



**HAL**  
open science

## The tooth, target organ of Bisphenol A, could be used as a biomarker of exposure to this agent

Katia Jedeon, Ariane Berdal, Sylvie Babajko

### ► To cite this version:

Katia Jedeon, Ariane Berdal, Sylvie Babajko. The tooth, target organ of Bisphenol A, could be used as a biomarker of exposure to this agent. Yann Gibert. Bisphenol A: Sources, Risks of Environmental Exposure and Human Health Effects, Nova Science Publishers, Inc, pp.205-225, 2015, Biochemistry Research Trends, 978-1-63463-210-2. hal-04199154

**HAL Id: hal-04199154**

**<https://hal.science/hal-04199154>**

Submitted on 7 Sep 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **The tooth, target organ of Bisphenol A, could be used as a biomarker of exposure to this agent**

Katia Jedeon<sup>1,2,3,4</sup>, Ariane Berdal<sup>1,2,3,4,5</sup>, Sylvie Babajko<sup>1,2,3,4</sup>

### **Affiliations of authors:**

<sup>1</sup>Centre de Recherche des Cordeliers, INSERM UMRS 1138, Laboratory of Molecular Oral Pathophysiology, Paris, F-75006 France.

<sup>2</sup>Université Paris-Descartes, Paris, F-75006 France.

<sup>3</sup>Université Pierre et Marie Curie-Paris, Paris, F-75006 France.

<sup>4</sup>Université Paris-Diderot, UFR d'Odontologie, F-75006, Paris, France.

<sup>5</sup>Centre de Référence des maladies rares de la face et de la cavité buccale MAFACE hôpital Rothschild, AP-HP, Paris, France.

### **Corresponding author:**

Sylvie Babajko

Centre de Recherche des Cordeliers, INSERM, UMRS 1138, Laboratory of Molecular Oral Pathophysiology, 15-21 rue de l'École de Médecine, 75270 Paris cedex 06, France

Tel: +33 (01) 44 27 55 79

Fax: +33 (01) 44 27 55 91

Email: [sylvie.babajko@crc.jussieu.fr](mailto:sylvie.babajko@crc.jussieu.fr)

## **Abstract**

Bisphenol A (BPA) is an endocrine disruptor chemical (EDC) whose effects on the development and function of reproductive organs and the mammary gland have been described for many years. In addition to these hormone-sensitive organs and cells, the tooth has been recently reported as an additional target organ of BPA (Jedeon *et al.*, 2013). The tooth comprises three compartments: the pulp cavity in its center, surrounded by dentin synthesized by odontoblasts, itself covered by enamel, the most mineralized tissue, synthesized by ameloblasts. After tooth eruption, ameloblasts disappear making any impact on enamel irreparable and thus irreversible. This property can be specifically exploited to relate the events that disturbed tooth development and especially the enamel mineralization taking place from the last third of fetal life to 4-5 years after birth in humans, precisely the time window of the highest susceptibility to EDCs. Experimental approaches consist to expose rats to low doses of BPA (alone or in combination with other enamel disrupting chemicals) and explore resulting enamel defects. Rats exposed to BPA present white opaque spots on their incisors reflecting enamel hypomineralization. These defects share many structural and biochemical characteristics with those described in the MIH (Molar Incisor Hypomineralization). MIH is an enamel disease that affects 15-18 % of children aged 6 to 9 years, diagnosed by white to brown lesions on their permanent first molars and sometimes on their permanent incisors. These data may suggest that BPA could be a causative agent of MIH and therefore teeth affected by MIH could be used as early markers of exposure to BPA and / or agents acting similarly. Finally, these fragile and often decayed teeth lead the dentist to use filling materials which contain BPA monomers and can constitute an additional source of BPA contamination, at least locally. The potential BPA contamination and consequent adverse health effects resulting from the use of dental materials is an important issue actively discussed.

## **BPA, an active endocrine disrupting chemical (EDC), is able to impact amelogenesis**

### *Oral contamination of BPA*

Bisphenol A (BPA), used in the manufacture of epoxy resins and polycarbonate plastics, is an exemplary widespread endocrine disruptor. BPA production has grown by 6 to 10% per year (Reuters, 2010) and is now estimated at 4 million tons. Indeed, over 95% of populations examined, in all parts of the world, are contaminated, mainly *via* food and food packaging (von Goetz et al., 2010; Pirard et al., 2012; Geens et al., 2012 for review; ANSES 2013). This contamination seems related to the population way of life since those limiting industrial goods, foods and habits are less contaminated (Martina *et al.*, 2012). 80% of whole BPA contamination result from oral exposure. Ingested BPA passes in blood circulation either by direct sublingual absorption (Gayrard et al., 2013) or by intestinal filtration. After hepatic detoxification and renal filtration, BPA-glucuronide is excreted in urine. Thus high urine and/or blood BPA concentrations may reflect high BPA exposure.

Epidemiological studies aim to associate high BPA urinary concentrations to adverse health effects including fertility troubles (Li et al., 2011; Ehrlich et al., 2012), obesity (Bhandari et al., 2013), diabetes (Lang et al., 2008; Silver et al., 2011 ; Shankar & Teppala, 2011 ; Sabanayagam et al., 2013 ; Kim et al., 2013), prostate cancer (Tarapore et al., 2014), increased risk of prematurity (Cantonwine et al., 2010), and behavior problems (Harley et al., 2013; Hong et al., 2013). Interestingly, BPA impact is not always evidenced on the whole population but more precisely on a selected group of individuals depending on age, sex, genetic background and the time of exposure.

### *Experimental studies carried out on animals*

Accordingly with epidemiological studies, experimental studies have shown that BPA impacts reproductive functions (Varayoud et al., 2011), prostate and mammary gland post-natal development (Taylor et al., 2011; Soto et al., 2013 for review). It also targets adipocyte, hepatic and pancreatic  $\beta$ -cells explaining metabolic effects of BPA leading to obesity, diabetes and other metabolic defects (Alonso-Magdalena et al., 2010; Vom Saal et al., 2012; Liu et al., 2013; Cabaton et

al., 2013). An important point that has to be underlined is that most of these data provide evidence for the importance of the perinatal exposure and its consequence during adulthood, as well as the preferential impacts mainly recorded in males. The importance of BPA perinatal exposure has been illustrated in other studies on behavioral impact (Braun et al., 2009; Anderson et al., 2013) and brain development (Poimenova et al., 2010).

Consistent with this, teeth have recently been found to be a target of BPA, where it disrupts amelogenesis leading to enamel hypomineralization (Jedeeon et al., 2013).

#### *How can BPA impact amelogenesis?*

The tooth comprises three compartments: the pulp cavity in its center, surrounded by dentin synthesized by odontoblasts, itself covered by enamel, the most mineralized tissue, synthesized by ameloblasts. Amelogenesis is a tightly time regulated process that includes ameloblast differentiation and enamel synthesis (Nancy, 2012), and it is time-coordinated with odontogenesis. Ameloblast differentiation originates in the dental cervical loop and begins with the secretory stage during which a partially mineralized enamel matrix is elaborated. The enamel matrix synthesized at that time is comprised of amelogenins, enamelin, ameloblastin and amelotin which are determinant for apatite deposition and for the first step of mineralization. When enamel mineralization is established and when crystal growth is determinant, matrix proteins are subject to extracellular processing by matrix metalloprotease-20 (MMP20). The matrix proteins are essential for the correct enamel formation as evidenced by the fact that mutations in any of the proteins involved leads to amelogenesis imperfecta (AI) (Wright, 2006; Chan et al., 2011). Once the full thickness of the enamel has been deposited, amelogenesis enters the maturation phase during which the serine protease kallikrein 4 (KLK4) degrades the enamel matrix proteins and calcium actively deposited. Abnormal retention of proteins or the presence of any extraneous proteins such as serum albumin lead to the eruption of hypomineralized enamel (Robinson et al., 1995). The rodent incisor provides a running record of how an impact on amelogenesis affects future development as the incisor continues to erupt (Berdal et

al., 1993; Lacruz et al., 2011) (**figure 1**). Once amelogenesis is complete, the ameloblasts and the overlying outer enamel epithelium cells degenerate and are ultimately lost through abrasion following tooth eruption. As a consequence, enamel defects are irreversible and provide a permanent record of any disturbances that occur during enamel development. This record of previous disturbances allows retrospective studies to be carried out and provides a means of temporally fixing a pathological event at some point during development. Among the different agents able to disturb amelogenesis, BPA has been recently shown to cause enamel hypomineralization (Jedeon et al. 2013).

#### *Enamel defects due to BPA exposure are similar to MIH*

BPA impact on amelogenesis was investigated by exposing Wistar rats daily by gavage to low doses of BPA (5 µg/kg/day) from conception until 30 or 100 days after birth (Jedeon et al., 2013). At 30 days, 75% of male rats (and 30% of females) presented white opaque spots on their incisors. In complete contrast, incisors from BPA treated rats at 100 days were no longer affected and were indistinguishable from controls indicating that rat amelogenesis has a window of susceptibility that does not extend to the period when the incisor enamel in P100 rats was being laid down (**figure 1**). BPA affected enamel exhibited asymmetrical hypomineralization area similar to human MIH.

MIH is diagnosed in children at around 6 to 8 years of age and presents as random white opacities on the enamel of first permanent molars and sometimes on permanent incisors also (Jälevik et al., 2001; Weerheijm & Merjare, 2003). Recent studies show that MIH prevalence turns around 15-18% of children affected whatever the region of the world considered (**table 1**). To date, MIH etiology remains unclear but the cause appears from environmental, even if enamelin gene polymorphism has been reported as a genetic factor associated with MIH development (Jeremias et al., 2013). Given that MIH affects those teeth that are undergoing mineralization around the time of birth, it is clear that the enamel forming ameloblasts are only sensitive to the causative agent(s) responsible for MIH in a specific time window. MIH is indicative of some adverse event(s) occurring during early

childhood that impact on enamel development (Jälevik, 2010). Diverse environmental conditions such as medication (amoxicillin), hypoxia, hypocalcaemia, dioxins, polychlorinated biphenyls (PCBs) and prolonged breast feeding have been associated with MIH (Alaluusua et al., 2002; Alaluusua, 2010). The white opaque opacities detected on 30 days-old rat incisors exposed to BPA provide a parallel between the etiology associated with the effect of BPA on rat amelogenesis and the etiology associated with human MIH. The perinatal window of sensitivity to BPA is precisely the same as the period of time corresponding to the mineralization of affected teeth, the first permanent molars and incisors (the first erupting permanent teeth). The mineralization of first permanent molars occurs between the 32th week of gestation and around four years after birth and that of permanent incisors between 3 months after birth and five years (AlQahtany, 2010). The mineralization of the other permanent teeth is delayed and they are very rarely affected by MIH.

Further investigations of both series of affected teeth by scanning electron microscopy showed that human MIH and BPA affected rat molars and incisors showed broken enamel in areas where the teeth occlude (Jedeon et al., 2013). In addition, the prismatic structure in human MIH enamel as well as BPA exposed enamel (of 30 days old rats) was obscured by a covering organic layer similar to the one reported previously (Jälevik et al., 2005). Among the main enamel matrix proteins (including amelogenins, enamelin and ameloblastin), enamelin expression (reported at the mRNA level as well as at the protein level) was higher in all BPA treated fractions. Enamelin amount is a central parameter for enamel synthesis as recently demonstrated by an experimental genetic approach (Hu et al., 2014). Too much or too little enamelin abolishes the production of enamel crystals and prism structure. Thus, BPA is able to specifically target one key gene involved in amelogenesis and affect enamel. BPA has also been shown to decrease KLK4 expression which is involved in the degradation of enamel matrix proteins that can remain during the maturation process of enamel and inhibit normal apatite crystal growth. This second event strengthens the first one by increasing much more the amount of enamelin. In such case, extraneous proteins as serum albumin are able to accumulate in the poor quality enamel (Farah et al., 2010) reinforcing the hypomineralization finally diagnosed by

the white opaque spots (Denis et al., 2013). All these events were reproduced *in vitro* in the rat ameloblastic cell line HAT-7 (Kawano et al., 2002) showing the BPA transcriptional regulation of enamelin and KLK4 gene expression (Jedeon et al., 2013).

The mechanism by which BPA impacts on ameloblast gene expression is still unclear. However BPA has been shown to interact with steroid/retinoid/thyroid nuclear receptor pathways (Paris et al., 2002; Delfosse et al., 2014). BPA has been shown to bind multiple receptors including estrogen receptors (ESRs) (Delfosse et al., 2012), GPR30 (Alonso-Magdalena et al., 2010; Chevalier et al., 2012), ERR gamma (Liu et al., 2012), but also retinoid receptor ROR gamma (Nishigori et al., 2012). It modulates androgen, thyroid and glucocorticoid receptor activities and expression levels of key-transcription factors such as CREB, CEBP, STAT3, THR, PPAR gamma or GATA-4. BPA may have a direct influence on ameloblasts by binding to a BPA sensitive receptor. The presence of ESR alpha, vitamin D and thyroid receptors in ameloblasts (Berdal et al., 1993; Ferrer et al., 2005) at the developmental stages where both enamelin and KLK4 expression is occurring maximally, invites the suggestion that BPA is affecting the expression of these genes through nuclear hormone pathway especially steroid receptors. The preferential enamel impact of BPA on male rats support this idea that should be further investigated.

#### *Combinatorial effects of BPA with other EDCs and enamel hypomineralizing factors*

Human and animal populations are exposed to many EDCs simultaneously. BPA certainly acts in combination with other EDCs or hypomineralizing agents. These molecules do not necessarily share same structural properties, and act through different signaling pathways and receptors that have not been exhaustively characterized. Consequently, the effects of combinations of EDCs are unpredictable: they can potentiate each other's actions or, on the contrary, suppress them. Combination of low doses of BPA with low doses of genistein and vinclozolin, two other EDCs, didn't lead to a greater phenotype (Jedeon et al., 2014). Whereas 6/8 rats exposed to BPA presented an enamel hypomineralization, only 3/8 presented similar white spots in the presence of combinations



of BPA with either genistein or vinclozolin or both. Such results were explained by the differential modulation of enamelin and KLK4 expression by these three agents, only BPA is able to increase enamelin expression (Jedeon et al., 2014). These results comfort the unpredictable impact of multiple agent combinations on a complex physiological function as amelogenesis.

In addition, one can suppose that BPA can have potentiated effects in the presence of hypomineralizing agents already described, especially dioxin that is also considered as a very active EDC (Alaluusua et al., 2004).

Among the different exogenous agents able to affect enamel mineralization, fluoride present in drinking water causes dental fluorosis by targeting mature ameloblasts, inhibiting enzymatic activity and interacting with calcium in hydroxyapatite crystal (Denbesten & Li, 2011; Perumal et al., 2013). The balance between beneficial effects on protection from caries and adverse effects leading to enamel hypomineralization occupies a narrow window of doses between 1 and 2 mg/day (Aoba & Fejerskov, 2002). Other chemicals are also able to lead to enamel hypomineralization as lead that exacerbates fluorosis (Leite et al., 2011), tetracyclin and amoxicillin (Tredwin et al., 2005; Antonini & Luder, 2011; Gottberg et al., 2014), and dioxin (Alaluusua et al., 2004; Gao et al., 2004). Interestingly, dioxin and amoxicillin exposures have been proposed as a causal factor of Molar Incisor Hypomineralisation (MIH) (Laisi et al., 2009; Alaluusua et al., 1999). It is noteworthy that both factors increase enamel hypomineralization in the presence of fluoride (Salmela et al., 2011; Sahlberg et al., 2013) and the importance of the perinatal exposure to these agents has been underlined (Alaluusua et al., 2002).

### **BPA may be leached from dental resins**

MIH affected teeth as well as any tooth with hypomineralized enamel (whatever the cause), are fragile and susceptible to caries. These decayed teeth lead the dentist to use filling materials which contain BPA monomers and can constitute an additional source of BPA contamination, at least

locally. The issue concerning the evaluation of BPA contamination due to dental materials and their possible adverse health effects are hotly debated.

#### *In vitro effects of BPA contained in dental materials*

Most dental sealants contain bisphenol-A diglycidyl methacrylate (bis-GMA), a common ingredient in restorative dental materials used since the 1960s. Olea and coworkers evidenced for the first time that dental resins leach BPA (present as an impurity) and related monomers bis-GMA, bis-DMA (bisphenol-A-diglycidyl-dimethacrylate) and BADGE (Bisphenol A diglycidyl ether) into saliva (Olea et al., 1996). This paper relates the highest amount of BPA ever detected in saliva (Rathee et al., 2012) but has the merit to raise the question of the safety around dental materials for patients but also for dentists. It triggered a lot of *in vitro* studies during the two last decades on potential effects of BPA (and related monomers) contained in dental resins, sealants, root canal sealers, orthodontic adhesives and polycarbonate brackets. These *in vitro* studies show that BPA leached from dental materials can present estrogen-like effects on hormone-sensitive cells, similar to other sources of BPA contaminations (Olea et al., 1996; Schafer et al., 1999). However, contrary to the first studies, the estrogen-like mitogenic activity of BPA related monomers, especially Bis-GMA, Bis-DMA and triethylene glycol dimethacrylate (TEGDMA) was not evidenced (Hashimoto et al., 2000, Nomura et al., 2003). In addition, *in vitro* tests on Chinese hamster lung cells show that contrary to BPA, nor Bis-GMA nor Bis-DMA activate on CYP3A4 and CYP3A7 (Hikage et al., 2003). However, as BPA monomers can be digested by esterases present in the saliva giving rise to free BPA and as BPA can be leached from dental resins due to incomplete polymerization, their presence must be checked despite the lack of their *in vitro* activity (Pulgar et al., 2000). In addition, some dental materials are still able to released bis-GMA at least during one year after *in vitro* elution (Polydorou et al., 2009) raising the issue of long-term effects and contamination.

At low doses, BPA is able to induce cell proliferation by up-regulating hTERT expression (Takahashi et al., 2004) whereas at high doses it presents cytotoxic effects by reducing oral epithelial cell number

(Seki et al., 2007). It has also been shown that high-doses ( $\mu\text{M}$  to  $\text{mM}$ ) of BPA monomers contained in dental resins present cytotoxic effects on gingival fibroblasts increasing cell apoptosis and DNA strand breakage (Engelman et al., 2004; Urcan et al., 2010). In summary, these *in vitro* data showed that BPA contained in dental materials shares similar properties with those of other plastic containers and that oral cells are directly susceptible to BPA. The dose of exposure is an important parameter to take into account as BPA present opposite effects depending on the concentration, mitogenic effects at low doses and apoptotic or cytotoxic effects at high doses.

#### *BPA contamination due to dental resins and composites*

Nevertheless, these *in vitro* investigations do not answer to the important issue of the potential BPA contamination nor to any adverse health effects resulting from BPA containing dental materials that may occur after orthodontic and/or restorative treatments. That's why many epidemiological studies on the impact of dental resins and sealants were reported in the literature since the publication of Olea and coworkers in 1996. The consensus idea that emerges from the literature is that BPA concentrations in saliva and urine increased one to three hours after treatment then return to basal levels (**Table 2**), suggesting that sufficient gargling after treatment can remove BPA from the oral cavity (Sasaki et al., 2005). Increased plasmatic levels of BPA were never reported whatever the study and the time of measurements after treatments. However, some recently published papers related that the number of dental composites could lead to a significant long-term BPA contamination determined by salivary and urinary BPA concentrations (Chung et al. 2012; Han et al., 2012; Kingman et al., 2012). The paper of Chung and co-workers presents the BPA urinary levels determined on 495 children aged 8-9 years and classified into four groups by the number of resin composites and sealant surfaces (0, 1-5, 6-10 and 11+). After adjusting for gender and age, the urinary BPA concentration in children with 11 or more resin composite surfaces was significantly higher than in the group without any resin composites. In the study of Han and co-workers, saliva of 124 children (62 controls with no dental sealant/resin and 62 cases with more than 4 tooth surfaces with dental sealant/resin) was

analyzed. The BPA level of control (0.40 µg/L) was significantly lower than that of case (0.92 µg/L). Kingman et al., included 172 participants receiving composite restorations in their study and showed that BPA levels in urine may remain increased until 30 hours after treatment. On the other hand, recent studies linking the number of dental resins composites and adverse health effects reported an increased level of anxiety, depression, social stress, interpersonal relation problems for children presenting more than two composites (Maserejian et al., 2012a). However, the number of composites is not associated to developmental troubles (Maserejian et al., 2012b; 2012c), nor to renal failure or damage (Trachtenberg et al., 2014; McGinley EL. 2014) or immune function alterations (Maserejian et al., 2013). These data point out the importance of the precise definition of the end-point and the difficulty to conclude of the non-effect of BPA in the absence of any positive control. They must be taken into account even if the number of subjects in cohorts is quite low to define further prospective studies including more BPA dosages during a longer period of time. Further studies are thus required before any conclusion on the possible BPA contamination by dental resins and possible adverse resulting effects.

#### *BPA contaminations due to orthodontic materials*

On the other hand, BPA and residual monomer leaching from orthodontic adhesive resins and polycarbonate brackets was evidenced *in vitro* as well as *in vivo* several months after treatment (Watanabe et al., 2001; Watanabe, 2004; Kloukos et al., 2013 for review). Even if their mitogenic activity was shown *in vitro* on the hormone-sensitive MCF-7 cell line (Gioka et al., 2009), it is not possible to evaluate their *in vivo* potential adverse effect (if any) as there are no available data relating urinary or salivary BPA concentration measurements long-term after orthodontic treatments, nor any adverse health effect described in children receiving such treatment.

It is however interesting to note that the studies carried on animals did not show any toxic effects of BPA-GMA (contained in dental materials) on reproductive function (Moilanen et al., 2013).

In conclusion, at the present time, long-term BPA contamination by dental materials is actively discussed and controversial because of the small size of cohorts and the imprecise duration of exposure (**table 2**, Akeroyd & Maserejian, 2013). Thus any reliable adverse effect is difficult/impossible to establish. Further studies are required to answer to these important issues that accelerate active research in the field of dental materials in order to find inert materials either resistant to enzymatic, mechanic and chemical degradation or devoid of any BPA related monomers. Taken all the published data together, the low dose effect (if any) of BPA leached by dental resins is probably negligible during adulthood. Nevertheless, more data are needed to evaluate the impact of orthodontic treatments often dispensed during childhood and the consequences on baby health of numerous resin composites carried by the mothers.

#### **The teeth are proposed as early biomarker of exposure to BPA**

Taking into account the huge amount of experimental data demonstrating BPA adverse impact on organ development and health, it is also important to lean on reliable biomarkers of exposure to BPA. The present data show that ameloblasts are susceptible to BPA and to other EDCs. Precise classification of the effects in each case would be valuable as these effects may not be as similar as may have been expected and specific enamel defects could reflect certain type of EDC combination. Considering the teeth as a complex organ resulting from epithelial and mesenchymal cell interactions, regrouping most of biological events and probably harboring many hormone receptors, the teeth or characterized *in vitro* models of enamel gene expression could constitute a novel marker of exposure to EDCs.

The data exposed in this paper underlie that BPA may be a causative agent in human MIH etiology. These fragile and often decayed teeth lead the dentist to use filling materials that may contain BPA based monomers thus maintaining pre-existing BPA contamination. MIH may represent a permanent record of exposure to BPA (or to EDCs sharing similar molecular effects) and could be easily used as a biomarker for retrospective analysis of infant exposure to EDCs and the impact of such exposure has

on health in later life. It is interesting to note that many toxicants mentioned in enamel hypomineralization including BPA, dioxin and genistein, are also suspected to be involved in mammary cancer development (Jenkins et al., 2012) reinforcing the idea of common mechanisms in teeth and other organs.

## References

Ahmadi R, Ramazani N, Nourinasab R. Molar incisor hypomineralization: a study of prevalence and etiology in a group of Iranian children. *Iran J Pediatr.* 2012;22(2):245-51.

Akeroyd JM, Maserejian NN. Composite restorations may lead to increased concentrations of salivary and urinary BPA. Article analysis and evaluation. *J Evid Base Dent Pract* 2013;13:64-66.

Alaluusua S. Aetiology of Molar-Incisor Hypomineralisation: A systematic review. *Eur Arch Paediatr Dent* 2010;11(2):53-58.

Alaluusua S, Calderara P, Gerthoux PM, Lukinmaa PL, Kovero O, Needham L, Patterson DG Jr, Tuomisto J, Mocarelli P. Developmental dental aberrations after the dioxin accident in Seveso. *Environ Health Perspect.* 2004;112(13):1313-8.

Alaluusua S, Kiviranta H, Leppäniemi A, Hölttä P, Lukinmaa PL, Lope L, Järvenpää AL, Renlund M, Toppari J, Virtanen H, Kaleva M, Vartiainen T. Natal and neonatal teeth in relation to environmental toxicants. *Pediatr Res.* 2002;52(5):652-5.

Alaluusua S, Lukinmaa PL, Torppa J, Tuomisto J, Vartiainen T. Developing teeth as biomarker of dioxin exposure. *Lancet.* 1999;353(9148):206.

AlQahtani SJ, Hector MP, Liversidge HM. Brief communication: The London atlas of human tooth development and eruption. *Am J Phys Anthropol.* 2010;142(3):481-90.

Alonso-Magdalena P, Ropero AB, Soriano S, García-Arévalo M, Ripoll C, Fuentes E, Quesada I, Nadal Á: Bisphenol-A acts as a potent estrogen via non-classical estrogen triggered pathways. *Mol Cell Endocrinol* 2012, 355:201-207.

Alonso-Magdalena P, Vieira E, Soriano S, Menes L, Burks D, Quesada I, Nadal A. Bisphenol A exposure during pregnancy disrupts glucose homeostasis in mothers and adult male offspring. *Environ Health Perspect.* 2010;118(9):1243-50.

Anderson OS, Peterson KE, Sanchez BN, Zhang Z, Mancuso P, Dolinoy DC. Perinatal bisphenol A exposure promotes hyperactivity, lean body composition, and hormonal responses across the murine life course. *FASEB J.* 2013;27(4):1784-92.

Anses - Evaluation du risque du bisphénol A pour la santé humaine. Avis de l'ANSES, rapport d'expertise collective. Édition scientifique. Mars 2013.

Antonini LG, Luder HU. Discoloration of teeth from tetracyclines--even today? *Schweiz Monatsschr Zahnmed.* 2011;121(5):414-31.

Aoba T, Fejerskov O. Dental fluorosis: chemistry and biology. *Critical reviews in oral biology and medicine* : an official publication of the American Association of Oral Biologists. 2002;13(2):155-70.

Arenholt-Bindslev D, Breinholt V, Preiss A, Schmalz G. Time-related bisphenol-A content and estrogenic activity in saliva samples collected in relation to placement of fissure sealants. *Clin Oral Investig.* 1999;3(3):120-5.

Balmer R, Toumba J, Godson J, Duggal M. The prevalence of molar incisor hypomineralisation in Northern England and its relationship to socioeconomic status and water fluoridation. *Int J Paediatr Dent.* 2012;22(4):250-7.

Berdal A, Hotton D, Pike JW, Mathieu H, Dupret JM: Cell- and stage-specific expression of vitamin D receptor and calbindin genes in rat incisor: regulation by 1,25-dihydroxyvitamin D3. *Dev Biol* 1993, 155:172-179.

Bhandari R, Xiao J, Shankar A. Urinary bisphenol A and obesity in U.S. children. *Am J Epidemiol.* 2013;177(11):1263-70.

Biondi AM, Cortese SG, Martínez K, Ortolani AM, Sebelli PM, Ienco M, Paván VH, Mendel N, Bertolino M, Hecht P. Prevalence of molar incisor hypomineralization in the city of Buenos Aires. *Acta Odontol Latinoam.* 2011;24(1):81-5.

Biondi AM, López Jordi Mdel C, Cortese SG, Alvarez L, Salveraglio I, Ortolani AM. Prevalence of molar-incisor hypomineralization (MIH) in children seeking dental care at the Schools of Dentistry of the University of Buenos Aires (Argentina) and University of la Republica (Uruguay). *Acta Odontol Latinoam.* 2012;25(2):224-30.

Braun JM, Yolton K, Dietrich KN, et al. Prenatal bisphenol A exposure and early childhood behavior. *Environ Health Perspect.* 2009;117(12):1945–1952.

Brogårdh-Roth S, Matsson L, Klingberg G. Molar-incisor hypomineralization and oral hygiene in 10-to-12-yr-old Swedish children born preterm. *Eur J Oral Sci.* 2011 Feb;119(1):33-9.

Cabaton NJ, Canlet C, Wadia PR, Tremblay-Franco M, Gautier R, Molina J, Sonnenschein C, Cravedi JP, Rubin BS, Soto AM, Zalko D. Effects of low doses of bisphenol A on the metabolome of perinatally exposed CD-1 mice. *Environ Health Perspect.* 2013;121(5):586-93.

Cantonwine D, Meeker JD, Hu H, Sánchez BN, Lamadrid-Figueroa H, Mercado-García A, Fortenberry GZ, Calafat AM, Téllez-Rojo MM. Bisphenol a exposure in Mexico City and risk of prematurity: a pilot nested case control study. *Environ Health.* 2010;9:62.

Chan HC, Estrella NM, Milkovich RN, Kim JW, Simmer JP, Hu JC. Target gene analyses of 39 amelogenesis imperfecta kindreds. *Eur J Oral Sci.* 2011;119 Suppl 1:311-23.

Chevalier N, Bouskine A, Fenichel P. Bisphenol A promotes testicular seminoma cell proliferation through GPER/GPR30. *Int J Cancer.* 2012;130(1):241-2.

Chung SY, Kwon H, Choi YH, Karmaus W, Merchant AT, Song KB, Sakong J, Ha M, Hong YC, Kang D. Dental composite fillings and bisphenol A among children: a survey in South Korea. *Int Dent J.* 2012;62(2):65-9.

Condò R, Perugia C, Maturo P, Docimo R. MIH: epidemiologic clinic study in paediatric patient. *Oral Implantol (Rome).* 2012 Apr;5(2-3):58-69.

da Costa-Silva CM, Jeremias F, de Souza JF, Cordeiro Rde C, Santos-Pinto L, Zuanon AC. Molar incisor hypomineralization: prevalence, severity and clinical consequences in Brazilian children. *Int J Paediatr Dent.* 2010 Nov;20(6):426-34.

Delfosse V, Grimaldi M, le Maire A, Bourguet W, Balaguer P. Nuclear receptor profiling of bisphenol-A and its halogenated analogues. *Vitam Horm.* 2014;94:229-51.

Delfosse V, Grimaldi M, Pons JL, Boulahtouf A, le Maire A, Cavailles V, Labesse G, Bourquet W, Balaguer P : Structural and mechanistic insights into bisphenols action provide guidelines for risk assessment and discovery of bisphenol A substitutes. *Proc Natl Acad Sci USA* 2012, 109:14930-14935.

Denbesten P, Li W. Chronic fluoride toxicity: dental fluorosis. *Monogr Oral Sci.* 2011;22:81-96.

Denis M, Atlan A, Vennat E, Tirlet G, Attal JP. White defects on enamel: diagnosis and anatomopathology: two essential factors for proper treatment (part 1). *Int Orthod.* 2013;11(2):139-65.

Engelmann J, Janke V, Volk J, Leyhausen G, von Neuhoff N, Schlegelberger B, Geurtsen W. Effects of BisGMA on glutathione metabolism and apoptosis in human gingival fibroblasts in vitro. *Biomaterials.* 2004;25(19) :4573-80.

Ehrlich S, Williams PL, Missmer SA, Flaws JA, Berry KF, Calafat AM, Ye X, Petrozza JC, Wright D, Hauser R. Urinary bisphenol A concentrations and implantation failure among women undergoing in vitro fertilization. *Environ Health Perspect.* 2012;120(7):978-83.

Farah RA, Monk BC, Swain MV, Drummond BK. Protein content of molar-incisor hypomineralisation enamel. *J Dent.* 2010;38(7):591-6.

Ferrer VL, Maeda T, Kawano Y: Characteristic distribution of immunoreactions for estrogen receptor alpha in rat ameloblasts. *Anat Rec* 2005, 284 :529–536.

Fung EY, Ewoldsen NO, St Germain HA Jr, Marx DB, Miaw CL, Siew C, Chou HN, Gruninger SE, Meyer DM. Pharmacokinetics of bisphenol A released from a dental sealant. *J Am Dent Assoc.* 2000;131(1):51-8.



Gao Y, Sahlberg C, Kiukkonen A, Alaluusua S, Pohjanvirta R, Tuomisto J, Lukinmaa PL. Lactational exposure of Han/Wistar rats to 2,3,7,8-tetrachlorodibenzo-p-dioxin interferes with enamel maturation and retards dentin mineralization. *J Dent Res.* 2004;83(2):139-44.

Garcia-Margarit M, Catalá-Pizarro M, Montiel-Company JM, Almerich-Silla JM. Epidemiologic study of molar-incisor hypomineralization in 8-year-old Spanish children. *Int J Paediatr Dent.* 2014 Jan;24(1):14-22.

Geens T, Aerts D, Berthot C, Bourguignon JP, Goeyens L, Lecomte P, Maghuin-Rogister G, Pironnet AM, Pussemier L, Scippo ML, Van Loco J, Covaci A. A review of dietary and non-dietary exposure to bisphenol-A. *Food Chem Toxicol.* 2012;50(10):3725-40.

Ghanim A, Morgan M, Mariño R, Bailey D, Manton D. Molar-incisor hypomineralisation: prevalence and defect characteristics in Iraqi children. *Int J Paediatr Dent.* 2011 Nov;21(6):413-21.

Gioka C, Eliades T, Zinelis S, Pratsinis H, Athanasiou AE, Eliades G, Kletsas D. Characterization and in vitro estrogenicity of orthodontic adhesive particulates produced by simulated debonding. *Dent Mater.* 2009;25(3):376-82.

Gottberg B, Berné J, Quiñónez B, Solórzano E. Prenatal effects by exposing to amoxicillin on dental enamel in Wistar rats. *Med Oral Patol Oral Cir Bucal.* 2014;19(1):e38-43.

Grošelj M, Jan J. Molar incisor hypomineralisation and dental caries among children in Slovenia. *Eur J Paediatr Dent.* 2013 Sep;14(3):241-5.

Han DH, Kim MJ, Jun EJ, Kim JB. Salivary bisphenol-A levels due to dental sealant/resin: a case-control study in Korean children. *J Korean Med Sci.* 2012;27(9):1098-104.

Harley KG, Gunier RB, Kogut K, Johnson C, Bradman A, Calafat AM, Eskenazi B. Prenatal and early childhood bisphenol A concentrations and behavior in school-aged children. *Environ Res.* 2013;126 :43-50.

Hashimoto Y, Nakamura M. Estrogenic activity of dental materials and bisphenol-A related chemicals in vitro. *Dent Mater J.* 2000;19(3):245-62.

Hikage S, Nakayama K, Saito T, Takahashi Y, Kamataki T, Suzuki S, Hongo T, Sato A. Cytotoxicity of bisphenol A glycidyl methacrylate on cytochrome P450-producing cells. *J Oral Rehabil.* 2003;30(5):544-9.

Hong SB, Hong YC, Kim JW, Park EJ, Shin MS, Kim BN, Yoo HJ, Cho IH, Bhang SY, Cho SC. Bisphenol A in relation to behavior and learning of school-age children. *J Child Psychol Psychiatry.* 2013;54(8):890-9.

Hu JC, Hu Y, Lu Y, Smith CE, Lertlam R, Wright JT, Suggs C, McKee MD, Beniash E, Kabir ME, Simmer JP. Enamelin is critical for ameloblast integrity and enamel ultrastructure formation. *PLoS One*. 2014;9(3):e89303.

Jälevik B. Enamel hypomineralization in permanent first molars. A clinical, histomorphological and biochemical study. *Swed Dent J Suppl* 2001;149:1-86.

Jälevik B, Dietz W, Norén JG. Scanning electron micrograph analysis of hypomineralized enamel in permanent first molars. *Int J Paediatr Dent* 2005;15(4):233-240.

Jedeon K, De La Dure-Molla M, Brookes SJ, Loiodice S, Marciano C, Kirkham J, Canivenc-Lavier M-C, Boudalia S, Bergès R, Harada H, Berdal A, Babajko S. Enamel defects reflect exposure to perinatal bisphenol A. *Am J Pathol*. 2013;183(1):108-18.

Jedeon K, Marciano C, Loiodice S, Boudalia S, Canivenc Lavier M-C, Berdal A, Babajko S. Enamel hypomineralization due to endocrine disruptors. *Connective Tissue Research*. 2014, in press

Jenkins S, Betancourt AM, Wang J, Lamartiniere CA. Endocrine-active chemicals in mammary cancer causation and prevention. *J Steroid Biochem Mol Biol*. 2012;129(3-5):191-200.

Jeremias F, de Souza JF, Silva CM, Cordeiro Rde C, Zuanon AC, Santos-Pinto L. Dental caries experience and Molar-Incisor Hypomineralization. *Acta Odontol Scand*. 2013;71(3-4):870-6.

Joskow R, Barr DB, Barr JR, Calafat AM, Needham LL, Rubin C. Exposure to bisphenol A from bis-glycidyl dimethacrylate-based dental sealants. *J Am Dent Assoc*. 2006;137(3):353-62.

Kang YG, Kim JY, Kim J, Won PJ, Nam JH. Release of bisphenol A from resin composite used to bond orthodontic lingual retainers. *Am J Orthod Dentofacial Orthop*. 2011 Dec;140(6):779-89.

Kawano S, Morotomi T, Toyono T, Nakamura N, Uchida T, Ohishi M, Toyoshima K, Harada H. Establishment of dental epithelial cell line (HAT-7) and the cell differentiation dependent on Notch signaling pathway. *Connect Tissue Res*. 2002;43(2-3):409-12.

Kim K, Park H. Association between urinary concentrations of bisphenol A and type 2 diabetes in Korean adults: a population-based cross-sectional study. *Int J Hyg Environ Health*. 2013;216(4):467-71.

Kingman A, Hyman J, Masten SA, Jayaram B, Smith C, Eichmiller F, Arnold MC, Wong PA, Schaeffer JM, Solanki S, Dunn WJ. Bisphenol A and other compounds in human saliva and urine associated with the placement of composite restorations. *J Am Dent Assoc*. 2012;143(12):1292-302.

Kloukos D, Pandis N, Eliades T. Bisphenol-A and residual monomer leaching from orthodontic adhesive resins and polycarbonate brackets: a systematic review. *Am J Orthod Dentofacial Orthop*. 2013;143(4 Suppl):S104-12.e1-2.

- Lacruz RS, Smith CE, Chen YB, Hubbard MJ, Hacia JG, Paine ML: Gene-expression analysis of early- and late-maturation-stage rat enamel organ. *Eur J Oral Sci* 2011, 119:149-157
- Laisi S, Ess A, Sahlberg C, Arvio P, Lukinmaa PL, Alaluusua S. Amoxicillin may cause molar incisor hypomineralization. *J Dent Res*. 2009;88(2):132-6.
- Lang IA, Galloway TS, Scarlett A, Henley WE, Depledge M, Wallace RB, Melzer D. Association of urinary bisphenol A concentration with medical disorders and laboratory abnormalities in adults. *JAMA*. 2008;300(11):1303-10.
- Leite GA, Sawan RM, Teófilo JM, Porto IM, Sousa FB, Gerlach RF. Exposure to lead exacerbates dental fluorosis. *Arch Oral Biol*. 2011;56(7):695-702.
- Li DK, Zhou Z, Miao M, He Y, Wang J, Ferber J, Herrinton LJ, Gao E, Yuan W. Urine bisphenol-A (BPA) level in relation to semen quality. *Fertil Steril*. 2011;95(2):625-30.e1-4.
- Liu J, Yu P, Qian W, Li Y, Zhao J, Huan F, Wang J, Xiao H. Perinatal bisphenol A exposure and adult glucose homeostasis: identifying critical windows of exposure. *PLoS One*. 2013;8(5):e64143.
- Liu X, Matsushima A, Nakamura M, Costa T, Nose T, Shimohigashi Y: Fine spatial assembly for construction of the phenol pocket to capture bisphenol A in the human nuclear receptor estrogen related receptor  $\gamma$ . *J Biochem* 2012, 151:403-415.
- Mahoney EK, Morrison DG. Further examination of the prevalence of MIH in the Wellington region. *N Z Dent J*. 2011 Sep;107(3):79-84.
- Martina CA, Weiss B, Swan SH. Lifestyle behaviors associated with exposures to endocrine disruptors. *Neurotoxicology*. 2012;33(6):1427-33.
- Martínez Gómez TP, Guinot Jimeno F, Bellet Dalmau LJ, Giner Tarrida L. Prevalence of molar-incisor hypomineralisation observed using transillumination in a group of children from Barcelona (Spain). *Int J Paediatr Dent*. 2012 Mar;22(2):100-9.
- Maserejian NN, Trachtenberg FL, Hauser R, McKinlay S, Shrader P, Tavares M, Bellinger DC. Dental composite restorations and psychosocial function in children. *Pediatrics*. 2012a;130(2):e328-38.
- Maserejian NN, Trachtenberg FL, Hauser R, McKinlay S, Shrader P, Bellinger DC. Dental composite restorations and neuropsychological development in children: treatment level analysis from a randomized clinical trial. *Neurotoxicology*. 2012c;33(5):1291-7.
- Maserejian NN, Hauser R, Tavares M, Trachtenberg FL, Shrader P, McKinlay S. Dental composites and amalgam and physical development in children. *J Dent Res*. 2012b;91(11):1019-25.

Maserejian NN, Shrader P, Brown OA, Trachtenberg FL, Soncini J, Hauser R, Shenker BJ. Dental sealants and composite restorations and longitudinal changes in immune function markers in children. *Int J Paediatr Dent*. 2013

McGinley EL. Summary of: dental composite materials and renal function in children. *Br Dent J*. 2014;216(2):80-1.

Mittal NP, Goyal A, Gauba K, Kapur A. Molar incisor hypomineralisation: prevalence and clinical presentation in school children of the northern region of India. *Eur Arch Paediatr Dent*. 2014 Feb;15(1):11-8.

Moilanen LH, Dahms JK, Hoberman AM. Reproductive toxicity evaluation of the dental resin monomer bisphenol a glycidyl methacrylate (CAS 1565-94-2) in mice. *Int J Toxicol*. 2013;32(6):415-25.

Nanci A. Enamel: Composition, Formation, and Structure. In: Ten Cate's Oral Histology Development, Structure, and Function / 8<sup>th</sup> edition United States of America: Elsevier mosby, 2012; p. 122-164.

Nishigori M, Nose T, Shimohigashi Y : Highly potent binding and inverse agonist activity of bisphenol A derivatives for retinoid-related orphan nuclear receptor ROR $\gamma$ . *Toxicol Lett* 2012;212:205-211.

Ng JJ, Eu OC, Nair R, Hong CH. Prevalence of molar incisor hypomineralization (MIH) in Singaporean children. *Int J Paediatr Dent*. 2014 Mar 7.

Nomura Y, Ishibashi H, Miyahara M, Shinohara R, Shiraishi F, Arizono K. Effects of dental resin metabolites on estrogenic activity in vitro. *J Mater Sci Mater Med*. 2003;14(4):307-10.

Olea N, Pulgar R, Pérez P, Olea-Serrano F, Rivas A, Novillo-Fertrell A, Pedraza V, Soto AM, Sonnenschein C. Estrogenicity of resin-based composites and sealants used in dentistry. *Environ Health Perspect*. 1996;104(3):298-305.

Parikh DR, Ganesh M, Bhaskar V. Prevalence and characteristics of Molar Incisor Hypomineralisation (MIH) in the child population residing in Gandhinagar, Gujarat, India. *Eur Arch Paediatr Dent*. 2012 Feb;13(1):21-6.

Paris F, Balaguer P, Térouanne B, Servant N, Lacoste C, Cravedi JP, Nicolas JC, Sultan C: Phenylphenols, biphenols, bisphenol-A and 4-tert-octylphenol exhibit alpha and beta estrogen activities and antiandrogen activity in reporter cell lines. *Mol Cell Endocrinol* 2002;193:43-49.

Perumal E, Paul V, Govindarajan V, Panneerselvam L. A brief review on experimental fluorosis. *Toxicol Lett*. 2013;223(2):236-51.

Pirard C, Sagot C, Deville M, Dubois N, Charlier C. Urinary levels of bisphenol A, triclosan and 4-nonylphenol in a general Belgian population. *Environ Int*. 2012;48 :78-83.

Poimenova A, Markaki E, Rahiotis C, Kitraki E. Corticosterone-regulated actions in the rat brain are affected by perinatal exposure to low dose of bisphenol A. *Neuroscience*. 2010;167(3):741-9.

Polydorou O, König A, Hellwig E, Kümmerer K. Long-term release of monomers from modern dental-composite materials. *Eur J Oral Sci.* 2009;117(1):68-75.

Pulgar R, Olea-Serrano MF, Novillo-Fertrell A, Rivas A, Pazos P, Pedraza V, Navajas JM, Olea N. Determination of bisphenol A and related aromatic compounds released from bis-GMA-based composites and sealants by high performance liquid chromatography. *Environ Health Perspect.* 2000;108(1):21-7.

Rathee M, Malik P, Singh J. Bisphenol A in dental sealants and its estrogen like effect. *Indian J Endocrinol Metab.* 2012;16(3):339-42.

Reuters. Experts demand European action on plastics chemical. 22 June 2010.

Robinson C, Kirkham J, Brookes SJ, Bonass WA: The chemistry of enamel development. *Int J Dev Biol* 1995;39:145-152.

Sabanayagam C, Teppala S, Shankar A. Relationship between urinary bisphenol A levels and prediabetes among subjects free of diabetes. *Acta Diabetol.* 2013;50(4):625-31.

Sahlberg C, Pavlic A, Ess A, Lukinmaa PL, Salmela E, Alaluusua S. Combined effect of amoxicillin and sodium fluoride on the structure of developing mouse enamel in vitro. *Arch Oral Biol.* 2013;58(9):1155-64.

Salmela E, Lukinmaa PL, Partanen AM, Sahlberg C, Alaluusua S. Combined effect of fluoride and 2,3,7,8-tetrachlorodibenzo-p-dioxin on mouse dental hard tissue formation in vitro. *Arch Toxicol.* 2011;85(8):953-63.

Sasaki N, Okuda K, Kato T, Kakishima H, Okuma H, Abe K, Tachino H, Tuchida K, Kubono K. Salivary bisphenol-A levels detected by ELISA after restoration with composite resin. *J Mater Sci Mater Med.* 2005;16(4):297-300.

Schafer TE, Lapp CA, Hanes CM, Lewis JB, Wataha JC, Schuster GS. Estrogenicity of bisphenol A and bisphenol A dimethacrylate in vitro. *J Biomed Mater Res.* 1999;45(3):192-7.

Seki K, Koshi R, Sugano N, Masutani S, Yoshinuma N, Ito K. Microarray analysis of bisphenol A-induced changes in gene expression in human oral epithelial cells. *Acta Biochim Biophys Sin (Shanghai).* 2007;39(11):879-84.

Shankar A, Teppala S. Relationship between urinary bisphenol A levels and diabetes mellitus. *J Clin Endocrinol Metab.* 2011;96(12):3822-6.

Silver MK, O'Neill MS, Sowers MR, Park SK. Urinary bisphenol A and type-2 diabetes in U.S. adults: data from NHANES 2003-2008. *PLoS One.* 2011;6(10):e26868.

Soto AM, Brisken C, Schaeberle C, Sonnenschein C. Does cancer start in the womb? altered mammary gland development and predisposition to breast cancer due to in utero exposure to endocrine disruptors. *J Mammary Gland Biol Neoplasia*. 2013;18(2):199-208.

Takahashi A, Higashino F, Aoyagi M, Kyo S, Nakata T, Noda M, Shindoh M, Kohgo T, Sano H. Bisphenol A from dental polycarbonate crown upregulates the expression of hTERT. *J Biomed Mater Res B Appl Biomater*. 2004;71(1):214-21.

Tarapore P, Ying J, Ouyang B, Burke B, Bracken B, Ho SM. Exposure to bisphenol a correlates with early-onset prostate cancer and promotes centrosome amplification and anchorage-independent growth in vitro. *PLoS One*. 2014 Mar 3;9(3):e90332.

Taylor JA, Richter CA, Ruhlen RL, vom Saal FS. Estrogenic environmental chemicals and drugs: mechanisms for effects on the developing male urogenital system. *J Steroid Biochem Mol Biol*. 2011;127(1-2):83-95.

Trachtenberg FL, Shrader P, Barregard L, Maserejian NN. Dental composite materials and renal function in children. *Br Dent J*. 2014;216(2):E4.

Tredwin CJ, Scully C, Bagan-Sebastian JV. Drug-induced disorders of teeth. *J Dent Res*. 2005;84(7):596-602.

Urcan E, Scherthan H, Styllou M, Haertel U, Hickel R, Reichl FX. Induction of DNA double-strand breaks in primary gingival fibroblasts by exposure to dental resin composites. *Biomaterials*. 2010;31(8):2010-4.

Varayoud J, Ramos JG, Bosquiaz VL, Lower M, Muñoz-de-Toro M, Luque EH. Neonatal exposure to bisphenol A alters rat uterine implantation-associated gene expression and reduces the number of implantation sites. *Endocrinology*. 2011;152(3):1101-11.

Vom Saal FS, Nagel SC, Coe BL, Angle BM, Taylor JA. The estrogenic endocrine disrupting chemical bisphenol A (BPA) and obesity. *Mol Cell Endocrinol*. 2012;354(1-2):74-84.

Von Goetz N, Wormuth M, Scheringer M, Hungerbühler K. Bisphenol a: how the most relevant exposure sources contribute to total consumer exposure. *Risk Anal*. 2010;30(3):473-87.

Watanabe M. Degradation and formation of bisphenol A in polycarbonate used in dentistry. *J Med Dent Sci*. 2004;51(1):1-6.

Watanabe M, Hase T, Imai Y. Change in the bisphenol A content in a polycarbonate orthodontic bracket and its leaching characteristics in water. *Dent Mater J* 2001;20:353-8.


Weerheijm KL, Mejàre I. Molar incisor hypomineralization: a questionnaire inventory of its occurrence in member countries of the European Academy of Paediatric Dentistry (EAPD). *Int J Paediatr Dent*. 2003;13(6):411-416.

Wright JT: The molecular etiologies and associated phenotypes of amelogenesis imperfecta. *Am J Med Genet A*. 2006;140:2547-2555.

Yamada H, Furuta I, Kato EH, et al. Maternal serum and amniotic fluid bisphenol A concentrations in the early second trimester. *Reprod Toxicol*. 2002;16(6):735–739.

Zawaideh FI, Al-Jundi SH, Al-Jaljoli MH. Molar incisor hypomineralisation: prevalence in Jordanian children and clinical characteristics. *Eur Arch Paediatr Dent*. 2011 Feb;12(1):31-6.

Zimmerman-Downs JM, Shuman D, Stull SC, Ratzlaff RE. Bisphenol A blood and saliva levels prior to and after dental sealant placement in adults. *J Dent Hyg*. 2010;84(3):145-50.



**Figure 1** Chronology of tooth development in human and rodent. Amelogenesis begins first with enamel matrix synthesis mainly constituted of amelogenin, enamelin and ameloblastin during the secretion stage (in green). After that, matrix proteins are degraded by MMP20 and KLK4, and mineralization takes place. In blue, mineralization period of affected teeth by exposure to BPA and in red, mineralization period of non affected teeth.



<b>Study</b>	<b>Number of children - Country</b>	<b>Age</b>	<b>MIH prevalence</b>
da Costa-Silva CM, <b>2010</b>	918 children - <b>Brazil</b>	6 to 12 years	<b>19,8%</b>
Brogårdh-Roth S <i>et al.</i> , <b>2011</b>	82 children - <b>Sweden</b>	10 to 12 years	<b>16%</b>
Biondi AM <i>et al.</i> , <b>2011</b>	1098 children - Buenos Aires <b>Argentina</b>	11,3 years	<b>15,9%</b>
Zawaideh FI <i>et al.</i> , <b>2011</b>	3666 children- <b>Jordan</b>	7 to 9 years	<b>13,7%</b>
Ghanim A <i>et al.</i> , <b>2011</b>	823 children - <b>Irak</b>	7 to 9 years	<b>18,6%</b>
Mahoney EK <i>et al.</i> , <b>2011</b>	235 children - Wellington, <b>New Zealand</b>	7 to 10 years	<b>18,8%</b>
Parikh DR <i>et al.</i> , <b>2012</b>	1366 children - <b>India</b>	8 to 12 years	<b>9,2%</b>
Balmer R <i>et al.</i> , <b>2012</b>	3233 children - North <b>England</b>	12 years	<b>15,9%</b>
Biondi AM <i>et al.</i> , <b>2012</b>	975 children - <b>Argentina / Uruguay</b>	11.6 years	<b>6.56%</b>
Ahmadi R <i>et al.</i> , <b>2012</b>	433 children - <b>Iran</b>	7-9 years	<b>12.7%</b>
Martínez Gómez TP <i>et al.</i> , <b>2012</b>	550 children - <b>Spain</b>	6 to 14 years	<b>17,8%</b>
Condò R <i>et al.</i> , <b>2012</b>	1500 children - <b>Italy</b>	4 to 15 years	<b>7,3%</b>
Jeremias F <i>et al.</i> , <b>2013</b>	1157 children - <b>Brazil</b>	6 to 12 years	<b>12,3%</b>
Grošelj M <i>et al.</i> , <b>2013</b>	558 children - <b>Slovenia</b>	6 to 11.5 years	<b>21.4%</b>
Garcia-Margarit M <i>et al.</i> , <b>2014</b>	840 children - <b>Spain</b>	8 years	<b>21,8%</b>
Mittal NP <i>et al.</i> , <b>2014</b>	1792 children - <b>India</b>	6 to 9 years	<b>6,31%</b>
Ng JJ <i>et al.</i> , <b>2014</b>	1083 children - <b>Singapore</b>	7.7 years	<b>12.5%</b>

**Table 2** Prevalence of MIH since 2010. The prevalence doesn't seem to depend on particular geographical region. Mean prevalence recorded the last five years turns around 15%.

Study	Number of subjects (age)	Time of measurements after treatments	Methods for BPA detection	Results	Conclusions
Olea N <i>et al.</i> , 1996	18 subjects (18-25 years)	1 h	HPLC and MS	In saliva: - Baseline: no detection - 1 h after treatment : 90-931 µg	The use of bis-GMA-based-resins in dentistry may represent an additional source of xenoestrogen exposure in humans.
Arenholt-Bindslev D <i>et al.</i> , 1999	8 subjects (20-23 years)	Immediately, 1 h and 24 h	HPLC	In saliva: - Baseline: no detection - Immediately after treatment: 0.3-2.8 ppm - 1 h and 24 h after treatment: no detection	Considerably lower amounts of BPA than previously reported, were detected in saliva samples collected immediately after but not 1 and 24 h after treatment.
Fung EY <i>et al.</i> , 2000	40 subjects (20-55 years)	1 h, 3 h, 1 day, 3 and 5 days	HPLC	In saliva: - 1 h and 3 h: 5.8-105.6 ppb in some specimens collected with a significant decrease from 1 to 3 h. In serum: - No detection beyond 3 h	BPA released orally from a dental sealant may not be absorbed or may be present in non detectable amounts in systemic circulation.
Sasaki N <i>et al.</i> , 2005	21 subjects	Immediately	ELISA	In saliva: - 10 to 100 ng/ml of BPA	Sufficient gargling after treatment is important for risk management.
Joskow R <i>et al.</i> , 2006	14 men	Immediately	MS	In saliva and urine: - 5.5 µg or 110 µg depending on the sealant used	Sealants should remain a useful part of routine preventive dental practice, especially those that leach negligible amounts of BPA.
Zimmerman-Downs JM <i>et al.</i> , 2010	30 adults (18 - 40 years)	1 h, 3 h and 24 h	ELISA	In saliva: - 0.07 to 6.00 ng/ml at baseline - Peak at 3 h: 3.98 ng/ml for one sealant group and 9.08 ng/ml for four sealant group - 24 h: same levels as baseline. In serum: - No detection	Dental sealant material used in this study do not influence the serum concentration levels of BPA.
Kang YG <i>et al.</i> , 2011	22 subjects	30 minutes, 1 day, 1 week and 1 month	HPLC / MS	In saliva: -20.889 ng/mL	The potential toxicity of BPA from placing lingual bonded retainer might be negligible.

Study	Number of subjects (age)	Time of measurements after treatments	Methods for BPA detection	Results	Conclusions
Maserejian NN <i>et al.</i> , 2012 (a)	534 children (6 - 10 years )	Comparison between the baseline and 5 years after treatments.	Psychosocial assessments by using validated instruments: Child Behavior Checklist (CBCL) parent-report, and Behavior Assessment for Children Self-Report (BASC-SR)		Greater exposure to bis-GMA based dental composite restorations was associated with impaired psychosocial function in children, whereas no adverse psychosocial outcomes were observed with greater UDMA-based compomer or amalgam treatment levels.
Maserejian NN <i>et al.</i> , 2012 (b)	534 children (6 - 10 years )	Comparison between the baseline and 5 years after treatments.	Tests of executive function, intelligence, memory, visual-spatial skills, verbal fluency, and problem-solving		Dental composite restorations had statistically insignificant associations of small magnitude with impairments in neuropsychological test change scores over 4- or 5-years of follow-up in this trial.
Maserejian NN <i>et al.</i> , 2012 (c)	218 boys and 256 girls (6- 10 years)	Comparison between the baseline and 5 years after treatments	changes in BMI-for-age z-scores, body fat percentage (BF%), and height velocity; exploratory analyses (n = 113) examined age at menarche.		Children with more treatment on primary teeth had greater increases in BF% regardless of material type. Girls assigned to composites had lower risk of menarche during follow-up.
Chung SY <i>et al.</i> , 2012	495 children (8-9 years)	Four groups based on the number of resin composites and sealant surfaces (0, 1-5, 6-10 and 11+)	HPLC-ESI-MS/MS	In urine: - 11 or more resin composite surfaces: 2.67 µg/g creatinine (creatinine-adjusted urinary BPA Concentrations)	Having many dental composite filling surfaces on teeth may increase the urinary BPA concentration in children.
Han DH <i>et al.</i> , 2012	124 children	62 controls with no dental sealant/resin on their tooth surfaces and 62 cases with more than 4 tooth surfaces with dental sealant/resin	BPA ELISA Kit	In saliva: -Control (0.40 µg/L) - Case (0.92 µg/L)	There may be a relationship between salivary BPA level and dental sealant/resin.
Kingman A <i>et al.</i> , 2012	172 subjects	Immediately, 30 h	LC/MS	- In saliva: increase immediately (within 1 h) and return to preresoration levels within 8 h after composite placement. - In urine: return to preresoration levels 9 to 30 h after restoration placement	Rubber dam use reduced BPA saliva concentrations but did not reduce the absorption of BPA (measured as BPA level in urine) during the study.