



**HAL**  
open science

# Effects of chaotropic salts on global proteome stability in halophilic archaea: Implications for life signatures on Mars

Lorenzo Carré, Daniel Gonzalez, Éric Girard, Bruno Franzetti

## ► To cite this version:

Lorenzo Carré, Daniel Gonzalez, Éric Girard, Bruno Franzetti. Effects of chaotropic salts on global proteome stability in halophilic archaea: Implications for life signatures on Mars. *Environmental Microbiology*, 2023, 25 (11), pp.2216-2230. 10.1111/1462-2920.16451 . hal-04199698

**HAL Id: hal-04199698**

**<https://hal.science/hal-04199698>**

Submitted on 24 Nov 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

# Effects of chaotropic salts on global proteome stability in halophilic archaea: Implications for life signatures on Mars

Lorenzo Carré<sup>1</sup> | Daniel Gonzalez<sup>2</sup> | Éric Girard<sup>1</sup> | Bruno Franzetti<sup>1</sup> 

<sup>1</sup>Université Grenoble Alpes, CNRS, CEA, IBS, Grenoble, France

<sup>2</sup>Université Grenoble Alpes, Grenoble, France

## Correspondence

Bruno Franzetti, Université Grenoble Alpes, CNRS, CEA, IBS, Grenoble, France.  
Email: [franzetti@ibs.fr](mailto:franzetti@ibs.fr)

## Funding information

French National Research Agency, Grant/Award Number: ANR-15-IDEX-02; Interdisciplinary Research Institute of Grenoble (IRIG, CEA)

## Abstract

Halophilic archaea thriving in hypersaline environments, such as salt lakes, offer models for putative life in extraterrestrial brines such as those found on Mars. However, little is known about the effect of the chaotropic salts that could be found in such brines, such as MgCl<sub>2</sub>, CaCl<sub>2</sub> and (per)chlorate salts, on complex biological samples like cell lysates which could be expected to be more representative of biomarkers left behind putative extraterrestrial life forms. We used intrinsic fluorescence to study the salt dependence of proteomes extracted from five halophilic strains: *Haloarcula marismortui*, *Halobacterium salinarum*, *Haloferax mediterranei*, *Halorubrum sodomense* and *Haloferax volcanii*. These strains were isolated from Earth environments with different salt compositions. Among the five strains that were analysed, *H. mediterranei* stood out as a results of its high dependency on NaCl for its proteome stabilization. Interestingly, the results showed contrasting denaturation responses of the proteomes to chaotropic salts. In particular, the proteomes of strains that are most dependent or tolerant on MgCl<sub>2</sub> for growth exhibited higher tolerance towards chaotropic salts that are abundant in terrestrial and Martian brines. These experiments bridge together global protein properties and environmental adaptation and help guide the search for protein-like biomarkers in extraterrestrial briny environments.

## INTRODUCTION

By shaping the range of conditions in which proteins retain their stability and solubility, amino acid substitutions and structural modifications influence the survivability of organisms in a given environment. Due to their profound long-evolved specialization, extremophiles have largely confirmed this link between molecular traits and adaptation to the environment (Ando et al., 2021; Carré et al., 2022; Reed et al., 2013). For example, *haloarchaea*, which thrive on Earth in hypersaline environments, cope with hyperosmolarity by intracellularly accumulating molar concentrations of K<sup>+</sup>. Yet, their cytosol remains an extreme environment in which non-halophilic proteins would lose stability and solubility. Thanks to several molecular traits, including

reduced hydrophobicity and increased frequency of negatively charged amino acids; proteins from halophiles have shown impressive tolerance and even dependency towards high concentrations of salts. Differences between salt-tolerance of homologous enzymes from different halophilic organisms have been pointed (Coquelle et al., 2010). In particular, the evolution of the model malate dehydrogenase (MDH) enzyme within the haloarchaeal class showed complex and non-linear trajectories leading to various adaptive strategies and responses to salt type and pH (Blanquart et al., 2021). These observations indicate that even among salt-accumulating halophiles, subtle differences in molecular traits could be related to dramatic differences in protein behaviour and salt-dependency.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2023 The Authors. *Environmental Microbiology* published by Applied Microbiology International and John Wiley & Sons Ltd.

While most terrestrial halophiles, including the archaeon *Halobacterium salinarum*, which is the most halophilic organism known to date, are found in Na<sup>+</sup> dominated brines, it is important to note that halophiles can also be found in highly chaotropic Mg<sup>2+</sup>-containing environments such as soda lakes, Dead Sea, cryobrines or deep hypersaline anoxic basins (DHABs). Chaotropicity is a concept that was first defined in 1962 (Hamaguchi & Geiduschek, 1962). It results from the activities that solutes exert on macromolecular systems, as well as from deviations of solvation water's behaviour from its bulk-like state (Ball & Hallsworth, 2015). Chaotropic cations such as Mg<sup>2+</sup> or Ca<sup>2+</sup> destabilize most proteins (Bye & Falconer, 2013; Tadeo et al., 2007), as predicted by Hofmeister series (Hofmeister, 1888). These ions strongly impair water structure and interact with side-chains of negatively charged amino acids, alleviating key interactions involved in protein structure (General et al., 2008; Lenton et al., 2017; Okur et al., 2017). Absolute determinations of kosmotropic/chaotropic activities have been measured for several compounds, leading to negative (kosmotropic) and similar values for NaCl and KCl, respectively  $-11.0$  and  $-11.3$  kJ kg<sup>-1</sup> mol<sup>-1</sup>, and positive (chaotropic) values for MgCl<sub>2</sub> and CaCl<sub>2</sub>, respectively  $+54.0$  and  $+92.2$  kJ kg<sup>-1</sup> mol<sup>-1</sup> (Cray et al., 2013). Along with thermodynamical approaches, the chaotropic or kosmotropic nature of various types of physiological solutes was also quantified biologically, by studying how they affect microbial cell growth and metabolism (Cray et al., 2013; de Lima Alves et al., 2015). It revealed that, even at concentrations that can be found in the natural environment, chaotropic solutes can inhibit the growth of bacteria and induce the accumulation of stress proteins responsible for stabilizing proteins and membranes (Hallsworth, Heim, & Timmis, 2003; Hallsworth, Prior, et al., 2003). These observations indicate that, besides hypersalinity, chaotropicity represents a major challenge limiting the window of life. However, cellular activities and cultivable microbes were obtained from a MgCl<sub>2</sub>-rich DHAB. Also, *Halorubrum sodomense* and *Haloferax volcanii*, two halophilic archaea isolated from the Dead Sea show impressively high tolerance to chaotropic ions (Oren, 1983; Pohlschroder & Schulze, 2019) and intracellular concentrations of Mg<sup>2+</sup> in *H. sodomense* has been reported to be notably higher than in other halophilic archaea (Oren, 2013). Finally studies on halophilic enzymes also indicated a peculiar role of Mg in enzyme stabilization (Blanquart et al., 2021; Madern & Zaccai, 1997, 2004). These observations indicated that extreme halophiles may have evolved specific mechanisms to cope with high chaotropicity.

Besides a better definition of life window with respect to hypersaline and chaotropic environments, halophilic organisms and their related biomolecules have also been proposed as models to test the

compatibility of biological processes and biomarker stability under simulated hypersaline Martian conditions (Bayles et al., 2020; Carré et al., 2022; Perl & Baxter, 2020; Sundarasami et al., 2019). Indeed, several studies have shown that metastable hypersaline liquid water bodies containing high amount of chaotropic ions could be present beneath the southern pole ice cap (Lauro et al., 2021; Orosei et al., 2018) or under the regolith in subsurface environments (Abotalib & Heggy, 2019; Clifford et al., 2010; McEwen et al., 2011; Rivera-Valentín et al., 2020). As confirmed by experimental studies (Fischer et al., 2014), such water systems would require to be hypersaline in order to remain liquid under the low temperatures of Martian environments (Brady, 1992). A notable feature of halophilic organisms is their capacity to survive for long periods when trapped in salt crystals or upon salt deliquescence (Crits-Christoph et al., 2016; Favreau, 2023; Favreau et al., 2022; Micheluz et al., 2022), some studies even suggesting a preservation over geological ages (Hallsworth, 2022; Schreder-Gomes et al., 2022). Moreover, halophilic enzymes were also shown to maintain exceptional stability when trapped in salt crystals (Tehei et al., 2002). It has also been shown that intracellular accumulation of chaotropic cations such as Mg<sup>2+</sup> or Ca<sup>2+</sup>, facilitates life at low temperatures as they counter the cold-induced stabilization of proteins (Chin et al., 2010; Cray et al., 2015). For these reasons, it has been proposed that traces of ancient halophilic life could be present in Martian halites (Hallsworth, 2020; Rivera-Valentín et al., 2020). Thus, the surface of the red planet should not be considered universally hostile to life (Hallsworth, 2021). In this context, extensive simulations have used members of the *Haloarchaea* class to test the effects of Martian-like simulated conditions such as UV irradiation (Fendrihan et al., 2009), desiccation-rehydration cycles (Vauclare et al., 2020), long-term desiccation and freezing cycles (Mancinelli et al., 2004) and low temperature (Laye & DasSarma, 2018; Reid et al., 2006). While such experimental approaches are useful to assess the capacity to tolerate Martian-like conditions at the organism level, they need to be complemented with studies on the biomolecules to get a complete picture of the molecular limits of life. In addition, Martian brines would display Mg<sup>2+</sup> and Ca<sup>2+</sup> as the dominant cations (Osterloo et al., 2008; Squyres et al., 2004) and high concentrations of chlorate and perchlorate anions (ClO<sub>3</sub><sup>-</sup> and ClO<sub>4</sub><sup>-</sup>), have been detected in situ (Hecht et al., 2009), especially in subsurface brines. As mentioned above, such chaotropic conditions represent a barrier to life (Hallsworth et al., 2007) and could make the Martian brines more extreme than their terrestrial analogues (Fox-Powell et al., 2016). At the molecular level, little is known about the effects of chaotropic ions on halophilic proteins and the existence of protein adaptation to chaotropic ions remains questionable. Such adaptation

would be crucial to cope with exotic conditions such as those present in Martian brines.

Recently, the effects of perchlorate salts on the reactivity of an halophilic enzyme (Laye & DasSarma, 2018) and on the stability and reactivity of a non-halophile enzyme have been investigated (Gault & Cockell, 2021). It should, however, be stressed that the conclusions arising from isolated model biomolecules may not reflect the limiting biological edifice required to cope with extreme conditions and, in the context of astrobiology, may not indicate what biomarker resulting from a putative extraterrestrial lifeform could be searched for (Fairén et al., 2020; Sánchez-García et al., 2020). Moreover, such approaches could be biased by the choice of the enzyme. In addition to variations between organisms, even between extreme halophiles (Coquelle et al., 2010), proteins from the same organism may display drastically different responses to salts, notably as a consequence of the composition of their amino acid sequence (Becker et al., 2014). Therefore, whole-proteome approaches to study the properties of a physiologically relevant mix of proteins may offer additional insights into global biomolecule adaptation to the environment. Various techniques, including neutron scattering (Zaccai, 2013), have been successfully used to unravel global protein adaptations in thermophiles, psychrophiles (Tehei et al., 2004), piezophiles (Martinez et al., 2016; Peters et al., 2014) and halophiles (Marty et al., 2013; Vauclare et al., 2015, 2020).

By using label-free differential scanning fluorimetry (nanoDSF), a method allowing analysis of thermal denaturation and thermal aggregation of proteins, we have recently revealed a difference between the stabilizing effect of KCl and NaCl on the proteome of the halophilic archaeon *Haloarcula marismortui* (Carré et al., 2021). In this study, we used this novel approach to explore the effects of combinations between kosmotropic (KCl, NaCl) and chaotropic salts ( $\text{MgCl}_2$ ,  $\text{CaCl}_2$ ,  $\text{NaClO}_3$ ,  $\text{NaClO}_4$ ,  $\text{Mg}(\text{ClO}_4)_2$  and  $\text{Ca}(\text{ClO}_4)_2$ ) on proteome stability in five different strains of *Haloarchaea*, including Mg-tolerant strains.

## EXPERIMENTAL PROCEDURES

### Experimental design

This study aimed to characterize response to various salt types and combinations at the scale of the proteome through nanoDSF analysis of cell lysates of halophilic archaea. After diluting them in concentrated salt solutions to achieve desired conditions, cell lysates were loaded in capillaries and analysed with a Prometheus NT.48 (NanoTemper). This allowed monitoring of denaturation and aggregation events throughout a temperature gradient. We focused on the shifts of dominant

denaturation peaks ( $T_{Pm}$ ) related to changes in salt types, combinations and concentrations. Relationships between  $T_{Pm}$  and salt concentration were fitted with a linear model whose slopes represented proteome sensitivity to the given salt.

### Strains and growth conditions

All strains were grown aerobically at 37°C under mild agitation by inoculating fresh medium with concentrated cell suspension stored in glycerol at -80°C. Initial stocks were purchased at the DSMZ-German Collection of Microorganisms (*H. mediterranei*, *H. sodomense*) or provided by A. Oren (*H. marismortui*, *H. salinarum*) or H. Myllykallio (*H. volcanii*). Growth medium was adapted from various references (Mevarech et al., 1976; Oesterhelt & Stoekenius, 1974; Oren, 1983; Rodriguez-Valera et al., 1980).

Composition of *H. marismortui* medium was 208 g L<sup>-1</sup> NaCl, 78 g L<sup>-1</sup>  $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ , 500 mg L<sup>-1</sup>  $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ , 125 mg L<sup>-1</sup>  $\text{MnCl}_2$ , 5 g L<sup>-1</sup>, NaBr 580 mg L<sup>-1</sup>, 10 g L<sup>-1</sup> yeast extract, pH adjusted to 7.5 with addition of 10 M NaOH after dissolution.

Composition of *H. salinarum* medium was 250 g L<sup>-1</sup> NaCl, 33.8 g L<sup>-1</sup>  $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ , 2 g L<sup>-1</sup> KCl, 125 mg L<sup>-1</sup>  $\text{MnCl}_2$ , NaBr 580 mg L<sup>-1</sup>, 3 g L<sup>-1</sup> yeast extract, 5 g L<sup>-1</sup> tryptone, pH adjusted to 7.2 with addition of 10 M NaOH after dissolution.

Composition of *H. mediterranei* medium was 195 g L<sup>-1</sup> NaCl, 84 g L<sup>-1</sup>  $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ , 500 mg L<sup>-1</sup> KCl, 1 g L<sup>-1</sup>  $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ , 125 mg L<sup>-1</sup>  $\text{MnCl}_2$ , NaBr 580 mg L<sup>-1</sup>, 5 g L<sup>-1</sup> yeast extract, pH adjusted to 7.2 with addition of 10 M NaOH after dissolution.

Composition of *H. sodomense* medium was 125 g L<sup>-1</sup> NaCl, 160 g L<sup>-1</sup>  $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ , 2.1 g L<sup>-1</sup> KCl, 130 mg L<sup>-1</sup>  $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ , 125 mg L<sup>-1</sup>  $\text{MnCl}_2$ , 5 g L<sup>-1</sup>, NaBr 580 mg L<sup>-1</sup>, 2 g L<sup>-1</sup> yeast extract, 1 g L<sup>-1</sup> peptone, 2 g L<sup>-1</sup> soluble starch, pH adjusted to 7.2 with addition of 10 M NaOH after dissolution.

Composition of *H. volcanii* medium was 125 g L<sup>-1</sup> NaCl, 62 g L<sup>-1</sup>  $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ , 3.7 g L<sup>-1</sup> KCl, 500 mg L<sup>-1</sup>  $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ , 125 mg L<sup>-1</sup>  $\text{MnCl}_2$ , 5 g L<sup>-1</sup>, NaBr 580 mg L<sup>-1</sup>, 3 g L<sup>-1</sup> yeast extract, 5 g L<sup>-1</sup> tryptone, pH adjusted to 7.2 with addition of 10 M NaOH after dissolution.

### Proteome extracts preparation

Sixty millilitres of a culture in mid or late exponential phase ( $\text{OD}_{600}$  between 0.4 and 1.0) were centrifugated at 20,000 g for 15 min. Supernatant was discarded by aspiration and pellet was resuspend in 3 mL of 4 M KCl or 5 M NaCl and cells were then lysed by sonication. For each condition, 50 µL of cell lysate was diluted with

water and concentrated salt solutions (4 M KCl, 5 M NaCl, 5 M MgCl<sub>2</sub>, 5 M CaCl<sub>2</sub>, 2 M NaClO<sub>3</sub>, 2 M NaClO<sub>4</sub>, 1 M Mg(ClO<sub>4</sub>)<sub>2</sub> or 1 M Ca(ClO<sub>4</sub>)<sub>2</sub>) to the final volume of 500 µL in order to reach the final desired concentration of salts.

### Characterization of effects of salts on *Haloarchaea* proteomes and *H. marismortui* MDH by differential scanning fluorimetry

Cell lysates extracted in various salt conditions were centrifugated at 20,000 g for 15 min and the supernatants (soluble fraction of cell lysate) were loaded in Prometheus™ NT.48 capillaries. Thermal denaturation curves were determined by measurements of protein intrinsic fluorescence. This analysis was performed using label-free, native differential scanning fluorimetry (nanoDSF; apparatus: Prometheus NT.48, NanoTemper). The tryptophan residues of the proteins in proteome extracts were excited at 280 nm and the fluorescence intensity was recorded at 330 and 350 nm. Excitation power was set at 80% or 100%. The temperature of the measurement compartment increased from 20 to 95°C at a rate of 2°C per minute. Light scattering was also measured as a way to detect aggregation events during heating.

To stress the inner variation of first derivative of ratio of 350/330 nm fluorescence in each sample, data were transformed using the highest and the lowest values within the 25–80°C range of temperature as boundaries by respectively setting them to 100 and 0.

### Statistical analysis

All graphs and analyses were made using GraphPad Prism 9 (GraphPad software Inc, California, USA).

Temperature associated with dominant denaturation peaks were automatically determined by the PR. ThermControl software (NanoTemper). When multiple peaks were detected, we only picked the dominant one with maximum height. When no temperature could be automatically determined, we used instead the temperature associated with maximum value of first derivative of 350/330 nm fluorescence ratio. Since events associated with abnormally high fluorescence or light scattering signals occasionally happened at highest temperatures in some individual experiments, the manual assignment of denaturation peaks was limited to temperatures between 25 and 80°C. Due to low signal over noise ratio, no denaturation peak and, hence,  $T_{Pm}$ , could be determined for several experiments, mostly with lysates of *H. salinarum*.

For each condition, linear regression was made by using the automatically and manually determined  $T_{Pm}$ .


Due to insufficient data, it could not be made for *H. salinarum* with the NaCl-only salt condition. For KCl-only and NaCl-only experiments, value of  $T_{Pm}$  measured at 0 M salt was excluded from the fit as this condition is actually particularly extreme for halophilic systems. This was not made with combinations since the proteome was always exposed to at least 1 M or counter-salt (KCl or NaCl). Slope of line fitting best  $T_{Pm}$  values was determined individually for each replicate. Statistical comparison of slopes in Figure 1A, was made using Brown–Forsythe and Welch ANOVA tests, comparing the mean of each column to the mean of every other column. All peak and slope values are given in Data S2–S5.

Since differences between slope values ranged over several orders of magnitude, data were transformed for Figure 2A. For each salt condition, the slope with highest absolute value was set to 100 or –100, depending on its sign, by multiplying it by a specific number and the other slopes were multiplied by the same number. This made comparisons between strains easier for every salt condition.


## RESULTS

### Response of the proteome of *H. marismortui* to different salts


To investigate proteome responses to KCl, NaCl, MgCl<sub>2</sub>, CaCl<sub>2</sub> and to combinations between them, we first used lysates of cells of *H. marismortui* in exponential phase. After dilution in various salt concentrations to reach the final desired composition and centrifugation, the supernatants, which contained the soluble fraction of the proteomes, were analysed using nanoDSF to derive proteome thermal denaturation and thermal aggregation curves as described in Carré et al. (2021). Untreated thermal aggregation and denaturation signals in various salt conditions are presented in supplementary material section (Figures S2–S5). Representative thermal denaturation curves for the proteome of *H. marismortui* are given in Figure 3 for solutions containing a single salt and for combinations between 1 M KCl or NaCl and chaotropic chloride or (per)chlorate salts. We first tested the effects of solutions containing one type of salt: KCl (0–4 M), NaCl (0–6 M), MgCl<sub>2</sub> (0–2 M) and CaCl<sub>2</sub> (Figure 3A). Then we studied combinations between KCl (1, 2 or 3 M) and chaotropic MgCl<sub>2</sub> or CaCl<sub>2</sub> (0–2 M) (Figure 3B) as well as combinations between NaCl (1, 2 or 3 M) and chaotropic chlorate/perchlorate salts: NaClO<sub>3</sub> (0–1 M), NaClO<sub>4</sub> (0–1 M), Mg(ClO<sub>4</sub>)<sub>2</sub> (0–0.5 M) or Ca(ClO<sub>4</sub>)<sub>2</sub> (0–0.5 M) (Figure 3C). The reason why we used NaCl rather than KCl with perchlorates is that immediate precipitation occurred when mixing (per)chlorate and KCl solutions, even at low concentrations.



**FIGURE 1** Proteome melting temperatures as a function of salt condition and proteome sensibility to various Earth-relevant salts in *Haloarchaea*.  $T_{Pm}$  values determined with KCl or NaCl only (A) and with combinations between KCl and  $MgCl_2$  (D). Corresponding slopes of line fitting  $T_{Pm}$  determined with KCl (B), NaCl (C) or combinations between KCl and  $MgCl_2$  (E). Statistical analysis was made using a Brown–Forsythe and Welch ANOVA test (\*\* $p < 0.0021$ ; \*\*\* $p < 0.0002$ ; \*\*\*\* $p < 0.0001$ ). Errors bars represent standard deviation.



**FIGURE 2** Proteome melting temperatures as a function of salt condition and proteome sensibility to various Mars-relevant salts in *Haloarchaea*.  $T_{Pm}$  values determined with combinations between NaCl and  $NaClO_3$  (A),  $NaClO_4$  (C),  $Mg(ClO_4)_2$  (E) or  $Ca(ClO_4)_2$  (G). Corresponding slopes of line fitting  $T_{Pm}$  determined with combinations between NaCl and  $NaClO_3$  (B),  $NaClO_4$  (D),  $Mg(ClO_4)_2$  (F) or  $Ca(ClO_4)_2$  (H).



**FIGURE 3** Representative thermal denaturation curves of *Haloarcula marismortui* proteome under various salt conditions. nanoDSF analysis was made with soluble fraction of lysates with only one type of salt (A), combinations between 1 M KCl and  $MgCl_2$  or  $CaCl_2$  (B) or with combinations between 1 M NaCl and (per)chlorate salts (C). Higher values (yellow) correspond to denaturation events. Curves were transformed to facilitate peak observation (see [Experimental Procedures](#) section). Occasionally, abnormally high signals were measured at the end of the heating ramp (in black). When signal over noise ratio was high, signal after 80°C may also appear in black due to the way colour range was set.

In the case of chaotropic salts, increasing the concentrations of both  $MgCl_2$  and  $CaCl_2$  delayed peak of the first derivative of the ratio of 350/330 nm

fluorescence and, hence, the dominant denaturation event (denaturation peak) reflecting a stabilizing effect of these two salts on the *H. marismortui* proteome in

spite of their chaotropicity (Figure 3A). However, the relationship between  $\text{CaCl}_2$  concentration and the temperature of the denaturation peak was non-linear.

Denaturation peaks followed a non-linear trend under KCl-MgCl<sub>2</sub> or KCl-CaCl<sub>2</sub> conditions. However, the denaturation events were less resolved than in single-salt conditions (Figure 3B). Yet, one dominant denaturation peak could still be observed with most KCl-MgCl<sub>2</sub> conditions. Several KCl-CaCl<sub>2</sub> conditions led to higher values of the first derivative of the ratio of 350/330 nm fluorescence above 80°C, which was used as the superior temperature boundary for determining the peak position in low-resolved curves. When excluding the events occurring above 80°C, dominant denaturation events could be observed at 45–65°C.

Combinations between NaCl and chlorate or perchlorate salts led to less resolved denaturation peaks similar to the KCl-MgCl<sub>2</sub> or KCl-CaCl<sub>2</sub> conditions (Figure 3C). The presence of chlorate or perchlorate salts caused denaturation peaks to occur at an even lower temperature, showing a stronger destabilizing effect compared to the KCl-MgCl<sub>2</sub> or KCl-CaCl<sub>2</sub> conditions. At low concentrations, chlorate or perchlorate salts do not seem to alleviate significantly the stabilizing effect of NaCl (Figure 3A).

Since the Prometheus NT.48, which we used for the nanoDSF assay, also measures light scattering, we used peaks of first derivative of light scattering as a proxy for aggregation events. However, aggregation peaks could only be observed at high concentrations of NaCl/KCl or with magnesium or calcium-containing chloride or perchlorate salts (Figure S1). KCl-MgCl<sub>2</sub> combinations allowed observation of two separated well-resolved aggregation peaks, the first happening at  $T < 75^\circ\text{C}$  and the second at higher temperatures. Both MgCl<sub>2</sub> and CaCl<sub>2</sub> promoted aggregation events since increasing their concentration lowered the temperatures at which these events were observed.

### Salt responses of the proteomes from *H. salinarum*, *Haloferax mediterranei*, *H. sodomense* and *H. volcanii* (compared to the one of *H. marismortui*)

To compare proteome response to salt conditions between species, the nanoDSF experiments were extended to cell lysates of four additional haloarchaeal strains: *H. salinarum*, *H. mediterranei*, *H. sodomense* and *H. volcanii*. These strains were isolated from environments with contrasting salt conditions and displayed different requirements for salt concentrations in their growth media (see Experimental Procedures section). In particular, *H. salinarum* is the most NaCl-dependent and NaCl-tolerant known organism, *H. sodomense* is the most Mg<sup>2+</sup>-tolerant organism known and *H. mediterranei* has the largest range of NaCl

concentrations compatible with growth. Details about salt requirements for growth and other microbiological information about these five strains from Bergey's Manual of Systematics of Bacteria and Archaea and is given in Table S1.

Thermal aggregation curves (Figure S2) were similar between strains either when only one type of salt was present or with salt combinations. Qualitative differences could only be seen between MgCl<sub>2</sub> and CaCl<sub>2</sub>, the latter promoting aggregation events at lower temperatures more drastically than MgCl<sub>2</sub>. Nonetheless, aggregation peaks were smaller at higher concentrations of KCl, especially at 3 M when combined with CaCl<sub>2</sub>. MgCl<sub>2</sub> favoured observation of two aggregation peaks at lower temperatures as its concentration increased. CaCl<sub>2</sub> leads to the observation of additional dominant aggregation events whose temperature was positively correlated with CaCl<sub>2</sub> concentration. Interestingly, this was not observed with proteomes of *H. marismortui* and *H. salinarum* the two most NaCl-dependent strains.

In contrast, the denaturation curves largely differed between strains. Figure S3 shows representative denaturation curves under one-salt conditions for all tested strains. Overall, proteomes showed similar behaviour, all being gradually stabilized by higher KCl or NaCl concentrations. Stabilization by MgCl<sub>2</sub> and CaCl<sub>2</sub> was also clearly observed with the proteomes of *H. mediterranei* and *H. sodomense* and, to a lesser extent, with *H. volcanii*. Cell lysates of *H. salinarum* were associated with the least resolved denaturation peaks, making their observation difficult or even impossible, especially when only MgCl<sub>2</sub> or CaCl<sub>2</sub> was present. Interestingly, thermal denaturation curves of lysates of *H. mediterranei* showed two clearly separated dominant denaturation events.

With salt combination experiments, increasing the concentration of chaotropic salt promoted observation of more than one noticeable denaturation event (Figure S4). As a reminder, variation of fluorescence signal is a function of exposure of aromatic amino acids to solvents upon protein denaturation.

This was observed with proteomes of *H. sodomense* and *H. volcanii* at 1 and 2 M KCl with increasing MgCl<sub>2</sub> concentrations. The first event to occur, which was observed at 1 and 2 M and unseen at 3 M KCl, was shifted towards low temperatures when MgCl<sub>2</sub> concentration increased. On the contrary, the second event to occur happened at a slightly higher temperature as MgCl<sub>2</sub> concentration increased from 0 to 2 M. This showed that destabilizing as well as stabilizing effects of MgCl<sub>2</sub> on both proteomes could be observed in the same conditions.

Denaturation peaks were less resolved with KCl-MgCl<sub>2</sub> combinations for proteomes of *H. marismortui* and *H. salinarum*. However, it could be confirmed for these strains that increasing concentration of MgCl<sub>2</sub>



lowered temperatures at which denaturation peaks could be observed at most KCl concentrations, showing an overall destabilizing effect of  $\text{MgCl}_2$ . Use of  $\text{CaCl}_2$  as the chaotropic salt reduced resolution of denaturation peaks for all strains, with the notable exception of *H. mediterranei* which maintained well-resolved denaturation peaks.

Effects of (per)chlorate-containing salts on proteome stability were similar between strains (Figure S5).  $\text{NaClO}_3$  was nearly always stabilizing while  $\text{NaClO}_4$  and  $\text{Mg}(\text{ClO}_4)_2$  had destabilizing effects. However, these effects were reduced by increasing the concentration of NaCl.  $\text{Ca}(\text{ClO}_4)_2$  had a complex non-linear effect on temperature associated with dominant denaturation events and generally reduced signal over noise ratio. As previously noted, denaturation curves obtained with proteomes of *H. marismortui* and *H. salinarum* were less resolved and, as with single-salt experiments, denaturation curves of proteome of *H. mediterranei* (Figure S5C) presented two distinct dominant denaturation events in most conditions.

## Statistical analyses to compare proteome melting temperatures and sensitivity

To enable statistical comparisons, experiments with KCl or NaCl only and with salt combinations (except KCl- $\text{CaCl}_2$ ) were independently replicated for the five strains: three additional experimental points with fewer salt concentrations for KCl or NaCl only and two additional experimental points with the same conditions for KCl- $\text{MgCl}_2$  and NaCl-(per)chlorate combinations. The temperatures associated with the peak value of the first derivative of ratio of 350/330 nm fluorescence associated with the main dominant denaturation events, which we previously defined as proteome melting temperatures ( $T_{\text{Pm}}$ ) (Carré et al., 2021), were retrieved. The slope of lines fitting  $T_{\text{Pm}}$  as a function of the concentration of the salt were also calculated.  $T_{\text{Pm}}$  and slope values are given in Figures 1 and 2.

The reproducibility of the measured  $T_{\text{Pm}}$  was good in spite of the complexity of the samples as shown by the low standard deviation values (Figures 1 and 2) except for *H. salinarum* for which it was not always possible to derive a  $T_{\text{Pm}}$  value. Variations in  $T_{\text{Pm}}$  values could be observed and were dependent on the strain and the salt type. The previously observed trends showed to be robust and reproducible. Increasing KCl or NaCl concentrations nearly always increased  $T_{\text{Pm}}$  in salt combinations experiments (Figures 1A,D and 2A,C,E,G). At every KCl concentration, the relationship between  $\text{MgCl}_2$  and  $T_{\text{Pm}}$  was complex and non-linear (Figure 1D).  $\text{NaClO}_3$  reproducibly stabilized proteomes and this effects linearly increased with its concentration (Figure 2A). In contrast,  $\text{NaClO}_4$  gradually decreased

$T_{\text{Pm}}$  in most of the conditions (Figure 2C). Effects of  $\text{Mg}(\text{ClO}_4)_2$  were similar to those due to  $\text{NaClO}_4$  (Figure 2E). Like  $\text{MgCl}_2$  in the presence of KCl,  $\text{Ca}(\text{ClO}_4)_2$  had complex non-linear effects on  $T_{\text{Pm}}$ , increasing stability of proteomes of *H. sodomense* and *H. volcanii* at low concentrations and then destabilizing them at higher concentrations (Figure 2G).


While not all salts lead to linear relationships between their concentration and the  $T_{\text{Pm}}$ , the slopes of the fitting line enabled the comparison of global proteome sensitivity to a given salt type, regardless of its intrinsic thermostability. No statistically significant differences in proteome stabilization by KCl were observed between the five tested strains (Figure 1B). In contrast, the proteome of *H. mediterranei* presented a statistically significant higher stabilization response to NaCl (Figure 1C), being more efficiently stabilized by this salt than the other strains (with the exclusion of *H. salinarum* whose proteome did not offer enough well-resolved denaturation peaks). In addition, the proteome of *H. salinarum* was more destabilized by  $\text{MgCl}_2$  (Figure 1E), less stabilized by  $\text{NaClO}_3$  (Figure 2B) and more destabilized by  $\text{NaClO}_4$  (Figure 2D). *H. sodomense* and *H. volcanii* presented strikingly similar proteome responses to all salt types, being the least sensitive to  $\text{NaClO}_4$  or  $\text{Mg}(\text{ClO}_4)_2$  and the only ones being stabilized by  $\text{Mg}(\text{ClO}_4)_2$  (Figure 2F) or  $\text{Ca}(\text{ClO}_4)_2$  (Figure 2H) in the presence of 3 M NaCl. Proteomes of *H. marismortui* and *H. mediterranei* showed intermediary behaviours between those observed for *H. salinarum* on the one hand and *H. volcanii*/*sodomense* on the other hand. In addition, both were very sensitive to  $\text{Mg}(\text{ClO}_4)_2$  (Figure 2F).

To assess possible correlations between sensitivity to the tested salt conditions, Pearson correlation coefficients were calculated using slope values from the five tested strains and the corresponding correlation matrix determined (Figure 4A). Sensitivities to  $\text{MgCl}_2$ ,  $\text{NaClO}_3$  and  $\text{NaClO}_4$  were positively correlated as were sensitivities to  $\text{Mg}(\text{ClO}_4)_2$  and  $\text{Ca}(\text{ClO}_4)_2$ . However, positive and negative correlations were dependent on the counter-salt (NaCl or KCl) concentration.

Our results clearly show that the proteome response to chaotropic salts differs between strains. This is summarized in Figure 4B, highlighting distinct behaviours, such as high sensitivity to NaCl in the case of *H. mediterranei* proteome and its peculiar response to Martian or Earth-relevant salts.

## DISCUSSION

Cellular proteomes contain different types of proteins with different biochemical and biophysical properties thus, studies related to single proteins cannot properly reflect the adaptative nature of the whole proteome. With this respect, nanoDSF is a valuable approach to



**FIGURE 4** Correlation between strains and sensitivities to salt. (A) Correlation matrix of sensitivities to salt over the five strains. Dotted lines separate conditions with same main salt with, when applicable, the conditions of counter-salt concentrations. (B) Radar chart of the normalized slopes of line fitting  $T_{Pm}$  values of proteomes of each haloarchaeal strain for every tested salt condition. Grey zone corresponds to negative slopes and, hence, destabilizing effects of salts whereas white zone corresponds to positive slopes and stabilizing effects.

probe molecular response applied to chemical conditions at the proteome scale. In the present work, we extended our proof of concept experiment (Carré et al., 2021) to the comparison of several halophilic extremophiles. Obviously, the intrinsic fluorescence signal depends on the existence of aromatic residues within the sample and comparison between strains may be biased if aromatic residue content differed. Thus, we compared frequencies of Phe/Tyr/Trp residues in the proteomes of the five strains and within a set of 93 representative proteins listed in Data S1, distribution being represented in Figure S6. While the genomes of both strains encode on average similar frequencies of the aromatic amino acids, *H. sodomense* showed a reproducibly strong nanoDSF signal in our experiments, unlike *H. salinarum*. Nonetheless, the frequency of the

aromatic amino acids in *H. salinarum* within the 93 representative proteins set is also close to the frequencies found in the genomes with *H. marismortui* and *H. sodomense*. Another aspect that may influence nanoDSF signal interpretation is the relative abundance of certain protein families within the proteome. However, our study relies only on *haloarchaea* strains, limiting the differences in proteome complexity and relative composition. Our exploration of the halophilic proteome response revealed the existence of limited protein denaturation and/or aggregation events happening at clearly distinct temperatures in several salt conditions, depending on the strain. This observation could be related to the fact that distinct protein populations behave differently with respect to salt nature and concentration.

The first important information provided by the nanoDSF experiments performed is that the proteome stability, with respect to salt, differed between the considered strains in spite of the fact that they are all strictly hyperhalophilic *Haloarchaea* accumulating molar concentrations of potassium ions in their cytosol. Of the five tested strains, *H. salinarum* and *H. mediterranei* showed the most distinctive behaviours. The proteome of *H. salinarum* gave the least resolved denaturation curves. This happened with every tested salt type, including the most chaotropic. This could be explained by a reduced frequency of aromatic residues contributing to the measured intrinsic protein fluorescence, hence reducing the peak height. As discussed, frequency of aromatic amino acids in the five considered strains is close. Therefore, scarcity of aromatic amino acids does not offer a reasonable explanation for the low-resolution denaturation peaks obtained with lysates of *H. salinarum*. Our observations and the similar aromatic residue frequency may indicate that proteins composing the cell lysates of *H. salinarum* may denature on a broader range of temperatures, thus limiting peak sharpness.

*H. mediterranei* stands out in another way: its proteome stability displayed significantly higher dependency on  $\text{Na}^+$  than the other strains. While all haloarchaea, including *H. mediterranei* (Pérez-Fillol & Rodríguez-Valera, 1986), intracellularly accumulate  $\text{K}^+$  rather than  $\text{Na}^+$ , the latter remains the dominant cation in most hypersaline environments. Most studies about intracellular salt composition in haloarchaea and its dependency on extracellular environment have essentially investigated intracellular  $\text{K}^+$  associated to medium with modified NaCl concentrations at constant  $\text{K}^+$  supply. In spite of the fact that no study has, to our knowledge, measured the intracellular concentration of  $\text{Na}^+$  as a function of  $\text{K}^+$  concentration in the growth medium, some studies with *H. marismortui* (Ginzburg et al., 1970; Jensen et al., 2015) and in *H. salinarum* (Lanyi & Silverman, 1972; Matheson et al., 1976) presented evidences of, respectively, decreased and increased accumulation of  $\text{K}^+$  and  $\text{Na}^+$  when  $\text{K}^+$  is scarce in the medium. This indicates that haloarchaea may accumulate  $\text{Na}^+$  instead when  $\text{K}^+$  is scarce. In such conditions, proteome of *H. mediterranei* could therefore benefit from  $\text{K}^+$  scarcity by having a proteome that can still benefit from  $\text{Na}^+$  abundancy. It should also be pointed out that *H. mediterranei* possess fewer carotenoid pigments and lacks both bacteriorhodopsin and halorhodopsin (Oren, 2017; Oren & Hallsworth, 2014), which are photodependent  $\text{H}^+$  and  $\text{Cl}^-$  pumps essential for  $\text{K}^+$  accumulation and for intracellular ion homeostasis in *Haloarchaea* (Oren, 1999). Moreover, *H. mediterranei* is the only strain among the five tested whose genome encodes the mechanosensitive transporter MscS protein, which promotes  $\text{K}^+$  effusion out of the cell, and all known haloarchaeal  $\text{Mg}^{2+}$  transporter

whereas genome of *H. volcanii* encodes only two of them (MgtA and MgtB) and *H. salinarum* and *H. marismortui* none (Table S2) (Becker et al., 2014).

Our study shows a positive correlation between the NaCl requirement for growth and proteome sensitivity to chaotropicity nature. Indeed, among the five studied strains, *H. salinarum* is the one requiring the highest amount of NaCl for optimal growth and is also the one that tolerates the highest NaCl concentration in its environment (Oren & Ventosa, 2017). Interestingly, *H. salinarum* proteome was destabilized by chaotropic salts,  $\text{MgCl}_2$  as well as (per)chlorate salts. By comparison, *H. sodomense* and *H. volcanii* optimally grow at lower NaCl concentrations (McGenity & Grant, 2015; Ventosa et al., 2015) and showed proteomes that are the least destabilized and even distinctively stabilized in some occasions by chaotropic salts. Moreover, *H. mediterranei* has the largest NaCl concentration range compatible with growth and tolerates both high and low NaCl concentrations (Oren, 2017; Oren & Hallsworth, 2014) and showed a proteome with intermediary behaviour in our measurements. Therefore, growth requirement for NaCl appears to be correlated with proteome sensitivity to  $\text{MgCl}_2$  and (per)chlorate salts, underlying the link between proteome intrinsic properties and adaptation to the environment. Moreover, our results indicate that proteome tolerance to  $\text{MgCl}_2$  is correlated to tolerance to perchlorate salts, suggesting the existence of common proteome traits allowing tolerance to general salt-induced chaotropicity. Whether this tolerance resulted from primary structure of proteins, compatible solutes accumulation, intracellular salinity, or expression of specific proteins such as chaperones or ion channels remains to be determined.

In cell lysates of *H. mediterranei*, two subpopulations of proteins denaturing at different temperatures were observed in NaCl or KCl single-salt conditions. One explanation could be that the proteins of this organism are, on average, more soluble over a larger range of KCl/NaCl conditions and, hence, more diverse with respect to the salt concentration effect. As this organism has one of the largest range of NaCl concentrations compatible with growth among haloarchaea (Oren, 2017; Oren & Hallsworth, 2014), this could reflect an adaptative trait allowing proteins to function over a broader range of environmental salt conditions. With *H. sodomense* and *H. volcanii*, which are two strains whose growths are exceptionally tolerant to  $\text{MgCl}_2$  (Mullakhanbhai & Larsen, 1975; Oren, 1983), increasing  $\text{MgCl}_2$  concentration at constant KCl concentration also segregated two subpopulations of proteins denaturing at different temperatures with one subpopulation stabilized by this divalent salt while the second is destabilized. These results further stress the complexity of the proteomes, which contains proteins with various response to high concentrations of  $\text{Mg}^{2+}$ .

Besides denaturation, multiple aggregation events were observed with all halophilic strains at reproducibly similar temperatures with combinations of KCl-MgCl<sub>2</sub> and NaCl-Mg(ClO<sub>4</sub>)<sub>2</sub>. Both populations of aggregates formed at lower temperatures as the concentration of Mg-containing salt increased. Mg<sup>2+</sup> hence segregates protein populations during thermal aggregation in a similar manner across strains. Interestingly, Hofmeister series predict instead that this salt should increase solubility and, hence, could limit aggregation. However, these predictions are mostly based on observations made with non-halophilic proteins, at a constant temperature, lower concentrations of chaotropic salts and generally for folded proteins. The thermal aggregation which happened during the experiments is made through hydrophobic interactions between residues exposed to solvent by the thermal unfolding. Since we observed, at least with some of the strains, two peaks in both denaturation and aggregation curves, it could be hypothesized that they are both formed by the same two distinct protein subpopulations. For example, a subpopulation more sensitive to Mg<sup>2+</sup> could denature and aggregate before the other, hence producing the first peaks, followed by the other subpopulation forming the latter peaks.

High concentrations of chaotropic salts such as MgCl<sub>2</sub> correspond to complex extreme conditions as they combine high salinity, high ionic strength, low water activity and high chaotropy. This is especially true with magnesium/calcium perchlorate salts which also add oxidative stress. Hence, it is commonly accepted that the effects of such salts on biomolecules are not well predictable and depend on the system. While the Hoffmeister series predict some of their effects, MgCl<sub>2</sub> and CaCl<sub>2</sub> increasing protein solubility and reducing protein stability, they do not indicate precisely the concentrations at which these salts become deleterious to biological systems.

We showed that increasing concentration of MgCl<sub>2</sub> reproducibly increased average thermostability of the proteomes, especially when the medium contained low concentrations of KCl. This effect continued to be observable at the highest MgCl<sub>2</sub> concentration values in *H. sodomense* and *H. volcanii*. This observation is astonishing as the concentrations of MgCl<sub>2</sub> or CaCl<sub>2</sub> tested in this study are considered to be particularly extreme at scales of both cellular life and biomolecular systems (Hallsworth et al., 2007; Yakimov et al., 2015), especially in the context of astrobiology and Martian brines (Fox-Powell et al., 2016; Hallsworth, 2021). For example, 2 M of CaCl<sub>2</sub> reduces by 20°C the *T<sub>m</sub>* of bovine pancreatic ribonuclease (Von Hippel & Wong, 1965) and the model MDH enzyme from *H. marismortui* has previously shown to be fully denatured by 2 M CaCl<sub>2</sub>.

Interestingly, in spite of being more chaotropic and oxidizing than chloride salts, Mars-relevant salts had

contrasting effects on proteome stability as illustrated by the *T<sub>Pm</sub>* values (Figure 4). NaClO<sub>3</sub> was always stabilizing all halophilic proteomes, except for *H. salinarum* at 3 M NaCl. NaClO<sub>4</sub> had a destabilizing effect, especially for *H. salinarum*, but only marginally for *H. sodomense* and *H. volcanii*. The trend remained the same for Mg(ClO<sub>4</sub>)<sub>2</sub> with even more destabilized proteomes of *H. marismortui*, *H. salinarum* and *H. mediterranei*, and again significantly less effect on *H. sodomense* and *H. volcanii*. In contrast, concentrations of Ca(ClO<sub>4</sub>)<sub>2</sub> below 0.5 M induced noticeable proteome stabilization with *H. sodomense* and *H. volcanii* at every NaCl concentration. The proteome of *H. mediterranei* was also stabilized by Ca(ClO<sub>4</sub>)<sub>2</sub> but only at 1 M NaCl. Since adding Ca(ClO<sub>4</sub>)<sub>2</sub> drastically diminished peak resolution with *H. salinarum*, *T<sub>Pm</sub>* determination may be even more biased with this strain. Interestingly, a recent study showed, with a similar thermostability-based approach, that the bovine α-chymotrypsin was stabilized by MgSO<sub>4</sub>, slightly stabilized by low concentrations of MgCl<sub>2</sub> and significantly destabilized by NaClO<sub>4</sub>, Mg(ClO<sub>4</sub>)<sub>2</sub> and Ca(ClO<sub>4</sub>)<sub>2</sub> (Gault & Cockell, 2021). Another recent study considered instead the effects of perchlorate salts on the β-galactosidase enzyme from *Halorubrum lacusprofundi*, an extremely cold-adapted halophilic archaeon (Laye & DasSarma, 2018), which is phylogenetically close to *H. sodomense* (Becker et al., 2014; Kamekura & Dyll-Smith, 1995). It has been found that the ClO<sub>4</sub><sup>-</sup> anion acted as a very weak non-competitive inhibitor at concentrations above 0.88 M NaClO<sub>4</sub>. Unlike the study on bovine α-chymotrypsin, the effects on stability were not investigated. In summary, over the five considered strains, *H. sodomense* and *H. volcanii* proteomes stand out with an outstanding tolerance to perchlorate salts such as those encountered on Mars, while *H. salinarum* proteome displays a high sensibility to these salts.

This work stresses the fact that simpler systems made with purified proteins may not represent the behaviour of the whole proteome. It has been previously shown that model MDH enzyme from *H. marismortui* and other *Haloarchaea* can be stabilized by low concentrations of MgCl<sub>2</sub> and CaCl<sub>2</sub>, which only become destabilizing above 0.5–1 M (Blanquart et al., 2021; Madern & Zaccai, 1997). However, each enzyme from a single organism may display a specific salt response. For example, various enzymes (Bräsen & Schönheit, 2005; Camacho et al., 2009; Cendrin et al., 1994; Goldman et al., 1990) and large molecular assemblies (Franzetti et al., 2002; Shevack et al., 1985) from *H. marismortui* are stable below 2 M KCl. Since these proteins may be significantly abundant in the proteomes, highly salt-sensitive enzymes like MDH may not represent properly the proteome as a whole. This is especially critical in the context of the search for biomarkers. Indeed, putative extraterrestrial

microorganisms would probably not leave behind purified systems but rather complex mixtures of their molecular components, based on their individual stability and eventual interactions. Hence, study of the biophysical effects of extraterrestrial-like simulated conditions on proteomes provides additional insights into how such conditions would shape habitability and the chances of finding biomarkers.

As pointed out by this study, halophilism, defined as NaCl-dependency, does not necessarily make the most relevant criterion for choosing a halophilic model for all astrobiological studies. Instead, organisms whose biomolecules present higher tolerance to MgCl<sub>2</sub>, such as *H. sodomense* and *H. volcanii*, should be considered as more relevant in the context of chaotropic extraterrestrial brines.

The fact that NaClO<sub>3</sub> stabilized proteomes of all tested *Haloarchaea* and that some displayed low sensitivity to NaClO<sub>4</sub> and Mg(ClO<sub>4</sub>)<sub>2</sub> and even some stabilization by MgCl<sub>2</sub> and Ca(ClO<sub>4</sub>)<sub>2</sub> in low concentrations of KCl and NaCl may also increase the chances of finding salt-adapted biomarkers in Martian brines where these two monovalent salts are scarcer. In particular, low amounts of NaCl/KCl, which are not expected to be abundant in Martian brines, are not a limit per se for halophilic proteome stability which can be increased by various other salts.

Further studies could also consider other factors which may compensate effects of Martian-relevant salts on halophilic proteomes. For example, high pressure, which reduces perchlorate enzymatic inhibition on bovine α-chymotrypsin (Gault et al., 2020) and deleterious effects of MgCl<sub>2</sub>, MgSO<sub>4</sub> and Mg(ClO<sub>4</sub>)<sub>2</sub> on ligand binding by tRNA (Jahmidi-Azizi et al., 2021). Moreover, low temperatures of the surface and upper subsurface of Mars would also counteract chaotropic effects of Martian salts (Chin et al., 2010; Cray et al., 2015). Other factors compensating the effects of Martian-relevant salts include trimethylamine N-oxide (TMAO), a common intracellular osmolyte, which mitigates chaotropic effect of Mg(ClO<sub>4</sub>)<sub>2</sub> on water structure (Laurent et al., 2020). Similarly, other kosmotropic salts, solutes and osmolytes counteract chaotropicity of chaotropic salts (de Lima Alves et al., 2015; Hallsworth et al., 2007; Stevenson et al., 2015; Yakimov et al., 2015).

Another important implication may arise from this work. While effects of pure water solutions on halophilic systems were out of the scope of this study, the denaturation curves obtained with only one type of salt show that all tested salts increased average thermostability in comparison to pure water conditions. This was seen with proteomes of all tested strains, except *H. salinarum* which should not always be considered as the most-used halophilic model for astrobiology. Indeed, sample returns are challenging goals of future Mars missions. Therefore, we stress the fact that extreme caution should be taken when analysing

samples from brines or salt-containing rocks. In particular, special attention should be paid to the composition of the hypersaline buffers that should be used to resuspend the Mars samples.

## AUTHOR CONTRIBUTIONS

**Lorenzo Carré:** Conceptualization (equal); data curation (equal); investigation (equal); methodology (lead); writing – original draft (equal). **Daniel Gonzalez:** Investigation (equal). **Éric Girard:** Conceptualization (equal); methodology (equal); supervision (equal). **Bruno Franzetti:** Conceptualization (lead); data curation (equal); funding acquisition (lead); investigation (equal); methodology (equal); project administration (lead); supervision (lead); validation (lead); writing – original draft (equal); writing – review and editing (equal).

## ACKNOWLEDGEMENTS

This work is supported by the French National Research Agency in the framework of the Investissements d'Avenir programme (ANR-15-IDEX-02), through the funding of the 'Origin of Life' project of the Université Grenoble-Alpes. IBS acknowledges integration into the Interdisciplinary Research Institute of Grenoble (IRIG, CEA).

## CONFLICT OF INTEREST STATEMENT

The authors declare that they have no competing interests.

## DATA AVAILABILITY STATEMENT

Datasets related to this article are available upon request.

## ORCID

Bruno Franzetti  <https://orcid.org/0000-0001-5323-0510>

## REFERENCES

- Abotalib, A.Z. & Heggy, E. (2019) A deep groundwater origin for recurring slope lineae on Mars. *Nature Geoscience*, 12, 235–241.
- Ando, N., Barquera, B., Bartlett, D.H., Boyd, E., Burnim, A.A., Byer, A.S. et al. (2021) The molecular basis for life in extreme environments. *Annual Review of Biophysics*, 50, 343–372.
- Ball, P. & Hallsworth, J.E. (2015) Water structure and chaotropicity: their uses, abuses and biological implications. *Physical Chemistry Chemical Physics*, 17, 8297–8305.
- Bayles, M., Belasco, B.C., Breda, A.J., Cahill, C.B., Silva, A.Z.D., Regan, M.J. et al. (2020) The Haloarchaea of Great Salt Lake as models for potential extant life on Mars. In: *Extremophiles as astrobiological models*. Chichester: John Wiley & Sons, Ltd, pp. 83–124. Available from: <https://doi.org/10.1002/9781119593096.ch4>
- Becker, E.A., Seitzer, P.M., Tritt, A., Larsen, D., Krusor, M., Yao, A.I. et al. (2014) Phylogenetically driven sequencing of extremely halophilic archaea reveals strategies for static and dynamic osmo-response. *PLoS Genetics*, 10, e1004784.
- Blanquart, S., Groussin, M., Le Roy, A., Szöllosi, G.J., Girard, E., Franzetti, B. et al. (2021) Resurrection of ancestral malate

- dehydrogenases reveals the evolutionary history of halobacterial proteins: deciphering gene trajectories and changes in biochemical properties. *Molecular Biology and Evolution*, 38, 3754–3774. Available from: <https://doi.org/10.1093/molbev/msab146>
- Brady, J.B. (1992) Does ice dissolve or does halite melt? *Journal of Geological Education*, 40, 116–118.
- Bräsen, C. & Schönheit, P. (2005) AMP-forming acetyl-CoA synthetase from the extremely halophilic archaeon *Haloarcula marismortui*: purification, identification and expression of the encoding gene, and phylogenetic affiliation. *Extremophiles*, 9, 355–365.
- Bye, J.W. & Falconer, R.J. (2013) Thermal stability of lysozyme as a function of ion concentration: a reappraisal of the relationship between the Hofmeister series and protein stability. *Protein Science*, 22, 1563–1570.
- Camacho, R.M., Mateos, J.C., González-Reynoso, O., Prado, L.A. & Córdova, J. (2009) Production and characterization of esterase and lipase from *Haloarcula marismortui*. *Journal of Industrial Microbiology and Biotechnology*, 36, 901–909.
- Carré, L., Girard, E. & Franzetti, B. (2021) Experimental study of proteome halophilicity using nanoDSF: a proof of concept. *Extremophiles*, 26(1), 1.
- Carré, L., Zaccai, G., Delfosse, X., Girard, E. & Franzetti, B. (2022) Relevance of Earth-bound extremophiles in the search for extraterrestrial life. *Astrobiology*, 22, 322–367. Available from: <https://doi.org/10.1089/ast.2021.0033>
- Cendrin, F., Jouve, H.M., Gaillard, J., Thibault, P. & Zaccai, G. (1994) Purification and properties of a halophilic catalase-peroxidase from *Haloarcula marismortui*. *Biochimica et Biophysica Acta (BBA)—Protein Structure and Molecular Enzymology*, 1209, 1–9.
- Chin, J.P., Megaw, J., Magill, C.L., Nowotarski, K., Williams, J.P., Bhaganna, P. et al. (2010) Solutes determine the temperature windows for microbial survival and growth. *Proceedings of the National Academy of Sciences of the United States of America*, 107, 7835–7840.
- Clifford, S.M., Lasue, J., Heggy, E., Boisson, J., McGovern, P. & Max, M.D. (2010) Depth of the Martian cryosphere: revised estimates and implications for the existence and detection of subpermafrost groundwater. *Journal of Geophysical Research*, 115, 1–17. Available from: <https://doi.org/10.1029/2009JE003462>
- Coquelle, N., Talon, R., Juers, D.H., Girard, É., Kahn, R. & Madern, D. (2010) Gradual adaptive changes of a protein facing high salt concentrations. *Journal of Molecular Biology*, 404, 493–505.
- Cray, J.A., Russell, J.T., Timson, D.J., Singhal, R.S. & Hallsworth, J.E. (2013) A universal measure of chaotropy and kosmotropy. *Environmental Microbiology*, 15, 287–296.
- Cray, J.A., Stevenson, A., Ball, P., Bankar, S.B., Eleutherio, E.C., Ezeji, T.C. et al. (2015) Chaotropy: a key factor in product tolerance of biofuel-producing microorganisms. *Current Opinion in Biotechnology*, 33, 228–259.
- Crits-Christoph, A., Gelsinger, D.R., Ma, B., Wierchos, J., Ravel, J., Davila, A. et al. (2016) Functional interactions of archaea, bacteria and viruses in a hypersaline endolithic community. *Environmental Microbiology*, 18, 2064–2077.
- de Lima Alves, F., Stevenson, A., Baxter, E., Gillion, J.L.M., Hejazi, F., Hayes, S. et al. (2015) Concomitant osmotic and chaotropy-induced stresses in *Aspergillus wentii*: compatible solutes determine the biotic window. *Current Genetics*, 61, 457–477.
- Fairén, A.G., Gómez-Elvira, J., Briones, C., Prieto-Ballesteros, O., Rodríguez-Manfredi, J.A., López Herederó, R. et al. (2020) The Complex Molecules Detector (CMOLD): a fluidic-based instrument suite to search for (bio)chemical complexity on Mars and icy moons. *Astrobiology*, 20, 1076–1096.
- Favreau, C. (2023). Réponse physiologique de l'haloarchée *Halobacterium salinarum* piégée au sein des cristaux d'halite: Développements méthodologiques et étude moléculaire ex situ. Sorbonne Université.
- Favreau, C., Tribondeau, A., Marugan, M., Guyot, F., Alpha-Bazin, B., Marie, A. et al. (2022) Molecular acclimation of *Halobacterium salinarum* to halite brine inclusions. *bioRxiv*. Available from: <https://doi.org/10.1101/2022.09.16.508090>
- Fendrihan, S., Bérces, A., Lammer, H., Musso, M., Rontó, G., Polacsek, T.K. et al. (2009) Investigating the effects of simulated Martian ultraviolet radiation on *Halococcus dombrowskii* and other extremely halophilic archaeobacteria. *Astrobiology*, 9, 104–112.
- Fischer, E., Martínez, G.M., Elliott, H.M. & Rennó, N.O. (2014) Experimental evidence for the formation of liquid saline water on Mars. *Geophysical Research Letters*, 41, 4456–4462.
- Fox-Powell, M.G., Hallsworth, J.E., Cousins, C.R. & Cockell, C.S. (2016) Ionic strength is a barrier to the habitability of Mars. *Astrobiology*, 16, 427–442.
- Franzetti, B., Schoehn, G., Garcia, D., Ruigrok, R.W.H. & Zaccai, G. (2002) Characterization of the proteasome from the extremely halophilic archaeon *Haloarcula marismortui*. *Archaea*, 1, 53–61.
- Gault, S. & Cockell, C.S. (2021) Perchlorate salts exert a dominant, deleterious effect on the structure, stability, and activity of  $\alpha$ -chymotrypsin. *Astrobiology*, 21, 405–412.
- Gault, S., Jaworek, M.W., Winter, R. & Cockell, C.S. (2020) High pressures increase  $\alpha$ -chymotrypsin enzyme activity under perchlorate stress. *Communications Biology*, 3, 1–9.
- General, I.J., Ascituro, E.K. & Madura, J.D. (2008) Structure of aqueous sodium perchlorate solutions. *The Journal of Physical Chemistry B*, 112, 15417–15425.
- Ginzburg, M., Sachs, L. & Ginzburg, B.Z. (1970) Ion metabolism in a Halobacterium. I. Influence of age of culture on intracellular concentrations. *The Journal of General Physiology*, 55, 187–207.
- Goldman, S., Hecht, K., Eisenberg, H. & Mevarech, M. (1990) Extracellular  $\text{Ca}^{2+}$ -dependent inducible alkaline phosphatase from extremely halophilic archaeobacterium *Haloarcula marismortui*. *Journal of Bacteriology*, 172, 7065–7070.
- Hallsworth, J.E. (2020) Salt deliquescence can support extraterrestrial life. *Nature Astronomy*, 4, 739–740.
- Hallsworth, J.E. (2021) Mars' surface is not universally biocidal. *Environmental Microbiology*, 23, 3345–3350. Available from: <https://doi.org/10.1111/1462-2920.15494>
- Hallsworth, J.E. (2022) Water is a preservative of microbes. *Microbial Biotechnology*, 15, 191–214.
- Hallsworth, J.E., Heim, S. & Timmis, K.N. (2003) Chaotropic solutes cause water stress in *Pseudomonas putida*. *Environmental Microbiology*, 5, 1270–1280.
- Hallsworth, J.E., Prior, B.A., Nomura, Y., Iwahara, M. & Timmis, K.N. (2003) Compatible solutes protect against chaotrope (ethanol)-induced, nonosmotic water stress. *Applied and Environmental Microbiology*, 69, 7032–7034.
- Hallsworth, J.E., Yakimov, M.M., Golyshin, P.N., Gillion, J.L.M., D'Auria, G., Alves, F.D.L. et al. (2007) Limits of life in  $\text{MgCl}_2$ -containing environments: chaotropy defines the window. *Environmental Microbiology*, 9, 801–813.
- Hamaguchi, K. & Geiduschek, E.P. (1962) The effect of electrolytes on the stability of the deoxyribonucleate helix. *Journal of the American Chemical Society*, 84, 1329–1338.
- Hecht, M.H., Kounaves, S.P., Quinn, R.C., West, S.J., Young, S.M.M., Ming, D.W. et al. (2009) Detection of perchlorate and the soluble chemistry of Martian soil at the Phoenix Lander site. *Science*, 325, 64–67.
- Hofmeister, F. (1888) Zur Lehre von der Wirkung der Salze. *Archiv für Experimentelle Pathologie und Pharmakologie*, 24, 247–260.
- Jahmidi-Azizi, N., Gault, S., Cockell, C.S., Oliva, R. & Winter, R. (2021) Ions in the deep subsurface of Earth, Mars, and icy moons: their effects in combination with temperature and pressure on tRNA–ligand binding. *International Journal of Molecular Sciences*, 22, 10861.
- Jensen, M.W., Matlock, S.A., Reinheimer, C.H., Lawlor, C.J., Reinheimer, T.A. & Gorrell, A. (2015) Potassium stress growth

- characteristics and energetics in the haloarchaeon *Haloarcula marismortui*. *Extremophiles*, 19, 315–325.
- Kamekura, M. & Dyall-Smith, M.L. (1995) Taxonomy of the family *Halobacteriaceae* and the description of two new genera *Halorubrobacterium* and *Natrialba*. *The Journal of General and Applied Microbiology*, 41, 333–350.
- Lanyi, J.K. & Silverman, M.P. (1972) The state of binding of intracellular K<sup>+</sup> in *Halobacterium cutirubrum*. *Canadian Journal of Microbiology*, 18, 993–995. Available from: <https://doi.org/10.1139/m72-154>
- Laurent, H., Soper, A.K. & Dougan, L. (2020) Trimethylamine N-oxide (TMAO) resists the compression of water structure by magnesium perchlorate: terrestrial kosmotrope vs. Martian chaotrope. *Physical Chemistry Chemical Physics*, 22, 4924–4937.
- Lauro, S.E., Pettinelli, E., Caprarelli, G., Guallini, L., Rossi, A.P., Mattei, E. et al. (2021) Multiple subglacial water bodies below the south pole of Mars unveiled by new MARSIS data. *Nature Astronomy*, 5, 63–70.
- Laye, V.J. & DasSarma, S. (2018) An Antarctic extreme halophile and its polyextremophilic enzyme: effects of perchlorate salts. *Astrobiology*, 18, 412–418.
- Lenton, S., Rhys, N.H., Towey, J.J., Soper, A.K. & Dougan, L. (2017) Highly compressed water structure observed in a perchlorate aqueous solution. *Nature Communications*, 8, 919.
- Madern, D. & Zaccai, G. (1997) Stabilisation of halophilic malate dehydrogenase from *Haloarcula marismortui* by divalent cations. Effects of temperature, water isotope, cofactor and pH. *European Journal of Biochemistry*, 249, 607–611.
- Madern, D. & Zaccai, G. (2004) Molecular adaptation: the malate dehydrogenase from the extreme halophilic bacterium *Salinibacter ruber* behaves like a non-halophilic protein. *Biochimie*, 86, 295–303.
- Mancinelli, R.L., Fahlen, T.F., Landheim, R. & Klovstad, M.R. (2004) Brines and evaporites: analogs for Martian life. *Advances in Space Research*, 33, 1244–1246.
- Martinez, N., Michoud, G., Cario, A., Ollivier, J., Franzetti, B., Jebbar, M. et al. (2016) High protein flexibility and reduced hydration water dynamics are key pressure adaptive strategies in prokaryotes. *Scientific Reports*, 6, 1–11.
- Marty, V., Jasnin, M., Fabiani, E., Vauclare, P., Gabel, F., Trapp, M. et al. (2013) Neutron scattering: a tool to detect in vivo thermal stress effects at the molecular dynamics level in micro-organisms. *Journal of the Royal Society Interface*, 10, 20130003.
- Matheson, A.T., Sprott, G.D., McDonald, I.J. & Tessier, H. (1976) Some properties of an unidentified halophile: growth characteristics, internal salt concentration, and morphology. *Canadian Journal of Microbiology*, 22, 780–786.
- McEwen, A.S., Ojha, L., Dundas, C.M., Mattson, S.S., Byrne, S., Wray, J.J. et al. (2011) Seasonal flows on warm Martian slopes. *Science*, 333, 740–743.
- McGenity, T.J. & Grant, W.D. (2015) *Halorubrum*. In: *Bergey's manual of systematics of archaea and bacteria*. Chichester: John Wiley & Sons, Inc., pp. 1–11. Available from: <https://doi.org/10.1002/9781118960608.gbm00487>
- Mevarech, M., Leicht, W. & Werber, M.M. (1976) Hydrophobic chromatography and fractionation of enzymes from extremely halophilic bacteria using decreasing concentration gradients of ammonium sulfate. *Biochemistry*, 15, 2383–2387.
- Micheluz, A., Pinzari, F., Rivera-Valentín, E.G., Manente, S. & Hallsworth, J.E. (2022) Biophysical manipulation of the extracellular environment by *Eurotium halophilicum*. *Pathogens*, 11, 1462.
- Mullakhanbhai, M.F. & Larsen, H. (1975) *Halobacterium volcanii* spec. nov., a Dead Sea Halobacterium with a moderate salt requirement. *Archives of Microbiology*, 104, 207–214.
- Oosterholt, D. & Stoeckenius, W. (1974) Isolation of the cell membrane of *Halobacterium halobium* and its fractionation into red and purple membrane. In: *Methods in enzymology*, Vol. 31. Amsterdam: Elsevier, pp. 667–678. Available from: <https://linkinghub.elsevier.com/retrieve/pii/0076687974310725>
- Okur, H.I., Hladílková, J., Rembert, K.B., Cho, Y., Heyda, J., Dzubiel, J. et al. (2017) Beyond the Hofmeister series: ion-specific effects on proteins and their biological functions. *The Journal of Physical Chemistry B*, 121, 1997–2014.
- Oren, A. (1983) *Halobacterium sodomense* sp. nov., a Dead Sea Halobacterium with an extremely high magnesium requirement. *International Journal of Systematic Bacteriology*, 33, 381–386.
- Oren, A. (1999) Bioenergetic aspects of halophilism. *Microbiology and Molecular Biology Reviews*, 63, 334–348.
- Oren, A. (2013) Life in magnesium- and calcium-rich hypersaline environments: salt stress by chaotropic ions. In: Seckbach, J., Oren, A. & Stan-Lotter, H. (Eds.) *Polyextremophiles. Volume 27 of cellular origin, life in extreme habitats and astrobiology*. Dordrecht: Springer Netherlands, pp. 215–232. Available from: [https://doi.org/10.1007/978-94-007-6488-0\\_8](https://doi.org/10.1007/978-94-007-6488-0_8)
- Oren, A. (2017) Why is n't *Haloferax mediterranei* more 'weed-like'? *FEMS Microbiology Letters*, 364. Available from: <https://doi.org/10.1093/femsle/fnx070>
- Oren, A. & Hallsworth, J.E. (2014) Microbial weeds in hypersaline habitats: the enigma of the weed-like *Haloferax mediterranei*. *FEMS Microbiology Letters*, 359, 134–142.
- Oren, A. & Ventosa, A. (2017) Halobacterium. In: Whitman, W.B., Rainey, F., Kämpfer, P., Trujillo, M., Chun, J., DeVos, P. et al. (Eds.) *Bergey's manual of systematics of archaea and bacteria*. Chichester: John Wiley & Sons, Ltd, pp. 1–12. Available from: <https://doi.org/10.1002/9781118960608.gbm00482.pub2>
- Orosei, R., Lauro, S.E., Pettinelli, E., Cicchetti, A., Coradini, M., Cosciotti, B. et al. (2018) Radar evidence of subglacial liquid water on Mars. *Science*, 361, 490–493.
- Osterloo, M.M., Hamilton, V.E., Bandfield, J.L., Glotch, T.D., Baldrige, A.M., Christensen, P.R. et al. (2008) Chloride-bearing materials in the southern highlands of Mars. *Science*, 319, 1651–1654.
- Pérez-Fillol, M. & Rodríguez-Valera, F. (1986) Potassium ion accumulation in cells of different halobacteria. *Microbiologia*, 2, 73–80.
- Perl, S.M. & Baxter, B.K. (2020) Great Salt Lake as an astrobiology analogue for ancient Martian hypersaline aqueous systems. In: Baxter, B.K. & Butler, J.K. (Eds.) *Great Salt Lake biology: a terminal lake in a time of change*. Cham: Springer International Publishing, pp. 487–514. Available from: [https://doi.org/10.1007/978-3-030-40352-2\\_16](https://doi.org/10.1007/978-3-030-40352-2_16)
- Peters, J., Martinez, N., Michoud, G., Cario, A., Franzetti, B., Oger, P. et al. (2014) Deep sea microbes probed by incoherent neutron scattering under high hydrostatic pressure. *Zeitschrift für Physikalische Chemie*, 228, 1121–1133. Available from: <https://doi.org/10.1515/zpch-2014-0547>
- Pohlschroder, M. & Schulze, S. (2019) *Haloferax volcanii*. *Trends in Microbiology*, 27, 86–87.
- Reed, C.J., Lewis, H., Trejo, E., Winston, V. & Evilia, C. (2013) Protein adaptations in archaeal extremophiles. *Archaea*, 2013, 1–14.
- Reid, I.N., Sparks, W.B., Lubow, S., McGrath, M., Livio, M., Valenti, J. et al. (2006) Terrestrial models for extraterrestrial life: methanogens and halophiles at Martian temperatures. *International Journal of Astrobiology*, 5, 89–97.
- Rivera-Valentín, E.G., Chevrier, V.F., Soto, A. & Martínez, G. (2020) Distribution and habitability of (meta)stable brines on present-day Mars. *Nature Astronomy*, 4, 756–761.
- Rodríguez-Valera, F., Ruiz-Berraquero, F. & Ramos-Cormenzana, A. (1980) Isolation of extremely halophilic bacteria able to grow in defined inorganic media with single carbon sources. *Microbiology*, 119, 535–538.
- Sánchez-García, L., Fernández-Martínez, M.A., Moreno-Paz, M., Carrizo, D., García-Villadangos, M., Machado, J.M. et al. (2020) Simulating Mars drilling mission for searching for life: ground-truthing lipids and other complex microbial biomarkers in the iron-sulfur rich Río Tinto analog. *Astrobiology*, 20, 1029–1047.

- Schreder-Gomes, S.I., Benison, K.C. & Bernau, J.A. (2022) 830-million-year-old microorganisms in primary fluid inclusions in halite. *Geology*, 50, 918–922.
- Shevack, A., Gewitz, H.S., Hennemann, B., Yonath, A. & Wittmann, H.G. (1985) Characterization and crystallization of ribosomal particles from *Halobacterium marismortui*. *FEBS Letters*, 184, 68–71.
- Squyres, S.W., Grotzinger, J.P., Arvidson, R.E., Bell, J.F., Calvin, W., Christensen, P.R. et al. (2004) In situ evidence for an ancient aqueous environment at Meridiani Planum, Mars. *Science*, 306, 1709–1714.
- Stevenson, A., Cray, J.A., Williams, J.P., Santos, R., Sahay, R., Neuenkirchen, N. et al. (2015) Is there a common water-activity limit for the three domains of life? *The ISME Journal*, 9, 1333–1351.
- Sundarasami, A., Sridhar, A. & Mani, K. (2019) Chapter 13—halophilic archaea as beacon for exobiology: recent advances and future challenges. In: Meena, S.N. & Naik, M.M. (Eds.) *Advances in biological science research*. Cambridge: Academic Press, pp. 197–214. Available from: <http://www.sciencedirect.com/science/article/pii/B9780128174975000136>
- Tadeo, X., Pons, M. & Millet, O. (2007) Influence of the Hofmeister anions on protein stability as studied by thermal denaturation and chemical shift perturbation. *Biochemistry*, 46, 917–923.
- Tehei, M., Franzetti, B., Madern, D., Ginzburg, M., Ginzburg, B.Z., Giudici-Ortoni, M.-T. et al. (2004) Adaptation to extreme environments: macromolecular dynamics in bacteria compared in vivo by neutron scattering. *EMBO Reports*, 5, 66–70.
- Tehei, M., Franzetti, B., Maurel, M.-C., Vergne, J., Hountondji, C. & Zaccai, G. (2002) The search for traces of life: the protective effect of salt on biological macromolecules. *Extremophiles*, 6, 427–430.
- Vauclare, P., Marty, V., Fabiani, E., Martinez, N., Jasnin, M., Gabel, F. et al. (2015) Molecular adaptation and salt stress response of *Halobacterium salinarum* cells revealed by neutron spectroscopy. *Extremophiles*, 19, 1099–1107.
- Vauclare, P., Natali, F., Kleman, J.P., Zaccai, G. & Franzetti, B. (2020) Surviving salt fluctuations: stress and recovery in *Halobacterium salinarum*, an extreme halophilic archaeon. *Scientific Reports*, 10, 1–10.
- Ventosa, A., de la Haba, R.R. & Sánchez-Porro, C. (2015) *Haloferax*. In: *Bergey's manual of systematics of archaea and bacteria*. Chichester: John Wiley & Sons, Inc., p. 1. Available from: <https://doi.org/10.1002/9781118960608>
- Von Hippel, P.H. & Wong, K.Y. (1965) On the conformational stability of globular proteins. The effects of various electrolytes and non-electrolytes on the thermal ribonuclease transition. *The Journal of Biological Chemistry*, 240, 3909–3923.
- Yakimov, M.M., La Cono, V., Spada, G.L., Bortoluzzi, G., Messina, E., Smedile, F. et al. (2015) Microbial community of the deep-sea brine Lake Kryos seawater–brine interface is active below the chaotropicity limit of life as revealed by recovery of mRNA. *Environmental Microbiology*, 17, 364–382.
- Zaccai, G. (2013) Ecology of protein dynamics. *Current Physical Chemistry*, 3, 9–16.

## SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**How to cite this article:** Carré, L., Gonzalez, D., Girard, É. & Franzetti, B. (2023) Effects of chaotropic salts on global proteome stability in halophilic archaea: Implications for life signatures on Mars. *Environmental Microbiology*, 25(11), 2216–2230. Available from: <https://doi.org/10.1111/1462-2920.16451>