



Mixing speed and stability of SRB measures through optimal transportation

Houssam Boukhecham, Benoît Kloeckner

► To cite this version:

Houssam Boukhecham, Benoît Kloeckner. Mixing speed and stability of SRB measures through optimal transportation. 2023. hal-04201539

HAL Id: hal-04201539

<https://hal.science/hal-04201539>

Preprint submitted on 10 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Mixing speed and stability of SRB measures through optimal transportation

Houssam Boukhecham ^{*} Benoît R. Kloeckner ^{*}

September 10, 2023

It is well-known that the SRB measure of a $C^{1+\alpha}$ Anosov diffeomorphism has exponential decay of correlations with respect to Hölder-continuous observables. We propose a new approach to this phenomenon, based on optimal transport. More precisely, we define a space of measures having absolutely continuous disintegrations with respect to some foliation close to the unstable foliation of the map, endowed with a variant of the Wasserstein metric where mass is only allowed to be transported along the diffeomorphism's stable foliation. We show that this metric is indeed finite on that space, and use that the construction makes the diffeomorphism act as a contraction to deduce two corollaries. First, the SRB measure has exponential decay of correlation with respect to pairs of observable that are only asked to be Hölder-continuous *in the stable, respectively unstable direction*, but can be discontinuous overall. Then, we prove quantitative statistical stability: the map sending a $C^{1+\alpha}$ Anosov diffeomorphism to its SRB measure is locally Hölder-continuous (using the C^1 metric for diffeomorphisms and the usual Wasserstein metric for measures).

Disclaimer. *This article is an advanced draft: some proofs are somewhat sketchy and the results should therefore be taken with a pinch of salt. Both authors have projects outside academia for the near future, and time did not permit to provide a more complete write-up. We still hope the idea is fruitful and the proofs detailed enough that the method can be used by others in many cases beyond Anosov diffeomorphism.*

^{*}Univ Paris Est Creteil, Univ Gustave Eiffel, CNRS, LAMA UMR8050, F-94010 Creteil, France

1 Introduction

Let $T : M \rightarrow M$ be a $C^{1+\alpha}$, topologically mixing Anosov diffeomorphism. The chaotic properties of T make it practically impossible to predict the future of an orbit at any given time, and at the same time enable one to predict accurately how such an orbit distributes over a long enough period of time, for Lebesgue-almost all starting point. This distribution is indeed well-known to be described by the *SRB measure* of T , which is also a *physical measure*. The “accuracy” of the distribution of orbits is quantified by the decay of correlations, which is exponential for Hölder observables. An important contemporary line of research consists in extending this to various classes of systems enjoying some flavor of non-uniformly hyperbolicity, see e.g. [Liv95, BV96, Dol98, You98, BY00, AdCJ04, Var08, AS12, DSL16, KKM19]. The purpose of the present article is to propose a novel method to study the decay of correlations, yielding some new results in the case of Anosov diffeomorphisms and hopefully simple enough to be extended in other settings.

The method is by a coupling argument, akin to the one commonly used in finite-states Markov Chains where one creates two realizations of the chain that are coupled one to another: they run independently until they meet at the same state, then they evolve along the same trajectory. Our idea is to do the same for orbits of T , with “meeting at the same state” replaced by “lying not too far away on the same stable leaf”.

The statement of our main result needs a few definitions which we introduce first.

1.1 Wasserstein metric and variants

To state our central result, let us first introduce the *Wasserstein metric* D_1 associated to the distance function $d(\cdot, \cdot)$ defined on M by any Riemannian metric : to any two probability measures μ_1, μ_2 , it associates the distance

$$D^1(\mu_1, \mu_2) := \inf_{\gamma \in \Gamma(\mu_1, \mu_2)} \int_{M \times M} d(x, y) \, d\gamma(x, y)$$

where $\Gamma(\mu_1, \mu_2)$ is the set of *coupling* (aka *transport plans*) between μ_1 and μ_2 , i.e. the set of probability measures γ on $M \times M$ having μ_1, μ_2 as marginals. This last condition can be written $p_{i*}\gamma = \mu_i$ for both $i = 1, 2$, where $p_i : M \times M \rightarrow M$ is the projection to the i th factor. “Kantorovich’s duality” enables one to rewrite this as

$$D^1(\mu_1, \mu_2) = \sup_{f \in \text{Lip}(d, 1)} \left| \int f \, d\mu_1 - \int f \, d\mu_2 \right|$$

where $\text{Lip}(d, k)$ denotes the space of functions $M \rightarrow \mathbb{R}$ that are Lipschitz of constant at most k with respect to the metric d . This metric induces the weak-* topology on the set of probability measure (this claim uses the compactness of M).

Now, the same concept can be used for other metrics than d . We will combine two variations: first, using $d(\cdot, \cdot)^\beta$ for some $\beta \in (0, 1)$, which is also a metric. The corresponding Wasserstein metric is

$$D^\beta(\mu_1, \mu_2) := \inf_{\gamma \in \Gamma(\mu_1, \mu_2)} \int_{M \times M} d(x, y)^\beta \, d\gamma(x, y),$$

it still metrizes the weak-* topology, and the duality will then be against β -Hölder functions with Hölder constant at most 1 (it is quite different from $D^1(\cdot, \cdot)^\beta$, since small movements of mass are disproportionately penalized compared to large ones). Second, we will use the extended metric $d_s : M \times M \rightarrow [0, +\infty]$ which gives the distance along the stable leaves: $d_s(x, y) = \infty$ when x, y are not in the same stable leaf, otherwise $d_s(x, y)$ is the least length of a smooth curve from x to y , constrained to stay on the same stable leaf. Then we write

$$D_s^1(\mu_1, \mu_2) := \inf_{\gamma \in \Gamma(\mu_1, \mu_2)} \int_{M \times M} d_s(x, y) \, d\gamma(x, y)$$

which can take the value $+\infty$ but is otherwise well-defined, and has a dual formulation against bounded Borel-measurable functions that are Lipschitz *in the stable direction*, i.e. with respect to d_s . Last, we combine both variations by introducing for all $\beta \in (0, 1]$

$$D_s^\beta(\mu_1, \mu_2) := \inf_{\gamma \in \Gamma(\mu_1, \mu_2)} \int_{M \times M} d_s(x, y)^\beta \, d\gamma(x, y).$$

Again, we have a duality and we introduce the corresponding space $\text{Hol}_s^\beta(M, T)$ of bounded, Borel-measurable functions $f : M \rightarrow \mathbb{R}$ such that for some $C \geq 0$ and all $x, y \in M$,

$$|f(x) - f(y)| \leq C \cdot d_s(x, y)^\beta.$$

The least such C is then denoted by $\text{Hol}_s^\beta(f)$, and we use on $\text{Hol}_s^\beta(M, T)$ the norm $\|\cdot\|_{s, \beta} = \|\cdot\|_\infty + \text{Hol}_s^\beta(\cdot)$, making it a Banach algebra. Then

$$D_s^\beta(\mu_1, \mu_2) = \sup_f \left| \int f \, d\mu_1 - \int f \, d\mu_2 \right|$$

where the supremum is over all $f \in \text{Hol}_s^\beta(M, T)$ such that $\text{Hol}_s^\beta(f) \leq 1$.

We define similarly Hol_W^β , where the stable foliation is replaced by any foliation W , and $\text{Hol}_u^\beta := \text{Hol}_{W^u}^\beta$.

1.2 Regularly Foliated Measures

The notion of SRB measure combines two aspects: the absolute continuity of the local disintegrations along the unstable foliation, and the T -invariance (the usual definition also includes existence of a positive Lyapunov exponent, which follows here from the Anosov assumption). We will consider non-invariant measures that satisfy the first condition for a foliation close to the unstable foliation of T .

To define “close”, we fix an open cone field $(C_x \subset T_x M)_{x \in M}$ containing a neighborhood of the unstable distribution E^u of T ; we will denote by U_x the corresponding neighborhood of E_x^u in the Grassmanian $G(k_u, T_x M)$ where $k_u := \dim E^u$; i.e. whenever E is a k_u -dimensional subspace of $T_x M$, we write indifferently $E \in U_x$ or $E \subset C_x$, and we say that E is *tangent to the cone field*. A k_u -dimensional submanifold N of M is said to be *tangent to the cone field* when $T_x N \in U_x$ for all $x \in N$, and a foliation is said to be tangent to the cone field whenever its leaves are. The cone field will be chosen

small enough that the action of T on the sections of $(U_x)_x$ is contracting (with E^u as its unique fixed point).

We use two parameters $\beta \in (0, 1]$ and $K > 0$ to bound the regularity of both the foliation and the densities of the disintegration. Foliations to be considered will be tangent to the cone field, and have their leaves written as β -Hölder graphs with constant at most K in some fixed finite atlas of M . Measures to be considered will have absolutely continuous local disintegrations with respect to such a foliation, with β -Hölder densities, and the log-densities will be asked to have Hölder constant at most K . The set of such probability measures is denoted by \mathcal{R}_K^β (the precise definition is given as Section 3.3.1).

1.3 Results

We are now in a position to state our central result.

Theorem 1.1. *Let $T : M \rightarrow M$ be a $C^{1+\alpha}$ Anosov diffeomorphism. For some $\beta_0 \in (0, 1)$ and for each $\beta \in (0, \beta_0)$ there exist $K_0, C > 0$ and $n_0 \in \mathbb{N}$ with the following properties:*

- i. $\mathcal{R}_{K_0}^\beta$ contains a unique T -invariant measure, which is the SRB measure μ_0 of T ,
- ii. $D_s^\beta(\mu_1, \mu_2) \leq C$ for all $\mu_1, \mu_2 \in \mathcal{R}_{K_0}^\beta$,
- iii. $T_*^{n_0} \mathcal{R}_{2K}^\beta \subset \mathcal{R}_K^\beta$ for all $K \geq K_0$.

In particular, for all $\mu_1, \mu_2 \in \mathcal{R}_{K_0}^\beta$ we can pair μ_1 and μ_2 by a coupling γ that is supported on the set of (x, y) where x and y are in the same stable leaf. Since the stable direction is contracting exponentially, denoting by $\lambda \in (0, 1)$ any constant such that T is λ -contracting along its stable foliation W^s , we get

$$D_s^\beta(T_*^n \mu_1, T_*^n \mu_2) \leq C \lambda^{\beta n} \quad \forall n \in \mathbb{N}; \quad (1)$$

and since D^1 is bounded above by $\text{diam}(M)^{1-\beta} D_s^\beta$, we also have exponential convergence in the usual Wasserstein metric. Given any $\mu \in \mathcal{R}_K^\beta$, the sequence $(T_*^n \mu)$ will converge to an invariant measure in $\mathcal{R}_{K_0}^\beta$, which must then be the SRB measure of T .

The following two corollaries are easily deduced from Theorem 1.1.

Corollary 1.2 (Exponential decay of correlations for stable/unstable-Hölder pairs of observables). *Let μ_0 be the SRB measure of T . For all $\beta \leq \beta_0$, there exist $C_\beta \geq 1$ such that for all $n \in \mathbb{N}$, all $f \in \text{Hol}_s^\beta$ and all $g \in \text{Hol}_u^\beta$,*

$$\left| \int f \circ T^n \cdot g \, d\mu_0 - \int f \, d\mu_0 \int g \, d\mu_0 \right| \leq C_\beta \|f\|_{s,\beta} \|g\|_{u,\beta} \lambda^{\beta n}$$

While exponential decay of correlations is long known for Anosov diffeomorphisms, this version applies to observables with only partial continuity requirement: f, g can exhibit discontinuities in the unstable, respectively stable, direction; the duality appearing between Hol_u^β and Hol_s^β seems very natural. One could also point out that the rate of decay is directly determined by the contraction factor λ of T , but beware that β_0 , while constructive, is not made completely explicit in the proof and can in principle be very small.

Corollary 1.3 (Hölder statistical stability). *There exist $\varepsilon > 0$, C' and β' such that for any $C^{1,\alpha}$ Anosov map $T_1 : M \rightarrow M$ which is ε -close from T in the C^1 topology, the SRB measures μ_0, μ_1 of T and T_1 are close in the Wasserstein distances:*

$$D_1(\mu_0, \mu_1) \leq C' \|T - T_1\|_{C^0}^{\beta'}$$

Here the main emphasis is on the explicit, Hölder modulus of continuity of the map sending an Anosov diffeomorphism to its SRB measure. Also note that the distance between measure is given in the quite strong Wasserstein metric, and that while we need T_1 to be C^1 -close to T , the control we get uses only the uniform distance

$$\|T - T_1\|_{C^0} := \max_{x \in M} d(T(x), T_1(x)).$$

2 Warm-up: expanding maps

Our goal here is to illustrate one strand of our method in the simplest possible setting by providing a short and elementary proof of the exponential decay of correlation for expanding maps.

The setting is as follows: $T : \mathbb{T}^1 \rightarrow \mathbb{T}^1$ is a uniformly expanding circle map of class $C^{1+\alpha}$ for some $\alpha \in (0, 1]$. Identify $\mathbb{T}^1 = \mathbb{R}/\mathbb{Z}$ with $[0, 1)$ and denote the usual circle distance by d . Since T is expanding, $\lambda := 1/\inf|T'| \in (0, 1)$. Since T is $C^{1+\alpha}$, there exist $H > 0$ such that $|T'(x)| \leq |T'(y)|e^{Hd(x,y)^\alpha}$ for all $x, y \in \mathbb{T}^1$. Let m be the Lebesgue measure; searching for an absolutely continuous invariant probability measure (ACIP), one as usual defines the transfer operator \mathcal{L} by $T_*(\rho m) = \mathcal{L}(\rho)m$, so that a non-negative eigenfunction of \mathcal{L} for the eigenvalue 1 is the density of an ACIP. By the change of variable formula, the transfer operator can be expressed as

$$\mathcal{L}\rho(x) := \sum_{z \in T^{-1}(x)} \frac{\rho(z)}{|T'(z)|}$$

and given any $x, y \in \mathbb{T}^1$, we can order $T^{-1}(x) =: \{x_i : 1 \leq i \leq k\}$ and $T^{-1}(y) =: \{y_i : 1 \leq i \leq k\}$ (where $k = |\deg T|$) in such a way that $d(x_i, y_i) \leq \lambda d(x, y)$ for all i . Such an ordering will always be assumed whenever we consider a pair of points and their inverse images. It will be convenient to have \mathcal{L} act on α -Hölder function, i.e. we view it as a linear, bounded operator on the Banach algebra $\text{Hol}_\alpha(\mathbb{T}^1)$ where the norm is

$$\|f\|_\alpha := \|f\|_\infty + \text{Hol}_\alpha(f); \quad \text{Hol}_\alpha(f) := \sup \left\{ \frac{|f(x) - f(y)|}{d(x, y)^\alpha} : x \neq y \in \mathbb{T}^1 \right\}.$$

Theorem 2.1 (Folklore¹). *There is an α -Hölder positive density $\rho_0 : \mathbb{T}^1 \rightarrow (0, +\infty)$ such that $\rho_0 m$ is T -invariant, and there exist $C \geq 1$ and $\theta \in (0, 1)$ such that for all α -Hölder positive density ρ and all $n \in \mathbb{N}$,*

$$\|\mathcal{L}^n \rho - \rho_0\|_\alpha \leq C \|\rho - \rho_0\|_\alpha \theta^n.$$

¹By this we mean that there are so many variants of such a result that we prefer not attribute it to a particular article or set of authors.

After the proof, we will briefly recall why this statements contains the decay of correlation, and its relation to a very strong convergence of probability measure, in the total variation norm. This convergence will inspire the metric to be used on measures in the Anosov case.

Our proof starts in a usual way, showing that the “stretching” effect stemming from the expansion hypothesis regularizes densities above some determined level. Then we phrase a simple coupling argument in term of densities: regular enough densities stay far from zero, so that any two of them “share” a definite amount of mass.

Proof. For each $K > 0$, consider the following set of Hölder densities:

$$\mathcal{H}_K^\alpha := \left\{ \rho : \mathbb{T}^1 \rightarrow (0, +\infty) \mid \int \rho \, dm = 1, \frac{\rho(x)}{\rho(y)} \leq e^{Kd(x,y)^\alpha} \ (\forall x, y \in \mathbb{T}^1) \right\}.$$

For all $\rho \in \mathcal{H}_K^\alpha$ and all $x, y \in \mathbb{T}^1$ with (suitably ordered) inverse images $\{x_i\}_i$ and $\{y_i\}_i$:

$$\begin{aligned} \frac{\rho}{|T'|}(x) &\leq \frac{\rho}{|T'|}(y) \cdot e^{(K+H)d(x,y)} \\ \frac{\rho}{|T'|}(x_i) &\leq \frac{\rho}{|T'|}(y_i) \cdot e^{(K+H)d(x_i, y_i)^\alpha} \\ \mathcal{L}\rho(x) &\leq \mathcal{L}\rho(y) \cdot e^{(K+H)\lambda^\alpha d(x,y)^\alpha} \end{aligned}$$

so that for all $K > 0$, $\mathcal{L}(\mathcal{H}_K^\alpha) \subset \mathcal{H}_{h(K)}^\alpha$ where $h(K) = (K + H)\lambda^\alpha$. For all K and $n \in \mathbb{N}$:

$$\begin{aligned} h^n(K) &= K\lambda^{n\alpha} + \lambda^\alpha H + \lambda^{2\alpha} H + \dots + \lambda^{n\alpha} H \\ &\leq K\lambda^{n\alpha} + \frac{\lambda^\alpha}{1 - \lambda^\alpha} H. \end{aligned}$$

Set $K_0 = 2\frac{\lambda^\alpha}{1 - \lambda^\alpha} H$ and let n_0 be the smallest integer such that $2\lambda^{n_0\alpha} \leq \frac{\lambda^\alpha}{1 - \lambda^\alpha}$, to get:

$$\mathcal{L}^{n_0}(\mathcal{H}_{2K_0}^\alpha) \subset \mathcal{H}_{K_0}^\alpha \quad \text{and} \quad \mathcal{L}(\mathcal{H}_{K_0}^\alpha) \subset \mathcal{H}_{K_0}^\alpha \quad (2)$$

Every $\rho \in \mathcal{H}_K^\alpha$, being a density, reaches a value at least 1 at some point x_0 , and each $x \in \mathbb{T}^1$ is at distance less than 1 from x_0 so that $\rho(x) \geq e^{-Kd(x, x_0)^\alpha} \geq e^{-K}$. For all $\tau \in (0, e^{-K_0})$ and all $\rho \in \mathcal{H}_{K_0}^\alpha$, we can decompose

$$\rho = \tau \mathbf{1} + (1 - \tau) \tilde{\rho}$$

where $\mathbf{1}$ is the constant function with value 1 and $\tilde{\rho} = (\rho - \tau \mathbf{1}) / (1 - \tau)$ is positive of integral 1 with respect to m . From now on we fix a value of τ small enough to further ensure that $\tilde{\rho} \in \mathcal{H}_{2K_0}^\alpha$ whenever $\rho \in \mathcal{H}_{K_0}^\alpha$. Then for all $\rho_1, \rho_2 \in \mathcal{H}_{K_0}^\alpha$ and all $n \geq n_0$:

$$\begin{aligned} \mathcal{L}^{n_0}(\rho_1 - \rho_2) &= \mathcal{L}^{n_0}(\tau \mathbf{1} + (1 - \tau) \tilde{\rho}_1 - \tau \mathbf{1} - (1 - \tau) \tilde{\rho}_2) \\ &= (1 - \tau) \mathcal{L}^{n_0}(\tilde{\rho}_1 - \tilde{\rho}_2) \\ \mathcal{L}^n(\rho_1 - \rho_2) &= (1 - \tau) \mathcal{L}^{n - n_0}(\mathcal{L}^{n_0} \tilde{\rho}_1 - \mathcal{L}^{n_0} \tilde{\rho}_2) \end{aligned} \quad (3)$$

where $\tilde{\rho}_i \in \mathcal{H}_{2K_0}^\alpha$ and $\mathcal{L}^{n_0} \tilde{\rho}_i \in \mathcal{H}_{K_0}^\alpha$. For each $n \in \mathbb{N}$, define

$$\gamma_n = \sup_{\rho_1, \rho_2 \in \mathcal{H}_{K_0}^\alpha} \|\mathcal{L}^n(\rho_1 - \rho_2)\|_\alpha \in [0, +\infty);$$

the above computation shows $\gamma_n \leq (1 - \tau)\gamma_{n-n_0}$ so that γ_n goes to 0, exponentially fast. The sequence of sets of functions $\mathcal{L}^n(\mathcal{H}_{K_0}^\alpha)$ thus converges to a point $\{\rho_0\} \subset \mathcal{H}_{K_0}^\alpha$ in the uniform norm (using compactness provided by the Azelà-Ascoli Theorem). Then ρ_0 must be a fixed point of \mathcal{L} and $\rho_0 m$ is an invariant measure of T . Being a fixed point, ρ_0 must belong to $\mathcal{H}_{K_0/2}^\alpha$, and $(\mathcal{L}^n \rho)_n$ converges to ρ_0 exponentially fast, uniformly over all $\rho \in \mathcal{H}_{K_0}^\alpha$. The statement follows. \square

Remark 2.2 (Decay of correlations). Denote by $\mu_0 = \rho_0 m$ the ACIP and let $f, g : \mathbb{T}^1 \rightarrow \mathbb{R}$ be two “observables”, with $f \in L^1(m)$ and g α -Hölder. For a small enough and $b = 1 - a\|g\|_\alpha^{-1} \int g \, d\mu_0$, the function $\rho = (a\|g\|_\alpha^{-1}g + b)\rho_0$ is a α -Hölder density with a g -uniform bound on $\|\rho - \rho_0\|_\alpha$, so that:

$$\begin{aligned} \int f \circ T^n \cdot g \, d\mu_0 &= \frac{\|g\|_\alpha}{a} \int f \circ T^n \cdot \left(\frac{\rho}{\rho_0} - b\right) d\mu_0 \\ &= \frac{\|g\|_\alpha}{a} \left(\int f \, dT_*^n(\rho m) - b \int f \, dT_*^n \mu_0 \right) \\ &= \frac{\|g\|_\alpha}{a} \int f (\mathcal{L}^n \rho - \rho_0) dm + \int g \, d\mu_0 \int f \, d\mu_0 \\ &= \int g \, d\mu_0 \int f \, d\mu_0 + O(\|g\|_\alpha \|f\|_{L^1(m)} \theta^n) \end{aligned}$$

i.e. we have exponential decay of correlations.

Remark 2.3. As seen in Remark 2.2, Theorem 2.1 implies that the measures $T_*^n(\rho m)$ converge to $\mu_0 := \rho_0 m$ in duality with $L^1(m)$ test functions; the convergence thus also holds against Borel bounded test functions, i.e. in the *total variation* distance. This can be rephrased as the existence of positive measures ν_n of total mass increasing to 1 such that $\nu_n \leq T_*^n(\rho m)$ and $\nu_n \leq \mu_0$ for all n . Such a strong convergence cannot be expected in the Anosov case, where the measures we will have to work with might be singular with respect to the limit measure. Seeing an expanding map as an Anosov endomorphism with stable dimension 0 gives a good guess on the natural replacement for the total variation distance: we shall use an optimal transportation distance where mass is only allowed to move along the stable direction. This is the second strand of proof we shall weave with the first one.

3 Anosov diffeomorphisms

3.1 Coupling measures and deduction of Theorem 1.1

For each $\beta \in (0, \alpha_0]$, we shall build a nested family of sets of probability measures $(\mathcal{R}_K^\beta)_K > 0$, where K is a regularity bound (the lesser, the more regular), where elements

of \mathcal{R}_K^β have absolutely continuous local disintegration with respect to some foliation close to the unstable foliation of T , and K bounds the Hölder regularity both of the foliation and of the densities on the leaves. For all $L > 0$ we define

$$\Delta_s(L) := \{(x, y) \in M \times M \mid d_s(x, y) \leq L\}$$

the set of pairs of points at distance at most L along the stable foliation. The slightly technical part of the article (detailed in Section 3.3) leads to the following properties:

Proposition 3.1. *There exist L_0 and $\tau \in (0, 1)$ (depending only on T), $K_0 > 0$ and $n_0 \in \mathbb{N}$ (both further depending on β) such that:*

- i. *The SRB measure of T lies in \mathcal{R}_K^β for some $K > 0$,*
- ii. *$\mathcal{R}_K^\beta \subset \mathcal{R}_{K'}^\beta$ for all $K' > K$,*
- iii. *for all $\mu \in \mathcal{R}_K^\beta$ associated with a foliation W and all positive $\rho \in \text{Hol}_W^\beta$ such that $\int \rho d\mu = 1$, the probability measure $\rho\mu$ lies in $\mathcal{R}_{K'}^\beta$, where $K' = K + \text{Hol}_W^\beta(\log \rho)$,*
- iv. *$T_*(\mathcal{R}_{K_0}^\beta) \subset \mathcal{R}_{K_0}^\beta$ and $T_*^{n_0}(\mathcal{R}_{2K}^\beta) \subset \mathcal{R}_K^\beta$ for all $K \geq K_0$,*
- v. *for all $\mu_1, \mu_2 \in \mathcal{R}_{K_0}^\beta$ we can find a probability measure η concentrated on $\Delta_s(L_0)$ and two probability measures $\mu'_1, \mu'_2 \in \mathcal{R}_{2K_0}$ such that*

$$\mu_i = \tau p_{i*} \eta + (1 - \tau) \mu'_i \quad (i \in \{1, 2\}).$$

Moreover, for any $\beta \in (0, \alpha_0)$ we can make the map $T \mapsto K_0$ upper semi-continuous in the C^1 topology, in particular there exist $\varepsilon > 0$ such that to any map T_1 that is ε -close to T in the C^1 topology we can apply the above with the same β and a constant $K_1 \leq 2K_0$ in the role of K_0 (and further constants L_1, n_1).

While item *iii* might seem difficult to apply, since W might not be known and Hol_W^β is thus quite abstract a space, it can be applied e.g. to a β -Hölder function ρ , and in the case of the SRB measure when $W = W_u$.

Proof of Theorem 1.1. We first prove that D_s^β is uniformly bounded over all $\mu_1, \mu_2 \in \mathcal{R}_{K_0}^\beta$, and that the SRB measure is the unique T -invariant measure in \mathcal{R}_K^β . For now β is fixed anywhere in $(0, \alpha_0]$, and we will see along the proof how we need to further restrict it.

We will apply item *v* of Proposition 3.1 repeatedly, defining sequences $(\mu_1^k, \mu_2^k)_{k \geq 0}$ and $(\eta^k)_{k \geq 0}$ as follows. First, $\mu_i^0 = \mu_i \in \mathcal{R}_{K_0}^\beta$ are the measures we start with. Given μ_1^k, μ_2^k , we apply to them Proposition 3.1 to get a “partial coupling” η^k and “residual measures” $(\mu_i^k)' \in \mathcal{R}_{2K_0}$. Then we set

$$\mu_i^{k+1} = T_*^{n_0}(\mu_i^k)' \in \mathcal{R}_{K_0}.$$

By induction, for both i and all k :

$$T_*^{kn_0} \mu_i = \sum_{j=0}^k (1 - \tau)^j \tau \left[T^{(k-j)n_0} \circ p_i \right]_* \eta_j + (1 - \tau)^{k+1} (\mu_i^k)'$$

Applying $T_*^{-kn_0}$ to this equality and letting $k \rightarrow \infty$, it follows that

$$\gamma := \tau \sum_{j \geq 0} (1 - \tau)^j (T^{-jn_0}, T^{-jn_0})_* \eta_j$$

is a coupling of (μ_1, μ_2) . Since η_j is concentrated on $\Delta_s(L_0)$, letting $\lambda_0 = \inf\{\|D_x T u\| : x \in M, u \in E_x^s, \|u\| = 1\}$ be the maximal contraction of T on the stable direction,

$$\begin{aligned} \int d_s(x, y)^\beta d\gamma(x, y) &= \tau \sum_{j \geq 0} (1 - \tau)^j \int d_s(T^{-jn_0} x, T^{-jn_0} y)^\beta d\eta_j \\ &\leq \tau \sum_{j \geq 0} (1 - \tau)^j \int \lambda_0^{-j\beta n_0} d_s(x, y)^\beta d\eta_j \\ &\leq \tau L_0^\beta \sum_{j \geq 0} \left(\frac{1 - \tau}{\lambda_0^{\beta n_0}} \right)^j \end{aligned}$$

Set $\beta_0 = \min\left\{\alpha_0, \frac{\log(1-\tau)}{n_0 \log \lambda_0}\right\}$; then whenever $\beta \in (0, \beta_0)$, the above series is convergent and provides the desired uniform bound.

Last, by Proposition 3.1 the SRB measure μ_0 lies in some \mathcal{R}_K^β . It must be that $\mu_0 \in \mathcal{R}_{K_0}^\beta$: if $K \leq K_0$ this is direct, otherwise it suffices to apply T sufficiently many times and use invariance of μ_0 . Uniqueness of the T -invariant measure in this set then follows from (3), which only depends on the part of Theorem 1.1 we already proved. \square

3.2 Proof of Corollaries 1.2 and 1.3

Let μ_0 be the SRB measure of T and consider any fixed $\beta \in (0, \beta_0)$. By Proposition 3.1, $\mu_0 \in \mathcal{R}_{K_0}^\beta$.

Proof of Corollary 1.2. Let $f \in \text{Hol}_s^\beta$ and $g \in \text{Hol}_u^\beta$. Much like in Remark 2.2, we take a small enough (but independent of g) and $b = 1 - a\|g\|_{u,\beta}^{-1} \int g d\mu_0$ ensuring that $\rho := a\|g\|_{u,\beta}^{-1} g + b$ is positive, that $\int \rho d\mu_0 = 1$, and that $\rho\mu_0 \in \mathcal{R}_{2K_0}^\beta$ (item *iii* of Proposition 3.1). Then

$$\begin{aligned} \int f \circ T^n \cdot g d\mu_0 &= \frac{\|g\|_\alpha}{a} \int f \circ T^n \cdot (\rho - b) d\mu_0 \\ &= \frac{\|g\|_\alpha}{a} \left(\int f dT_*^n(\rho\mu_0) - b \int f dT_*^n \mu_0 \right) \\ &= \frac{\|g\|_\alpha}{a} \left(\int f dT_*^n(\rho\mu_0) - \int f d\mu_0 \right) + \int g d\mu_0 \int f d\mu_0 \\ &= \int g d\mu_0 \int f d\mu_0 + O\left(\|g\|_{u,\beta} \|f\|_{s,\beta} D_s^\beta(T_*^n(\rho\mu_0), \mu_0)\right) \end{aligned}$$

Using Theorem 1.1 and its consequence (1), since $T_*^{n_0}(\rho\mu_0) \in \mathcal{R}_{K_0}^\beta$, for all $n \in \mathbb{N}$:

$$D_s^\beta(T_*^n(\rho\mu_0), \mu_0) = D_s^\beta(T_*^{n-n_0}(T_*^{n_0}\rho\mu_0), T_*^{n-n_0}\mu_0) \leq C\lambda^{\beta(n-n_0)}.$$

\square

Proof of Corollary 1.3. We shall denote with an index 1 instead of 0 the quantities related to T_1 instead of T (e.g. μ_1, α_1, K_1 , etc.) with implied dependency on T_1, β , etc.

Fix any $\beta < \beta_0$; taking ε small enough ensures that the unstable distribution E_1^u of T_1 is tangent to the cone field fixed around E^u to define the $(\mathcal{R}_K^\beta)_K$, and that $\alpha_1 > \beta$. By Proposition 3.1 we can further assume that $K_1 \leq 2K_0$. Observe that whenever Proposition 3.1 holds for some value of K_0 , it also holds for all larger values. Up to enlarging K_0 , we can thus assume that $K_1 \leq K_0$ and that the unstable foliation of T_1 is β -Hölder of constant K_0 . This enlargement really depends on ε only, uniformly on all T_1 . We thus have $\mu_1 \in \mathcal{R}_{K_0}^\beta$.

Normalize the Riemannian metric to ensure the diameter of M is ≤ 1 . Let $A = \|T\|_{C^1} + \varepsilon$, so that both T and T_1 are A -Lipschitz. Set $D = D_1(\mu_0, \mu_1)$, and fix $n \in \mathbb{N}$ large enough to ensure that $D_\beta^s(T_*^n \mu_1, T_*^n \mu_0) \leq \frac{1}{2} D_1(\mu_0, \mu_1)$; we can take $n \simeq \log 1/D$, so that $A^n \simeq D^{-p}$ for some $p > 0$ independent of T_1 . Then

$$\begin{aligned} D = D_1(\mu_1, \mu_0) &\leq D_1(\mu_1, T_*^n \mu_1) + D_1(T_*^n \mu_1, \mu_0) \\ &= D_1(T_{1*}^n \mu_1, T_*^n \mu_1) + D_1(T_*^n \mu_1, T_*^n \mu_0) \\ &\leq \|T_1^n - T^n\|_{C^0} + D_\beta^s(T_*^n \mu_1, T_*^n \mu_0) \end{aligned}$$

We bound $\|T_1^n - T^n\|_{C^0}$ by induction: for all $x \in M$,

$$\begin{aligned} d(T_1^n x, T^n x) &\leq d(T_1(T_1^{n-1} x), T_1(T^{n-1} x)) + d(T_1(T^{n-1} x), T(T^{n-1} x)) \\ &\leq A d(T_1^{n-1} x, T^{n-1} x) + \|T_1 - T\|_{C_0} \\ &\leq (1 + A + \dots + A^{n-1}) \|T_1 - T\|_{C_0} \\ &\lesssim A^n \|T_1 - T\|_{C_0} \end{aligned}$$

Plugging this into the previous computation, we get

$$\begin{aligned} D &\lesssim A^n \|T_1 - T\|_{C^0} + \frac{D}{2} \\ \frac{D}{2} &\lesssim D^{-p} \|T_1 - T\|_{C^0} \\ D &\lesssim \|T_1 - T\|_{C^0}^{\frac{1}{1+p}}. \end{aligned}$$

□

3.3 Technical part of the proof: $(\mathcal{R}_K^\beta)_K$ and its properties

We start by recalling classical facts on Anosov diffeomorphisms and setting up some notations.

Recall that we assume $T : M \rightarrow M$ to be a $C^{1+\alpha}$, topologically mixing Anosov diffeomorphism. We denote by E^s, E^u the *stable* and *unstable* distributions, which are T -invariant, of constant dimensions k_s, k_u , and such that $E_x^s \oplus E_x^u = T_x M$ for all $x \in M$. By choosing an adapted Riemannian metric on M , we can assume that there exist $\lambda \in (0, 1)$ such that for all $x \in M, u \in E_x^s, v \in E_x^u$ and $n \in \mathbb{N}$:

$$\|D_x(T^n)(u)\| \leq \lambda^n \|u\|, \quad \|D_x(T^{-n})(v)\| \leq \lambda^n \|v\|.$$

The stable and unstable distribution can be integrated into the *stable* and *unstable foliations* $(W_x^s)_x, (W_x^u)_x$.

The Riemannian metric on M induces a canonical Riemannian metric on each Grassmannian $G(k, T_x M)$ (the space of all k -dimensional subspaces of $T_x M$), we denote the induced distance function by $d_{G(k, T_x M)}$. We can in particular define for all continuous k -dimensional distributions E, F :

$$d(E, F) := \max_{x \in M} d_{G(k, T_x M)}(E_x, F_x);$$

the Anosov property in particular implies that T_* acts on $G(k_u, T_x M)$ with E^u as an exponentially attracting fixed point: for some $\eta \in (0, 1)$ and some $\varepsilon > 0$, for all $x \in M$ and all $E_x \in G(k_u, T_x M)$ such that $d_{G(k_u, T_x M)}(E_x, E_x^u) \leq \varepsilon$,

$$d_{G(k_u, T_{T(x)} M)}(D_x T(E_x), E_{T(x)}^u) \leq \eta \cdot d_{G(k_u, T_x M)}(E_x, E_x^u).$$

We fix a field of open cones $(C_x)_{x \in M}$ in the tangent bundle TM containing E_u , strongly preserved by T (meaning $\overline{T(C_x)} \subset C_{T(x)}$ for all $x \in M$) and in such a way that the above contraction holds on each $U_x := \{E \in G(k_u, T_x M) \mid E \subset C_x\}$.

To compare subspaces of $T_x M$ at different nearby base points x , we simply fix a finite smooth atlas of M ; we say that a k -dimension distribution E is β -Hölder if for some constant K , in all charts in the fixed atlas, E is β -Hölder with that constant K . The exponent β does not depend on the choice of atlas, but the constant does.

There exist some $\alpha_0 \in (0, \alpha)$ such that:

- E^s, E^u are α_0 -Hölder continuous, and in particular the leafs of W^s, W^u are $C^{1+\alpha_0}$ submanifolds;
- the holonomies $\pi_s^{N_1 \rightarrow N_2}$ and $\pi_u^{N_1 \rightarrow N_2}$ of W^s, W^u between C^{1, α_0} transversals N_1, N_2 to E^s (or E^u) are α_0 -Hölder and absolutely continuous with α_0 -Hölder jacobian; moreover the α_0 -Hölder constants can be chosen uniform over all (N_1, N_2) that are bounded in diameter, are written as graphs of bounded C^{1, α_0} norm, and are at bounded distance one from the other along W^s (or W^u).

While α_0 depends on T , it does so only through a few parameters (α , the α -Hölder constants of E^s and E^u , λ , a uniform bound on DT , etc.) and it can be chosen such that every $C^{1+\alpha}$ maps T_1 that is sufficiently close to T in the C^1 topology shares the same α_0 .

From now on we fix any $\beta \in (0, \alpha_0)$.

3.3.1 Définition of $(\mathcal{R}_K^\beta)_K$

While elements of $(\mathcal{R}_K^\beta)_K$ are probability measures, their crucial defining property is a relation with some foliation, which we thus have to define beforehand. We will need to define a regularity constant to control certain submanifolds; to this end we will again use a specified atlas, but with some additionnal features.

We consider smooth charts $\varphi : U_\varphi \subset M \mapsto V_\varphi \subset \mathbb{R}^k$ (where k is the dimension of M) whose range V_φ are of the form $B^{k_u}(0, 1) \times B^{k_s}(0, 1) \subset \mathbb{R}^k$, where $B^\bullet(p, r)$ denotes the ball of center p and radius r in a factor \mathbb{R}^\bullet of \mathbb{R}^k , with respect to the canonical Euclidean metric. A typical point of V will be denoted by (p, q) , with $p \in B^{k_u}(0, 1)$, $q \in B^{k_s}(0, 1)$. We denote by $x_\varphi := \varphi^{-1}(0, 0)$ the “center” of the chart, and we request that the “slices” $\varphi^{-1}(B^{k_u}(0, 1) \times \{q\})$ are C^1 -close to $W_{\varphi^{-1}(0, q)}^u$, in particular their tangent space shall be contained in the cone field C at all point; and similarly the $\varphi^{-1}(\{p\} \times B^{k_s}(0, 1))$ shall be C^1 -close to $W_{\varphi^{-1}(p, 0)}^s$, in particular their tangent space shall avoid the closure $\bar{C} = (\bar{C}_x)_x$ of the cone field at all point. We can further assume (taking the domains U_φ somewhat tall in the stable direction and thin in the unstable direction) that the image φ_*C of the cone field is very narrow; in particular that every k_u -dimensional C^1 submanifold N whose tangent space is everywhere contained in C and that meets $\text{Core}_\varphi := \varphi^{-1}(B^{k_u}(0, 1/2) \times B^{k_s}(0, 1/2))$ at some point x , can be written locally around x as $\varphi^{-1}(G)$ where G is a graph of a map $B^{k_u}(0, 1/2) \rightarrow B^{k_s}(0, 1)$. In particular, every foliation whose tangent distribution is everywhere contained in C admits a local trivialization that cover the core of φ .

From now on we fix a finite atlas of such charts, where the Core_φ cover M .

Definition 3.2. Let us say that a k_u -dimensional foliation W of M is K -adapted (to T, C, β) when the leafs of W are $C^{1+\beta}$ and their tangent space is inside c at all point, and in each chart of the specified atlas the leafs are graphs of β -Hölder with constant at most K .

Definition 3.3. Let \mathcal{R}_K^β be the set of probability measure μ on M such that there exist a K -adapted foliation W with respect to which μ has absolutely continuous local disintegrations with positive β -Hölder densities, and each of those densities ρ satisfy

$$\frac{\rho(x)}{\rho(y)} \leq e^{K d_W(x, y)^\beta} \quad (4)$$

where d_W denotes the distance along the leafs of W . We say that K is a *log constant* of *log β -Hölder constant* for ρ , or for μ .

Assume U_1, U_2 are two intersecting open sets on which W can be trivialized and N_1, N_2 are transversals; i.e. $U_i = \cup_{y \in N_i} W_y \cap U_i \simeq N_i \times B^{k_u}(0, 1)$ and consider $x \in U_1 \cap U_2$. Then there are two different densities ρ_1, ρ_2 corresponding to the disintegrations of the restrictions of μ to U_1, U_2 with respect to N_1 and N_2 ; but this densities are proportional to each other along leaves where they are both defined. Indeed, denoting by $p_i : U_i \rightarrow N_i$ the holonomic projection to the transversal, $\rho_1(x)/\rho_2(x)$ is the Radon derivative of $\pi_*^{N_1 \rightarrow N_2} p_{1*} \mu$ with respect to $p_{2*} \mu$ at $y = p_2(x)$. In particular, equation (4) holds for both ρ_i simultaneously, with the same constant.

3.3.2 Proof of Proposition 3.1

We consider $\beta \in (0, \alpha_0)$ and the above specified atlas fixed. For each $r > 0$ and $x \in M$, set

$$B_\times(x, r) = \{y \in M \mid \exists z \in M, d_s(x, z) < r, d_u(z, y) < r\}.$$

This “product balls” form a basis of the topology of M , and we can find $\delta_-, \delta_+ > 0$ such that for every chart φ , $B_\times(x_\varphi, \delta_-) \subset \text{Core}_\varphi \subset U_\varphi \subset B_\times(x_\varphi, \delta_+)$. Up to refining the atlas, we can moreover ensure that δ_+ is small enough that both the stable and unstable foliations of T are topologically trivial at scale δ_+ (i.e. the $B_\times(x, \delta_+)$ are domains of trivializing charts). Since T is assumed to be topologically mixing, its stable leafs are dense and we can find \tilde{L}_0 such that for every pair of charts φ, ψ , there is a stable path of length at most \tilde{L}_0 from $\varphi^{-1}(0, 0)$ to a point $\delta_-/100$ -close to x_ψ in the distance d_u . We set $L_0 = \tilde{L}_0 + 2\delta_+$.

To see that the SRB measure μ_0 lies in some \mathcal{R}_K^β , it suffices to use the usual expression for the densities of its local disintegrations as an infinite product involving the unstable jacobian of T , see section 9.3 in [BP07].

By definition, we immediately get $\mathcal{R}_K^\beta \subset \mathcal{R}_{K'}^\beta$, whenever $K' > K$. Moreover if $\mu \in \mathcal{R}_K^\beta$ and $\rho \in \text{Hol}_W^\beta$ is a positive density with respect to μ , then $\rho\mu$ has absolutely continuous disintegrations with respect to the same K -adapted foliation W as μ ; if ρ_1 is a local leaf density of μ , then the local leaf density of $\rho\mu$ is proportional to $\rho\rho_1$, hence $\rho\mu \in \mathcal{R}_{K'}^\beta$ for $K' = K + \|\log \rho\|_{W, \beta}$.

As in Section 2, there is $H > 0$ and $\lambda' \in (0, 1)$ such that whenever $\mu \in \mathcal{R}_K^\beta$, $T_*\mu \in \mathcal{R}_{\lambda'(K+H)}^\beta$; the constant λ' is β th power of the contraction factor of T on every submanifold whose tangent space is contained in C at all point, and H accounts for the unstable jacobian of T and the way T and change of charts distort β -Hölder graphs. We can thus define K_0 and n_0 by

$$K_0 = 2 \frac{\lambda'}{1 - \lambda'} H, \quad 2\lambda'^{n_0} \leq \frac{\lambda'}{1 - \lambda'}$$

to obtain that $T_*(\mathcal{R}_{K_0}^\beta) \subset \mathcal{R}_{K_0}^\beta$ and $T_*^{n_0}(\mathcal{R}_{2K}^\beta) \subset \mathcal{R}_K^\beta$ for all $K \geq K_0$.

We now prove the core coupling property, item *v*. Let $\mu_1, \mu_2 \in \mathcal{R}_{K_0}^\beta$ and denote by W^1, W^2 their K_0 -adapted foliations. Denote by N the number of charts in the chosen atlas and recall that the $\text{Core}_\varphi := \varphi^{-1}(B^{k_u}(0, 1/2) \times B^{k_s}(0, 1/2))$ cover M . We can thus find two charts φ_1, φ_2 such that $\mu_i(\text{Core}_{\varphi_i}) \geq 1/N$ for both i ; and since the cores are covered by trivializing charts of the W^i , we can locally disintegrate restriction $\tilde{\mu}_i$ of the μ_i through the holonomic projections p_i to local stable leaves $D_i \subset W_{x_{\varphi_i}}^s$:

$$\mu_i = \int_{D_i} \rho_{i,x}(y) \, \text{dvol}_{W_x^i}(y) \, \text{d}\eta_i(x)$$

where η_i are positive measures on D_i of mass at least $1/N$ and ρ_i are β -Hölder with log constant at most K_0 . Consider a $1/N$ -partial coupling $\tilde{\gamma}$ of (η_1, η_2) , i.e. γ is a positive measure of mass $1/N$ on $M \times M$ with marginals $\leq \eta_i$; for concreteness, let us take $\tilde{\gamma} = \frac{1}{N} \eta_1 \otimes \eta_2$. For each pair of leafs W_x^1, W_y^2 we construct a partial coupling of the $\rho_i \, \text{dvol}_{W_x^i}$ as follows. Given $h > 0$, Let $f = f_{x,h} : W_x^1 \rightarrow \mathbb{R}_+$ be defined by $f(z) = \max(0, h - \frac{10h}{\delta_-} d_{W_x^1}(x, z))$. Let $g = g_{y,h}$ be defined as the composition of f with the holonomy $\pi_{x,y}^s$ of W^s , along a curve of length at most L_0 . By choosing $h \leq h_0$ with h_0 small enough, we can ensure that both $\rho_{1,x} - f_{x,h}$ and $\rho_{1,x} - g_{y,h}$ are positive and β -Hölder with log constant at most $2K_0$ (indeed, f is Lipschitz with constant only

depending on δ_- , hence on T , and g is α_0 -Hölder with constant depending further on L_0 and the stable holonomy Hölder constant, thus still only depending on T). Now all considered submanifold in the foliations are $C^{1,\beta}$ with bounded Hölder constant in the specified charts, so that there exist $\tilde{\tau} > 0$ such that the function f_{x,h_0} has integral at least $\tilde{\tau}$ with respect to $\text{vol}_{W_x^1}$. By choosing $h = h_x \leq h_0$ depending on x , we can make $f_{x,h}$ of integral exactly $\tilde{\tau}$. finally, our partial coupling of μ_1, μ_2 is given by

$$\gamma = \int (\text{Id}, \pi_{x,y}^s)_* (f_{x,h_x} \text{vol}_{W_x^1}) \, d\tilde{\gamma}(x, y).$$

It has mass $\tau := \tilde{\tau}/N$, and its marginals are $\leq \mu_i$; moreover $\mu'_i := \mu_i - p_{i*}\gamma$ have local disintegrations with respect to W^i that are β -Hölder with log constant at most $2K_0$, hence $\mu'_i \in \mathcal{R}_{2K_0}^\beta$.

3.3.3 Final remarks

A similar method can be used to handle hyperbolic attractors instead of Anosov maps, but Theorem 1.1 does not hold anymore in this setting (stable leafs are no longer dense, and most measures to be considered will no longer be at finite D_s^β distance, for any β). What can be done is to pair iterates of measures $T_*^{n_1}\mu_1, T_*^{n_1}\mu_2$ where n_1 is chosen to ensure that a significant part of the mass of μ_1, μ_2 are at small stable distance one from the other.

If we relax uniform hyperbolicity, we can hope to keep weaker results in the same vein. In the case of lack of uniform hyperbolicity in the stable direction, it might be necessary to use $\omega \circ d_s$ instead of d_s^β , where ω is a concave function increasing quicker than any Hölder function at 0. In the case of lack of uniform hyperbolicity in the unstable direction, it might be necessary to use densities and observables with an adapted regularity. In both cases, weaker speed for the decay of correlation is of course expected.

It might be possible to adapt the present method to maps of flows with discontinuities, given it relies on a relatively flexible coupling argument: pair together along the stable direction part of any two measures that are regular enough in the unstable direction, then apply the dynamics long enough to recover the initial regularity for the remaining, uncoupled parts. Rince and repeat.

References

- [AdCJ04] A. ARMANDO DE CASTRO JÚNIOR – “Fast mixing for attractors with a mostly contracting central direction”, *Ergodic Theory Dynam. Systems* **24** (2004), no. 1, p. 17–44. [1](#)
- [AS12] J. F. ALVES et M. SOUFI – “Statistical stability and limit laws for Rovella maps”, *Nonlinearity* **25** (2012), no. 12, p. 3527–3552. [1](#)
- [BP07] L. BARREIRA et Y. PESIN – *Nonuniform hyperbolicity*, Encyclopedia of Mathematics and its Applications, vol. 115, Cambridge University Press,

Cambridge, 2007, Dynamics of systems with nonzero Lyapunov exponents.
[3.3.2](#)

- [BV96] V. BALADI et M. VIANA – “Strong stochastic stability and rate of mixing for unimodal maps”, *Ann. Sci. École Norm. Sup. (4)* **29** (1996), no. 4, p. 483–517.
[1](#)
- [BY00] M. BENEDICKS et L.-S. YOUNG – “Markov extensions and decay of correlations for certain Hénon maps”, no. 261, 2000, Géométrie complexe et systèmes dynamiques (Orsay, 1995), p. xi, 13–56. [1](#)
- [Dol98] D. DOLGOPYAT – “On decay of correlations in Anosov flows”, *Ann. of Math. (2)* **147** (1998), no. 2, p. 357–390. [1](#)
- [DSL16] J. DE SIMOI et C. LIVERANI – “Statistical properties of mostly contracting fast-slow partially hyperbolic systems”, *Invent. Math.* **206** (2016), no. 1, p. 147–227. [1](#)
- [KKM19] A. KOREPANOV, Z. KOSLOFF et I. MELBOURNE – “Explicit coupling argument for non-uniformly hyperbolic transformations”, *Proc. Roy. Soc. Edinburgh Sect. A* **149** (2019), no. 1, p. 101–130. [1](#)
- [Liv95] C. LIVERANI – “Decay of correlations”, *Ann. of Math. (2)* **142** (1995), no. 2, p. 239–301. [1](#)
- [Var08] P. VARANDAS – “Correlation decay and recurrence asymptotics for some robust nonuniformly hyperbolic maps”, *J. Stat. Phys.* **133** (2008), no. 5, p. 813–839. [1](#)
- [You98] L.-S. YOUNG – “Statistical properties of dynamical systems with some hyperbolicity”, *Ann. of Math. (2)* **147** (1998), no. 3, p. 585–650. [1](#)