

#### Rapid screening of epidemiologically important Salmonella enterica subsp. enterica serovars by Matrix-Assisted Laser Desorption Ionization—Time of Flight (MALDI-TOF) mass spectrometry

Frédérique Moury, Benoît Gassilloud, Jean-Sébastien Py, Wilhelm Amandine, Antonin Ribeyre, Christine Piquet, Viviane Morel, Renaud Lailler, Sabrina Cadel Six

#### ▶ To cite this version:

Frédérique Moury, Benoît Gassilloud, Jean-Sébastien Py, Wilhelm Amandine, Antonin Ribeyre, et al.. Rapid screening of epidemiologically important Salmonella enterica subsp. enterica serovars by Matrix-Assisted Laser Desorption Ionization—Time of Flight (MALDI-TOF) mass spectrometry. Societé Française de Microbiologie (SFM), Mar 2016, Paris, France. https://na.eventscloud.com/ehome/134460/www.sfm-microbiologie.org. hal-04205250

HAL Id: hal-04205250

https://hal.science/hal-04205250

Submitted on 14 Sep 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Rapid screening of epidemiologically important *Salmonella* enterica subsp. enterica serovars by Matrix-Assisted Laser Desorption Ionization—Time of Flight (MALDI-TOF) mass spectrometry.



Frédérique Moury<sup>1</sup>, Benoît Gassilloud<sup>2</sup>, Jean-Sébastien Py<sup>2</sup>, Wilhelm Amandine<sup>2</sup>, Antonin Ribeyre<sup>2</sup>, Christine Piquet<sup>1</sup>, Viviane Morel<sup>1</sup>, Renaud Lailler<sup>1</sup> and Sabrina Cadel Six<sup>1</sup>

1 Université PARIS-EST; ANSES; Food Safety Laboratory; Listeria, Salmonella, E. coli unit; Maisons Alfort, France.

2 MALDI-TOF Platform; Nancy Laboratory for Hydrology; ANSES, Nancy, France.



12<sup>e</sup> Congrès Nationale de la Société Française de Microbiologie 22 et 23 mars 2016, Paris - France

Salmonella spp. is one of the most widely distributed foodborne pathogens in the European Union (EU) resulting in significant morbidity, mortality and social and economic cost. Accurate and rapid typing of this pathogen is essential for effective surveillance nevertheless conventional serotyping, which enables to type the 2600 serovars known, is a laborious procedure, time consuming and expensive. Private and public laboratories implied in surveillance need a rapid and not expensive prescreening method for at least the 25 serovars, the most frequently isolated [1].


MALDI-TOF mass spectrometry could represent an alternative. In the last decade this technique was considered as revolutionary allowing making the diagnosis of pathogens more easily and quickly compared to conventional methods however even if its ability to discriminate pathogens at genus and specie level was well established, at serovars level it remains scare and needs to be explored.


The **aim of this study** was to evaluate the potential of MALDI-TOF mass spectrometry as prescreening method for typing of the 5 Salmonella serovars considered to be of public health significance under EU regulations (the regulated serovars): Enteritidis, Hadar, Virchow, Infantis, Typhimurium and its variant S.4,[5],12:i:-

## Results:

Statistical analysis via ClinProTools:


 Global ACP from 36 raw spectra:
the 5 serovars are distinct with the exception of the variant S.4,[5],12:i:-






Characteristics
 biomarkers via
 alignment of
 spectra were found
 for the 5 serovars

 A mapping of the specific mass for each regulated serovars was realized



## Method employed:

The strains were grown 24 h at 36 °C over Drigalski medium. The mass spectrometry MALDITOF identification was realized with a Microflex LT® (Bruker Daltonics) after proteomic extraction with ethanol, formic acid and acetonitrile.



## Conclusion and outlook:

MALDI-TOF/MS seems to be a possible methodology which could be use for the prescreening of the 5 regulated serovars.

Nevertheless the specificity of each mass, selected for the characterization, must be evaluated compared with other serovars and additional robustness tests are essential before using these methodology routinely.

Both these tests are actually in process in our laboratory onto 20 others serovars, the most frequently isolated [1].

#### References:

[1] Frédérique Moury, Sylvine Frémy, Claude Oudart, Christine Piquet, Catherine Pires Gomes, Corinne Danan, Sophie Granier, Renaud Lailler, Anne Brisabois. Inventaire des *Salmonella* d'origine non humaine - Réseau *Salmonella* 2011. Edité en décembre 2013.

anses.fr