



HAL
open science

Revisiting β -dicyanovinyl substituted calix[4]pyrrole: Toward the chemodosimetric detection of hydrazine in solution

Amaury Kasprowiak, Ishfaq Ahmad Rather, Rashid Ali, Pierre-Edouard Danjou

► To cite this version:

Amaury Kasprowiak, Ishfaq Ahmad Rather, Rashid Ali, Pierre-Edouard Danjou. Revisiting β -dicyanovinyl substituted calix[4]pyrrole: Toward the chemodosimetric detection of hydrazine in solution. *Journal of Molecular Structure*, 2023, 1287, pp.135694. <10.1016/j.molstruc.2023.135694>. <hal-04210185>

HAL Id: hal-04210185

<https://hal.science/hal-04210185v1>

Submitted on 6 Jan 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Revisiting β -dicyanovinyl substituted calix[4]pyrrole : toward the chemodosimetric detection of hydrazine in solution

Amaury Kasprowiak^a, Ishfaq Ahmad Rather^b, Rashid Ali^{b*}, Pierre-Edouard Danjou^{c*}

Email : [rali1@jmi.ac.in](mailto:r.ali1@jmi.ac.in) ; danjou@univ-littoral.fr

Affiliation

^a Département de Chimie, Université du Littoral Côte d'Opale, 220 avenue de l'université, 59140 Dunkerque, France

^b Organic and Supramolecular Functional Materials Research Laboratory, Department of Chemistry, Jamia Millia Islamia, Jamia Nagar, Okhla, New Delhi-110025, India

^c Unité de Chimie Environnementale et Interactions sur le Vivant, UR 4492, Université du Littoral Côte d'Opale, 145 Avenue Maurice Schumann, MREI 1, 59140 Dunkerque, France

Abstract

Calix[4]pyrrole derivatives are usually used to recognize charged species or polar guests through non-covalent interactions, but the chemodosimetric approach remains scarce in the literature. In this study, the selective chemodosimetric detection and quantification of hydrazine, a hazardous pollutant commonly used in industry, was performed using UV-spectroscopy with a repurposed β -dicyanovinyl substituted calix[4]pyrrole sensor. The selectivity of the chemodosimeter towards hydrazine was evaluated in acetonitrile with various nucleophiles (nitrogen-containing compounds and a thiol). In addition, the influence of several parameters (time, water content, and temperature) on hydrazine detection by the sensor was evaluated. This study allows for the sensing of hydrazine with a limit of detection (LOD) of 1.3 mg/L and a linear response in the 10-1000 μ M range. The ability to detect hydrazine with the naked eye has also been demonstrated. This paper reports one of the first chemodosimetric approach employed with calix[4]pyrrole to detect and quantify a neutral molecule, namely hydrazine.

Introduction

Hydrazine, a small covalent molecule, is regarded as a hazardous environmental pollutant and being a carcinogenic chemical has proved to be highly toxic for both humans and animals.¹ The continuous exposure to hydrazine leads to nose, throat, and eye irritation, dizziness, nausea, pulmonary edema, etc., which in turn proves fatal for kidneys, liver and central nervous system in human beings.² As a matter of the fact, U.S based Environmental Protection Agency (EPA) has marked hydrazine as potential carcinogen and fixed an Immediately Dangerous to Life and Health (IDLH) value of 50 ppm in air.³ By virtue of

widespread use of hydrazine in industries (pharmaceuticals, polymers, dyes etc.)⁴, hydrazine pollution in water and air has become a major concern globally. Indeed, it was reported that upon exposure to about 0.1 ppm hydrazine in air, the urine of Japanese workers contains an average of 2.7 ppm of hydrazine.⁵ It is also expected to find concentrations above 10 ppm in urine in case of an event-level exposure.⁶ Thus, there arises a need to efficiently monitor the level of hydrazine in air, water, human fluids and other related systems. To date, numerous conventional analytical methods have been employed for the detection of hydrazine like chromatography, spectrophotometric flow injection, and electroanalytical methods.⁷⁻⁹ However, naked eye and UV-visible detection are particularly attractive because they are generally cheap, fast, selective, and reliable, even if the detection limit is not as low as other methods such as molecular fluorescence spectroscopy¹⁰⁻¹³. Analytical methods utilizing optical signatures and supramolecular sensors/receptors have been of great interest to the scientific community in recent years.^{14,15} Among the diverse supramolecular receptors available, the tetrapyrrolic macrocyclic receptor known as calix[4]pyrrole (C4P) has shown a promising role in the recognition of diverse analytes¹⁶. C4P possess a built-in recognition site able to recognize anions or polar guests, while its functionalization at the level of meso-carbon or pyrrole allow to achieve cation or ion-pair recognition through non-covalent interactions.¹⁷⁻¹⁹ To date, the use of calix[4]pyrrole scaffold for chemodosimetric detection (i.e. detection with significant chemical transformation involving both breaking and forming of the covalent bonds²⁰) of non-charged molecules is still scarce²¹. In this study, we repurposed a β -dicyanovinyl substituted C4P preliminary designed for the recognition of cyanide anion in order to achieve the naked eyed and spectroscopic detection of hydrazine in acetonitrile (ACN) and acetonitrile-water mixture.

Material and Methods

NMR spectra were recorded on a Bruker Ascend 400 spectrometer operating at 400 MHz and 101 MHz for ¹H and ¹³C respectively. Traces of residual solvent were used as the internal standard. High-resolution mass spectrometry spectra (HRMS) were recorded in the positive mode using an Agilent 6540 Accurate-Mass Quadrupole Time-of-Flight (Q-TOF) LC/MS system equipped with an Agilent Jet Stream dual electrospray source. UV-Vis measurements were performed at a fixed temperature using a Varian Cary 3500 UV-Vis spectrophotometer equipped with a compact Peltier apparatus. A quartz suprasil 100-QS TO 10 mm pathlength cuvette was used.

Synthetic procedure for compound 3: To a clear low melting mixture (5.00 g) of *L*-(+)-tartaric acid (TA) and *N,N'*-dimethyl urea (DMU) in the ratio of 3:7 at 70 °C, was added freshly distilled pyrrole (5.00 ml, 72.07 mmol, 1 equiv) and acetone (26.69 ml, 360.34 mmol, 5 equiv). The reaction mixture was stirred at 70 °C for 4-5 h and was monitored by TLC. After the completion of reaction, the warm reaction mixture was diluted with water (20 ml). The reaction mixture was cooled to room temperature, filtered through a sintered glass funnel and the solid material was washed with water (4 x 20 mL). The crude product was purified by silica gel column chromatography using (5% EtOAc:petroleum ether). The first eluted fraction was the desired compound **3** (4.64 g, 60% yield, white solid). ¹H NMR (400 MHz, CDCl₃) δ 7.06 (s, 4H), 5.93 (d, *J* = 2.4 Hz, 8H), 1.54 (s, 24H). ¹³C NMR (101 MHz, CDCl₃) δ 138.67, 102.93,

35.61, 29.09; HRMS (ESI, Q-TOF) m/z : calculated for $C_{28}H_{36}N_4$ $[M+H]^+$ 429.3013, found: 429.3021. The spectral data is in accordance with the literature²².

Synthetic procedure for compound 4: In a dried two-necked 50 ml round bottomed flask, $POCl_3$ (0.52 ml, 5.60 mmol) was added dropwise at 0 °C under an inert nitrogen atmosphere to dry DMF (0.43 ml, 5.60 mmol) with vigorous stirring. After some time, dry CH_2Cl_2 (5 ml) was added to this reaction mixture and left for stirring at 0 °C for 45 minutes under inert atmosphere to obtain a Vilsmeier reagent. This prepared reagent was later on added dropwise at 0° C under inert atmosphere to a stirred solution of compound 3 (2.40 g, 5.60 mmol) in dry CH_2Cl_2 (100 ml) prepared in another two-necked 250 ml round bottomed flask. The solution turns dark red and was left for stirring at 0 °C for 2 h. After the completion of the reaction, the reaction mixture was thoroughly washed with saturated aqueous Na_2CO_3 and NaCl. The organic contents were extracted in CH_2Cl_2 , dried over Na_2SO_4 , filtered, and concentrated over vacuum to afford crude red solid. This crude mixture was later on subjected to column chromatography (SiO_2 , CH_2Cl_2 followed by 5% CH_2Cl_2 :ethyl acetate) to get pure **4** (0.537 g, 21% yield, white solid). 1H NMR (500 MHz, $CDCl_3$) δ 10.09 (s, 1H), 7.35 (s, 2H), 6.96 (s, 1H), 6.47 (s, 1H), 6.0 – 5.83 (m, 7H), 1.73 (s, 6H), 1.53 (d, s, 12H), 1.49 (s, 6H). The spectral data is in accordance with the literature²³.

Synthetic procedure for chemodosimeter 5: In a dried two-necked 50 ml round bottomed flask, compound **4** (0.50 g, 1.10 mmol) and malononitrile (89 μ l, 1.60 mmol) were dissolved in 18 ml of toluene at room temperature. To this stirred solution, piperidine (0.15 ml, 1.52 mmol) was added and left for stirring at room temperature for 6 h. After the reaction gets completed, the organic contents were extracted in CH_2Cl_2 , dried over anhydrous Na_2SO_4 , and concentrated over vacuum to afford the crude mixture. This crude mixture was then subjected to column chromatography (SiO_2 , pure CH_2Cl_2) to afford pure sensor **5** (381 mg, 69% yield) as a yellowish solid. 1H NMR (400 MHz, $CDCl_3$) δ 7.95 (s, 1H), 7.42 (bs, 1H), 7.26 (bs, 1H), 7.20 (bs, 1H), 6.95 (d, $J=2.7$ Hz, 1H), 6.88 (bs, 1H), 5.95-6.02 (m, 4H), 5.91 (m, 1H), 5.85 (m, 1H), 1.68 (s, 3H), 1.55 (s, 3H), 1.53 (s, 3H), 1.50 (s, 3H). The spectral data is in accordance with the literature²⁴.


UV-spectroscopy analysis : For UV spectroscopy analysis, a stock solution of sensor **5** was prepared in pure acetonitrile at 1.10^{-3} M and then diluted to 10 μ M (10 mL). Stock solutions of nucleophilic compounds were also prepared in pure acetonitrile at 1.10^{-1} M. An aliquot of the nucleophilic compounds (100 equivalents) was added to the 10 μ M solution of sensor **5** and then UV spectra were recorded every 10 minutes during 2 hours. For calibration curves, 7 solutions of compound **5** at a concentration of 10 μ M in acetonitrile were incubated in a water bath at 25 °C. Every two minutes, a known concentration of hydrazine (i.e. 10 equiv. for solution 2, 20 equiv. for solution 3, 40 equiv for solution 4 etc.) was added to each individual solution. The two minutes interval between hydrazine addition allow the operator to fill the UV cuvette and record UV spectra at known time interval of 1, 2 and 4 hours.

Results and discussion:

1) synthesis and characterization

The synthesis of sensor **5** is described in Scheme 1. First, octamethylcalix[4]pyrrole **3** was obtained from the condensation of pyrrole with acetone following a green procedure employing

a deep eutectic solvent.²² Then compound **4** and **5** were obtained according to previously published procedures.^{23,24} The ¹H NMR spectra in CDCl₃ is in accordance with the literature (Figures S1-2).



Scheme 1. Synthesis of β -vinyl substituted calix[4]pyrrole (sensor **5**) and its reaction with hydrazine

2) NMR investigation

Prior to UV-spectroscopy studies, the reaction of sensor **5** with hydrazine was investigated by ¹H NMR spectroscopy. The addition of an excess hydrazine to a solution of sensor **5** in CD₃CN led after a few minutes to a change in color from yellow to almost colorless. The aromatic part of the ¹H NMR spectrum of sensor **5** and its reaction product with hydrazine are presented in Figure 1. It can be seen that the reaction produce a single new product, presenting a significant upfield shift of the proton present on the pyrrole bearing the sensing arm ($\Delta\delta = 0.69$ ppm) and a slight shielding of vinylidene singlet proton at 8.12 ppm ($\Delta\delta = 0.008$) ppm. These observations are in agreement with the previously published literature employing the dicyanovinyl group as sensing unit.²⁵ Moreover, the proposed structure of compound **6** was consistent with the high resolution mass spectrometry experiment (Figure S4).




Figure 1 : Partial ^1H NMR spectra (400 MHz, CD_3CN , 298K) of sensor **5** before (top) and after (bottom) addition of an excess of hydrazine. (full spectra are provided in Figure S5)

3) Study of selectivity

The recognition behavior of sensor **5** was firstly studied by UV-spectroscopy in pure acetonitrile towards various nucleophilic compounds (hydrazine, ammonia, propylamine, diethylamine, triethylamine, pyrrole, aniline, and 2-mercaptoethanol). The UV spectrum of sensor **5** presents a maximum absorption at 373 nm, which is similar to previously published data²⁴ recorded in an ACN-DMSO (3%) mixture and possesses a molar extinction coefficient of $19303 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$ (Figure S6). This value should allow the observation of reasonably low concentration of analytes.²⁶

As expected from the NMR experiments, the addition of 100 equivalents of hydrazine to a solution of sensor **5** at $10 \mu\text{M}$ in ACN led to a decrease in absorbance at 373 nm (Figure 2) with the emergence of an isobestic point at 298 nm. It is important to note that the reaction between the dicyanovinyl group of sensor **5** and hydrazine was time-dependant. To take into account the dependency of the UV response on time, UV spectra were acquired every 10 minutes for 2 hours for each nitrogen-containing compounds (Figures S7-13). To our delight, the addition of 100 equivalents of other nucleophilic compounds did not significantly modify the absorption spectra of sensor **5** after 60 or 120 minutes (Figure 3A), indicating that sensor **5** was highly selective towards hydrazine. A competition experiment was also performed (Figure 3B). In this experiment, Sensor **5** was exposed to 100 equivalents of the 7 competing nucleophiles (ammonia, propylamine, diethylamine, triethylamine, pyrrole, aniline, and 2-mercaptoethanol) and a UV spectra was recorded after an incubation time of 2h. A modest decrease in absorbance intensity of 8% was observed. Then 100 equivalents of hydrazine

were added to the previous solution and allow to react for 2 more hours. As expected, a drastic decrease in absorbance intensity was observed, indicating that sensor **5** is selective towards hydrazine. This selectivity could be attributed to the stronger nucleophilicity of hydrazine compared to other tested nucleophiles²⁷. Kinetic studies indicate that the reaction between hydrazine and calixpyrrole **5** is pseudo-first order (Figure S14)




Figure 2: UV absorption of sensor **5** ($1 \cdot 10^{-5} \text{M}$, 25°C) upon addition of 100 equivalents of hydrazine in acetonitrile. Spectra acquired every 10 min. Inset: variation of absorbance at 373 nm




Figure 3: A) Variation of UV absorption at 373 nm of sensor **5** ($1 \cdot 10^{-5} \text{M}$, 25°C) upon addition of 100 equivalents of different competing nucleophiles in pure acetonitrile. B) UV spectra of sensor **5** alone, then 2h after addition of 100 equivalents of each competing nucleophiles and finally 2h after addition of 100 equivalent of hydrazine.

This great selectivity was also demonstrated at a higher concentration (Figure 4). Addition of 100 equivalents hydrazine to a 1 mM solution of sensor **5** lead to a complete bleaching after less than 90 seconds that can be perceived by naked eye.




Figure 4: Color change of sensor **5** (1 mM) in acetonitrile in the presence of 100 equivalent of nucleophiles. 2-mercaptoethanol (1), pyrrole (2), aniline (3), triethylamine (4), diethylamine (5), propylamine (6), ammonia (7), hydrazine (8), sensor **5** alone (9). Picture was taken 90s after hydrazine addition.

4) Study of the influence of water

The influence of water on the detection capacity of sensor **5** was evaluated by measuring the maximum absorbance of a solution of calixpyrrole at 10 μM in the presence of 100 equivalents of hydrazine with different percentages of water at 25 °C. The addition of water led to a slight red shift in the maximum absorption, reaching 376 nm for 40% of water. It can be observed on Figure 5 (and Figure S15-20) that the response decreases with an increasing amount of water. This can be explained by the lower nucleophilicity of hydrazine in water compared to acetonitrile.²⁸ The same type of response decrease can be observed to a lesser degree by adding a polar protic solvent such as ethanol in acetonitrile instead of water (Figure S21). This lower decrease could be explained by the fact that ethanol is a less effective hydrogen bond donor than water. Nevertheless, the detection capacity of sensor **5** in presence of water can be easily restored by enhancing the reaction rate by increasing the temperature to 50 °C (Figure 5). This finding would allow the use of sensor **5** for the detection of environmental samples on aqueous media.




Figure 5: Variation of UV absorption at maximum absorption of sensor **5** ($1 \cdot 10^{-5}\text{M}$, 25°C or 50°C) upon addition of 100 equivalents of hydrazine in acetonitrile containing increasing water content.

5) Analytical performance towards hydrazine

In order to evaluate the analytical performance of sensor **5**, calibration curves were constructed by monitoring the absorbance at 373 nm with increasing amounts of hydrazine (0 to 100 equiv.). To take into account the time dependency of the reaction between calixpyrrole and hydrazine, 7 individual solutions of compound **5** at a concentration of 10 μM were prepared and incubated in a water bath at 25 $^{\circ}\text{C}$. Then, an increasing amount of hydrazine was added to each solution with a time interval of 2 minutes in order to allow the operator to fill the UV cuvette and record the UV spectra. A series of solutions were analyzed after 1, 2, and 4 hours. The calibration curves are shown in Figure 6 A and Figure S22.

One can observe that good correlation coefficients were obtained after 1 and 2 hours. After 4h, high hydrazine concentrations (i.e. 80 and 100 equiv.) start to reach a plateau that derive from linearity and therefore do not allow accurate quantification of hydrazine. However, the first 5 points of the calibration curve remained aligned, with a good correlation coefficient. In our hand, the best result was achieved after 2 hours of incubation (Figure 6 A) with a correlation coefficient of 0.997. The same methodology was employed in an ACN-water 40% mixture with an incubation time of 2 hours at 50 $^{\circ}\text{C}$. A representative calibration curve is depicted on figure 6B, which shows a good correlation coefficient of 0.994.

The limit of detection (LOD) and limit of quantification (LOQ) were deduced from the calibration curves according to the following equations²⁹:

$$\text{LOD} = 3.3 \times \sigma / \text{slope}$$

$$\text{LOQ} = 10 \times \sigma / \text{slope}$$

with σ = the standard deviation of y-intercept of the regression line




Figure 6 : Calibration curves for hydrazine monitoring in the range 10-1000 μM obtained after 2h of incubation. A) in acetonitrile, 25 $^{\circ}\text{C}$ B) ACN-Water 40%, 50 $^{\circ}\text{C}$

LOD was evaluated at 4.10^{-5} M (1.3 mg/L) and LOQ at $1.2.10^{-4}$ M (3.9 mg/L) in 100% acetonitrile solution. In 40% water solution, LOD and LOQ were evaluated at 2.5 mg/L and 7.5 mg/L respectively.

Conclusion

In conclusion, we have successfully repurposed a β -dicyanovinyl-substituted calix[4]pyrrole chemodosimetric sensor initially designed for the sensing of cyanide anion in order to achieve the spectroscopic detection of hydrazine. It is one of the first time that chemodosimetric approach was used with calixpyrrole for the detection of a covalent analyte. Sensor **5** proved to be highly selective towards hydrazine among other competing aliphatic or aromatic amino

compounds (ammonia, propylamine, diethylamine, triethylamine, pyrrole, aniline) and 2-mercaptoethanol. The sensing capacities were evaluated by UV-spectroscopy in pure acetonitrile and in acetonitrile-water mixtures with up to 40% water. Calibration curves obtained after an incubation time of 2 hours at 25°C in pure acetonitrile or 2 hours at 50°C for ACN-water 40% mixture proved to be linear in the 10-1000 µM range. Limit of detection of sensor toward hydrazine were estimated at 4.10^{-5} M (1.3 mg/L) and $7.5.10^{-5}$ M (2.5 mg/L) in pure acetonitrile and ACN-water 40% mixture respectively. Finally, sensor **5** was employed for the naked-eye detection of hydrazine, which is the only compound studied that induce a total bleaching of the yellow solution. The chemodosimetric approach used in this study could be exploited in the future for the design of multitopic receptors based on calix[4]pyrroles thanks to the presence of a responsive side arm and the pyrrole cavity.

Acknowledgement

The authors are grateful to DST-SERB, New-Delhi for the financial support (Project File no. ECR/2017/000821), and Jamia Millia Islamia for providing the wonderful infrastructure. I.A.R also thanks CSIR, New-Delhi for the SRF fellowship award. Continuous financial support from the University of Littoral Côte d'Opale is also warmly acknowledged (P.-E.D & A.K)

Conflicts of interest

There are no conflicts to declare.

Notes and references :

- (1) Garrod, S.; Bollard, M. E.; Nicholls, A. W.; Connor, S. C.; Connelly, J.; Nicholson, J. K.; Holmes, E. Integrated Metabonomic Analysis of the Multiorgan Effects of Hydrazine Toxicity in the Rat. *Chem Res Toxicol* **2005**, *18* (2), 115–122. <https://doi.org/10.1021/tx0498915>.
- (2) Ameen, S.; Akhtar, M. S.; Shin, H. S. Hydrazine Chemical Sensing by Modified Electrode Based on in Situ Electrochemically Synthesized Polyaniline/Graphene Composite Thin Film. *Sensors and Actuators B: Chemical* **2012**, *173*, 177–183. <https://doi.org/10.1016/j.snb.2012.06.065>.
- (3) CDC - Immediately Dangerous to Life or Health Concentrations (IDLH): Hydrazine - NIOSH Publications and Products. <https://www.cdc.gov/niosh/idlh/302012.html> (accessed 2022-07-04).
- (4) Maiti, A.; Manna, S. K.; Halder, S.; Mandal, M.; Karak, A.; Banik, D.; Jana, K.; Mahapatra, A. K. A Benzothiazole-Based Dual Reaction Site Fluorescent Probe for the Selective Detection of Hydrazine in Water and Live Cells. *Org. Biomol. Chem.* **2022**. <https://doi.org/10.1039/D2OB00709F>.
- (5) Koizumi, A.; Nomiya, T.; Tsukada, M.; Wada, Y.; Omae, K.; Tanaka, S.; Miyauchi, H.; Imamiya, S.; Sakurai, H. Evidence on N-Acetyltransferase Allele-Associated Metabolism of Hydrazine in Japanese Workers. *Journal of Occupational and Environmental Medicine* **1998**, *40* (3), 217–222.
- (6) Isenberg, S. L.; Carter, M. D.; Crow, B. S.; Graham, L. A.; Johnson, D.; Beninato, N.; Steele, K.; Thomas, J. D.; Johnson, R. C. Quantification of Hydrazine in Human Urine

- by HPLC-MS/MS. *J Anal Toxicol* **2016**, *40* (4), 248–254. <https://doi.org/10.1093/jat/bkw015>.
- (7) Ensafi, A. A.; Chamjangali, M. A. Flow Injection Spectrophotometric Determination of Trace Amounts of Hydrazine by the Inhibition of the Pyrogallol Red-Iodate Reaction. *Journal of Analytical Chemistry* **2004**, *59* (2), 129–133. <https://doi.org/10.1023/B:JANC.0000014738.38853.c6>.
 - (8) Oh, J.-A.; Shin, H.-S. Simple and Sensitive Determination of Hydrazine in Drinking Water by Ultra-High-Performance Liquid Chromatography–Tandem Mass Spectrometry after Derivatization with Naphthalene-2,3-Dialdehyde. *Journal of Chromatography A* **2015**, *1395*, 73–78. <https://doi.org/10.1016/j.chroma.2015.03.051>.
 - (9) Christova, R.; Ivanova, M.; Novkirishka, M. Indirect Potentiometric Determination of Arsenite, Sulphite, Ascorbic Acid, Hydrazine and Hydroxylamine with an Iodide-Selective Electrode. *Analytica Chimica Acta* **1976**, *85* (2), 301–307. [https://doi.org/10.1016/S0003-2670\(01\)84695-X](https://doi.org/10.1016/S0003-2670(01)84695-X).
 - (10) Wang, M.; Wang, X.; Li, X.; Yang, Z.; Guo, Z.; Zhang, J.; Ma, J.; Wei, C. A Coumarin-Fused “off-on” Fluorescent Probe for Highly Selective Detection of Hydrazine. *Spectrochim Acta A Mol Biomol Spectrosc* **2020**, *230*, 118075. <https://doi.org/10.1016/j.saa.2020.118075>.
 - (11) Song, Y.; Chen, G.; Han, X.; You, J.; Yu, F. A Highly Sensitive Near-Infrared Ratiometric Fluorescent Probe for Imaging of Mitochondrial Hydrazine in Cells and in Mice Models. *Sensors and Actuators B: Chemical* **2019**, *286*, 69–76. <https://doi.org/10.1016/j.snb.2019.01.116>.
 - (12) Zhang, X.-Y.; Yang, Y.-S.; Wang, W.; Jiao, Q.-C.; Zhu, H.-L. Fluorescent Sensors for the Detection of Hydrazine in Environmental and Biological Systems: Recent Advances and Future Prospects. *Coordination Chemistry Reviews* **2020**, *417*, 213367. <https://doi.org/10.1016/j.ccr.2020.213367>.
 - (13) Roy, B.; Bandyopadhyay, S. The Design Strategies and Mechanisms of Fluorogenic and Chromogenic Probes for the Detection of Hydrazine. *Anal. Methods* **2018**, *10* (10), 1117–1139. <https://doi.org/10.1039/C7AY02866K>.
 - (14) Sasaki, Y.; Lyu, X.; Tang, W.; Wu, H.; Minami, T. Supramolecular Optical Sensor Arrays for On-Site Analytical Devices. *Journal of Photochemistry and Photobiology C: Photochemistry Reviews* **2022**, *51*, 100475. <https://doi.org/10.1016/j.jphotochemrev.2021.100475>.
 - (15) You, L.; Zha, D.; Anslyn, E. V. Recent Advances in Supramolecular Analytical Chemistry Using Optical Sensing. *Chem. Rev.* **2015**, *115* (15), 7840–7892. <https://doi.org/10.1021/cr5005524>.
 - (16) Rather, I. A.; Danjou, P.-E.; Ali, R. Aryl- and Superaryl-Extended Calix[4]Pyrroles: From Syntheses to Potential Applications. *Top Curr Chem (Z)* **2023**, *381* (1), 7. <https://doi.org/10.1007/s41061-022-00419-0>.
 - (17) Rather, I. A.; Wagay, S. A.; Hasnain, M. S.; Ali, R. New Dimensions in Calix[4]Pyrrole: The Land of Opportunity in Supramolecular Chemistry. *RSC Adv.* **2019**, *9* (66), 38309–38344. <https://doi.org/10.1039/C9RA07399J>.
 - (18) Kim, A.; Ali, R.; Park, S. H.; Kim, Y.-H.; Park, J. S. Probing and Evaluating Anion– π Interaction in Meso-Dinitrophenyl Functionalized Calix[4]Pyrrole Isomers. *Chem. Commun.* **2016**, *52* (74), 11139–11142. <https://doi.org/10.1039/C6CC04562F>.
 - (19) Lee, J. Y.; Root, H. D.; Ali, R.; An, W.; Lynch, V. M.; Bähring, S.; Kim, I. S.; Sessler, J. L.; Park, J. S. Ratiometric Turn-On Fluorophore Displacement Ensembles for Nitroaromatic Explosives Detection. *J. Am. Chem. Soc.* **2020**, *142* (46), 19579–19587. <https://doi.org/10.1021/jacs.0c08106>.
 - (20) Kaur, K.; Saini, R.; Kumar, A.; Luxami, V.; Kaur, N.; Singh, P.; Kumar, S. Chemodosimeters: An Approach for Detection and Estimation of Biologically and Medically Relevant Metal Ions, Anions and Thiols. *Coordination Chemistry Reviews* **2012**, *256* (17), 1992–2028. <https://doi.org/10.1016/j.ccr.2012.04.013>.
 - (21) Rather, I. A.; Sofi, F. A.; Bhat, M. A.; Ali, R. Synthesis of Novel One-Walled Meso-Phenylboronic Acid-Functionalized Calix[4]Pyrrole: A Highly Sensitive Electrochemical

- Sensor for Dopamine. *ACS Omega* **2022**, 7 (17), 15082–15089.
<https://doi.org/10.1021/acsomega.2c00926>.
- (22) Rather, I. A.; Ali, R. A Catalytic and Solvent-Free Approach for the Synthesis of Diverse Functionalized Dipyrromethanes (DPMs) and Calix[4]Pyrroles (C4Ps). *Green Chem.* **2021**, 23 (16), 5849–5855. <https://doi.org/10.1039/D1GC01515J>.
- (23) Anzenbacher, Pavel; Jursíková, K.; Shriver, J. A.; Miyaji, H.; Lynch, V. M.; Sessler, J. L.; Gale, P. A. Lithiation of Meso-Octamethylcalix[4]Pyrrole: A General Route to C-Rim Monosubstituted Calix[4]Pyrroles. *J. Org. Chem.* **2000**, 65 (22), 7641–7645.
<https://doi.org/10.1021/jo005610w>.
- (24) Hong, S.-J.; Yoo, J.; Kim, S.-H.; Kim, J. S.; Yoon, J.; Lee, C.-H. β -Vinyl Substituted Calix[4]Pyrrole as a Selective Ratiometric Sensor for Cyanide Anion. *Chem. Commun.* **2009**, No. 2, 189–191. <https://doi.org/10.1039/B815326D>.
- (25) Bu, L.; Rémond, M.; Colinet, P.; Jeanneau, E.; Le Bahers, T.; Chaput, F.; Andraud, C.; Bretonnière, Y. Sensitive 1,1-Dicyanovinyl Push-Pull Dye for Primary Amine Sensing in Solution by Fluorescence. *Dyes and Pigments* **2022**, 202, 110258.
<https://doi.org/10.1016/j.dyepig.2022.110258>.
- (26) Nishiyabu, R.; Anzenbacher, P. 1,3-Indane-Based Chromogenic Calixpyrroles with Push–Pull Chromophores: Synthesis and Anion Sensing. *Org. Lett.* **2006**, 8 (3), 359–362. <https://doi.org/10.1021/ol0521782>.
- (27) Hou, J.-T.; Wang, B.; Wang, S.; Wu, Y.; Liao, Y.-X.; Ren, W. X. Detection of Hydrazine via a Highly Selective Fluorescent Probe: A Case Study on the Reactivity of Cyano-Substituted CC Bond. *Dyes and Pigments* **2020**, 178, 108366.
<https://doi.org/10.1016/j.dyepig.2020.108366>.
- (28) Nigst, T. A.; Antipova, A.; Mayr, H. Nucleophilic Reactivities of Hydrazines and Amines: The Futile Search for the α -Effect in Hydrazine Reactivities. *J. Org. Chem.* **2012**, 77 (18), 8142–8155. <https://doi.org/10.1021/jo301497g>.
- (29) ICH-Guidelines Q2(R1), Validation of Analytical Procedures: Text and Methodology. <https://Database.Ich.Org/Sites/Default/Files/Q2%28R1%29%20Guideline.Pdf>.
<https://database.ich.org/sites/default/files/Q2%28R1%29%20Guideline.pdf>.