



## **Extraction of soluble proteins by bead milling from Tetraselmis chui in two different physiological states**

Hugo Mear, P. Gillon, I. Gifuni, L. Lavenant, A. Poidevin, Estelle Couallier

### **► To cite this version:**

Hugo Mear, P. Gillon, I. Gifuni, L. Lavenant, A. Poidevin, et al.. Extraction of soluble proteins by bead milling from *Tetraselmis chui* in two different physiological states. European Phycological Congress, Aug 2023, Brest, France. <hal-04211525>

**HAL Id: hal-04211525**

**<https://hal.science/hal-04211525v1>**

Submitted on 19 Sep 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

## **Extraction of soluble proteins by bead milling from *Tetraselmis chui* in two different physiological states**

Mear H.<sup>1,2\*</sup>, Gillon P.<sup>2</sup>, Gifuni I.<sup>1</sup>, Lavenant L.<sup>2</sup>, Poidevin A.<sup>2</sup>, Couallier E.<sup>2</sup>

\*Main speaker

<sup>1</sup>Algosource, France

<sup>2</sup> Nantes Université, Oniris, CNRS, GEPEA, UMR 6144, France

\*mear.hugo@laposte.net

*Tetraselmis* is an interesting microalga for food products because of its high protein content and surface productivity as well as the possibility to scale-up the culture easily. However, this specie is yet understudied compared to other algae such as *Chlorella*. Different batches of *Tetraselmis chui* were received from the same supplier, with different morphological properties, probably associated with a difference in physiology.

The batches were characterized biochemically and the rupturability was tested to characterize their behaviour to cell disruption, with one batch more fragile than the other. The bead milling of the two batches also showed striking differences in terms of cell disruption kinetic, as well as in dry matter repartition after centrifugation and in protein recovery yields within the supernatant. Nearly no differences in protein yield were observed after bead milling in comparison to freeze-thawing for the resistant batch, with a final yield of 11% of the initial proteins. However, a clear protein release was observed for the fragile batch along the milling residence time, with a final protein yield of 32%. The electrophoresis of the supernatants as a function of the milling showed very different protein profiles.

The results underlined that a difference in physiology, which may come from a different life cycle stage or a different cell cycle stage of *Tetraselmis chui* had a significant impact on the following biorefinery, affecting directly the cell disruption and the protein recovery yields. The valorization of the whole microalgae thus necessitates the finest coupling between upstream and downstream.