



HAL
open science

Response to PEEP in COVID-19 ARDS patients with and without extracorporeal membrane oxygenation. A multicenter case–control computed tomography study

Jean-Christophe Richard, Florian Sigaud, Maxime Gaillet, Maciej Orkisz, Sam Bayat, Emmanuel Roux, Touria Ahaouari, E.E. Dávila Serrano, Loic Bousset, Gilbert Ferretti, et al.

► To cite this version:

Jean-Christophe Richard, Florian Sigaud, Maxime Gaillet, Maciej Orkisz, Sam Bayat, et al.. Response to PEEP in COVID-19 ARDS patients with and without extracorporeal membrane oxygenation. A multicenter case–control computed tomography study. *Critical Care*, 2022, 26 (1), pp.195. 10.1186/s13054-022-04076-z . hal-04212435

HAL Id: hal-04212435

<https://hal.science/hal-04212435v1>

Submitted on 20 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RESEARCH

Open Access



Response to PEEP in COVID-19 ARDS patients with and without extracorporeal membrane oxygenation. A multicenter case–control computed tomography study

Jean-Christophe Richard^{1,2,3*}, Florian Sigaud⁴, Maxime Gaillet¹, Maciej Orkisz², Sam Bayat^{5,6}, Emmanuel Roux², Touria Ahaouari⁵, Eduardo Davila², Loic Bousset^{2,7}, Gilbert Ferretti⁸, Hodane Yonis¹, Mehdi Mezidi^{1,3}, William Danjou¹, Alwin Bazzani¹, Francois Dhelft^{1,3}, Laure Folliet¹, Mehdi Girard¹, Matteo Pozzi⁹, Nicolas Terzi^{4,10} and Laurent Bitker^{1,2,3}

Abstract

Background: PEEP selection in severe COVID-19 patients under extracorporeal membrane oxygenation (ECMO) is challenging as no study has assessed the alveolar recruitability in this setting. The aim of the study was to compare lung recruitability and the impact of PEEP on lung aeration in moderate and severe ARDS patients with or without ECMO, using computed tomography (CT).

Methods: We conducted a two-center prospective observational case–control study in adult COVID-19-related patients who had an indication for CT within 72 h of ARDS onset in non-ECMO patients or within 72 h after ECMO onset. Ninety-nine patients were included, of whom 24 had severe ARDS under ECMO, 59 severe ARDS without ECMO and 16 moderate ARDS.

Results: Non-inflated lung at PEEP 5 cmH₂O was significantly greater in ECMO than in non-ECMO patients. Recruitment induced by increasing PEEP from 5 to 15 cmH₂O was not significantly different between ECMO and non-ECMO patients, while PEEP-induced hyperinflation was significantly lower in the ECMO group and virtually nonexistent. The median [IQR] fraction of recruitable lung mass between PEEP 5 and 15 cmH₂O was 6 [4–10]%. Total superimposed pressure at PEEP 5 cmH₂O was significantly higher in ECMO patients and amounted to 12 [11–13] cmH₂O. The hyperinflation-to-recruitment ratio (i.e., a trade-off index of the adverse effects and benefits of PEEP) was significantly lower in ECMO patients and was lower than one in 23 (96%) ECMO patients, 41 (69%) severe non-ECMO patients and 8 (50%) moderate ARDS patients. Compliance of the aerated lung at PEEP 5 cmH₂O corrected for PEEP-induced recruitment ($C_{\text{BABY LUNG}}$) was significantly lower in ECMO patients than in non-ECMO patients and was linearly related to the logarithm of the hyperinflation-to-recruitment ratio.

Conclusions: Lung recruitability of COVID-19 pneumonia is not significantly different between ECMO and non-ECMO patients, with substantial interindividual variations. The balance between hyperinflation and recruitment

*Correspondence: j-christophe.richard@chu-lyon.fr

¹ Service de Médecine Intensive Réanimation, Hôpital de la Croix Rousse, Hospices Civils de Lyon, 103 Grande Rue de la Croix Rousse, 69004 Lyon, France

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

induced by PEEP increase from 5 to 15 cmH₂O appears favorable in virtually all ECMO patients, while this PEEP level is required to counteract compressive forces leading to lung collapse. $C_{\text{BABY LUNG}}$ is significantly lower in ECMO patients, independently of lung recruitability.

Keywords: ARDS, Computed tomography, ECMO, PEEP, COVID-19

Background

Lung recruitability of COVID-19 acute respiratory distress syndrome (ARDS) remains to date a matter of debate, as conflicting results have been reported using several techniques derived from change in respiratory mechanics and/or gas exchange in response to positive end-expiratory pressure (PEEP) increase [1–5]. Various factors may explain the heterogeneity of literature reports regarding recruitability of COVID-19 ARDS: time from ARDS onset, ARDS severity, reliability of these methods to compute recruitment, ventilatory management before evaluation, body mass index (BMI), among others. Furthermore, COVID-19 ARDS lungs may be intrinsically less recruitable than “classical” ARDS, as non-aerated lung regions in COVID-19 patients may represent alveolar spaces substituted by fibrosis, cellular debris and necrotic tissue rather than atelectasis [6].

Two small studies have been published so far using computed tomography (CT) in the context of COVID-19 ARDS and provided conflicting results [7, 8]. One study performed during the later phase of the disease identified most patients as being non-recruiters by PEEP [7], while median recruitability in the other study [8] performed within 72 h of ARDS onset was substantially higher than in non-COVID ARDS historical studies [9].

No study has assessed to date the recruitability of severe COVID-19 ARDS under veno-venous extracorporeal membrane oxygenation (ECMO). In non-COVID-19 ARDS patients, median lung recruitability was substantially higher in 47 patients under ECMO [10] than in the seminal study of Gattinoni performed on non-ECMO ARDS patients [9], suggesting that higher PEEP levels should be beneficial in ARDS patients under ECMO. Whether these results apply to COVID-19 ARDS patients under ECMO remains to date unknown, as well as optimal PEEP level in this subset of patients. We hypothesized that lung recruitability is influenced by time from ARDS onset and may be lower in ECMO patients cannulated more than 72 h after ARDS onset. Furthermore, we hypothesized that compliance of the baby lung (i.e., aerated lung at PEEP 5 cmH₂O) may explain poor lung recruitability, as it may be an early marker of fibrosis.

The primary aim of the study was to compare lung recruitability assessed with CT in moderate and severe ARDS patients with or without ECMO. The secondary

aim of the study was to identify the mechanisms underlying lung recruitability in COVID-19 ARDS.

Methods

Study design and setting

The study is a multicenter prospective observational case–control study, performed in 2 intensive care units located in university hospitals, and was conducted in accordance with the amended declaration of Helsinki. The study was approved by a local independent ethics committee (Comité Scientifique et Ethique des Hospices Civils de Lyon, 20_194) and complied to the STROBE criteria for observational studies [11]. Patients were enrolled between November 1, 2020, and June 16, 2021, in center#1 and between November 1, 2020, and December 31, 2021, in center#2. Consent for data utilization was sought from the patients or their representative, and follow-up lasted 90 days. The primary endpoint of the study was the amount of recruitable lung between PEEP 5 and 15 cmH₂O on CT.

Patients and protocol

Eligible participants were ARDS patients [12] aged 18 years or older, under invasive mechanical ventilation, who had a COVID-19 pneumonia with a positive SARS-CoV-2 reverse transcription polymerase chain reaction, and an indication for CT.

Exclusion criteria were ARDS onset > 72 h in non-ECMO patients, ECMO onset > 72 h, contra-indication to transport to the imaging facility (ratio of oxygen arterial partial pressure to inspired oxygen fraction ($\text{PaO}_2/\text{FiO}_2$) < 60 Torr, mean arterial pressure < 65 mmHg, or intracranial hypertension), chronic obstructive pulmonary disease, pneumothorax or bronchopleural fistula, previous inclusion in the present study, the presence of intrathoracic metallic devices, pregnancy, patient under a legal protective measure, and refusal to participate by patient and/or relative.

Non-ECMO patients were ventilated with a tidal volume (VT) of 4 to 6 mL.kg⁻¹ of predicted body weight (PBW) to keep plateau pressure ($P_{\text{Plat,rs}}$) below 30 cmH₂O, with recommendation to use a PEEP-FiO₂ table to adjust PEEP [13]. ECMO patients were ventilated with a VT of 1 mL.kg⁻¹ PBW, with PEEP adjusted to target a $P_{\text{Plat,rs}}$ approximating 20–22 cmH₂O.

Respiratory measurements and arterial blood gas analysis were performed at least 1 h after adjustment of ventilatory settings. Patients were then transferred to the imaging facility using a transport ventilator (MONNAL T60—Air Liquide Medical Systems, Antony, France) with unchanged ventilatory settings. The endotracheal tube was transiently occluded with a Kocher clamp during ventilator change to avoid derecruitment.

Data analysis

Total PEEP ($PEEP_{tot,rs}$) and $P_{Plat,rs}$ were measured at the end of 3-s end-expiratory and end-inspiratory pauses. Airway driving pressure (ΔP_{rs}) was computed as $P_{Plat,rs}$ minus $PEEP_{tot,rs}$. Elastance of the respiratory system was computed as ΔP_{rs} divided by VT.

Low-dose CT acquisitions were performed in the supine position with an iCT 256 Ingenuity CT (Philips Healthcare, Eindhoven, The Netherlands), or a GE Optima CT scan (GE Medical Systems, Milwaukee, USA) using the following settings: voltage 140 kVp, slice thickness 1 mm, and matrix size 512×512 . Field of view, pixel size, and mAs were adapted for each patient. Lung scanning was performed from apex to base during end-expiratory and end-inspiratory pauses at the PEEP level set by the clinician ($CT_{Expi-Inspi}$), and during end-expiratory pauses at PEEP 15 and 5 cmH₂O ($CT_{PEEP5-15}$). Lack of respiratory efforts during the pauses was visually checked on the ventilator pressure–time curves. Image reconstruction was performed using a smooth filter (kernel B). The lungs were interactively segmented with a CreaTools-based software [14], excluding pleural effusions, hilar and mediastinal structures. Segmented lung volumes were analyzed using MATLAB (MathWorks, Natick, MA).

Voxel tissue and gas fraction were computed as previously described [15]. Tissue and gas volumes were computed as the product of their respective fractions times voxel volume times number of voxels in segmented lung volume, respectively.

Lung parenchyma was classified into four compartments, according to CT number: non-inflated (density between +100 and –100 Hounsfield units (HU)), poorly inflated (density between –101 and –500 HU), normally inflated (density between –501 and –900 HU), and hyperinflated tissue (density ≤ -901 HU).

Total lung weight and weight of each compartment were estimated using lung tissue volume, assuming a tissue density of 1 g.mL⁻¹ [16].

VT was assessed on CT (VT_{CT}) by subtracting the volume of gas at end-inspiration and at end-expiration in segmented lungs.

The amount of recruitable lung between PEEP 5 and 15 cmH₂O ($\Delta PEEP_{5-15}$ -induced recruitment) was computed as the weight of the non-inflated compartment at

PEEP 5 cmH₂O minus its weight at PEEP 15 cmH₂O, and standardized to total lung weight.

Tidal recruitment of the non-inflated compartment was defined as the weight of the non-inflated compartment at end-expiration minus its weight at end-inspiration [17], and standardized to total lung weight.

Change in lung aerated volume induced by PEEP change from 5 to 15 cmH₂O ($PEEP_{volume}$) was computed as the difference in the total volume of gas within the lungs between PEEP 15 and 5 cmH₂O.

The hyperinflation-to-recruitment ratio was computed as the difference between hyperinflated compartment total volume at PEEP 15 minus its value at PEEP 5 cmH₂O, over the difference between non-inflated compartment total volume at PEEP 5 minus its value at PEEP 15 cmH₂O [8].

Tidal hyperinflation was computed as the volume of the hyperinflated compartment at end-inspiration minus its volume at end-expiration [17], and standardized to predicted body weight.

The total superimposed pressure in the most dorsal parts of the lung was computed as previously described [18].

The lung inhomogeneity extent was measured by comparing the inflation of neighboring lung regions as previously described [19, 20] and was defined as the percentage of lung volume presenting an inflation ratio of neighboring regions greater than 1.61 (i.e., the 95th percentile of a control population) [19].

We finally developed a method to estimate elastic properties of the already aerated lung at PEEP 5 cmH₂O with CT ($C_{BABY LUNG}$, Additional file 1). Classical computation of compliance between PEEP 5 and 15 cmH₂O (i.e., change in lung aerated volume divided by change in PEEP) overestimates $C_{BABY LUNG}$ as recruited alveoli account partly for the change in aeration. As recruitment assessed by CT is computed as the difference in non-aerated lung compartment weight between PEEP levels, a computation of recruited aerated volume ($Rec_{Aer vol}$) from recruited lung weight was performed using the methodology proposed by Paula and coworkers [21], assuming that recruitable alveoli would remain aerated at PEEP 5 cmH₂O and have equilibrated to a level of expansion equivalent to that of other already open alveoli at PEEP 5 [21]. $C_{BABY LUNG}$ between PEEP 5 and 15 cmH₂O was finally computed as: $(PEEP_{volume} - Rec_{Aer vol})/\Delta PEEP$ (i.e., 10 cmH₂O).

A quality control was performed on both couples of CT images ($CT_{Expi-Inspi}$, $CT_{PEEP5-15}$). Images couples with segmented lung weight differing by more than 5% were excluded. $CT_{Expi-Inspi}$ in which VT_{CT} differed from VT set on the ventilator by more than 60 mL were also excluded (Additional file 2).

Statistical analysis

Statistical analysis was performed using R version 4.1.1 [22] with packages multcomp [23], lme4 [24], lmerTest [25], and interactions [26]. A p value ≤ 0.05 was chosen for statistical significance.

Data were expressed as count (percentage) or median [interquartile range] and compared between groups with the Fisher's exact test for categorical variables and ANOVA for continuous variables. Multiple comparisons between groups were made using the Holm–Sidak procedure. Comparisons of variables involving the same individual were made with linear mixed models, using patient as a random effect.

Multivariate analyses were performed using linear models, by incorporating variables with p values < 0.2 in univariate analysis and stepwise backward selection.

Estimation of sample size was not computed as the study is exploratory, and data collection stopped with the control of COVID-19 fifth wave in our geographic area.

Missing data were not imputed owing to the low rate of missingness for variables included in the multivariate models (Additional file 3).

Results

Flowchart of the study is presented in Additional file 2. Ninety-nine patients were included, of whom 24 were severe ARDS under ECMO, 59 severe ARDS without ECMO and 16 moderate ARDS. Three non-ECMO patients with delay between ARDS onset > 72 h (4 to 6 days after ARDS onset) were erroneously included and remained in the analysis. Patients' characteristics at inclusion are reported in Table 1, and respiratory mechanics and blood gas data are presented in Table 2.

Lung CT parameters at PEEP 5 cmH₂O

CT images obtained in 4 representative ECMO and non-ECMO patients are provided in Fig. 1. Lung weight assessed on CT was not significantly different between groups (Fig. 2). Non-inflated lung at PEEP 5 cmH₂O was significantly greater in ECMO than in non-ECMO patients (Fig. 2), and in severe ARDS non-ECMO patients vs. moderate ARDS patients. End-expiratory aerated lung volume (EELV) at PEEP 5 cmH₂O was significantly lower in ECMO patients. Total superimposed pressure at PEEP 5 cmH₂O (Fig. 3, left panel) was significantly higher

Table 1 Patient characteristics

Variables	Whole dataset (n = 99)	Moderate ARDS without ECMO (n = 16)	Severe ARDS without ECMO (n = 59)	Severe ARDS on ECMO (n = 24)
Sex male—no. (%)	75 (76%)	14 (88%)	42 (71%)	19 (79%)
Age—yr	62 [54–71]	61 [55–70]	66 [57–73]	57 [50–61] ^b
BMI—kg.m ⁻²	30 [26–37]	29 [25–32]	30 [26–37]	32 [27–39]
Delay between hospital admission and CT—day	5 [3–8]	5 [3–6]	4 [3–7]	10 [6–12] ^b
Delay between ICU admission and CT—day	3 [1–6]	3 [1–5]	2 [1–3]	7 [5–8] ^{a,b}
Delay between ARDS onset and CT—day	1 [1–3]	1 [1, 2]	1 [1, 2]	5 [3–7] ^{a,b}
Delay between IMV onset and CT—day	1 [1–3]	1 [1, 2]	1 [1, 2]	5 [3–7] ^{a,b}
SAPS 2 at ICU admission	39 [30–45]	41 [28–45]	40 [31–47]	36 [30–42]
SOFA score at inclusion	7 [5–8]	6 [5–7]	7 [4–8]	7 [5–9]
Prone position in the 24 h preceding CT—no. (%)	81 (82%)	14 (88%)	46 (78%)	21 (88%)
iNO in the 24 h preceding CT—no. (%)	19 (19%)	0 (0%)	12 (20%)	7 (29%)
NMBA in the 24 h preceding CT—no. (%)	96 (96%)	15 (94%)	57 (97%)	24 (100%)
RRT in the 24 h preceding CT—no. (%)	4 (4%)	1 (6%)	2 (3%)	1 (4%)
Inotropes in the 24 h preceding CT—no. (%)	1 (1%)	0 (0%)	1 (2%)	0 (0%)
Vasopressor in the 24 h preceding CT—no. (%)	67 (68%)	13 (81%)	37 (63%)	17 (71%)
ICU mortality—no. (%) ^c	49 (49%)	3 (19%)	32 (54%)	14 (58%)
Day-90 mortality—no. (%) ^c	48 (49%)	3 (19%)	32 (55%)	13 (54%)
Ventilator-free days at day 90—day	0 [0–71]	59 [13–83]	0 [0–70]	0 [0–56]
ICU length of stay—day	25 [16–40]	24 [13–45]	23 [16–36]	36 [21–48]
Hospital length of stay—day	37 [23–58]	44 [28–78]	35 [21–49]	41 [29–59]

Values are median [1st quartile–3rd quartile] or count (percentage)

^a $p < 0.05$ vs. moderate ARDS without ECMO, ^b $p < 0.05$ vs. severe ARDS without ECMO, ^c $p < 0.05$ between groups

ARDS acute respiratory distress syndrome, BMI body mass index, CT computed tomography, ECMO extracorporeal membrane oxygenation, ICU intensive care unit, IMV invasive mechanical ventilation, iNO inhaled nitric oxide, NMBA neuromuscular blocking agents, no number, RRT renal replacement therapy, SAPS2 Simplified Acute Physiology Score

Table 2 Respiratory mechanics and arterial blood gas

Variables	Whole dataset (n = 99)	Moderate ARDS without ECMO (n = 16)	Severe ARDS without ECMO (n = 59)	Severe ARDS on ECMO (n = 24)
PEEP—cmH ₂ O	10 [5–14]	9 [5–10]	10 [5–10]	15 [13–15] ^{ab}
VT—mL.kg ⁻¹ PBW	5.9 [3.9–6.0]	6.0 [5.9–6.0]	6.0 [5.9–6.0]	1.0 [1.0–1.0] ^{ab}
RR—min ⁻¹	22 [16–28]	23 [20–28]	25 [22–29]	5 [5–10] ^{ab}
PEEP _{tot,rs} —cmH ₂ O	10 [7–15]	9 [6–10]	10 [6–11]	15 [14–15] ^{ab}
P _{plat,rs} —cmH ₂ O	20 [17–22]	18 [17–24]	20 [16–23]	20 [19–21]
P _{peak} —cmH ₂ O	29 [25–35]	30 [28–34]	32 [27–37]	22 [21–24] ^{ab}
ΔP _{rs} —cmH ₂ O	9 [7–12]	10 [9–13]	10 [9–12]	5 [3–8] ^{ab}
E _{rs} —cmH ₂ O. L ⁻¹	33 [25–44]	29 [22–37]	28 [23–35]	70 [53–95] ^{ab}
pH	7.38 [7.32–7.44]	7.37 [7.33–7.42]	7.37 [7.31–7.41]	7.46 [7.39–7.47] ^{ab}
PaO ₂ —Torr	72 [65–80]	78 [67–89]	74 [66–80]	69 [62–74] ^a
FiO ₂ or FmO ₂ – %	90 [60–100]	50 [44–53]	90 [65–100] ^a	100 [80–100] ^a
PaCO ₂ —Torr	49 [41–54]	42 [37–51]	50 [43–58]	49 [41–51]
Bicarbonates—mmol.L ⁻¹	28 [24–32]	26 [22–28]	27 [24–31]	32 [29–34] ^{ab}
Lactate—mmol.L ⁻¹	1.7 [1.3–2.2]	1.5 [1.3–1.9]	1.6 [1.3–2.2]	1.9 [1.6–2.4]

Values are median [1st quartile–3rd quartile]. ^a $p < 0.05$ vs. moderate ARDS without ECMO. ^b $p < 0.05$ vs. severe ARDS without ECMO

ARDS acute respiratory distress syndrome, ΔP_{rs} driving pressure of the respiratory system, ECMO extracorporeal membrane oxygenation, E_{rs} elastance of the respiratory system, FiO₂ inspired fraction of oxygen, FmO₂ ECMO membrane oxygen fraction, PaCO₂ carbon dioxide partial pressure in arterial blood, PaO₂ oxygen partial pressure in arterial blood, PBW predicted body weight, PEEP positive end-expiratory pressure, PEEP_{tot,rs} total PEEP of the respiratory system, P_{peak} peak airway pressure, P_{plat,rs} plateau pressure of the respiratory system, RR respiratory rate, VT tidal volume

in ECMO patients (12 [11–13] cmH₂O), as compared to both moderate (10 [8–11] cmH₂O) and severe non-ECMO patients (9 [7–13] cmH₂O).

(Additional file 5 and 6). ΔPEEP₅₋₁₅-induced recruitment significantly decreased over time from ARDS onset in non-ECMO patients, but not in ECMO patients (Fig. 5).

Lung recruitability

Left panel of Fig. 4 shows the frequency distribution of patients according to lung recruitability between PEEP 5 and 15 cmH₂O. The median amount of recruitable lung between PEEP 5 and 15 cmH₂O was 6 [4–10]%, and was not significantly different between groups. PEEP_{volume} was significantly lower in ECMO patients, while ΔPEEP₅₋₁₅-induced hyperinflation was significantly lower in the ECMO group and virtually inexistent (Fig. 2). A sensitivity analysis excluding the 3 patients with exclusion criteria yielded identical results (Additional file 4). The hyperinflation-to-recruitment ratio (a trade-off index of the adverse effects and benefits of PEEP increase from 5 to 15 cmH₂O) was significantly lower in ECMO patients, as compared to both severe non-ECMO ARDS patients and moderate ARDS patients (Fig. 3, right panel). This ratio was lower than 1 in 8 (50%) moderate ARDS patients, 41 (69%) severe non-ECMO patients, and in 23 (96%) ECMO patients, i.e., virtually all ECMO patients responded to PEEP 15 cmH₂O by recruited volume outweighing hyperinflation volume.

Multivariate analysis identified poorly inflated lung at PEEP5, EELV at PEEP5 and the interaction of ECMO status and delay between ARDS onset and CT as independent predictors of ΔPEEP₅₋₁₅-induced recruitment

Tidal hyperinflation and tidal recruitment

The frequency distribution of patients according to tidal hyperinflation is reported in Fig. 4 (right panel). Tidal hyperinflation amounted to 0.3 [0.1–1.0] mL.kg⁻¹ PBW. Tidal hyperinflation greater than 1 mL.kg⁻¹ PBW was observed in 19 (25%) non-ECMO patients and 0 (0%) ECMO patients. As expected, tidal hyperinflation was significantly lower in ECMO patients, while tidal recruitment was significantly higher in non-ECMO severe ARDS patients (Fig. 2).

Lung inhomogeneity

The extent of lung inhomogeneity decreased significantly from PEEP 5 to PEEP 15 cmH₂O in non-ECMO patients, but not in ECMO patients (Fig. 6), with substantial interindividual variations. Lung inhomogeneity extent decreased between PEEP5 and 15 cmH₂O in 94% of moderate ARDS patients, 97% of severe ARDS patients, and 62% of severe ECMO patients.

Compliance of the aerated lung at PEEP 5 cmH₂O

C_{BABY LUNG} was significantly lower in ECMO patients than in both severe ARDS patients without ECMO and




Fig. 1 CT images acquired at the mid-chest level in 4 representative patients. ECMO patients are presented in the uppermost quadrants, while non-ECMO patients are presented in the lowermost quadrants. Patients with ΔPEEP_{5-15} -induced recruitment below the median value of the whole population (deemed as low recruiters) are presented in the leftmost quadrants, while patients with PEEP_{5-15} -induced recruitment above the median value of the whole population (deemed as high recruiters) are presented in the rightmost quadrants. In each quadrant, the upper two images were acquired at PEEP 5 cmH_2O (one raw CT image on the left and one quantitative parametric CT on the right), and the lower two were acquired at PEEP 15 cmH_2O . The color code used for parametric images is provided in the figure legend. CT, computed tomography, ΔPEEP_{5-15} -induced recruitment, amount of recruitable lung between PEEP 5 and 15 cmH_2O , ECMO, extracorporeal membrane oxygenation, PEEP, positive end-expiratory pressure



moderate ARDS (Fig. 7, left panel). $C_{\text{BABY LUNG}}$ was linearly related to the logarithm of the hyperinflation-to-recruitment ratio (Fig. 7, right panel).


Multivariate analysis identified BMI, ECMO, and EELV at PEEP 5 cmH_2O as independent variables significantly associated with $C_{\text{BABY LUNG}}$ (Additional file 7), while $C_{\text{BABY LUNG}}$ was not independently associated with lung recruitability (Additional file 5 and 6).

Discussion

The main findings of the study are the following: 1—lung recruitability between PEEP 5 and 15 cmH_2O was not significantly different between categories of ARDS severity, 2—the trade-off between hyperinflation and recruitment induced by a PEEP change from 5 to 15 cmH_2O was favorable in virtually all ECMO patients, 3—compliance of the aerated lung at PEEP 5 was significantly lower in ECMO patients, independently of lung recruitability, and may be a protective factor against PEEP-induced hyperinflation under ECMO.

Patient characteristics at inclusion were similar in the present study and in the largest French epidemiological study on critically ill adults with COVID-19 [27]. ECMO patients were older in the present study than in the ELSO [28] and the Paris–Sorbonne University Hospital Network [29] registries, but with similar BMI. However, PEEP level under ECMO was higher (15 vs 10 and 12 cmH_2O , respectively), while VT, ΔP_{rs} , and $P_{\text{plat,rs}}$ were substantially lower in the present study [28, 29]. In non-ECMO-patients, the amount of non-aerated lung at PEEP 5 cmH_2O was similar in the present study and in 3 previous studies on COVID-19 patients under mechanical ventilation (approximately 40%) [5, 7, 8]. On the other hand, median lung recruitability between PEEP 5 and 15 cmH_2O was substantially lower in the present study as compared to the study by Protti et al. (6% vs. 11% of lung weight) [8], with a relatively similar CT protocol and delay between ARDS onset and CT. In addition to differences between pre-intubation management and case-mix, this may be explained by CT acquisition at higher PEEP levels (10 and 20 cmH_2O , respectively) in some





obese patients in the latter study. In ECMO patients, the amount of non-aerated lung at PEEP 5 cmH₂O was similar in our study and in non-COVID-19 ECMO patients (approximately 80%) [10].

An important finding of our study is that the delay from ARDS onset is an important confounding factor of lung recruitability and that lung recruitability decreases as early as during the first 3 days after ARDS onset. Surprisingly, this effect was not identified in ECMO patients and the following potential reasons may be hypothesized to explain this finding: patient selection for ECMO eligibility, lower sample size of the ECMO subgroup, beneficial impact of fluid administration required to maintain ECMO flow on lung recruitability, or preventive effect of ECMO on fibrotic changes potentially involved in the progressive loss of recruitability observed under standard ventilation [30], although this remains speculative.

To date, there is no undisputable threshold defining low or high recruiter patients with CT. Most studies defined high recruiters as patients with an amount of recruited lung between 5 and 45 cmH₂O above the median value of the population [9, 10]. As approximately 50% of the recruitment between PEEP 5 and 45 cmH₂O is achieved between PEEP 5 and 15 cmH₂O [8, 9], and as the threshold defining high recruiter amounted to 10% of total lung weight in the seminal study by Gattinoni and coworkers [9], it can be speculated that the 6% median lung

recruitment value identified in our study suggests that COVID-19 and non-COVID-19 ARDS are roughly similar regarding lung recruitability. Remarkably, recruitability was similar among categories of ARDS severity in our study, despite a stepwise increase in non-inflated lung compartment as severity increases, in striking contrast with Gattinoni's study [9], suggesting that the underlying mechanisms responsible for lung consolidation are different between COVID-19 and non-COVID-19 ARDS. However, we failed to relate decreased compliance of the aerated lung to poor lung recruitability, suggesting that the mechanical characteristics of the aerated and non-aerated lung may be unrelated.

Another striking result of the present study is that the balance between hyperinflation and recruitment induced by a PEEP increase from 5 to 15 cmH₂O is favorable in virtually all ECMO patients, as opposed to COVID-19 ARDS patients without ECMO. Interestingly, most of the patients included in large ECMO cohort studies on COVID-19 ARDS were ventilated with substantially lower PEEP levels [28, 29]. Our results suggests that the lower compliance of the aerated lung at PEEP 5 cmH₂O during ECMO (a consequence of lower EELV) may prevent occurrence of PEEP-induced hyperinflation (Figs. 2 and 7). Hyperinflation may have nonetheless been undetected in ECMO patients as it may occur at a level below the resolution of CT, or since the decrease in CT density




Fig. 5 Relationship between ΔPEEP_{5-15} -induced recruitment and delay between ARDS onset and CT, according to ECMO status. Each point is partial residuals of the multivariate model with ΔPEEP_{5-15} -induced recruitment as the dependent variable, and the following independent variables: EELV at PEEP 5 cmH₂O, poorly inflated lung at PEEP5 and the interaction between ECMO status x delay between CT and ARDS onset. Lines are regression lines according to ECMO status. Shaded areas are 95% confidence interval for each regression line. ARDS, acute respiratory distress syndrome, CT, computed tomography, ECMO, extracorporeal membrane oxygenation, ΔPEEP_{5-15} -induced recruitment, amount of recruitable lung between PEEP 5 and 15 cmH₂O, EELV, end-expiratory lung volume, NS, non-statistically significant, PEEP, positive end-expiratory pressure





Fig. 6 Lung inhomogeneities and PEEP. The extent of lung inhomogeneity of the individual patients with moderate ARDS (left panel), severe ARDS without ECMO (middle panel), and severe ARDS with ECMO (right panel) is reported at PEEP 5 (red circles) and 15 cmH₂O (blue circles) * $p < 0.05$ between PEEP 5 and 15 cmH₂O of the same category of severity. Lung inhomogeneities are registered to the following patent: WO 2013/088336, and agreement to use this patent for research purposes was obtained from the owner (Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico). ARDS, acute respiratory distress syndrome, ECMO, extracorporeal membrane oxygenation, MODERATE, moderate ARDS, PEEP, positive end-expiratory pressure, SEVERE, severe ARDS without ECMO, SEVERE ECMO, severe ARDS under ECMO.



due to hyperinflation can be masked by the increased tissue mass in severe ARDS [31]. Hyperinflation was hence not detected during PEEP increase from 5 to 45 cmH₂O in a previous study on ECMO patients [10]. Furthermore, we cannot exclude that overdistension without hyperinflation occurred at the interface between non-aerated and aerated lung units in some patients (see below) [19, 20]. Finally, the impact of reduced lung compliance on overdistension occurrence during PEEP increase in ECMO patients remains unknown.

A beneficial impact of PEEP increase on lung inhomogeneity extent was only identified in non-ECMO patients. As lung inhomogeneities act as stress raisers, this suggests that the energy load was more evenly distributed within the lung parenchyma at PEEP 15 in this subgroup of patients. This effect was not observed in ECMO patients, although with substantial interindividual variations, as was previously observed in non-COVID ARDS [20]. Moreover, PEEP increase was expected to be detrimental on stress raisers in 38% of ECMO patients, favoring individualization of PEEP settings.

Some limitations of the present study should be acknowledged. First, potentially recruitable lung at PEEP 45 cmH₂O was not assessed, as this PEEP level was deemed excessive by our team [32] and others in COVID-19 ARDS [7] and higher PEEP levels may have led to different results. ECMO patients were studied a median of 4 days later than non-ECMO patients, and this delay may have impacted lung recruitability under

ECMO, although this effect was ruled out by our multivariate analysis (Additional file 6, Fig. 5). Occurrence of complete airway closure [1] was not assessed in the present study, and this effect may have biased measurement of $C_{\text{BABY LUNG}}$. Approximately 70% of the screened population (i.e., patients with COVID-19 pneumonia under invasive mechanical ventilation) lacked eligibility criteria, mostly because a CT was performed before intensive care admission, and the study population may be a biased subset of the initial population. Variable selection for multivariate analysis was performed on the basis of bivariable association, and this may have increased the risks of detecting spurious statistical associations [33]. Finally, the observational design of the study precludes any firm conclusion to be drawn regarding optimal ventilation settings in ARDS COVID-19 patients.

Nevertheless, the study has the following strengths. Lung evaluation was performed using CT, i.e., the reference method to perform a quantitative analysis of lung aeration and recruitment [15, 31, 34]. External validity of the study is expected to be good owing to the low rate of screen failure despite the context of pandemics, the substantial sample size for a CT study, and the multicenter design. Furthermore, the study is the first to assess lung recruitability in both ECMO and non-ECMO patients, in a homogeneous time frame (i.e., at the early phase of ARDS), thus minimizing potential confounding effects related to ventilator-induced lung injury or ventilator-associated pneumonia.

According to our results, a PEEP level around 13–15 cmH₂O is required to overcome the superimposed hydrostatic pressure and prevent alveolar collapse in the majority of COVID-19 ARDS patients under ECMO. Higher PEEP levels may be required in patients with elevated chest wall to respiratory system elastance ratio, such as superobese patients, as part of this pressure level may be dissipated in the chest wall. In addition, a PEEP level of 15 cmH₂O is associated with a favorable balance ratio between hyperinflation and recruitment induced by PEEP in virtually all COVID-19 ECMO patients. However, this PEEP level may be detrimental on stress raisers in a subset of ECMO patients favoring individualization of PEEP setting. Furthermore, the rationale for high PEEP use in patients under ECMO may be questionable as it relies on a putative protective effect on derecruitment related to the use of ultra-low tidal volume, while it may overdistend the lung, alter hemodynamics or increase the mechanical power transmitted to the lung from the ventilator. Finally, lung recruitability should be reassessed early in the course of COVID-19 ARDS without ECMO, as it decreases sharply over time.

Conclusion

Lung recruitability of COVID-19 pneumonia is not significantly different among categories of ARDS severity, with substantial interindividual variations favoring individualization of PEEP setting. The balance between hyperinflation and recruitment induced by PEEP increase from 5 to 15 cmH₂O appears to be favorable in virtually all ECMO patients, although lung inhomogeneities acting as stress raisers were not significantly improved by PEEP increase in this group of patients. A PEEP level of 12–15 cmH₂O is required to counteract compressive forces leading to lung collapse in most ECMO patients. In non-ECMO patients, lung recruitability decreases steadily during the first 3 days after ARDS onset and justifies reevaluation of the PEEP setting on a daily basis.

Abbreviations

ARDS: Acute respiratory distress syndrome; BMI: Body mass index; $C_{\text{BABY LUNG}}$: Compliance of the aerated lung at PEEP 5 cmH₂O, corrected for PEEP-induced recruited aerated volume; CT: Computed tomography; $CT_{\text{ExpI-Inspi}}$: CT acquired during end-expiratory and end-inspiratory pauses at the PEEP level set by the clinician; $CT_{\text{PEEP5-15}}$: CT acquired during end-expiratory pauses at PEEP 5 and 15 cmH₂O; ΔPEEP_{5-15} : Change in PEEP from 5 to 15 cmH₂O; ΔP_{rs} : Airway driving pressure; ECMO: Extracorporeal membrane oxygenation; EELV: End-expiratory aerated lung volume; FiO_2 : Inspired oxygen fraction; HU: Hounsfield units; PaO_2/FiO_2 : Ratio of oxygen arterial partial pressure to inspired oxygen fraction; PBW: Predicted body weight; PEEP: Positive end-expiratory pressure; $PEEP_{\text{tot,rs}}$: Total PEEP of the respiratory system; $PEEP_{\text{volume}}$: Change in lung aerated volume induced by PEEP change from 5 to 15 cmH₂O; $P_{\text{plat,rs}}$: Plateau pressure of the respiratory system; $Rec_{\text{Aer Vol}}$: Recruited aerated volume between PEEP5 and 15 cmH₂O; VT: Tidal volume; VT_{CT} : Tidal volume was assessed on CT.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13054-022-04076-z>.

Additional file 1: Computation of compliance of the already aerated lung at PEEP 5 ($C_{\text{BABY LUNG}}$)

Additional file 2: Flow chart of the study

Additional file 3: Missing data per variable

Additional file 4: Sensitivity analysis

Additional file 5: Univariate analysis of variables associated with lung recruitment

Additional file 6: Multivariate analysis of variables associated with PEEP-induced lung recruitment

Additional file 7: Univariate and multivariate analyses of variables associated with $C_{\text{BABY LUNG}}$

Acknowledgements

The authors wish to thank Loredana Baboi for her help with inclusion of the patients in the study and data acquisition. The authors also wish to thank the following residents for their help with patient transport to the imaging facility: Hugo Dumargne, Claire Guerin, Mathilde Legros, Anaëlle Romagny, Hippolyte Lequain, Maxime Billon, Alexandre Pinede, Bastien Berger, Mathieu Chivot, Alexandre Berland, Thomas Foddis, Mathieu Sadoul, Gabriel Sarciron, Amaury Gausiome, Ali Chour, Vincent Alcazer, Martin Deslais, Caroline Abou-Zeid, Amaury Berrou, Camille Chaffard, Victor Dechaene, Florian Genty, Malik Haoutar, Julien Perrone, Benjamin Pequignot, and Louis Auffray. The authors wish to thank the intensive care unit nurses for their help with patients transport to the imaging facility, and patients and their relatives for agreeing to participate in the study. The authors wish to thank the radiology technicians for their help in organizing and acquiring the CT scans

Author contributions

JCR and LBitker made substantial contributions to study design, to data acquisition, study analysis, and interpretation of data, drafted the manuscript, approved the version to be published, and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. FS, MO, SB, ER, LBoussel, and NT made substantial contributions to study design, data acquisition, study analysis, and interpretation of data, revised the manuscript critically for important intellectual content, approved the version to be published, and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. MGaillet, TA, GF, HY, MM, WD, AB, FD, LF, and MGirard made substantial contributions to data acquisition, study analysis, and interpretation of data, revised the manuscript critically for important intellectual content, approved the version to be published, and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. ED made substantial contributions to study analysis and interpretation of data, revised the manuscript critically for important intellectual content, approved the version to be published, and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. MP made substantial contributions to interpretation of data, revised the manuscript critically for important intellectual content, approved the version to be published, and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. All authors have read and approved the final manuscript.

Funding

The study was funded by the Hospices Civils de Lyon. The funding body did not take part in the design of the study, nor in data collection, analysis, or interpretation, and was not involved in writing the manuscript.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

The authors state that the study has been approved by a local independent ethics committee (Comité Scientifique et Ethique des Hospices Civils de Lyon, 20_194). Consent for data utilization was sought from the patients or their representative.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Service de Médecine Intensive Réanimation, Hôpital de la Croix Rousse, Hospices Civils de Lyon, 103 Grande Rue de la Croix Rousse, 69004 Lyon, France. ²INSA-Lyon, CNRS, INSERM, CREATIS UMR 5220, Univ Lyon, Université Claude Bernard Lyon 1, U1294 Villeurbanne, France. ³Université de Lyon, Université Claude Bernard Lyon 1, Villeurbanne, France. ⁴Service de Médecine-Intensive Réanimation, CHU Grenoble-Alpes, Grenoble, France. ⁵Synchrotron Radiation for Biomedicine Laboratory (STROBE), INSERM UA07, Univ. Grenoble Alpes, Grenoble, France. ⁶Department of Pulmonology and Physiology, Grenoble University Hospital, Grenoble, France. ⁷Service de Radiologie, Hôpital De La Croix Rousse, Hospices Civils de Lyon, Lyon, France. ⁸Grenoble, France Service de Radiologie Diagnostique Et Interventionnelle, Université Grenoble Alpes, CHU Grenoble-Alpes, Grenoble, France. ⁹Service de Chirurgie Cardiaque, Hôpital Louis Pradel, Hospices Civils de Lyon, Lyon, France. ¹⁰INSERM, U1042, University Grenoble Alpes, HP2 Grenoble, France.

Received: 28 April 2022 Accepted: 27 June 2022

Published online: 02 July 2022

References

- Chen L, Del Sorbo L, Grieco DL, Junhasavasdikul D, Rittayamai N, Soliman I, et al. Potential for lung recruitment estimated by the recruitment-to-inflation ratio in acute respiratory distress syndrome. *Am J Respir Crit Care Med.* 2019;201(2):178.
- Pan C, Chen L, Lu C, Zhang W, Xia J-A, Sklar MC, et al. Lung recruitability in SARS-CoV-2 associated acute respiratory distress syndrome: a single-center, observational study. *Am J Respir Crit Care Med.* 2020;201:1294–7.
- Sang L, Zheng X, Zhao Z, Zhong M, Jiang L, Huang Y, et al. Lung recruitment, individualized PEEP, and prone position ventilation for COVID-19-associated severe ARDS: A single center observational study. *Front Med (Lausanne).* 2020;7: 603943.
- Beloncle FM, Pavlovsky B, Desprez C, Fage N, Olivier P-Y, Asfar P, et al. Recruitability and effect of PEEP in SARS-Cov-2-associated acute respiratory distress syndrome. *Ann Intensive Care.* 2020;10:55.
- Chiumello D, Busana M, Coppola S, Romitti F, Formenti P, Bonifazi M, et al. Physiological and quantitative CT-scan characterization of COVID-19 and typical ARDS: a matched cohort study. *Intensive Care Med.* 2020;46(12):2187.
- Nunes MC, Hale MJ, Mahtab S, Mabena FC, Dlodlu N, Baillie VL, et al. Clinical characteristics and histopathology of COVID-19 related deaths in South African adults. *PLoS ONE.* 2022;17: e0262179.
- Ball L, Robba C, Maiello L, Herrmann J, Gerard SE, Xin Y, et al. Computed tomography assessment of PEEP-induced alveolar recruitment in patients with severe COVID-19 pneumonia. *Crit Care.* 2021;25:81.
- Protti A, Santini A, Pennati F, Chiurazzi C, Cressoni M, Ferrari M, et al. Lung response to a higher positive end-expiratory pressure in mechanically ventilated patients with COVID-19. *Chest.* 2022;161:979–88.
- Gattinoni L, Caironi P, Cressoni M, Chiumello D, Ranieri VM, Quintel M, et al. Lung recruitment in patients with the acute respiratory distress syndrome. *N Engl J Med.* 2006;354:1775–86.
- Camporota L, Caricola EV, Bartolomeo N, Di Mussi R, Wyncoll DLA, Meadows CIS, et al. Lung recruitability in severe acute respiratory distress syndrome requiring extracorporeal membrane oxygenation. *Crit Care Med.* 2019;47:1177–83.
- von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, et al. The strengthening of reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet.* 2007;370:1453–7.
- Ranieri VM, Rubenfeld GD, Thompson BT, Ferguson ND, Caldwell E, et al. Acute respiratory distress syndrome: the Berlin definition. *JAMA.* 2012;307(23):2526–33.
- Brower RG, Matthay MA, Morris A, Schoenfeld D, Thompson BT, et al. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med.* 2000;342:1301–8.
- Davila Serrano E, Dhelft F, Bitker L, Richard JC, Orkisz M. Software for CT-image analysis to assist the choice of mechanical-ventilation settings in acute respiratory distress syndrome. Warsaw, Poland; Leszek J Chmielewski, Ryszard Kozera, Arkadiusz Orłowski Eds., Springer, 2020. p. 48–58.
- Gattinoni L, Pesenti A, Avalli L, Rossi F, Bombino M. Pressure-volume curve of total respiratory system in acute respiratory failure. computed tomographic scan study. *Am Rev Respir Dis.* 1987;136(3):730–6.
- Gattinoni L, Pesenti A, Bombino M, Baglioni S, Rivolta M, Rossi F, et al. Relationships between lung computed tomographic density, gas exchange, and PEEP in acute respiratory failure. *Anesthesiology.* 1988;69:824–32.
- Terragni PP, Rosboch G, Tealdi A, Corno E, Menaldo E, Davini O, et al. Tidal hyperinflation during low tidal volume ventilation in acute respiratory distress syndrome. *Am J Respir Crit Care Med.* 2007;175:160–6.
- Pelosi P, D'Andrea L, Vitale G, Pesenti A, Gattinoni L. Vertical gradient of regional lung inflation in adult respiratory distress syndrome. *Am J Respir Crit Care Med.* 1994;149:8–13.
- Cressoni M, Cadringer P, Chiurazzi C, Amini M, Gallazzi E, Marino A, et al. Lung inhomogeneity in patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med.* 2014;189:149–58.
- Cressoni M, Chiumello D, Algieri I, Brioni M, Chiurazzi C, Colombo A, et al. Opening pressures and atelectrauma in acute respiratory distress syndrome. *Intensive Care Med.* 2017;43:603–11.
- da Paula LFS, C, Wellman TJ, Winkler T, Spieth PM, Güldner A, Venegas JG, et al. Regional tidal lung strain in mechanically ventilated normal lungs. *J Appl Physiol.* 2016;121(6):1335–47.
- R Core Team. R: A language and environment for statistical computing [Internet]. Vienna, Austria: R Foundation for statistical Computing, 2020. Available from: <http://www.R-project.org>
- Hothorn T, Bretz F, Westfall P. Simultaneous inference in general parametric models. *Biom J.* 2008;50:346–63.
- Bates D, Maechler M, Bolker B, Walker S. Fitting linear mixed-effects models using lme4. *J Stat Softw.* 2015;67:1–48.
- Kuznetsova A, Brockhoff PB, Christensen RHB. lmerTest package: tests in linear mixed effects models. *J Stat Softw.* 2017;82:1–26.
- Long JA. Interactions: Comprehensive, user-friendly toolkit for probing interactions [Internet]. 2019. Available from: <https://cran.r-project.org/package=interactions>
- COVID-ICU Group on behalf of the REVA Network and the COVID-ICU Investigators. Clinical characteristics and day-90 outcomes of 4244 critically ill adults with COVID-19: a prospective cohort study. *Intensive Care Med.* 2020;47:60–73.
- Barbaro RP, MacLaren G, Boonstra PS, Iwashyna TJ, Slutsky AS, Fan E, et al. Extracorporeal membrane oxygenation support in COVID-19: an international cohort study of the extracorporeal life support organization registry. *Lancet.* 2020;396:1071–8.
- Schmidt M, Hajage D, Lebreton G, Monsel A, Voiriot G, Levy D, et al. Extracorporeal membrane oxygenation for severe acute respiratory distress syndrome associated with COVID-19: a retrospective cohort study. *Lancet Respir Med.* 2020;8(11):1121.
- Araos J, Alegria L, Garcia P, Cruces P, Soto D, Erranz B, et al. Near-apneic ventilation decreases lung injury and fibroproliferation in an acute respiratory distress syndrome model with extracorporeal membrane oxygenation. *Am J Respir Crit Care Med.* 2019;199:603–12.

31. Gattinoni L, Caironi P, Pelosi P, Goodman LR. What Has computed tomography taught us about the acute respiratory distress syndrome? *Am J Respir Crit Care Med.* 2001;164:1701–11.
32. Chauvelot L, Bitker L, Dhelft F, Mezidi M, Orkisz M, Davila Serrano E, et al. Quantitative-analysis of computed tomography in COVID-19 and non COVID-19 ARDS patients: A case-control study. *J Crit Care.* 2020;60:169–76.
33. Leisman DE, Harhay MO, Lederer DJ, Abramson M, Adjei AA, Bakker J, et al. Development and reporting of prediction models: guidance for authors from editors of respiratory, sleep, and critical care journals. *Crit Care Med.* 2020;48:623–33.
34. Chiumello D, Marino A, Brioni M, Cigada I, Menga F, Colombo A, et al. Lung recruitment assessed by respiratory mechanics and computed tomography in patients with acute respiratory distress syndrome what is the relationship? *Am J Respir Crit Care Med.* 2016;193(11):1254–63.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

