



**HAL**  
open science

## **Immune effector cell–associated hematotoxicity: EHA/EBMT consensus grading and best practice recommendations**

Kai Rejeski, Marion Subklewe, Mahmoud Aljurf, Emmanuel Bachy, Adriana Balduzzi, Pere Barba, Benedetto Bruno, Reuben Benjamin, Matteo Carrabba, Christian Chabannon, et al.

### ► **To cite this version:**

Kai Rejeski, Marion Subklewe, Mahmoud Aljurf, Emmanuel Bachy, Adriana Balduzzi, et al.. Immune effector cell–associated hematotoxicity: EHA/EBMT consensus grading and best practice recommendations. *Blood*, 2023, 142 (10), pp.865-877. 10.1182/blood.2023020578 . hal-04212662

**HAL Id: hal-04212662**

**<https://hal.science/hal-04212662>**

Submitted on 3 Oct 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Immune Effector Cell-Associated Hematotoxicity (ICAHT):  
EHA/EBMT Consensus Grading and Best Practice Recommendations**

Kai Rejeski<sup>\*1</sup>, Marion Subklewe<sup>\*1</sup>, Mahmoud Aljurf<sup>2</sup>, Emmanuel Bachy<sup>3</sup>, Adriana Balduzzi<sup>4</sup>, Pere Barba<sup>5</sup>, Benedetto Bruno<sup>6</sup>, Reuben Benjamin<sup>7</sup>, Matteo G. Carrabba<sup>8</sup>, Christian Chabannon<sup>9</sup>, Fabio Ciceri<sup>8</sup>, Paolo Corradini<sup>10</sup>, Julio Delgado<sup>11</sup>, Roberta Di Blasi<sup>12</sup>, Raffaella Greco<sup>8</sup>, Roch Houot<sup>13</sup>, Gloria Iacoboni<sup>5</sup>, Ulrich Jäger<sup>14</sup>, Marie José Kersten<sup>15</sup>, Stephan Mielke<sup>16</sup>, Arnon Nagler<sup>17</sup>, Francesco Onida<sup>18</sup>, Zinaida Peric<sup>19</sup>, Claire Roddie<sup>20</sup>, Annalisa Ruggeri<sup>8</sup>, Fermín Sánchez-Guijo<sup>21</sup>, Isabel Sánchez-Ortega<sup>22</sup>, Dominik Schneidawind<sup>23</sup>, Maria-Luisa Schubert<sup>24</sup>, John A. Snowden<sup>25</sup>, Catherine Thieblemont<sup>12</sup>, Max Topp<sup>26</sup>, Pier-Luigi Zinzani<sup>27</sup>, John G. Gribben<sup>28</sup>, Chiara Bonini<sup>29</sup>, Anna Sureda<sup>30</sup>, Ibrahim Yakoub-Agha<sup>31</sup>

\* Participated equally to this work

1-Department of Medicine III – Hematology/Oncology, University Hospital, Ludwig-Maximilians- University, Munich, Germany

2-Oncology Center, King Faisal Specialist Hospital and Research Center, Riyadh, Saudi Arabia

3-Department of Hematology, Hospices Civils de Lyon and University Claude Bernard Lyon 1, Lyon, France

4- Pediatric Transplantation Unit, Department of Medicine and Surgery, University of Milano-Bicocca - Fondazione IRCCS San Gerardo dei Tintori

5- Department of Hematology, Vall d'Hebron University Hospital, Experimental Hematology, Vall d'Hebron Institute of Oncology (VHIO), Vall d'Hebron Barcelona Hospital Campus, Barcelona; Department of Medicine, Universitat Autònoma de Barcelona, Bellaterra

6-Division of Hematology and Cell Therapy Unit, Department of Molecular Biotechnology and Health Sciences, University of Torino, Torino, Italy

7- School of Cancer & Pharmaceutical Sciences, King's College London

8- Unit of Hematology and Bone Marrow Transplantation, IRCCS San Raffaele Hospital, Vita-Salute San Raffaele University, Milan, Italy

9- Institut Paoli-Calmettes Comprehensive Cancer Centre & module Biothérapies du Centre d'Investigations Cliniques de Marseille, Inserm – Aix-Marseille Université – AP-HM – IPC, CBT-1409

10- Division of Hematology & Stem Cell Transplantation, Fondazione IRCCS Istituto Nazionale dei Tumori, University of Milano

11- Oncoimmunotherapy Unit, Department of Hematology, Hospital Clinic de Barcelona, University of Barcelona, FCRB-IDIBAPS, Barcelona, Spain

12- Université de Paris ; Assistance Publique – Hôpitaux de Paris, Service d'hémo-oncologie, DMU DHI, Paris, France

13- Department of Hematology, CHU Rennes, University of Rennes, INSERM U1236, Rennes, France

14- Medical University of Vienna, Department of Medicine I, Division of Hematology and Hemostaseology, Vienna, Austria

15- Department of Hematology, Amsterdam University Medical Centers, location University of Amsterdam, Cancer Center Amsterdam, Amsterdam, LYMMCARE, the Netherlands

16- Department of Cellular Therapy and Allogenic Stem Cell Transplantation (CAST), Department of Laboratory Medicine and Medicine Huddinge, Karolinska University Hospital and Institutet, Karolinska Comprehensive Cancer Center, Stockholm, Sweden

17- Division of Hematology and CBB, Chaim Sheba Medical Center, Tel Aviv University, Tel-Hashomer, Israel.

18- Hematology and BMT Unit, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, University of Milan, Milan, Italy.

19- Department of Hematology, University Hospital Centre Zagreb and School of Medicine, University of Zagreb, Zagreb, Croatia.

20-University College London Cancer Institute, London, UK; Department of Hematology, University College London Hospital, London, UK.

21-University of Salamanca, IBSAL-University Hospital of Salamanca, Salamanca, Spain

22-EBMT, Executive Office, Barcelona, Spain

23-University Hospital Zurich, Department of Medical Oncology and Hematology, Zurich, Switzerland

24-Department of Medicine V, University Hospital Heidelberg, Heidelberg, Germany

25-Department of Haematology, Sheffield Teaching Hospitals NHS Foundation Trust & Department of Oncology and Metabolism, University of Sheffield, United Kingdom.  
26-Medizinische Klinik und Poliklinik II, Universitätsklinikum Würzburg, Würzburg, Germany  
27-IRCCS Azienda Ospedaliero-Universitaria di Bologna, Istituto di Ematologia "Seràgnoli," Dipartimento di Medicina Specialistica, Diagnostica e Sperimentale, Università di Bologna, Bologna, Italy.  
28-Barts Cancer Institute, Queen Mary, University of London, London UK  
29-Experimental Hematology Unit, Division of Immunology, Transplantation and Infectious Disease, IRCCS San Raffaele Hospital, Vita-Salute San Raffaele University, Milan, Italy.  
30-Clinical Hematology Department, Institut Català d'Oncologia-L'Hospitalet, Barcelona. Institut d'Investigació Biomèdica de Bellvitge (IDIBELL), Universitat de Barcelona (UB), Spain  
31-CHU de Lille, Univ Lille, INSERM U1286, Infinite, 59000 Lille, France

**Corresponding authors:****Marion Subklewe, M.D.**

LMU – Klinikum der Universität München  
Department of Hematology & Oncology  
Marchioninistrasse 15, 81377 München, Germany  
Email: marion.subklewe@med.uni-muenchen.de

**Prof. Ibrahim Yakoub-Agha, M.D, P.H.D**

UAM allogreffes de CSH, CHRU, F-59037 Lille CEDEX, France  
Tel: +33(0)3.20.44.55.51  
Email: ibrahim.yakoubagha@chu-lille.fr

Running head: Hematotoxicity following CAR T-cell therapy

Keywords: immune effector cell-associated hematotoxicity, ICAHT, growth factors, hematopoietic cell boost, chimeric antigen receptor (CAR) T-cell therapy

**1 Abstract**

2 Hematological toxicity represents the most common adverse event following chimeric antigen  
3 receptor (CAR) T-cell therapy. Cytopenias can be profound, long-lasting, and can predispose for  
4 severe infectious complications. In a recent worldwide survey, we demonstrated that there  
5 remains considerable heterogeneity in regards to current practice patterns. Here, we sought to  
6 build consensus on the grading and management of Immune Effector Cell Associated Hemato-  
7 Toxicity (ICAHT) following CAR-T therapy. For this purpose, a joint effort between the European  
8 society for Blood and Marrow Transplantation (EBMT) and the European Hematology Association  
9 (EHA) involved an international panel of 36 CAR-T experts who met in a series of virtual  
10 conferences, culminating in a 2-day meeting in Lille, France. On the basis of these deliberations,  
11 best practice recommendations were developed. For the grading of ICAHT, a classification  
12 system based on depth and duration of neutropenia was developed for early (day 0-30) and late  
13 cytopenia (after day +30). Detailed recommendations on risk factors, available pre-infusion  
14 scoring systems (e.g. CAR-HEMATOTOX score), and diagnostic work-up are provided. A further  
15 section focuses on identifying hemophagocytosis in the context of severe hematotoxicity. Finally,  
16 we review current evidence and provide consensus recommendations for the management of  
17 ICAHT, including growth factor support, anti-infectious prophylaxis, transfusions, autologous  
18 hematopoietic cell boost, and allogeneic hematopoietic cell transplantation. In conclusion, we  
19 propose ICAHT as a novel toxicity category following immune effector cell therapy, provide a  
20 framework for its grading, review literature on risk factors, and outline expert recommendations  
21 for the diagnostic work-up and short- and long-term management.

**22 Introduction and state-of-the-art**

23 The last decade has firmly established chimeric antigen receptor (CAR) T-cell therapy as a  
24 practice-changing immunotherapy platform for an increasing number of refractory B-cell  
25 malignancies.<sup>1-7</sup> While durable remissions can be achieved, this comes with the caveat of a  
26 unique spectrum of side effects ranging from Cytokine Release Syndrome (CRS), to Immune  
27 Effector Cell Associated Neurotoxicity Syndrome (ICANS), and Immune Effector Cell Associated  
28 Hemophagocytic Lymphohistiocytosis-like Syndrome (IEC-HS).<sup>8-11</sup> Real-world evidence has  
29 underlined the growing importance of hematological toxicity as the most frequent Common  
30 Terminology Criteria for Adverse Events (CTCAE) grade  $\geq 3$  adverse event following CAR T-cell  
31 therapy.<sup>12-14</sup> Similarly high rates of cytopenias have been reported for other T-cell based  
32 immunotherapies such as bispecific antibodies.<sup>15-19</sup> Notably, profound and often long-lasting  
33 cytopenias can add to the immunosuppression conferred by B-cell aplasia and consecutive  
34 hypogammaglobulinemia.<sup>20</sup> Importantly, severe infections are a major driver of both morbidity and  
35 non-relapse mortality (NRM) following CAR T-cell therapies.<sup>21-23</sup>

36

37 Hematological side effects have been described after CAR T-cell therapy regardless of the target  
38 antigen (e.g., CD19 vs. CD22 vs. BCMA) and across various disease entities (e.g., LBCL, BCP-  
39 ALL, MCL, MM, FL).<sup>3-5,24-29</sup> Several features underline the unique nature of CAR-T related  
40 hematotoxicity. First, cytopenias can persist long after the resolution of clinical CRS, and have  
41 been reported as long as months to years following CAR T-cell infusion.<sup>30</sup> Hematopoietic count  
42 recovery often follows a biphasic trajectory, with intermittent recovery followed by second, or  
43 multiple, dips.<sup>12,13</sup> Second, patients can develop very severe bone marrow (BM) aplasia that is  
44 often refractory to therapeutic measures such as growth factor support.<sup>13,31,32</sup> Finally, the  
45 underlying pathophysiology remains to be elucidated, although recent evidence points towards  
46 the importance of both baseline hematopoietic reserve and the systemic inflammatory state of the  
47 host.<sup>13</sup> Moreover, the inflammatory stress conferred by severe CRS and the associated  
48 alterations in cytokine patterns can exert myelosuppressive effects.<sup>33-35</sup>

49

50 In a recent international survey led by EHA and EBMT, we identified a high degree of  
51 heterogeneity both in regards to the grading and management of cytopenias.<sup>36</sup> Current grading

52 systems such as the CTCAE describe cytopenias predominantly in quantitative terms by  
53 assigning severity grades according to the depth of cytopenia. However, they are difficult to apply  
54 in daily practice and fail to capture the distinct nature of post-CAR-T hematopoietic reconstitution,  
55 such as the biphasic and/or delayed course. Furthermore, the cumulative risk of secondary  
56 complications (e.g., infections, bleeding) primarily increases with the respective duration of  
57 observed cytopenia.<sup>22,37</sup> Classification systems that were developed for cytopenia following  
58 classic cytotoxic chemotherapies may not apply to patients receiving novel T-cell based  
59 immunotherapies. To accommodate these unique features of hematological side effects in adult  
60 patients receiving such therapies, we herein introduce the concept of Immune Effector Cell  
61 Associated Hemato-Toxicity, or ICAHT. Based on a novel framework for grading, we outline  
62 expert recommendations for its diagnostic work-up and management.

63

#### 64 **Methodology**

65 This workshop is based on the EBMT PH & G committee method.<sup>38</sup> In September 2022, KR and  
66 MS proposed to set up a workshop to issue European recommendations regarding the grading  
67 and management of ICAHT, particularly following autologous CAR T-cell therapy. As a first step,  
68 an international survey on current practices at >50 global CAR-T centers was sent out and results  
69 were analyzed.<sup>36</sup> Experts from different countries and belonging to EBMT and EHA were  
70 subsequently invited to join the workshop. As a second step, several teleconferences took place  
71 to discuss and advance the first draft. Along with the results of the international survey, a  
72 comprehensive literature review was carried out by the workshop participants within each  
73 subgroup, which served as the basis for the discussions. The third step consisted of a two-day  
74 face-to-face meeting which took place in Lille, France on March 2<sup>nd</sup> and 3<sup>rd</sup>, 2023.

75

76 These recommendations are intended to be general in scope and applicable to all diseases and  
77 types of autologous CAR T-cell therapies or other T-cell based immunotherapies (e.g., bispecific  
78 antibody constructs) adopted as standard clinical practice. They are intended to reflect current  
79 best practices in this new and rapidly evolving field and aim to help clinicians and other  
80 healthcare professionals in providing consistent, high-quality patient care. These  
81 recommendations were created due to the growing number of autologous CAR T-cell therapies

82 currently available outside clinical trials for the treatment of hematological malignancies. Given  
83 the lack of high-quality evidence from randomized trials in this area (expected Evidence Levels 3-  
84 5, Oxford Centre for Evidence Based Medicine), the decision was made not to grade these  
85 recommendations. They therefore represent the consensus point of view of the authors. When  
86 administering CAR T-cell therapies within clinical trials, physicians are advised to follow  
87 respective trial protocols.

88

## 89 **Consensus recommendations**

### 90 **1. ICAHT Grading**

91 On the basis of the results of the international survey on behalf of EHA and EBMT, the expert  
92 panel defined early ICAHT as cytopenia occurring during the first 30 days after CAR T-cell  
93 infusion. Conversely, late ICAHT was classified as cytopenia observed beyond day +30. The  
94 expert panel resolved that the main clinical action points of post-CAR-T cytopenias concerned  
95 profound and/or prolonged neutropenia, and that isolated thrombocytopenia or anemia represent  
96 rare occurrences. Concomitantly, a grading system based on neutropenia was pursued. For early  
97 ICAHT (day 0-30), a grading system based on both depth and duration of neutropenia was  
98 defined due to the associated clinical sequelae (**Table 1**, top). Late ICAHT was graded based on  
99 the elapsed time from CAR T-cell infusion (e.g., occurring after day +30) with the severity (grade  
100 I-IV) defined by the depth of neutropenia (**Table 1**, bottom). For anemia and thrombocytopenia,  
101 the expert panel refers to existing grading systems and recommends that institutional guidelines  
102 should be followed, as further outlined in **Section 6** and **Table 4** (see *transfusions*).

103

### 104 **2. Risk factors for developing post-CAR-T cytopenias**

105 The overall incidence of hematological toxicity in the key registrational trials for CAR T-cell  
106 products endorsed by the European Medicines Agency (EMA) are outlined in the **Supplemental**  
107 **Table 1**. Furthermore, we performed an extensive literature review of prominent real-world  
108 studies with a specific focus on correlative studies and potential risk factors (**Supplemental**  
109 **Table 2**). Overall, a plethora of factors contribute to the development of cytopenias after CAR-T,  
110 some of which remain incompletely understood. Broadly, they relate to the underlying disease  
111 and its previous treatments, baseline risk factors (e.g., hematopoietic reserve, BM infiltration,

112 systemic inflammation), as well as CAR-T product features and CRS-related inflammatory  
113 patterns (summarized in **Table 2** and the **Supplementary Text**).<sup>12,13,23,30,33,34,39-56</sup>

114

### 115 **3. What scoring systems to use**

116 Based on several of the risk factors delineated above, the CAR-HEMATOTOX score was  
117 developed to identify patients at high risk for prolonged neutropenia, and especially the  
118 development of the aplastic phenotype of neutrophil recovery.<sup>13</sup> An online calculator can be found  
119 on the website of the German Lymphoma Alliance (GLA): [https://www.german-lymphoma-](https://www.german-lymphoma-alliance.de/Scores.html)  
120 [alliance.de/Scores.html](https://www.german-lymphoma-alliance.de/Scores.html)). The score incorporates factors related to hematopoietic reserve  
121 (absolute neutrophil count [ANC], hemoglobin, platelet count) and baseline inflammatory state  
122 (CRP, ferritin) and was validated for a primary endpoint of severe neutropenia (ANC <500/ $\mu$ L)  
123 lasting longer than 14 days during the first 60 days after CAR-T infusion. Importantly, the CAR-  
124 HEMATOTOX score is determined prior to lymphodepleting chemotherapy and thus enables  
125 early risk-stratification into a high vs. low risk of developing severe hematotoxicity after CAR T-  
126 cell treatment (**Figure 1**). In subsequent studies, the score also identified patients at risk for  
127 severe infections and poor treatment outcomes across multiple disease entities (e.g., LBCL,  
128 MCL, MM).<sup>22,42-44,57</sup> However, it is important to note that the score remains to be validated  
129 prospectively and for adult and pediatric BCP-ALL patients. Furthermore, the test characteristics  
130 (high sensitivity, lower specificity) indicate a lower positive predictive value, meaning that not all  
131 patients deemed high-risk will develop severe hematotoxicity. Conversely, the high negative  
132 predictive value suggests that the score is particularly helpful in ruling out patients at risk for  
133 severe hematotoxicity.

134

### 135 **4. Assessment and diagnostic work-up of ICAHT**

136 In patients with a high-risk profile for developing ICAHT (**Table 2, Figure 1**), baseline BM  
137 studies (prior to apheresis or lymphodepletion) should be considered to risk-stratify patients for  
138 hematological toxicity and identify underlying marrow infiltration as a pertinent risk factor.  
139 Cryopreservation of the BM aspirate and/or peripheral blood mononuclear cells (PBMCs) is  
140 optional, but may provide useful information in case the patient develops secondary BM failure  
141 (e.g., presence of CHiP clone).



142

143 In case of cytopenia that persists beyond the expected reconstitution of lymphodepleting  
144 chemotherapy (typically following week 2-3 after CAR-T infusion), the first step in the work-up  
145 comprises defining the differential diagnosis, which can include drug-induced cytopenia, vitamin  
146 deficiencies, infectious causes, sustained inflammatory stressors, relapse and/or active BM  
147 disease. The expert panel recommends performing an incremental diagnostic-work-up, with an  
148 initial tier 1 assessment comprising standard diagnostic tests that should be performed in all  
149 cases of severe, or grade  $\geq$ III, ICAHT (**Figure 2**). In case the tier 1 results are inconclusive and  
150 cytopenias persist and/or are G-CSF refractory (absence of count recovery despite  $\geq$ 5 days of  
151 G-CSF support), a subsequent tier 2 diagnostic work-up can be pursued. Importantly, this  
152 includes extended viral studies, as well as BM aspiration and biopsy. The expert panel would  
153 reserve cytogenetics and next-generation sequencing to rule out an underlying myeloid  
154 malignancy to either cases of profound, long-lasting marrow aplasia (e.g., no count recovery  
155 above an ANC  $\geq$ 500/ $\mu$ L by day +30, pancytopenia), or new-onset pancytopenia that is refractory  
156 to therapeutic measures late after CAR-T infusion.

157

### 158 **5. Hemophagocytosis associated with severe hematotoxicity after CAR T-cell therapy**

159 Hemophagocytic lymphohistiocytosis (HLH) represents a hyper-inflammatory condition  
160 resulting from abnormal immune activation, which is associated with high fever,  
161 hyperferritinemia, prolonged cytopenia and eventually multi-organ failure. HLH remains a  
162 diagnostic quandary as unique biomarkers are still lacking and/or not readily available. In the  
163 context of CAR T-cell therapy, the incidence of HLH-like symptoms ranges from 1% to  
164 3.4%.<sup>10,58</sup> Two entities, CRS/MAS and IEC-HS, can be distinguished according to time of  
165 onset and presence of concomitant CRS/ICANS symptoms.<sup>29,59-61</sup> In patients with severe  
166 ICAHT that present with aplastic neutrophil recovery and rising serum ferritin, the diagnosis of  
167 HLH should be considered, as both can present with profound immune dysregulation and  
168 increased IFN signaling.<sup>42,54</sup> A comprehensive work-up is recommended in order to identify  
169 additional abnormalities such as new-onset hepatosplenomegaly, hypertriglyceridemia,  
170 coagulopathy and hypofibrinogenemia, as well as hemophagocytosis features on BM biopsy  
171 or in other tissues (**Fig. 2**). Existing scoring systems that can guide the diagnosis of HLH in

172 the context of severe ICAHT include HLH-2004 criteria, the H-score, and the OHI index.<sup>62-64</sup>  
173 Additionally, **Table S3** outlines the MD Anderson criteria<sup>58</sup>, EBMT/EHA/JACIE  
174 recommendations<sup>59</sup> and IEC-HS criteria<sup>60</sup>, which were deemed more specific to CAR-T  
175 therapy by the expert panel. In patients in whom ICAHT manifests in the form of HLH, anti-  
176 inflammatory measures should be promptly initiated to mitigate cytokine storm and its clinical  
177 sequelae. Patients should be treated with anakinra, a recombinant humanized IL-1 receptor  
178 antagonist, in combination with high-dose corticosteroids (**Figure S1**). In refractory cases,  
179 ruxolitinib, cytokine adsorption, and emapalumab (IFN- $\gamma$  inhibitor) can be considered, albeit  
180 data remains scarce.<sup>65-67</sup>

181

## 182 **6. Management of cytopenias**

183 The management of ICAHT can broadly be separated into an initial phase which addresses the  
184 (expected) early cytopenias and aims to mitigate the risk of infections and/or other complications,  
185 as well as a later phase that is initiated in case of persistent and/or therapy-refractory  
186 cytopenias. An overview of the expert recommendations for early ICAHT management is  
187 provided in **Table 3**.

188

### 189 **Transfusions**

190 Due to the frequent nature of severe anemia and thrombocytopenia after CAR-T therapy,  
191 transfusions are an essential part of supportive care and include either packed red blood cell  
192 concentrates (pRBCs) or platelet concentrates (PCs). Transfusion-associated GvHD (ta-GvHD)  
193 is a rare complication of transfusion wherein viable donor T lymphocytes in cellular blood  
194 products mount an immune response against the recipient.<sup>68</sup> Considering the high mortality rate  
195 (>90%), prevention of ta-GvHD is recommended, though there is no internationally agreed upon  
196 consensus on the duration of the use of irradiated blood products across cellular therapies. In  
197 the setting of HCT, standard practice is to use irradiated blood for (1) at least 2 weeks prior to  
198 stem cell collection until at least 3 months after auto-HCT, and (2) starting with conditioning at  
199 the latest until at least 6 months after allo-HCT, or until immune reconstitution.<sup>69</sup> In the context  
200 of CAR-T therapy, the expert panel recommended the irradiation of blood products from 7 days  
201 prior to leukapheresis until at least 90 days post-CAR-T infusion unless conditioning, disease or

202 previous treatment determine indefinite duration (**Table 3**). Of note, the use of the purine  
203 analogue fludarabine as a component of lymphodepletion prior to CAR-T infusion may impact  
204 local guidance for irradiated blood products.<sup>69</sup> Given its relative rarity, we recommend reporting  
205 cases of ta-GVHD following CAR-T to regulatory authorities.

206

### 207 **Growth factor support**

#### 208 *Granulocyte-macrophage colony-stimulating factor (GM-CSF)*

209 GM-CSF is typically elevated in CAR-T patients with CRS and ICANS. The use of GM-CSF as a  
210 growth factor for patients with low blood counts should be avoided as it may promote  
211 inflammatory toxicity and induce neuroinflammation following CAR-T therapy.<sup>70,71</sup>

212

#### 213 *Granulocyte colony-stimulating factor (G-CSF)*

214 Due to the concerns for the use of GM-CSF and the hypothesized, but largely unknown risks of  
215 exacerbating toxicities, early guidance suggested generally deferring G-CSF until resolution of  
216 acute CAR T-cell related immunotoxicity (typically week 3). However, several recent reports  
217 question this as a general rule and point towards an acceptable safety profile for the early use of  
218 G-CSF, with no increase of high-grade ( $\geq 3^\circ$ ) CRS/ICANS.<sup>72-76</sup> In the largest retrospective analysis  
219 by Miller and colleagues (n=197), prophylactic G-CSF *before* CAR-T (mostly pegylated G-CSF)  
220 was associated with faster neutrophil recovery, comparable treatment outcomes, and similar  
221 rates of severe ICANS.<sup>75</sup> While prophylactic G-CSF was associated with a higher rate of grade  $\geq 2$   
222 CRS, this observation did not extend to the clinically relevant grade  $\geq 3$  CRS. In a subgroup  
223 analysis, the authors found that G-CSF did not worsen severity of CRS in patients who *already*  
224 present with low-grade ( $1^\circ$ ) toxicity. In a further study by Lievin et al, early G-CSF administration  
225 (from day +2) in neutropenic patients was associated with a reduced risk of febrile neutropenia  
226 without increasing the risk of severe CRS or ICANS.<sup>74</sup> Notably, G-CSF was also safe in  
227 maintaining CAR T-cell expansion kinetics and anti-lymphoma activity, without any deleterious  
228 impact on the quality of response and outcomes.<sup>73,74</sup> Appraising the above evidence and  
229 weighing the benefits and risks, early G-CSF administration on day +2 can be considered in high-  
230 risk patients to shorten the length of expected severe neutropenia (see **Table 2** and **Figure 1**).  
231 Therapeutic G-CSF in case of prolonged severe neutropenia (ANC  $<500/\mu\text{L}$ ) can also be

232 considered, and can be of diagnostic benefit for identifying the aplastic neutrophil recovery  
233 phenotype<sup>13,32</sup>, which is often G-CSF unresponsive. The large majority of CAR-T patients (>80%)  
234 ultimately respond to growth factor support with count recovery.<sup>32,34</sup> However, recurrent neutrophil  
235 dips (biphasic course) can necessitate intermittent application of therapeutic G-CSF (**Figure 3**).  
236 Finally, a uniform consensus was reached on the necessity of *prospective*, and ideally  
237 multicenter, clinical trials that evaluate the safety and optimal treatment protocol for G-CSF  
238 (prophylactic vs. early / pegylated vs. non-pegylated) in the context of CAR-T therapy and across  
239 disease entities (B-ALL vs. B-NHL vs. MM).

240

#### 241 *Thrombopoietin (TPO) agonists*

242 TPO agonists (e.g., eltrombopag, romiplostim) are considered primarily in patients with  
243 prolonged and late thrombocytopenia, with the thrombocytopenic nadir typically occurring in the  
244 2<sup>nd</sup> month after CAR-T therapy.<sup>12,13</sup> Data supporting the use of TPO agonists in the CAR-T  
245 setting are extremely limited and are restricted to a few case series from single centers with  
246 limited patient numbers.<sup>77-79</sup> In these limited reports, improvement in platelets and also  
247 hemoglobin and ANC was noted, with some patients becoming transfusion independent both  
248 for platelets and pRBCs similar to improvement in hematopoiesis observed with TPO agonist  
249 use in cases of acquired BM failure.<sup>80,81</sup> Due to the limited available data, the expert panel  
250 advises that the use of TPO agonists should parallel the practice for HCT.<sup>82</sup> They can also be  
251 utilized in G-CSF refractory cases of ICAHT (**Figure 3**).

252

#### 253 **Infection prophylaxis**

254 Regarding the administration of anti-infectious prophylaxis during cytopenia, the expert panel  
255 broadly recommends adherence to the general EHA/EBMT/JACIE guidelines for patients  
256 receiving CAR T-cell therapy.<sup>59</sup> The following specific recommendations were issued (**Table 3**):

- 257     ▪ Adherence to current EHA/EBMT guidelines regarding anti-viral and anti-pneumocystis  
258         pneumonia (PCP) prophylaxis, as well as intravenous immunoglobulin (IVIG) substitution  
259         for post-CAR-T hypogammaglobulinemia.<sup>59</sup>
- 260     ▪ The expert panel does not recommend the use of a neutropenic diet to reduce the risk of  
261         infection in neutropenic CAR-T patients.<sup>83-85</sup>

- 262     ▪ **Antibacterial prophylaxis:** the panel proposes a risk-adapted strategy based on the  
263     patient-individual risk profile for infections including the expected incidence rate of  
264     protracted, profound neutropenia (ANC <100/ $\mu$ L for  $\geq$ 7 days), in line with the consensus  
265     American Society of Clinical Oncology (ASCO)/ Infectious Diseases Society of America  
266     (IDSA) recommendations for adult cancer patients.<sup>86</sup> Antibacterial prophylaxis with a  
267     fluoroquinolone (e.g. levofloxacin, ciprofloxacin) is not recommended in patients who are at  
268     a low risk of severe (grade  $\geq$ III) ICAHT (**Table 1, Table 3**) and should be avoided due to  
269     fluoroquinolone-specific side effects, the potential emergence of resistant strains, and  
270     selection for *C. difficile* and *enterococci*.<sup>37,87-90</sup> Furthermore, recent publications have  
271     demonstrated that antibiotic exposure prior to CAR T-cell therapy reduces microbiome  
272     diversity and is associated with inferior outcomes, potentially due to the multifunctional and  
273     immunomodulatory role of the gut microbiome.<sup>91-94</sup> On the other hand, antibacterial  
274     prophylaxis can be considered in high-risk patients once the ANC falls below <500/ $\mu$ L to  
275     mitigate the risk of severe infections. The CAR-HEMATOTOX score may be useful for  
276     guidance and identification of high-risk candidates.<sup>13</sup> In a large retrospective analysis of  
277     LBCL patients receiving CD19 CAR-T, a significant reduction of severe bacterial infections  
278     with fluoroquinolone prophylaxis was observed in CAR-HEMATOTOX<sup>high</sup> but not CAR-  
279     HEMATOTOX<sup>low</sup> patients, supporting a risk-adapted approach. Importantly, the panel  
280     recommends adherence to institutional guidelines that take into account local epidemiology  
281     and resistance patterns. In this context, monitoring for multi-drug resistant gram-negative  
282     bacteria (MDR GNB) colonization (i.e., active surveillance through rectal swab culture) may  
283     be useful both for baseline risk assessment and during prolonged neutropenia.
- 284     ▪ **Antifungal prophylaxis:** To reduce the risk of invasive fungal disease (IFD), anti-mold  
285     prophylaxis (e.g., micafungin or posaconazole) can be considered in patients at high risk  
286     for severe ICAHT (grade  $\geq$ III) once the ANC falls below <500/ $\mu$ L (**Table 3**). Additional risk  
287     factors to consider are prior allo-HCT, prior invasive aspergillosis and receipt of  
288     corticosteroids (either long-term  $\geq$ 72h or high-dose, e.g., greater than 10 mg of  
289     dexamethasone or equivalent). The low overall incidence rate for IFD in the context of  
290     CAR-T should be taken into account<sup>95</sup>, although fungal infections represent a frequent  
291     cause of fatal infectious complications.<sup>22,96</sup> Systemic primary antifungal prophylaxis should

292 be continued until stable count recovery (ANC >500/ $\mu$ L over 3 days) and discontinuation of  
293 steroids for CRS/ICANS management.

294

295 *Hematopoietic cell boost*

296 Patients who are unresponsive and/or refractory to G-CSF beyond day +14 after CAR-T infusion  
297 represent a clinically challenging subgroup of patients at high risk for severe and even fatal  
298 infectious complications. While the evidence remains limited, TPO agonists can be offered in this  
299 setting, especially in cases of associated thrombocytopenia.<sup>79</sup> In cases of severe ICAHT in which  
300 an inflammatory stressor is deemed contributory (severe CRS/ICANS, CRS/MAS), anti-  
301 inflammatory strategies such as pulse-dose corticosteroids and/or anti-cytokine therapies (e.g.,  
302 tocilizumab, anakinra) should be used. A promising strategy pertains to the use of cryopreserved  
303 autologous or allogeneic CD34<sup>+</sup> hematopoietic cells from prior collection (either prior auto- or allo-  
304 HCT).<sup>97-99</sup> Three recent case series shed light on both the safety and clinical feasibility of this  
305 approach across a broad population of pediatric and adult patients (summarized in  
306 **Supplemental Table 4**). High rates of sustained neutrophil and platelet engraftment were noted  
307 across studies. While hematopoietic cell boost (HCB) has been successfully applied during active  
308 infection<sup>100</sup>, clinicians should be aware of the possibility of immune reconstitution inflammatory  
309 syndrome (IRIS) in patients with prolonged bone marrow aplasia.<sup>31</sup> As the earlier application of an  
310 *available* HCB was associated with superior survival outcomes,<sup>99</sup> the expert panel recommends  
311 considering the application of a HCB without prior conditioning chemotherapy for grade  $\geq$ III  
312 ICAHT beyond day +14 if (1) a boost is readily available and (2) G-CSF refractoriness has been  
313 established. At the same time, the survey results highlighted that even when HCB were  
314 considered a viable treatment option in a patient with prior auto-HCT, they were often not  
315 available. While prophylactic collection in high-risk candidates has been proposed as a potential  
316 mitigating strategy, the panel cautioned that the collection process may add to the already high  
317 logistic burden of CAR T-cell therapy (e.g., coordination of apheresis slots and storage capacity),  
318 which could negatively impact vein-to-vein times in a state of high disease burden. Furthermore,  
319 the process could incur unnecessary collection- and storage-associated costs.<sup>101,102</sup> Ultimately, it  
320 was concluded that further research is needed to assess the number needed to treat for  
321 prophylactic stem cell collection.

322

323 *Allogeneic hematopoietic cell transplantation*

324 If the above options remain ineffective or elusive and grade IV ICAHT persists beyond day +30,  
325 the expert panel recommends initiating a donor search for a potential allo-HCT as a last resort  
326 (ultima ratio). In such cases of life-threatening ICAHT, the benefit and risks of allo-HCT need to  
327 be carefully weighed and aligned with the patient's goals-of-care. Furthermore, the possibility of  
328 spontaneous count recovery needs to seriously be considered.<sup>34,103,104</sup> Accordingly, the expert  
329 panel suggested that the ultimate trigger for allo-HCT needs to be discussed on a case-by-case  
330 basis. Month 3-6 post CAR-T infusion was deemed a reasonable time frame to balance both the  
331 risk of infection and possibility of spontaneous count recovery. Once the decision for allo-HCT  
332 has been made, details regarding donor selection, conditioning regimens and  
333 immunosuppression have to be discussed. Experience and evidence are very limited and only  
334 general considerations can be reviewed here. As for every allo-HCT, the same basic principles  
335 should apply keeping in mind that the primary indication is severe and persistent cytopenia  
336 although basically all patients currently receive commercially available CAR-T cells to treat  
337 malignant lymphoid disorders. Most importantly, salvage allo-HCT is also capable to provide  
338 tumor control through the conditioning regimen and graft-versus-tumor effects and current  
339 standard procedures will most likely lead to eradication of CAR-T cells at the latest when full  
340 donor chimerism has been established. Therefore, remission status must be determined prior to  
341 allo-HCT and may guide the choice of conditioning regimen and the taper of immunosuppression.  
342 As usual, performance status, comorbidities, prior therapies and expected anti-tumor activity  
343 should be carefully considered when discussing the transplantation modalities, donor choice and  
344 selection.

345

## 346 **7. Conclusions and Outlook**

347 Much progress has been made in the last years in defining hematological toxicity as a distinct  
348 toxicity entity of CAR T-cell therapy. While the underlying pathophysiology remains incompletely  
349 understood, growing evidence points towards critical interactions between host hematopoiesis  
350 and CAR T-cell function and efficacy. By defining ICAHT and delineating a specific grading

351 system, we herein provide a nomenclature that enables cross-trial comparisons and invites  
352 severity-based management strategies.

353

354 In this international consensus guidelines document, we have proposed a structured approach  
355 to diagnosis, grading/staging and clinical management of ICAHT. This endeavor has also set  
356 the stage for areas of future development that will require collaboration between various  
357 European and non-European stakeholders involved in CAR T-cell therapy. Structured sample  
358 collection across multiple centers represents the basis for translational projects that delineate  
359 the underlying mechanisms of ICAHT by leveraging novel technologies such as multi-omics and  
360 single-cell approaches. One area of particular interest lies in identifying early determinants of  
361 ICAHT by studying the peripheral blood immune contexture and/or the local BM  
362 microenvironment from pre-CAR-T samples. Furthermore, large retrospective real-world  
363 analyses may shed light on some of the differences in the clinical management of ICAHT that  
364 were identified by the EHA/EBMT survey. Residual questions relate to the optimal timing of G-  
365 CSF initiation as well as the optimal protocol to employ (e.g., prophylactic vs. early G-CSF). The  
366 question of prophylactic collection of CD34+ hematopoietic cells in high-risk candidates and the  
367 optimal trigger time point for both HCB and allo-HCT represent unresolved issues that warrant  
368 further systematic study. Ultimately, prospective clinical trials will be needed that determine the  
369 potential benefits and evidence-base of treatment strategies that mitigate ICAHT.



370 **Acknowledgements**

371 We are grateful for the support by EHA and EBMT without whom this work would not have been  
372 possible.

373

374 **Competing interests**

375 **K.R.** Kite/Gilead: Research Funding and travel support; Novartis: Honoraria; BMS/Celgene:  
376 Consultancy, Honoraria

377 **M.S.** receives industry research support from Amgen, BMS/Celgene, Kite/Gilead, Janssen,  
378 Miltenyi Biotec, Novartis, Roche, Seattle Genetics and Takeda and serves as a  
379 consultant/advisor to AvenCell, CDR-Life, Ichnos Sciences, Incyte Biosciences, Janssen,  
380 Miltenyi Biotec, Molecular Partners, Novartis, Pfizer and Takeda. She serves on the speakers'  
381 bureau at Amgen, AstraZeneca, BMS/Celgene, Gilead, GSK, Janssen, Novartis, Pfizer, Roche  
382 and Takeda.

383 **E.B.** Honorarium/Advisory Board: Kite/Gilead, BMS/Celgene, Novartis, Pfizer, Incyte, ADC  
384 Therapeutics, Roche, Takeda. Personal fees: Kite/Gilead, BMS/Celgene, Novartis, Pfizer.  
385 Research funding: Amgen, BMS.

386 **A.B.** Speaker's bureau from Novartis, Amgen, Medac; meeting support from Novartis, Amgen,  
387 Medac, Neovii

388 **P.B.** Honoraria and travel support from Allogene, Amgen, BMS, Kite/Gilead, Incyte, Jazz  
389 Pharmaceuticals, Miltenyi Biomedicine, Nektar, Novartis and Pierre Fabre

390 **R.B.** Research funding from Servier and Allogene. Participated in advisory boards for  
391 BMS/Celgene, Janssen, Allogene, Takeda, Enara Bio, Securabio, Menarini Stemline, Arovella,  
392 Kite/Gilead

393 **M.C.** Consultancy: Novartis, Janssen.

394 **C.C.** Consultant advisor: Bellicum Pharmaceuticals, BMS, Kite/Gilead, Janssen  
395 Pharmaceuticals, Novartis. Speaker's bureau: BMS, Kite/Gilead, Janssen Pharmaceuticals,  
396 Jazz, Miltenyi Biotec, Novartis. Travel support: BMS, Kite/Gilead

397 **P.C.** Honoraria: Abbvie, Amgen, Celgene, Daiichi Sankyo, Janssen, Kite/Gilead, Novartis,  
398 Roche, Sanofi, Servier, Takeda (Consulting, Advisory role or Lecturer); travel and

399 accommodations paid by for-profit health care companies during the past 2 years: Novartis,  
400 Janssen, BMS/Celgene, Takeda, Kite/Gilead, Roche  
401 **R.D.** Novartis: Scientific Advisory Board and Conference speaker. Kite/Gilead: Scientific  
402 Advisory Board, Conference speaker, Travel accommodation. Janssen: Scientific Advisory  
403 Board. Pfizer: Conference speaker. BMS/Celgene: Scientific Advisory Board. Abbvie :  
404 Conference speaker. Incyte : Conference speaker  
405 **R.G.** discloses speaking honoraria from Biotest, Pfizer, Medac, and Magenta.RH:  
406 **R.H.** honoraria from Kite/Gilead, Novartis, Incyte, Janssen, MSD, Takeda and Roche; and  
407 consultancy at Kite/Gilead, Novartis, Bristol-Myers Squibb/Celgene, ADC Therapeutics, Incyte,  
408 Miltenyi.  
409 **G.I.** Honoraria: BMS/Celgene, Novartis, Janssen, AstraZeneca, Abbvie, Miltenyi, Roche.  
410 **U.J.** Consultancy: Novartis, Janssen, Roche; Honoraria: Gilead, BMS/CELGENE, Miltenyi  
411 **M.J.K.** honoraria from and consulting/advisory role for Bristol Myers Squibb, Celgene, Kite, a  
412 Gilead Company, Miltenyi Biotech, Novartis, and Roche; research funding from Kite, Roche,  
413 Takeda, and Celgene; and travel support from Kite, Miltenyi Biotech, Novartis, and Roche.  
414 **S.M.** Via Institution: Celgene/BMS: Speaker's bureau; Novartis: Speaker's bureau; Janssen:  
415 Speaker's bureau; Gilead/KITE: Travel support, Expert panel; Miltenyi: Data safety monitoring  
416 board; Immunicum/Mendes: Data safety monitoring board; SWECARNET: Founder/Leadership;  
417 ScientifyResearch: Founder (spouse).  
418 **F.O.** Travel, Accommodations, Expenses: Takeda, Kyowa Kirin International, Medac  
419 **C.R.** Honoraria BMS, Kite/Gilead, Novartis, Amgen  
420 **F.S.G.** Research support from Novartis, Gilead. Honorary from Novartis, Gilead, Pfizer, BMS-  
421 Celgene,  
422 **D.S.** Advisory Board with Autolus  
423 **M.L.S.** Advisory board / consultancy: Novartis, Kite-Gilead, Takeda, Janssen; Honoraria:  
424 Kite/Gilead  
425 **J.S.** received speaker fees from Gilead and consultancy from Kiadis, Medac and Vertex.  
426 **C.T.** Board with Novartis, Kite/Gilead, BMS, Roche, Takeda, Incyte, Beigene  
427 MT: Ad Board: KITE/Gilead, BMS, Janssen, Novartis, Roche, Incyte, Astra Zeneca. Research  
428 funding: KITE/Gilead, Takeda, Incyte, Roche, Regeneron

429 **P.L.Z.** Consultancy: MSD, Eusapharma, Novartis; Advisory Board: Securia Bio, Celltrion,  
430 Kite/Gilead, Janssen, BMS/Celgene, Servier, Sandoz, MSD, AstraZeneca, Takeda, Roche,  
431 Eusapharma, Kyowa Kirin, Novartis, ADC therap., Incyte, Belgene. Speakers' Bureau: Celltrion,  
432 Kite/Gilead, Janssen, BMS/Celgene, Servier, MSD, AstraZeneca, Takeda, Roche, Eusapharma,  
433 Kyowa Kirin, Novartis, Incyte, Belgene.

434 **J.G.G.** Received honoraria from Amgen, BMS Gilead/Kite, Janssen, Novartis, and received  
435 research funding from AZ, BMS and Janssen.

436 **C.B.** inventor of different patents on cancer immunotherapy and genetic engineering; member of  
437 Advisory Boards and Consultant for, Intellia, Kite/Gilead, Miltenyi, Kiadis, QuellTx, Janssen,  
438 Chroma, Genyo, Pancancer-T, Alia, and received research support from Intellia Therapeutics.

439 **A.S.** Honoraria: Takeda, Bristol Myers Squibb, Merck Sharp & Dohme, Celgene, Janssen, Sanofi,  
440 Roche, Novartis, Gilead Sciences, Janssen-Cilag; Consulting or Advisory Role: Takeda, Bristol  
441 Myers Squibb, Gilead Sciences, Celgene, Janssen, Novartis; Speakers' Bureau: Takeda; Travel,  
442 Accommodations, Expenses: Kite/Gilead

443 **I.Y-A.** Recived honoraria from Novartis, Gilead/Kite, BMS and Janssen.

444

445 The remaining authors have nothing to declare. None of the mentioned conflicts of interest were  
446 related to financing of the content of this manuscript.

447

#### 448 **Author contributions**

449 *Conceptualization: KR, MS;*

450 *Investigation: KR, MS, MA, EB, AB, PB, BB, RB, MGC, CC, FC, PC, JD, RDB, RG, RH, GI, UJ,*  
451 *MJK, SM, AN, FO, ZP, CR, AR, FSG, ISO, DS, MLS, JAS, CT, MT, PLZ, JGG, CB, AS, IYA;*

452 *Formal Analysis and Visualization: KR, MS, RG, RH, MJK, SM, CR, FSG, DS, MLS, CT, AS, IYA*

453 *Methodology: KR, MS, ISO, AS, IYA*

454 *Writing Original Draft: KR, MS*

455 *Writing Review and Editing: KR, MS, MA, EB, AB, PB, BB, RB, MGC, CC, FC, PC, JD, RDB, RG,*  
456 *RH, GI, UJ, MJK, SM, AN, FO, ZP, CR, AR, FSG, ISO, DS, MLS, JAS, CT, MT, PLZ, JGG, CB,*  
457 *AS, IYA*

458 *All authors read and approved the final manuscript.*

459 **References**

- 460  
461 1. Locke FL, Miklos DB, Jacobson CA, et al. Axicabtagene Ciloleucel as Second-Line  
462 Therapy for Large B-Cell Lymphoma. *N Engl J Med.* 2022;386(7):640-654.  
463 2. Abramson JS, Solomon SR, Arnason JE, et al. Lisocabtagene maraleucel as second-  
464 line therapy for large B-cell lymphoma: primary analysis of phase 3 TRANSFORM  
465 study. *Blood.* 2022.  
466 3. Shah BD, Ghobadi A, Oluwole OO, et al. KTE-X19 for relapsed or refractory adult B-cell  
467 acute lymphoblastic leukaemia: phase 2 results of the single-arm, open-label,  
468 multicentre ZUMA-3 study. *Lancet.* 2021;398(10299):491-502.  
469 4. Wang M, Munoz J, Goy A, et al. KTE-X19 CAR T-Cell Therapy in Relapsed or  
470 Refractory Mantle-Cell Lymphoma. *N Engl J Med.* 2020;382(14):1331-1342.  
471 5. Raje N, Berdeja J, Lin Y, et al. Anti-BCMA CAR T-Cell Therapy bb2121 in Relapsed or  
472 Refractory Multiple Myeloma. *N Engl J Med.* 2019;380(18):1726-1737.  
473 6. Snowden JA, Sanchez-Ortega I, Corbacioglu S, et al. Indications for haematopoietic cell  
474 transplantation for haematological diseases, solid tumours and immune disorders:  
475 current practice in Europe, 2022. *Bone Marrow Transplant.* 2022;57(8):1217-1239.  
476 7. Passweg JR, Baldomero H, Chabannon C, et al. Hematopoietic cell transplantation and  
477 cellular therapy survey of the EBMT: monitoring of activities and trends over 30 years.  
478 *Bone Marrow Transplant.* 2021;56(7):1651-1664.  
479 8. Shimabukuro-Vornhagen A, Godel P, Subklewe M, et al. Cytokine release syndrome. *J*  
480 *Immunother Cancer.* 2018;6(1):56.  
481 9. Karschnia P, Jordan JT, Forst DA, et al. Clinical presentation, management, and  
482 biomarkers of neurotoxicity after adoptive immunotherapy with CAR T cells. *Blood.*  
483 2019;133(20):2212-2221.  
484 10. Sandler RD, Tattersall RS, Schoemans H, et al. Diagnosis and Management of  
485 Secondary HLH/MAS Following HSCT and CAR-T Cell Therapy in Adults; A Review of  
486 the Literature and a Survey of Practice Within EBMT Centres on Behalf of the  
487 Autoimmune Diseases Working Party (ADWP) and Transplant Complications Working  
488 Party (TCWP). *Front Immunol.* 2020;11:524.  
489 11. Hines MR, Knight TE, McNerney KO, et al. Immune Effector Cell associated  
490 Hemophagocytic Lymphohistiocytosis-like Syndrome (IEC-HS). *Transplant Cell Ther.*  
491 2023.  
492 12. Fried S, Avigdor A, Bielora B, et al. Early and late hematologic toxicity following CD19  
493 CAR-T cells. *Bone Marrow Transplant.* 2019;54(10):1643-1650.  
494 13. Rejeski K, Perez A, Sesques P, et al. CAR-HEMATOTOX: a model for CAR T-cell-  
495 related hematologic toxicity in relapsed/refractory large B-cell lymphoma. *Blood.*  
496 2021;138(24):2499-2513.  
497 14. Wudhikarn K, Pennisi M, Garcia-Recio M, et al. DLBCL patients treated with CD19 CAR  
498 T cells experience a high burden of organ toxicities but low nonrelapse mortality. *Blood*  
499 *Adv.* 2020;4(13):3024-3033.  
500 15. Dickinson MJ, Carlo-Stella C, Morschhauser F, et al. Glofitamab for Relapsed or  
501 Refractory Diffuse Large B-Cell Lymphoma. *N Engl J Med.* 2022;387(24):2220-2231.  
502 16. Thieblemont C, Phillips T, Ghesquieres H, et al. Epcoritamab, a Novel, Subcutaneous  
503 CD3xCD20 Bispecific T-Cell-Engaging Antibody, in Relapsed or Refractory Large B-  
504 Cell Lymphoma: Dose Expansion in a Phase I/II Trial. *J Clin Oncol.* 2022;JCO2201725.  
505 17. Chari A, Minnema MC, Berdeja JG, et al. Talquetamab, a T-Cell-Redirecting GPRC5D  
506 Bispecific Antibody for Multiple Myeloma. *N Engl J Med.* 2022;387(24):2232-2244.  
507 18. Moreau P, Girgis S, Goldberg JD. Teclistamab in Relapsed or Refractory Multiple  
508 Myeloma. Reply. *N Engl J Med.* 2022;387(18):1722-1723.  
509 19. Goebeler ME, Knop S, Viardot A, et al. Bispecific T-Cell Engager (BiTE) Antibody  
510 Construct Blinatumomab for the Treatment of Patients With Relapsed/Refractory Non-  
511 Hodgkin Lymphoma: Final Results From a Phase I Study. *J Clin Oncol.*  
512 2016;34(10):1104-1111.  
513 20. Hill JA, Seo SK. How I prevent infections in patients receiving CD19-targeted chimeric  
514 antigen receptor T cells for B-cell malignancies. *Blood.* 2020;136(8):925-935.  
515 21. Nastoupil LJ, Jain MD, Feng L, et al. Standard-of-Care Axicabtagene Ciloleucel for  
516 Relapsed or Refractory Large B-Cell Lymphoma: Results From the US Lymphoma CAR  
517 T Consortium. *J Clin Oncol.* 2020;38(27):3119-3128.

- 518 22. Rejeski K, Perez A, Iacoboni G, et al. The CAR-HEMATOTOX risk-stratifies patients for  
519 severe infections and disease progression after CD19 CAR-T in R/R LBCL. *J*  
520 *Immunother Cancer*. 2022;10(5).
- 521 23. Bethge WA, Martus P, Schmitt M, et al. GLA/DRST real-world outcome analysis of  
522 CAR-T cell therapies for large B-cell lymphoma in Germany. *Blood*. 2022.
- 523 24. Berdeja JG, Madduri D, Usmani SZ, et al. Ciltacabtagene autoleucl, a B-cell  
524 maturation antigen-directed chimeric antigen receptor T-cell therapy in patients with  
525 relapsed or refractory multiple myeloma (CARTITUDE-1): a phase 1b/2 open-label  
526 study. *Lancet*. 2021;398(10297):314-324.
- 527 25. Munshi NC, Anderson LD, Jr., Shah N, et al. Idecabtagene Vicleucl in Relapsed and  
528 Refractory Multiple Myeloma. *N Engl J Med*. 2021;384(8):705-716.
- 529 26. Jacobson CA, Chavez JC, Sehgal AR, et al. Axicabtagene ciloleucl in relapsed or  
530 refractory indolent non-Hodgkin lymphoma (ZUMA-5): a single-arm, multicentre, phase  
531 2 trial. *Lancet Oncol*. 2022;23(1):91-103.
- 532 27. Iacoboni G, Rejeski K, Villacampa G, et al. Real-world evidence of brexucabtagene  
533 autoleucl for the treatment of relapsed or refractory mantle cell lymphoma. *Blood Adv*.  
534 2022.
- 535 28. Maude SL, Laetsch TW, Buechner J, et al. Tisagenlecleucl in Children and Young  
536 Adults with B-Cell Lymphoblastic Leukemia. *N Engl J Med*. 2018;378(5):439-448.
- 537 29. Lichtenstein DA, Schischlik F, Shao L, et al. Characterization of HLH-like manifestations  
538 as a CRS variant in patients receiving CD22 CAR T cells. *Blood*. 2021;138(24):2469-  
539 2484.
- 540 30. Cordeiro A, Bezerra ED, Hirayama AV, et al. Late Events after Treatment with CD19-  
541 Targeted Chimeric Antigen Receptor Modified T Cells. *Biol Blood Marrow Transplant*.  
542 2020;26(1):26-33.
- 543 31. Rejeski K, Kunz WG, Rudelius M, et al. Severe *Candida glabrata* pancolitis and fatal  
544 *Aspergillus fumigatus* pulmonary infection in the setting of bone marrow aplasia after  
545 CD19-directed CAR T-cell therapy - a case report. *BMC Infect Dis*. 2021;21(1):121.
- 546 32. Jain T, Olson TS, Locke FL. How I Treat Cytopenias after CAR T-cell Therapy. *Blood*.  
547 2023.
- 548 33. Juluri KR, Wu V, Voutsinas JM, et al. Severe cytokine release syndrome is associated  
549 with hematologic toxicity following CD19 CAR T-cell therapy. *Blood Adv*. 2021.
- 550 34. Jain T, Knezevic A, Pennisi M, et al. Hematopoietic recovery in patients receiving  
551 chimeric antigen receptor T-cell therapy for hematologic malignancies. *Blood Adv*.  
552 2020;4(15):3776-3787.
- 553 35. Rejeski K, Wu Z, Blumenberg V, et al. Oligoclonal T-cell expansion in a patient with  
554 bone marrow failure after CD19 CAR-T for Richter transformed DLBCL. *Blood*. 2022.
- 555 36. Rejeski K, Greco R, Onida F, et al. An International Survey on Grading, Diagnosis, and  
556 Management of Immune Effector Cell-Associated Hematotoxicity (ICAHT) Following  
557 CAR T-cell Therapy on Behalf of the EBMT and EHA. *Hemasphere*. 2023;7(5):e889.
- 558 37. Taplitz RA, Kennedy EB, Bow EJ, et al. Antimicrobial Prophylaxis for Adult Patients  
559 With Cancer-Related Immunosuppression: ASCO and IDSA Clinical Practice Guideline  
560 Update. *J Clin Oncol*. 2018;36(30):3043-3054.
- 561 38. Yakoub-Agha I, Greco R, Onida F, et al. Practice harmonization workshops of EBMT:  
562 an expert-based approach to generate practical and contemporary guidelines within the  
563 arena of hematopoietic cell transplantation and cellular therapy. *Bone Marrow*  
564 *Transplant*. 2023.
- 565 39. Xia Y, Zhang J, Li J, et al. Cytopenias following anti-CD19 chimeric antigen receptor  
566 (CAR) T cell therapy: a systematic analysis for contributing factors. *Ann Med*.  
567 2022;54(1):2951-2965.
- 568 40. Brudno JN, Natrakul D, Lam N, Dulau-Florea A, Yuan CM, Kochenderfer JN. Acute and  
569 delayed cytopenias following CAR T-cell therapy: an investigation of risk factors and  
570 mechanisms. *Leuk Lymphoma*. 2022;63(8):1849-1860.
- 571 41. Roddie C, Neill L, Osborne W, et al. Effective bridging therapy can improve CD19 CAR-  
572 T outcomes while maintaining safety in patients with large B-cell lymphoma. *Blood Adv*.  
573 2023.
- 574 42. Rejeski K, Perez A, Iacoboni G, et al. Biphasic Neutrophil Recovery after CD19  
575 CART in R/R LBCL Is Associated with Superior PFS/OS, Robust CAR T-Cell Expansion  
576 in Relation to Baseline Tumor Volume, and a Decrease of Systemic Inflammation over  
577 Time. *Blood*. 2022;140(Supplement 1):4549-4551.

- 578 43. Rejeski K, Wang Y, Albanyan O, et al. The CAR-Hematotox Score Identifies Patients at  
579 High Risk for Hematological Toxicity, Infections and Poor Clinical Outcomes Following  
580 Brexucabtagene Autoleucel in Relapsed/Refractory Mantle Cell Lymphoma. *Blood*.  
581 2022;140(Supplement 1):651-653.
- 582 44. Rejeski K, Hansen DK, Bansal R, et al. The CAR-Hematotox Score As a Prognostic  
583 Model of Toxicity and Response in Patients Receiving BCMA-Directed CAR-T for  
584 Relapsed/Refractory Multiple Myeloma. *Blood*. 2022;140(Supplement 1):7506-7508.
- 585 45. Liu Y, Derkach A, Lewis N, et al. Clonal hematopoiesis in diffuse large B-cell lymphoma:  
586 clinical impact and genetic relatedness to lymphoma and therapy-related myeloid  
587 neoplasm. *Haematologica*. 2023;108(3):917-922.
- 588 46. Saini NY, Swoboda DM, Greenbaum U, et al. Clonal Hematopoiesis Is Associated with  
589 Increased Risk of Severe Neurotoxicity in Axicabtagene Ciloleucel Therapy of Large B-  
590 Cell Lymphoma. *Blood Cancer Discov*. 2022;3(5):385-393.
- 591 47. Miller PG, Sperling AS, Brea EJ, et al. Clonal hematopoiesis in patients receiving  
592 chimeric antigen receptor T-cell therapy. *Blood Adv*. 2021;5(15):2982-2986.
- 593 48. Jain MD, Zhao H, Wang X, et al. Tumor interferon signaling and suppressive myeloid  
594 cells are associated with CAR T-cell failure in large B-cell lymphoma. *Blood*.  
595 2021;137(19):2621-2633.
- 596 49. Bachy E, Le Gouill S, Di Blasi R, et al. A real-world comparison of tisagenlecleucel and  
597 axicabtagene ciloleucel CAR T cells in relapsed or refractory diffuse large B cell  
598 lymphoma. *Nat Med*. 2022;28(10):2145-2154.
- 599 50. Luo W, Li C, Zhang Y, et al. Adverse effects in hematologic malignancies treated with  
600 chimeric antigen receptor (CAR) T cell therapy: a systematic review and Meta-analysis.  
601 *BMC Cancer*. 2022;22(1):98.
- 602 51. Kawalekar OU, RS OC, Fraietta JA, et al. Distinct Signaling of Coreceptors Regulates  
603 Specific Metabolism Pathways and Impacts Memory Development in CAR T Cells.  
604 *Immunity*. 2016;44(3):712.
- 605 52. Zhao Z, Condomines M, van der Stegen SJC, et al. Structural Design of Engineered  
606 Costimulation Determines Tumor Rejection Kinetics and Persistence of CAR T Cells.  
607 *Cancer Cell*. 2015;28(4):415-428.
- 608 53. Wang Y, Song Z, Geng Y, et al. The risk factors and early predictive model of  
609 hematotoxicity after CD19 chimeric antigen receptor T cell therapy. *Front Oncol*.  
610 2022;12:987965.
- 611 54. Li X, Deng Q, Henderson J, et al. Targetable Cellular Etiology of Prolonged Cytopenia  
612 Following CD19 CAR T-Cell Therapy. *Blood*. 2022;140(Supplement 1):4502-4503.
- 613 55. de Bruin AM, Demirel O, Hooibrink B, Brandts CH, Nolte MA. Interferon-gamma impairs  
614 proliferation of hematopoietic stem cells in mice. *Blood*. 2013;121(18):3578-3585.
- 615 56. Morales-Mantilla DE, King KY. The Role of Interferon-Gamma in Hematopoietic Stem  
616 Cell Development, Homeostasis, and Disease. *Curr Stem Cell Rep*. 2018;4(3):264-271.
- 617 57. Rejeski K, Blumenberg V, Iacoboni G, et al. Identifying Early Infections in the Setting of  
618 CRS With Routine and Exploratory Serum Proteomics and the HT10 Score Following  
619 CD19 CAR-T for Relapsed/Refractory B-NHL. *Hemasphere*. 2023;7(4):e858.
- 620 58. Neelapu SS, Tummala S, Kebriaei P, et al. Chimeric antigen receptor T-cell therapy -  
621 assessment and management of toxicities. *Nat Rev Clin Oncol*. 2018;15(1):47-62.
- 622 59. Hayden PJ, Roddie C, Bader P, et al. Management of adults and children receiving  
623 CAR T-cell therapy: 2021 best practice recommendations of the European Society for  
624 Blood and Marrow Transplantation (EBMT) and the Joint Accreditation Committee of  
625 ISCT and EBMT (JACIE) and the European Haematology Association (EHA). *Ann*  
626 *Oncol*. 2022;33(3):259-275.
- 627 60. Hines MR, Knight TE, McNerney KO, et al. Immune Effector Cell associated  
628 Hemophagocytic Lymphohistiocytosis-like Syndrome (IEC-HS). *Transplantation and*  
629 *Cellular Therapy* 2023.
- 630 61. Cutini I, Puccini B, Fabbri A, et al. Late haemophagocytic lymphohistiocytosis in a  
631 patient treated with Axicabtagene ciloleucel. *Transpl Immunol*. 2022;75:101719.
- 632 62. Henter JI, Horne A, Arico M, et al. HLH-2004: Diagnostic and therapeutic guidelines for  
633 hemophagocytic lymphohistiocytosis. *Pediatr Blood Cancer*. 2007;48(2):124-131.
- 634 63. Fardet L, Galicier L, Lambotte O, et al. Development and validation of the HScore, a  
635 score for the diagnosis of reactive hemophagocytic syndrome. *Arthritis Rheumatol*.  
636 2014;66(9):2613-2620.

- 637 64. Zoref-Lorenz A, Murakami J, Hofstetter L, et al. An improved index for diagnosis and  
638 mortality prediction in malignancy-associated hemophagocytic lymphohistiocytosis.  
639 *Blood*. 2022;139(7):1098-1110.
- 640 65. McNerney KO, DiNofia AM, Teachey DT, Grupp SA, Maude SL. Potential Role of  
641 IFN $\gamma$  Inhibition in Refractory Cytokine Release Syndrome Associated with CAR  
642 T-cell Therapy. *Blood Cancer Discov*. 2022;3(2):90-94.
- 643 66. Rainone M, Ngo D, Baird JH, et al. Interferon-gamma blockade in CAR T-cell therapy-  
644 associated macrophage activation syndrome/hemophagocytic lymphohistiocytosis.  
645 *Blood Adv*. 2023;7(4):533-536.
- 646 67. La Rosee P, Horne A, Hines M, et al. Recommendations for the management of  
647 hemophagocytic lymphohistiocytosis in adults. *Blood*. 2019;133(23):2465-2477.
- 648 68. Kopolovic I, Ostro J, Tsubota H, et al. A systematic review of transfusion-associated  
649 graft-versus-host disease. *Blood*. 2015;126(3):406-414.
- 650 69. Foukaneli T, Kerr P, Bolton-Maggs PHB, et al. Guidelines on the use of irradiated blood  
651 components. *Br J Haematol*. 2020;191(5):704-724.
- 652 70. Giavridis T, van der Stegen SJC, Eyquem J, Hamieh M, Piersigilli A, Sadelain M. CAR  
653 T cell-induced cytokine release syndrome is mediated by macrophages and abated by  
654 IL-1 blockade. *Nat Med*. 2018;24(6):731-738.
- 655 71. Sterner RM, Sakemura R, Cox MJ, et al. GM-CSF inhibition reduces cytokine release  
656 syndrome and neuroinflammation but enhances CAR-T cell function in xenografts.  
657 *Blood*. 2019;133(7):697-709.
- 658 72. Barreto JN, Bansal R, Hathcock MA, et al. The impact of granulocyte colony stimulating  
659 factor on patients receiving chimeric antigen receptor T-cell therapy. *Am J Hematol*.  
660 2021;96(10):E399-E402.
- 661 73. Galli E, Allain V, Di Blasi R, et al. G-CSF does not worsen toxicities and efficacy of  
662 CAR-T cells in refractory/relapsed B-cell lymphoma. *Bone Marrow Transplant*.  
663 2020;55(12):2347-2349.
- 664 74. Lievin R, Di Blasi R, Morin F, et al. Effect of early granulocyte-colony-stimulating factor  
665 administration in the prevention of febrile neutropenia and impact on toxicity and  
666 efficacy of anti-CD19 CAR-T in patients with relapsed/refractory B-cell lymphoma. *Bone  
667 Marrow Transplant*. 2022.
- 668 75. Miller KC, Johnson PC, Abramson JS, et al. Effect of granulocyte colony-stimulating  
669 factor on toxicities after CAR T cell therapy for lymphoma and myeloma. *Blood Cancer  
670 J*. 2022;12(10):146.
- 671 76. Ma S, Li H, Zhou D, et al. Associations of granulocyte colony-stimulating factor with  
672 toxicities and efficacy of chimeric antigen receptor T-cell therapy in relapsed or  
673 refractory multiple myeloma. *Cytotherapy*. 2023.
- 674 77. Baur R, Jitschin R, Kharboutli S, et al. Thrombopoietin receptor agonists for acquired  
675 thrombocytopenia following anti-CD19 CAR-T-cell therapy: a case report. *J Immunother  
676 Cancer*. 2021;9(7).
- 677 78. Beyar-Katz O, Perry C, On YB, et al. Thrombopoietin receptor agonist for treating bone  
678 marrow aplasia following anti-CD19 CAR-T cells-single-center experience. *Ann  
679 Hematol*. 2022;101(8):1769-1776.
- 680 79. Drillet G, Lhomme F, De Guibert S, Manson G, Houot R. Prolonged thrombocytopenia  
681 after CAR T-cell therapy: the role of thrombopoietin receptor agonists. *Blood Adv*.  
682 2023;7(4):537-540.
- 683 80. Drexler B, Passweg J. Current evidence and the emerging role of eltrombopag in  
684 severe aplastic anemia. *Ther Adv Hematol*. 2021;12:2040620721998126.
- 685 81. Peffault de Latour R, Kulasekararaj A, Iacobelli S, et al. Eltrombopag Added to  
686 Immunosuppression in Severe Aplastic Anemia. *N Engl J Med*. 2022;386(1):11-23.
- 687 82. Bento L, Bastida JM, Garcia-Cadenas I, et al. Thrombopoietin Receptor Agonists for  
688 Severe Thrombocytopenia after Allogeneic Stem Cell Transplantation: Experience of  
689 the Spanish Group of Hematopoietic Stem Cell Transplant. *Biol Blood Marrow  
690 Transplant*. 2019;25(9):1825-1831.
- 691 83. Ball S, Brown TJ, Das A, Khera R, Khanna S, Gupta A. Effect of Neutropenic Diet on  
692 Infection Rates in Cancer Patients With Neutropenia: A Meta-analysis of Randomized  
693 Controlled Trials. *Am J Clin Oncol*. 2019;42(3):270-274.
- 694 84. Sonbol MB, Jain T, Firwana B, et al. Neutropenic diets to prevent cancer infections:  
695 updated systematic review and meta-analysis. *BMJ Support Palliat Care*.  
696 2019;9(4):425-433.

- 697 85. Stella F, Marasco V, Levati GV, et al. Non-Restrictive Diet Does Not Increase Infections  
698 in Patients with Neutropenia after Stem Cell Transplantation: Final Analysis of the  
699 Neurodiet Multicenter, Randomized Trial. *Blood*. 2022;140(Supplement 1):417-419.
- 700 86. Teipel R, Kroschinsky F, Kramer M, et al. Prevalence and variation of CHIP in patients  
701 with aggressive lymphomas undergoing CD19-directed CAR T-cell treatment. *Blood*  
702 *Adv*. 2022;6(6):1941-1946.
- 703 87. Spellberg B, Doi Y. The Rise of Fluoroquinolone-Resistant *Escherichia coli* in the  
704 Community: Scarier Than We Thought. *J Infect Dis*. 2015;212(12):1853-1855.
- 705 88. Lautenbach E, Metlay JP, Bilker WB, Edelstein PH, Fishman NO. Association between  
706 fluoroquinolone resistance and mortality in *Escherichia coli* and *Klebsiella pneumoniae*  
707 infections: the role of inadequate empirical antimicrobial therapy. *Clin Infect Dis*.  
708 2005;41(7):923-929.
- 709 89. Trecarichi EM, Tumbarello M, Spanu T, et al. Incidence and clinical impact of extended-  
710 spectrum-beta-lactamase (ESBL) production and fluoroquinolone resistance in  
711 bloodstream infections caused by *Escherichia coli* in patients with hematological  
712 malignancies. *J Infect*. 2009;58(4):299-307.
- 713 90. Bow EJ. Fluoroquinolones, antimicrobial resistance and neutropenic cancer patients.  
714 *Curr Opin Infect Dis*. 2011;24(6):545-553.
- 715 91. Schubert ML, Rohrbach R, Schmitt M, Stein-Thoeringer CK. The Potential Role of the  
716 Intestinal Microenvironment and Individual Microbes in the Immunobiology of Chimeric Antigen  
717 Receptor T-Cell Therapy. *Front Immunol*. 2021;12:670286.
- 718 92. Blumenberg V, Busch G, Baumann S, et al. High Bacterial Abundances of *Dorea* and  
719 *Pediococcus* in the Gut Microbiome Linked to Expansion, Immune Checkpoint  
720 Expression and Efficacy of CD19-Directed CAR T-Cells in Patients with r/r DLBCL.  
721 *Blood*. 2021;138(Supplement 1):2792-2792.
- 722 93. Smith M, Dai A, Ghilardi G, et al. Gut microbiome correlates of response and toxicity  
723 following anti-CD19 CAR T cell therapy. *Nat Med*. 2022;28(4):713-723.
- 724 94. Stein-Thoeringer CK, Saini NY, Zamir E, et al. A non-antibiotic-disrupted gut  
725 microbiome is associated with clinical responses to CD19-CAR-T cell cancer  
726 immunotherapy. *Nat Med*. 2023.
- 727 95. Little JS, Aleissa MM, Beluch K, et al. Low incidence of invasive fungal disease  
728 following CD19 chimeric antigen receptor T-cell therapy for non-Hodgkin lymphoma.  
729 *Blood Adv*. 2022;6(16):4821-4830.
- 730 96. Haidar G, Dorritie K, Farah R, Bogdanovich T, Nguyen MH, Samanta P. Invasive Mold  
731 Infections After Chimeric Antigen Receptor-Modified T-cell Therapy: A Case Series,  
732 Review of the Literature, and Implications for Prophylaxis. *Clin Infect Dis*. 2019.
- 733 97. Rejeski K, Burchert A, Iacoboni G, et al. Safety and feasibility of stem cell boost as a  
734 salvage therapy for severe hematotoxicity after CD19 CAR T-cell therapy. *Blood Adv*.  
735 2022;6(16):4719-4725.
- 736 98. Mullanfiroze K, Lazareva A, Chu J, et al. CD34+-selected stem cell boost can safely  
737 improve cytopenias following CAR T-cell therapy. *Blood Adv*. 2022;6(16):4715-4718.
- 738 99. Gagelmann N, Wulf GG, Duell J, et al. Hematopoietic stem cell boost for persistent  
739 neutropenia after CAR T-cell therapy: a GLA/DRST study. *Blood Adv*. 2023;7(4):555-  
740 559.
- 741 100. Lipsitt A, Beattie L, Harstead E, et al. Allogeneic CD34(+) selected hematopoietic stem  
742 cell boost following CAR T-cell therapy in a patient with prolonged cytopenia and active  
743 infection. *Pediatr Blood Cancer*. 2023;70(3):e30166.
- 744 101. Chhabra S, Thapa B, Szabo A, et al. Utilization and Cost Implications of Hematopoietic  
745 Progenitor Cells Stored for a Future Salvage Autologous Transplantation or Stem Cell  
746 Boost in Myeloma Patients. *Biol Blood Marrow Transplant*. 2020;26(11):2011-2017.
- 747 102. Liang EC, Muffly LS, Shiraz P, et al. Use of Backup Stem Cells for Stem Cell Boost and  
748 Second Transplant in Patients with Multiple Myeloma Undergoing Autologous Stem Cell  
749 Transplantation. *Transplant Cell Ther*. 2021;27(5):405 e401-405 e406.
- 750 103. Logue JM, Peres LC, Hashmi H, et al. Early cytopenias and infections after standard of  
751 care idacabtagene vicleucel in relapsed or refractory multiple myeloma. *Blood Adv*.  
752 2022;6(24):6109-6119.
- 753 104. Logue JM, Zucchetti E, Bachmeier CA, et al. Immune reconstitution and associated  
754 infections following axicabtagene ciloleucel in relapsed or refractory large B-cell  
755 lymphoma. *Haematologica*. 2020.



- 756 105. Alizadeh D, Wong RA, Yang X, et al. IL15 Enhances CAR-T Cell Antitumor Activity by  
757 Reducing mTORC1 Activity and Preserving Their Stem Cell Memory Phenotype.  
758 *Cancer Immunol Res.* 2019;7(5):759-772.
- 759 106. Pascutti MF, Erkelens MN, Nolte MA. Impact of Viral Infections on Hematopoiesis: From  
760 Beneficial to Detrimental Effects on Bone Marrow Output. *Front Immunol.* 2016;7:364.
- 761 107. Porter TJ, Lazarevic A, Ziggas JE, et al. Hyperinflammatory syndrome resembling  
762 haemophagocytic lymphohistiocytosis following axicabtagene ciloleucel and  
763 brexucabtagene autoleucel. *Br J Haematol.* 2022;199(5):720-727.
- 764

765 **Main Tables and Table Legends**766 **Table 1: ICAHT Grading**

Grading	I	II	III	IV
<i>Early ICAHT (day 0-30)</i>				
<b>ANC ≤ 500/μL</b>	<7 days	7-13 days	≥14 days	Never above 500/μL
<b>ANC ≤ 100/μL</b>	-	-	≥7 days	≥14 days
<i>Late ICAHT (after day +30)*</i>				
<b>ANC ≤ 1500/μL</b>				
<b>ANC ≤ 1000/μL</b>				
<b>ANC ≤ 500/μL</b>				
<b>ANC ≤ 100/μL</b>				

767 \*measured ≥2 time points, or non-transient neutropenia

768

769 **Table 2: Risk factors associated with an increased risk of post-CAR-T cytopenias**

	Risk Factors	Comments	References
<b>Disease-related features</b>	Underlying disease (ALL > B-NHL)	Evidence concerning the rate of cytopenias in multiple myeloma patients still emerging	Xia et al. <sup>39</sup>
	Disease burden at CAR-T infusion (progressive disease, high LDH)	Specially BM disease burden	Wudhikarn et al. <sup>14</sup> Logue et al. <sup>103</sup>
<b>Prior therapies</b>	Number of prior therapy lines	Associated with baseline hematopoietic function	Xia et al. <sup>39</sup>
	Prior hematopoietic stem cell transplantation (HSCT)		Fried et al. <sup>105</sup>
	Bridging Therapy		Roddie et al. <sup>41</sup>
<b>Baseline Marrow Status</b>	Bone marrow infiltration		Rejeski et al. <sup>42</sup> Brudno et al. <sup>40</sup>
	Pre-existing cytopenias	Particularly pre-existing thrombocytopenia	Rejeski et al. <sup>13</sup> Juluri et al. <sup>33</sup>
	Clonal hematopoiesis of indeterminate potential (CHIP)?	Has been linked to increased inflammation, potential emerging risk factor	Saini et al. <sup>46</sup> Miller et al. <sup>47</sup> Teipel et al. <sup>86</sup>
<b>Baseline Inflammatory Status</b>	Increased Serum CRP		Rejeski et al. <sup>13</sup>
	Increased Serum Ferritin		Rejeski et al. <sup>13</sup>
<b>CAR-T Product and post-infusion risk factors</b>	Co-stimulatory molecule (CD28>41BB)	May also reflect differences in lymphodepletion dosing (cyclophosphamide dosing)	Xia et al. <sup>39</sup>
	Type of construct (Tandem > single target)		Xia et al. <sup>39</sup>
	Severe CRS		Juluri et al. <sup>33</sup> Jain et al. <sup>34</sup>
	Sustained increased inflammatory markers		Juluri et al. <sup>33</sup>
	Oligoclonal T-cell expansion	In select patients; the success of auto-HCT boost argues against this as a general mechanism	Rejeski et al. <sup>35</sup>
	Active Infection	Mainly viral or in case of concomitant sepsis	Pascutti et al. <sup>106</sup>
	CRS/MAS or IEC-HS	Cytopenia as overlapping symptomology	Sandler et al. <sup>10</sup> Hines et al. <sup>11</sup> Porter et al. <sup>107</sup>

770 **Table 3: Short-term management of cytopenias**

	When	How	Precautions	Comments
<b>Packed red blood cell (pRBC)/ Platelet transfusions</b>	As per institutional standards, based on patient risk profile	As per institutional standards  For pRBC: consider using 1 product per time to reduce iron overload <sup>69</sup>	Irradiation of blood products; Start 7 days prior to leukapheresis until at least 90 days post CAR-T	Due to the use of fludarabine
<b>G-CSF</b>	<b>Prophylactic G-CSF:</b> On day +2 in patients with a high-risk profile for ICAHT (e.g. high CAR-HEMATOTOX score and risk profile according to <b>Table 2</b> )	Based on individual risk profile: Consider early G-CSF administration (from day +2) as prophylaxis in high risk for ICAHT  Dosing: 5 µg/kg once daily	In patients at low risk for ICAHT, G-CSF probably not necessary*	Reduced risk of febrile neutropenia (without increasing the risk of severe, or grade ≥3, CRS nor ICANS).  No detrimental effect on CAR-T expansion kinetics or treatment outcomes <sup>74,75</sup>
	<b>Therapeutic G-CSF:</b> Severe neutropenia (ANC <500/µL) neutropenia with or without infectious complications	In case of prolonged neutropenia with/without infectious complications.  Dosing: 5 µg/kg once daily, consider increasing dose in case of non-response		Patients with intermittent neutrophil recovery often rapidly respond to G-CSF stimulation, while aplastic patients are often G-CSF unresponsive
<b>Antibacterial prophylaxis</b>	In patients with a low risk for ICAHT, not recommended.  In patients with a high-risk profile for ICAHT, prophylaxis may be considered once ANC <500/µL.	As per institutional standards (e.g. levofloxacin or ciprofloxacin).	Warning in case of colonization by MDR pathogens.	Look at local bacterial epidemiology. High local prevalence of MDR GNB might prevent the use of antibacterial prophylaxis
<b>Anti-viral</b>	All patients	Start from LD conditioning until 1-year post-CAR T-cell infusion AND/OR until CD4+ count >0.2 × 10 <sup>9</sup> /l  Valaciclovir 500 mg bid or aciclovir 800 mg bid		
<b>Anti-pneumocystis</b>	All patients	To start from LD conditioning until 1-year post-CAR-T cell infusion AND/OR until CD4+ count >0.2x10 <sup>9</sup> G/l  Co-trimoxazole 480 mg once daily or 960 mg three times each week	In case of co-trimoxazole allergy, pentamidine inhalation (300 mg once every month), dapsonsone 100 mg daily or atovaquone 1500 mg once daily can be considered	Can be started later depending on center guidelines
<b>Systemic primary anti-fungal prophylaxis</b>	Prophylaxis may be considered in severe neutropenia (ANC <500) with a high-risk profile for ICAHT (e.g. CAR HEMATOTOX score and risk profile according to <b>Table 2</b> ) and/or prolonged neutropenia	Mold-active prophylaxis for 1-3 months (depending on the duration of neutropenia and use of steroids): posaconazole (300 mg/day) or micafungin (50 mg i.v./day)		In patients with prior allo-HCT, prior invasive aspergillosis and those receiving corticosteroids (long-term >72 h, or high-dose), prophylaxis is recommended

771

772 **Figure Legends**773 **Figure 1. The CAR-HEMATOTOX score as a risk-stratification tool**

774

775 **Figure 2. Step-by-step diagnostic work-up depending on ICAHT severity**776 \* In case of elevated ferritin and clinical suspicion of MAS, see **Table S3** and **Figure S1**

777

778 **Figure 3. Treatment algorithm for Immune Effector Cell Associated Neutropenia**779 \*High-risk defined as prior history of hematopoietic stem cell transplantation, baseline cytopenia,  
780 high tumor burden and systemic inflammation, presence of BM infiltration.781 \*\*Anti-fungal prophylaxis particularly recommended in patients with prior IFD, prior allo-HCT, and  
782 receiving corticosteroids (long-term >72h or high-dose). Decision for/against anti-bacterial  
783 prophylaxis should incorporate local bacterial epidemiology (e.g. prevalence for MDR GNB); not  
784 recommended for patients with a low-risk profile for ICAHT

785 † Also extends to late ICAHT if these criteria are met

# Figure 1

Prior to lymphodepleting chemotherapy (day -5)

➔ Determine patient-individual risk of heme-tox and infections using the **CAR-HEMATOTOX score**

- Leniency time period for lab values: 3 days

Features	0 Point	1 Point	2 Points
Platelet Count	> 175.000/μl	75.000 - 175.000/μl	< 75.000/μl
Absolute Neutrophil Count (ANC)	> 1200/μl	< 1200/μl	-
Hemoglobin	> 9.0 g/dl	< 9.0 g/dl	-
C-reactive protein (CRP)	< 3.0 mg/dl	> 3.0 mg/dl	-
Ferritin	< 650 ng/ml	650-2000 ng/ml	> 2000 ng/ml

Low: 0-1 High: ≥ 2

Low Risk (HT 0-1)

High Risk (HT 2-7)

Risk Profile

	LBCL (n=235)	MCL (n=103)	MM (n=113)
Median duration of severe neutropenia (ANC<500/μL, D0-60)	5.5 days (95% CI 5-8 days)	6 days (95% CI 5-7 days)	3 days (95% CI 2-5 days)
Aplastic Phenotype	2.6%	0%	3%
Severe Infection Rate	8%	5%	5%
Severe Bacterial Infection Rate	0.9%	5%	3%

	LBCL (n=235)	MCL (n=103)	MM (n=113)
Duration of severe neutropenia (ANC <500/μL day 0-60)	12 days (95% CI 10-16 days)	14 days (95% CI 9-18 days)	9 days (95% CI 7-13 days)
Aplastic Phenotype	36%	47%	32%
Severe Infection Rate	40%	30%	40%
Severe Bacterial Infection Rate	27%	28%	34%

**Figure 2**

	Categories	Putative causes	Test	Time points	Comments from expert panel
<b>TIER 1</b>	<b>Lower threshold to perform – minimal workup</b>				
	<b>Poor bone marrow reserve</b>	Prior treatments including allo-HCT, fludarabine, marrow infiltration	Complete blood count (CBC), reticulocyte production index (RPI), peripheral blood smear	routinely	Recommended
	<b>Medication – drug side effects</b>	Check for concomitant myelosuppressive medications		routinely	
	<b>Vitamin deficiencies</b>	Vitamin B12, Folic acid	Serum levels	routinely	Recommended
	<b>Rule out infections</b>	Bacterial/ Viral/Fungal infections	Blood cultures, CMV PCR, Procalcitonin CD4+ T-cell, IgG, B-cell levels	routinely	Recommended
	<b>Rule out macrophage-activation syndrome*</b>	CRS/MAS or IEC-HS	Ferritin, triglycerides	routinely	Recommended
<b>TIER 2</b>	<b>Subsequent work-up – In case of G-CSF refractory state, if tier 1 results are negative and/or risk factors are present</b>				
	<b>Viral PCR considering the clinical presentation</b>	Parvovirus	Parvovirus B19 PCR	In case of prolonged anemia	Recommended
		HHV6, JC	HHV6, JC PCR blood/CSF	In case of neurologic symptoms	Recommended
		EBV, adeno, HSV	PCR	In case of HLH	Recommended
	<b>Bone marrow disease</b>	(MDS/AML/myelofibrosis) or relapse	BM aspirate, biopsy, Flow cytometry, immunohistochemistry, cytogenetics, NGS	In case of prolonged cytopenia	Recommended
		Relapse of leukemia/lymphoma	Flow cytometry peripheral blood / bone marrow, With B-cell panel	routinely	Recommended
	<b>Other causes</b>	Other rare hematologic diseases, myeloid diseases, PNH, autoimmune processes	Myeloid panel, PI-linked structures, Direct Antiglobulin Test (DAT)	In case of suspected MPN/PNH/ autoimmune processes	Recommended

**Figure 3**

