



**HAL**  
open science

# The application of a time exponential integrator to the wave equations, oriented to seismic imaging

Fernando Valdés, Pedro Peixoto, Martin Schreiber

## ► To cite this version:

Fernando Valdés, Pedro Peixoto, Martin Schreiber. The application of a time exponential integrator to the wave equations, oriented to seismic imaging. LACIAM 2023 - First Latin American Congress of Industrial and Applied Mathematics, Jan 2023, Valparaíso, Chile. pp.1-1. hal-04215838

**HAL Id: hal-04215838**

**<https://hal.science/hal-04215838>**

Submitted on 22 Sep 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License




# The application of a time exponential integrator to the wave equations, oriented to seismic imaging

Fernando Valdés\*, Pedro Peixoto & Martin Schreiber  
Institute of Mathematics and Statistic of the University of São Paulo  
fernandr@ime.usp.br



## Introduction

The production of images from the subsurface using elastic waves is an important and challenging problem in geophysics [1]. A key stage is the computation of the solution of the wave propagation equations with absorbing boundary conditions (ABC). Encouraging a continuously development of innovative numerical methods to approximate its solution, trying to increase the accuracy and efficiency.



A particular class of methods, known as time exponential integrators, have been shown to outperform classical schemes in several physics-related differential equations [2]. However, their application in hyperbolic systems with ABC, like the ones arising in seismic imaging, still lacks theoretical and experimental investigations. For these equations, we study one of the most promising exponential integrators schemes, which is based on Faber polynomials.

## Background

Exponential integrators are a class of time integrating methods used to solve ordinary differential equations of first order in time,

$$\frac{du(t)}{dt} = Hu(t) + f(t, u(t)), \quad u(t_0) = u_0.$$

where  $u(t), u_0 \in \mathbb{C}^n$ ,  $H \in \mathbb{C}^{n \times n}$ ,  $f: \mathbb{R} \times \mathbb{C}^n \rightarrow \mathbb{C}^n$ ; through the approximation of the semi-analytic solution of the constants variation formula,

$$u(t) = e^{(t-t_0)H}u_0 + \int_{t_0}^t e^{(t-\tau)H}f(\tau, u(\tau))d\tau. \quad (1)$$

Eq. (1) can be further transformed to the expression

$$u(t) = [I_{n \times n} \ 0] e^{(t-t_0)H} \begin{bmatrix} u_0 \\ e_p \end{bmatrix},$$

where  $e_p \in \mathbb{R}^p$  is a canonic vector with one in its last element.

**Faber polynomials:** given a degree  $j$ , and a square matrix  $H$ , Faber's polynomials are defined as  $F_j(H)$ , with

$$\begin{aligned} F_0(H) &= I_{n \times n}, & F_1(H) &= H/\gamma - c_0 I_{n \times n}, \\ F_2(H) &= F_1(H)F_1(H) - 2c_1 I_{n \times n}, \\ F_j(H) &= F_1(H)F_{j-1}(H) - c_1 F_{j-2}(H), & j &\geq 3, \end{aligned}$$

with

$$\gamma = (a+b)/2, \quad c_0 = d/\gamma, \quad c_1 = c_f^2/(4\gamma^2),$$

where the parameters  $a, b, c_f$ , and  $d$ , are set according to the spectrum of the operator  $H$ .


Acoustic wave equations with ABC in two dimensions  $((x, y) \in \Omega = [a_1, a_2] \times [b_1, b_2], t > t_0)$

$$\frac{\partial}{\partial t} \begin{pmatrix} u \\ v \\ w_x \\ w_y \end{pmatrix} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -\beta_x \beta_y + c^2 \left( \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) & -(\beta_x + \beta_y) & c^2 \frac{\partial}{\partial x} & c^2 \frac{\partial}{\partial y} \\ (\beta_y - \beta_x) \frac{\partial}{\partial x} & 0 & -\beta_x & 0 \\ (\beta_x - \beta_y) \frac{\partial}{\partial y} & 0 & 0 & -\beta_y \end{pmatrix} \begin{pmatrix} u \\ v \\ w_x \\ w_y \end{pmatrix} + \begin{pmatrix} 0 \\ f \\ 0 \\ 0 \end{pmatrix},$$


where  $f(t) = (1 - f_0^2 \pi^2 (t - t_0)^2) e^{-f_0^2 \pi^2 (t - t_0)^2}$  is a Ricker wavelet in time.

## Von Neumann stability and dispersion

Stability and dispersion analysis for the acoustic 2D equations with PML in a homogeneous medium using different spatial discretization orders and equations formulations.




(a) CFL number,  $c_{CFL} = c \Delta t_{max} / \Delta x$ .



(b) Maximum  $\alpha_{max} = c \Delta t_{max} / \Delta x$ , with phase change less than  $10^{-5}$ .


High polynomial degrees render larger  $c_{CFL}$  and  $\alpha_{max}$ .

## Corner model



When the degree of the Faber polynomial (FA) increases, the error calculated over the black line in Subfigure (e) diminishes by orders of magnitude (Subfigure (f)).

## Convergence and efficiency



Higher polynomial degrees allows larger time steps, and they are more efficient than using low degrees with small time-steps.

## Conclusions and future research

- The exponential integrator method based on Faber polynomials can be used to obtain accurate solutions of the wave equations with PML.
- When higher polynomials degrees are used there is an improvement in the stability and dispersion of the scheme.
- Higher degrees permit selecting larger time steps, without losing accuracy, and this strategy is computationally more efficient.

In the future...

- We will compare this method with other time exponential integrators, high-order Runge-Kuttas with strong stability preserving properties, and lower order schemes.
- We will use realistic velocity fields to asses the performance of the methods in real-life scenarios.

## References

- [1] Ikelle LT, Amundsen L (2018). *Introduction to petroleum seismology*. Society of Exploration Geophysicists.
- [2] Loffeld J, Tokman M (2013). *Comparative performance of exponential, implicit, and explicit integrators for stiff systems of odes*. Journal of Computational and Applied Mathematics 241:45-67

**Acknowledgment** This research was carried out in association with the ongoing RD project registered as ANP 20714-2, Software technologies for modelling and inversion, with applications in seismic imaging (University of Sao Paulo / Shell Brasil / ANP)- Desenvolvimento de tecnicas numericas e software para problemas de inversao com aplicacoes em processamento sismico, sponsored by Shell Brasil under the ANP RD levy as "Compromisso de Investimentos com Pesquisa e Desenvolvimento".