



**HAL**  
open science

## Impact of meteorological forcing on seasonal phytoplankton succession in a peri-urban lake

Nicolas Clercin, Brigitte Vinçon-Leite, Francesco Piccioni, Philippe Dubois,  
Mohamed Saad

► **To cite this version:**

Nicolas Clercin, Brigitte Vinçon-Leite, Francesco Piccioni, Philippe Dubois, Mohamed Saad. Impact of meteorological forcing on seasonal phytoplankton succession in a peri-urban lake. Symposium for European Freshwater Sciences - SEFS12, European Freshwater Science Association, Jul 2021, Dublin ( virtual ), Ireland. hal-04216130

**HAL Id: hal-04216130**

**<https://hal.science/hal-04216130>**

Submitted on 23 Sep 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Impact of meteorological forcing on seasonal phytoplankton succession in a peri-urban lake

Nicolas Clercin<sup>1</sup>, Brigitte Vinçon-Leite<sup>1</sup>, Francesco Piccioni<sup>1</sup>, Philippe Dubois<sup>1</sup>, Mohamed Saad<sup>1</sup>

<sup>1</sup> Leesu, Ecole des Ponts, Univ Paris Est Creteil, Marne-la-Vallée, France

## Sessions

RS22 Urban freshwaters

RS11 Fundamental and applied freshwater (rivers, lakes & wetlands) ecology

## Abstract

Located in the Great Paris peri-urban area (France), Lake Champs-sur-Marne is a former shallow sandpit whose water source is primarily groundwater and now, a eutrophic recreational water body. The water quality must comply with European Union bathing water directive. The monitoring of water quality has been set up since the late 2000's with the deployment of autonomous submersible sondes recording for physics, biology and chemistry parameters whereas phytoplankton identification and enumeration have been implemented since April 2019. Unlike phosphorus, nitrogen was found to be the limiting nutrient for substantial algal growth. During the summer 2019, a massive bloom of nitrogen-fixing cyanobacteria *Aphanizomenon flos-aquae* was observed in parallel with nitrogen depletion in water. Although the same pattern of nitrogen was observed in 2020, no cyanobacteria bloomed. Despite similar average air temperatures between summers 2019 and 2020, the lack of a strong thermal stratification was observed throughout the summer 2020. This phenomenon led to a dominance of chlorophycean and later cryptophycean species with total algal biovolumes much lower than those developed by cyanobacteria the previous year. As commonly assumed, global warming will increase water temperatures, strengthen water column thermal stratifications and subsequently lead to more frequent and more severe cyanobacterial blooms. However, complex interplays between meteorological conditions and phytoplankton traits may result in different species assemblages, potentially outcompeting cyanobacteria.