



HAL
open science

Macdonald duality and the proof of the quantum Q-system conjecture

Philippe Di Francesco, Rinat Kedem

► **To cite this version:**

Philippe Di Francesco, Rinat Kedem. Macdonald duality and the proof of the quantum Q-system conjecture. 2023. hal-04216442

HAL Id: hal-04216442

<https://hal.science/hal-04216442>

Preprint submitted on 24 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MACDONALD DUALITY AND THE PROOF OF THE QUANTUM Q-SYSTEM CONJECTURE

PHILIPPE DI FRANCESCO AND RINAT KEDEM

ABSTRACT. The $SL(2, \mathbb{Z})$ -symmetry of Cherednik’s spherical double affine Hecke algebras in Macdonald theory includes a distinguished generator which acts as a discrete time evolution of Macdonald operators, which can also be interpreted as a torus Dehn twist in type A . We prove for all twisted and untwisted affine algebras of type $ABCD$ that the time-evolved q -difference Macdonald operators, in the $t \rightarrow \infty$ q -Whittaker limit, form a representation of the associated discrete integrable quantum Q-systems, which are obtained, in all but one case, via the canonical quantization of suitable cluster algebras. The proof relies strongly on the duality property of Macdonald and Koornwinder polynomials, which allows, in the q -Whittaker limit, for a unified description of the quantum Q-system variables and the conserved quantities as limits of the time-evolved Macdonald operators and the Pieri operators, respectively. The latter are identified with relativistic q -difference Toda Hamiltonians. A crucial ingredient in the proof is the use of the “Fourier transformed” picture, in which we compute time-translation operators and prove that they commute with the Pieri operators or Hamiltonians. We also discuss the universal solutions of Koornwinder-Macdonald eigenvalue and Pieri equations, for which we prove a duality relation, which simplifies the proofs further.

1. INTRODUCTION

1.1. Overview. The purpose of this paper is to determine, in a uniform manner, a certain discrete integrable structure associated with the finite-dimensional representations of quantum affine algebras or Yangians of affine algebras \mathfrak{g} , and to understand the relation of this structure with Koornwinder-Macdonald theory and spherical double affine Hecke algebras (sDAHA) for all \mathfrak{g} of non-exceptional type.

Specifically, the characters of a special subset of representations of these algebras, known as KR-modules [CM06, KR87], satisfy recursion relations which can be viewed as discrete time evolution equations. These evolutions, called Q-systems [HKO⁺99, HKO⁺02], are known to have an integrable structure for a subset of algebras \mathfrak{g} . When $\mathfrak{g} = A_{N-1}^{(1)}$, the integrable structure was shown to be a Coxeter-Toda system [DFK10, GSV11], and this was generalized [Wil15] to $\mathfrak{g} = D_N^{(1)}, A_{2N-1}^{(2)}, D_{N+1}^{(2)}$ using the refactorization map [HKKR00]. In type $\mathfrak{g} = B_N^{(1)}$ the system was shown to be integrable [Vic18] using a Goncharov-Kenyon type dimer model [GK13]. Generally, discrete integrability was a conjecture and does seem to be related to refactorization maps.

The combinatorial structure of Q-system evolution equations is such that, in almost all cases, they can be interpreted as mutations in a cluster algebra of geometric type [Ked08, DFK09]. This structure can be encoded in the form of a quiver, which, in type $A_{N-1}^{(1)}$ appears in various contexts such as K -theoretic Coulomb branches [SS19], gauge theories [CNV], factorization maps [Wil15] and shifted quantum affine algebras or Yangians [BFN19, FT19, DFK19].

Cluster algebras have a natural quantization [BZ05], and in [DFK14, Lin21] it was shown that this quantization is associated to the grading of the so-called fusion product of KR-modules [FL99]. That is, the character of the graded fusion product can be expressed as a constant term identity involving the corresponding solutions of the quantum Q-system.

The Hamiltonians associated with the quantum Q-system in type $A_{N-1}^{(1)}$ are the q -difference (or relativistic) quantum Toda Hamiltonians of [Eti99], and therefore the graded characters of fusion products of fundamental representations in this type can be identified with q -Whittaker functions of $U_q(\mathfrak{sl}_N)$ [DFK18]. More generally, the quantum Q-system generators can be identified with the algebra of creation and annihilation operators of generalized q -Whittaker functions or graded characters. This algebra was identified in [DFK18, DFK17a, DFK19] with the q -Whittaker limit $t \rightarrow \infty$ of the spherical double affine Hecke algebra of type A [Che05] as follows. There is a natural $SL(2, \mathbb{Z})$ -symmetry of the sDAHA, and the generator $\tau_+ \in SL(2, \mathbb{Z})$ acts on the functional representation as conjugation by Cherednik's Gaussian. The subset of the sDAHA generators consisting of all τ_+ -translates of the commuting Macdonald difference operators can be identified, in the q -Whittaker limit, as the set of solutions of the quantum Q-system. This gives a functional representation of the quantum Q-system as an algebra of difference operators. In particular, single τ_+ -translates of Macdonald operators are the q -Whittaker limits of the Kirillov-Noumi raising operators for Macdonald polynomials [KN99].

There are generalizations of DAHAs and their functional representation to the other classical affine root systems [Che05, Che95a]. The corresponding Macdonald operators have generalized Macdonald polynomials as common eigenfunctions. The latter can also be obtained as specializations of the Koornwinder polynomials, which depend on several extra parameters (a, b, c, d) [Koo92, YY21, vD95] (see Table 1). These are eigenfunctions of the Koornwinder difference operator, the first in a family of N commuting difference operators [vD95], where N is the rank of the finite sub-algebra R . These are elements in the functional representation [Nou95] of a DAHA depending on (a, b, c, d) .

Quantum Q-systems can be defined for all affine algebras. It is natural to expect that some selected commuting family of difference operators together with their τ_+ -translates will, in the q -Whittaker limit, satisfy these non-commutative evolution equations. For a subset of algebras \mathfrak{g} , we presented a conjectural family of such difference operators in [DFK21]. In this paper, we generalize and prove these conjectures for all non-exceptional \mathfrak{g} using duality and the Fourier transform.

In type $A_{N-1}^{(1)}$, we present a simpler, alternative proof to that of [DFK18], that the τ_+ -translates of the Macdonald difference operators, in the q -Whittaker limit, satisfy the quantum Q-system. An essential ingredient in the simplification of the proof is the use of the Macdonald duality property [Mac95]. The Macdonald difference operators are elements of $\mathbb{C}_t(x)[\Gamma_i]$, written in terms of the generators of the quantum torus

$$(1.1) \quad \mathbb{T}_x := \langle x_i, \Gamma_i \rangle_{i=1, \dots, N}, \quad \Gamma_i x_j = q^{\delta_{ij}} x_j \Gamma_i$$

acting on functions of $x = (x_1, \dots, x_N)$. The monic, symmetric Macdonald polynomials $P_\lambda(x)$ are the common eigenfunctions of the Macdonald difference operators, depending on two sets of parameters, the variables x as well as integer partitions $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_N)$. Encoding the partitions using the variables $s = (s_1, \dots, s_N)$ with $s_i = \Lambda_i t^{\rho_i}$, $\Lambda_i = q^{\lambda_i}$ and ρ is the half-sum of positive roots ($\rho_i = N - i$), the duality of Macdonald theory is a symmetry property of the properly normalized eigenfunctions of the Macdonald operators under the interchange of the variables x and s [Mac95, vD96, Che95b].

This symmetry can be used to relate the Macdonald operator eigenvalue equations to the Pieri rules, which express the effect of multiplication of the Macdonald polynomial by elementary symmetric functions, as the action of q -difference operators in the variables s or Λ . These difference operators are Pieri operators, and can be written in terms of the quantum torus

$$(1.2) \quad \mathbb{T}_\Lambda := \langle \Lambda_i, T_i \rangle_{i=1, \dots, N}, \quad T_i \Lambda_j = q^{\delta_{ij}} \Lambda_j T_i.$$

The eigenvalue equations for Macdonald operators with eigenfunctions $P_\lambda(x)$ can be thought of as a special case of the sDAHA “Fourier transform”, which is a map from operators $f(x)$ to operators $\bar{f}(\Lambda)$, such that

$$(1.3) \quad f(x) P_\lambda(x) = \bar{f}(\Lambda) P_\lambda(x)$$

for all λ . In particular, the Fourier transforms of the Macdonald operators are the elementary symmetric functions in the variable s , and the transform of the Pieri operators are the elementary symmetric functions in x .

In the q -Whittaker limit, switching to the “Fourier transformed” \mathbb{T}_Λ picture results in drastic simplifications. Indeed, the transforms of the Macdonald operators in the q -Whittaker limit are the limits of the elementary symmetric function in s , which are the leading monomials in Λ_i due to the dependence of s on t . Because of this, the τ_+ -translates of the Fourier-transformed operators are simpler. In the functional representation as difference operators in x -variables, the action of τ_+ is expressed as conjugation by the Cherednik Gaussian $\gamma(x)$ [Che05], which is independent of t . In the Fourier-transformed picture in Λ -variables, we find the explicit operator $g(\Lambda)$ which represents the τ_+ -translation, generates the discrete time evolution of the quantum Q-system, and commutes with the Hamiltonians.

The operator $g(\Lambda)$ is interpreted as a Dehn twist [SS18] from a quantum higher Teichmüller theory interpretation in type A , and can be expressed as a product of quantum

dilogarithms. It was shown to be a specialization of the Baxter Q -operator of [SS18] for the Q -system quiver of type $A_{N-1}^{(1)}$. The Baxter Q -operator is the generating function of commuting q -difference Toda Hamiltonians [Eti99], which coincide with the q -Whittaker limit of the Macdonald Pieri rules.

The above discussion for type $A_{N-1}^{(1)}$ can be extended to the affine algebras, twisted or untwisted, of the form $\mathfrak{g} = X_N^{(r)}$, with X a classical Lie algebra of type $ABCD$ and $r = 1, 2$, in the notation of [Kac90, HKO⁺02]. This includes the cases presented as conjectures in [DFK21]. In this paper, we prove the relation between the q -Whittaker limit of the $SL(2, \mathbb{Z})$ -translates of appropriately chosen q -difference Koornwinder-Macdonald operators, for each algebra \mathfrak{g} , with the solutions of the type \mathfrak{g} quantum Q -system systems.

At the same time, we find a uniform description of the integrable structure of the quantum Q -systems as discrete dynamical systems with commuting integrals of motion. These are versions of the q -difference Toda Hamiltonians for the various root systems. The classical Q -systems associated with the affine root systems of types $A_N^{(1)}, D_N^{(1)}, A_{2N-1}^{(2)}, D_{N+1}^{(2)}$ are known [Wil15] to be the evolution equations of the refactorization maps of types A, D, C, B , respectively. As such, their integrable structure is given by a classical Toda-type lattice [HKKR00, Res03]. The integrable structure of the Q -systems for $\mathfrak{g} = B_N^{(1)}, C_N^{(1)}$ and $A_{2N}^{(2)}$, also given by a Toda-type Hamiltonian, but do not arise from factorization dynamics.

Here, we show that the unifying structure in the case of the quantum Q -systems is given, instead, by the Koornwinder-Macdonald operators [vD96, Sah99, Mac01] via the duality relating the eigenvalue equations to the Pieri rules, which become q -difference Toda equations [Eti99, GT21] in the q -Whittaker limit $t \rightarrow \infty$.

In this limit, the symmetry in the duality relations is broken, and as a result the Fourier transform of the Macdonald operators acting on Λ -space is greatly simplified. The rationale of our proofs is to identify the Fourier transforms of the $SL(2, \mathbb{Z})$ -translates of the Macdonald operators in the q -Whittaker limit and to transform the relations occurring in Λ -space back to x -space.

We also use another formulation of duality, based on the notion of the “universal solution” $P(x; s)$ to the Koornwinder-Macdonald eigenvalue equations (c.f. [Che95b, Che09] and the asymptotically-free basic Harish-Chandra series of [S14]). In [NS12] such solutions are explicitly computed in type $A_{N-1}^{(1)}$. In the Koornwinder case and its specializations, this suggests the definition of series solutions for the eigenvalue equation in the formal variables (x, s) . These truncate to the polynomial eigenfunctions when λ is specialized to \mathfrak{g} -partitions corresponding to dominant integral weights of \mathfrak{g} , where s is a discrete variable. Duality relates the universal solution $P(x; s)$ of the eigenvalue equation, to the universal solution $Q(s; x)$ of the Pieri equations, and formalizes the Fourier transform.

1.2. Summary of the main results. All the results here refer to all affine algebras \mathfrak{g} of type $A_{N-1}^{(1)}, B_N^{(1)}, C_N^{(1)}, D_N^{(1)}, A_{2N-1}^{(2)}, D_{N+1}^{(2)}$ and $A_{2N}^{(2)}$. The subsequent labeling of theorems

refers to both Section 2 (dedicated to the case of $A_{N-1}^{(1)}$ for pedagogical reasons) and Section 4 which addresses the cases $B_N^{(1)}, C_N^{(1)}, D_N^{(1)}, A_{2N-1}^{(2)}, D_{N+1}^{(2)}$ and $A_{2N}^{(2)}$.

For each such algebra \mathfrak{g} , we define a preferred set of N commuting q -difference operators $\{D_a^{(\mathfrak{g})}(x; q, t), a \in [1, N]\}$, chosen via specializations of linear combinations of van Diejen's higher Koornwinder operators, augmented by Rains and Macdonald operators for each algebra. The operators are chosen so that their eigenvalues form a basis for the space spanned by the fundamental characters of the underlying finite subalgebras R^* listed in Table 1, with $R^* = A_{N-1}$ in the case of $\mathfrak{g} = A_{N-1}^{(1)}$.

Once an appropriate family of commuting difference operators is constructed, we define the q -Whittaker limit $t \rightarrow \infty$ of the family, $\{D_a^{(\mathfrak{g})}(x; q)\}$, as well as their $\tau_+^n \in SL(2, \mathbb{Z})$ -translates, $\{D_{a;n}^{(\mathfrak{g})}(x; q), n \in \mathbb{Z}\}$. These operators satisfy the \mathfrak{g} -type quantum Q-systems, (2.13), (2.14) or (4.2)–(4.4). The main results of this paper may be stated as the following Theorems:

Theorems 2.3, 4.3. *For each \mathfrak{g} , the family of q -difference operators $\{D_{a;n}^{(\mathfrak{g})}(x; q), a \in [1, N], n \in \mathbb{Z}\}$ defined in Section 3.3.3 satisfy the corresponding quantum Q-system relations.*

These operators can be viewed as generalized raising or lowering operators, and in particular, we prove

Theorem 4.4. *The q -difference operators $\{D_{a;i}^{(\mathfrak{g})}(x; q) : i = 0, 1\}$ satisfy eigenvalue and raising operator properties:*

$$\begin{aligned} D_{a;0}^{(\mathfrak{g})}(x; q) \Pi_\lambda^{(\mathfrak{g})}(x) &= \Lambda^{\omega_a^*} \Pi_\lambda^{(\mathfrak{g})}(x), \\ D_{a;1}^{(\mathfrak{g})}(x; q) \Pi_\lambda^{(\mathfrak{g})}(x) &= \Lambda^{\omega_a^*} \Pi_{\lambda+\omega_a}^{(\mathfrak{g})}(x), \end{aligned}$$

where ω_a, ω_a^* are fundamental weights of R, R^* the finite root lattices of $\mathfrak{g}, \mathfrak{g}^*$.

The proof uses the $t \rightarrow \infty$ limit of the ‘‘Fourier transform’’ (1.3) (sometimes called q -Whittaker transform), which relies on the completeness of eigenfunctions of the q -Whittaker difference operators. Let $\Pi_\lambda^{(\mathfrak{g})}(x)$ be the set of common eigenfunctions of the operators $D_a^{(\mathfrak{g})}(x) = D_{a;0}^{(\mathfrak{g})}(x)$, i.e. limits of Macdonald polynomials. The Fourier transform relates operators $f(x)$ in \mathbb{T}_x to operators $f(\Lambda)$ in \mathbb{T}_Λ via $f(x) \Pi_\lambda^{(\mathfrak{g})}(x) = \bar{f}(\Lambda) \Pi_\lambda^{(\mathfrak{g})}(x)$. Then we define the q -difference operators acting on the index variables Λ , $\bar{D}_{a;n}^{(\mathfrak{g})}(\Lambda)$, starting from the initial (Macdonald eigenvalue) condition $D_{a;0}^{(\mathfrak{g})}(x) \Pi_\lambda^{(\mathfrak{g})}(x) = \bar{D}_{a;0}^{(\mathfrak{g})}(\Lambda) \Pi_\lambda^{(\mathfrak{g})}(x)$, and further constrained to obey the \mathfrak{g} quantum Q-system with the opposite multiplication. Finally we set out to prove that the operators $\bar{D}_{a;n}^{(\mathfrak{g})}(\Lambda)$ are the Fourier transforms of the operators $D_{a;n}^{(\mathfrak{g})}(x)$.

To this end, we first find the explicit time-translation operators $n \mapsto n + 1$ (or $n + 2$) acting on the operators $\bar{D}_{a;n}^{(\mathfrak{g})}$ for each \mathfrak{g} :

Theorems 2.7, 4.9. *There is a unique, up to scalar multiple, function $g = g^{(\mathfrak{g})}(\Lambda)$ in the generators of the quantum torus \mathbb{T}_Λ , such that*

$$\bar{D}_{a,n+t_a} = q^{-\omega_a^* \cdot \omega_{at_a}/2} g \bar{D}_{a,n} g^{-1},$$

where $t_a = 1$ for all labels a except for the so-called short labels N in type $B_N^{(1)}$, $[1, N-1]$ in type $C_N^{(1)}$ and $[1, N]$ in type $A_{2N}^{(2)}$, in which case $t_a = 2$. Here ω_a (ω_a^*) are fundamental weights of R (R^*).

The g operators are listed in (2.27) and (4.12). They can be expressed as explicit products of quantum dilogarithms. In the case of $A_{N-1}^{(1)}$ the operator g^{-1} is a particular evaluation of the Baxter Q -operator of [SS18].

Next, we use the duality in Koornwinder-Macdonald theory, which relates the eigenvalue equation of the Koornwinder-Macdonald operators for \mathfrak{g} to the Pieri rules for the dual \mathfrak{g}^* , where all cases are self-dual except $B_N^{(1)*} = C_N^{(1)}$ and $C_N^{(1)*} = B_N^{(1)}$. In the q -Whittaker limit, the Pieri operators are interpreted as (relativistic) q -difference Toda Hamiltonians, and the polynomials $\Pi_\lambda^{(\mathfrak{g})}(x)$ as q -Whittaker functions. The First Pieri operators for $\Pi_\lambda^{(\mathfrak{g})}(x)$ are listed in (2.12) and (3.38). They express the effect of multiplying $\Pi_\lambda^{(\mathfrak{g})}(x)$ with the first elementary symmetric function in x , $e_1(x)$ in type $A_{N-1}^{(1)}$ or $\hat{e}_1(x) = \sum_{i=1}^N (x_i + x_i^{-1})$ for all other \mathfrak{g} . The N commuting Pieri operators $H_a^{(\mathfrak{g})}(\Lambda)$, which can be thought of as q -difference Toda Hamiltonians, commute with the time-translation operator $g^{(\mathfrak{g})}(\Lambda)$. Thus, we have

Corollaries 2.13, 4.14. *The Pieri operators $\{H_a^{(\mathfrak{g})}(\Lambda), a \in [1, N]\}$ are algebraically independent conserved quantities of the opposite quantum Q -system for type \mathfrak{g} .*

The origin of the operators $g^{(\mathfrak{g})}(\Lambda)$ is clarified by the following theorem:

Theorems 2.11, 4.12. *The operators $g^{(\mathfrak{g})}(\Lambda)$ are the Fourier transforms of Cherednik's Gaussian operator $\gamma(x)^{t_1}$, with $\gamma(x)$ as in Equation (2.16) the functional representation of the $SL(2, \mathbb{Z})$ generator τ_+ :*

$$\gamma(x)^{t_1} \Pi_\lambda(x) = g^{(\mathfrak{g})}(\Lambda) \Pi_\lambda(x), \quad t_1 = \begin{cases} 2 & \text{if } \mathfrak{g} = A_{2N}^{(2)}, C_N^{(1)}, \\ 1 & \text{otherwise.} \end{cases}$$

Using these properties, and a uniqueness argument involving series solutions of the first Pieri rule, we finally prove

Theorems 2.6, 4.15. *For all \mathfrak{g} ,*

$$D_{a;n}^{(\mathfrak{g})}(x) \Pi_\lambda^{(\mathfrak{g})}(x) = \bar{D}_{a;n}^{(\mathfrak{g})}(\Lambda) \Pi_\lambda^{(\mathfrak{g})}(x), \quad (n \in \mathbb{Z})$$

valid for any \mathfrak{g} -partition λ and any root label $a \in [1, N]$.

In other words, the operators $\bar{D}_{a;n}^{(\mathfrak{g})}(\Lambda)$ are the Fourier transforms of the Whittaker limits of Macdonald operators $D_{a;n}^{(\mathfrak{g})}(x)$. This completes the proof of the main claim of this paper

(Theorems 2.3, 4.3 and 4.4), and the conjectures of [DFK21], that the q -difference operators $D_{a,n}^{(\mathfrak{g})}(x; q)$ satisfy the quantum Q-system of type \mathfrak{g} .

We consider duality in terms of the universal solutions of the eigenvalue and Pieri equations, $P(x; s)$ and $Q(s; x)$. These are continuations of the Koornwinder or Macdonald polynomials to arbitrary values of the parameters λ , with $s = q^\lambda t^\rho$ and $x = q^\mu t^{\rho^*}$ with ρ a function of (a, b, c, d) as in Section 3.1.4. The series have the form $P(x; s) = q^{\lambda \cdot \mu} \sum_{\beta \in Q_+} c_\beta(s) x^{-\beta}$ and $Q(s; x) = q^{\lambda \cdot \mu} \sum_{\beta \in Q_+^*} \bar{c}_\beta(x) s^{-\beta}$, with normalizations $c_0(s) = \bar{c}_0(x) = 1$, and Q_+, Q_+^* are the positive root lattices of the root systems R, R^* . Under specialization of the parameters (a, b, c, d) , universal solutions of the Koornwinder equation specialize \mathfrak{g} -Macdonald solutions, and whenever λ corresponds to any dominant integral R -weight, the series truncates to a polynomial. They also specialize to Chalykh's Baker-Akhiezer quasi-polynomials [Cha02] when parameters a, b, c, d, t (resp. t) are specialized to negative (half-)integer (resp. integer) powers of q , while λ remains arbitrary.

We establish the following relation between the series $P(x; s)$ and $Q(x; s)$, and a simple subsequent reformulation of duality.

Theorem 5.3, Corollary 5.4, Theorem 5.5. *The universal Koornwinder functions $Q^{(a,b,c,d)}(s; x)$ and $P^{(a,b,c,d)}(x; s)$ and their \mathfrak{g} -specializations are related via*

$$Q^{(a,b,c,d)}(s; x) = \frac{P^{(a,b,c,d)}(x; s)}{\Delta^{(a,b,c,d)}(x)}, \quad Q^{(\mathfrak{g})}(s; x) = \frac{P^{(\mathfrak{g})}(x; s)}{\Delta^{(\mathfrak{g})}(x)}$$

with $\Delta^{(a,b,c,d)}$ as in (3.14), and $\Delta^{(\mathfrak{g})}$ as in (3.27). Moreover the duality of Koornwinder and \mathfrak{g} -Macdonald polynomials extends to universal solutions as follows

$$Q^{(a^*, b^*, c^*, d^*)}(x; s) = Q^{(a,b,c,d)}(s; x), \quad Q^{(\mathfrak{g}^*)}(x; s) = Q^{(\mathfrak{g})}(s; x).$$

1.3. Outline of the paper. This paper is organized as follows.

We first revisit the A type in Section 2 as an illustration of the concepts used for other types in the remainder of the paper. After recalling the definition of Macdonald operators and polynomials, and showing how eigenvalue equations relate to Pieri rules via duality, we discuss the q -Whittaker limit $t \rightarrow \infty$ and the quantum Q-system (Sects. 2.1-2.4). Our main character is the time translation operator, whose adjoint action allows to advance Macdonald operators $D_a(x)$ in discrete time, thus producing operators $D_{a,n}(x)$. In x space it takes the form of the (scalar) Gaussian operator $\gamma(x)$ (Sect. 2.5). To prove that the Macdonald operators obey the quantum Q-system we switch to Λ space: in Sect. 2.6, using the eigenvalue $\bar{D}_{a;0}(\Lambda)$ of the Macdonald operator $D_a(x)$ as initial data, we *define* the candidate Fourier duals $\bar{D}_{a;n}(\Lambda)$ of the translated Macdonald operators as solutions of the opposite quantum Q-system for this initial data. In Sect. 2.7 we derive the associated explicit time translation operator $g(\Lambda)$. Finally the latter is shown to be the Fourier transform of the Gaussian $\gamma(x)$ in Sect. 2.10 leading to the proof of the main Theorem 2.3 for type A by identifying $D_{a,n}(x)$ with the Fourier transform of $\bar{D}_{a;n}(\Lambda)$. A key ingredient

is the commutation of $g(\Lambda)$ with the first Pieri operator $H_1(\Lambda)$, shown in Sect. 2.8. This allows to identify the Pieri operators as the commuting conserved quantities of the quantum Q-system, also known as the relativistic Toda Hamiltonians of type A (Sect. 2.12). Finally Section 2.13 is devoted to a reformulation of the duality properties, Fourier transform, and proofs in terms of the universal solution of the Macdonald eigenvalue equation considered by [NS12].

The remainder of the paper focusses on the other types. In Section 3, we define suitable families of commuting Macdonald operators, borrowing from various existing constructions. The first approach uses the known specialization scheme of the Koornwinder operator to Macdonald operators, completed by van Diejen and Rains into a set of commuting difference operators (Sects. 3.1 and 3.2, and details in Appendix A). The duality of Koornwinder polynomials descends to a duality between Macdonald operators and Pieri rules for dual types. Sect. 3.3 describes the q -Whittaker limit $t \rightarrow \infty$, and provides detailed definitions of the translated limiting Macdonald operators $D_{a;n}^{(\mathfrak{g})}$ for all types. Some new subtleties arise for non- A types, in particular the distinction between long and short labels a for which the time translation has to be defined separately (type A only has long labels). Section 4 defines the quantum Q-systems for all types and presents the main results of this paper, Theorems 4.3 and 4.4 (Sect. 4.1), and their proof (Sect. 4.2 complemented by Appendices B and C), along the same lines as for the case of type A . We define candidate Fourier transforms $\bar{D}_{a;n}^{(\mathfrak{g})}(\Lambda)$ of the translated Macdonald operators by use of the opposite \mathfrak{g} -quantum Q-system, and construct the time translation operators $g^{(\mathfrak{g})}(\Lambda)$ explicitly. The latter commute with the Toda Hamiltonians, identified as the conservation laws of the quantum Q-systems, thus allowing us to identify $g^{(\mathfrak{g})}(\Lambda)$ with the Fourier transform of the Gaussian operator $\gamma^{(\mathfrak{g})}(x)$. Finally in Sect. 4.3 we prove that the $\bar{D}_{a;n}^{(\mathfrak{g})}(\Lambda)$ are the Fourier transforms of the translated Macdonald operators $D_{a;n}^{(\mathfrak{g})}(x)$, from which the main results follow.

Section 5 is an extension of the constructions of Section 2.13. In Sect. 5.1 we introduce universal solutions $P(x, s)$ to the Koornwinder-Macdonald eigenvalue equations in the form of power series of the variables $x^{-\alpha_i}, s^{-\alpha_i^*}$ where α_i, α_i^* are the simple roots of suitable lattices. $P(x, s)$ has the remarkable property that it reduces to Koornwinder-Macdonald polynomials upon specializing $s = q^{\lambda t^{\rho}}$ for λ a (\mathfrak{g}) -partition. In Sect. 5.2 we extend the duality of Koornwinder-Macdonald polynomials to some relation between $P(x, s)$ and its dual $P^*(s, x)$. Finally in Sect. 5.3 we show how this new formulation allows to rewrite the proofs of this paper more elegantly.

We finally gather some concluding remarks in Section 6. We comment on the three additional “companion” quantum Q-systems obtained as a by-product of our study and the associated representation by q -difference operators, which were not part of the original setting (Sect. 6.1). We discuss path models for the various universal functions of the paper in Sect. 6.2. We interpret the universal function results in terms of q -Whittaker functions in Sect. 6.3. Sect. 6.4 summarizes our results and lists related open questions.

Acknowledgments. We thank I. Cherednik, G. Schrader, A. Shapiro, J. Shiraishi and C. Stroppel for useful discussions, and G. Barraquand for pointing out reference [Rai05]. We are also thankful to the referee who raised a number of interesting issues and suggested further important references. This work was supported by the following grants: National Science Foundation grants DMS 18-02044 and DMS-1937241; NSF Grant No. 1440140 while the authors were in residence at the Mathematical Sciences Research Institute in Berkeley, California in 2021; Simons Foundation Fellowship grants 613580 and 617036 and Simons Foundation grants MP-TSM-00002262 and MP-TSM-00001941. PDF is supported by the Morris and Gertrude Fine Endowment. RK thanks the Institut Henri Poincaré and the Institut de Physique Théorique-CEA Paris Saclay for their hospitality.

2. DUALITY PROPERTY IN TYPE $A_{N-1}^{(1)}$

The main result of [DFK18] is a theorem which states that the q -Whittaker limit of the A-type Macdonald operators and their τ_+ -translates satisfy the quantum Q-system of type $A_{N-1}^{(1)}$. To prove this, we used the explicit form of these difference operators in x -space. In this section we will give a different method for proving the same theorem by using Macdonald's duality relation and the Fourier transform of these operators.

We use the notation for the roots and weights of \mathfrak{gl}_N as follows. Let $\{e_1, \dots, e_N\}$ be the standard basis of \mathbb{R}^N , then the simple roots are $\{\alpha_i = e_i - e_{i+1}; i = 1, \dots, N-1\}$, the \mathfrak{gl}_N fundamental weights (including ω_N) are $\{\omega_i = e_1 + \dots + e_i, i = 1, \dots, N\}$, the positive roots are $R_+ = \{e_i - e_j, 1 \leq i < j \leq N\}$, and $\rho = \sum_{i=1}^N (N-i)e_i$. For $u, v \in \mathbb{R}^N$, we use the notation $u \cdot v$ for the standard scalar product. We consider $q \in \mathbb{C}^*$, $|q| < 1$.

2.1. Macdonald operators and polynomials. Let $x = (x_1, \dots, x_N)$ be formal variables, and define the q -difference operators $\mathcal{D}_a(x; q, t)$ be the a -th Macdonald q -difference operator in type $A_{N-1}^{(1)}$ [Mac95]:

$$(2.1) \quad \mathcal{D}_a(x; q, t) = \sum_{\substack{I \subset \{1, \dots, N\} \\ |I|=a}} \prod_{\substack{j \notin I \\ i \in I}} \left(\frac{tx_i - x_j}{x_i - x_j} \right) \prod_{i \in I} \Gamma_i, \quad a \in \{1, \dots, N\}, \quad \Gamma_i x_j = q^{\delta_{ij}} x_j \Gamma_i.$$

These difference operators form a commuting family and preserve the space of symmetric polynomials in x . The common eigenvectors are the monic symmetric Macdonald polynomials $P_\lambda(x)$, where $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_N \geq 0)$ is an integer partition:

$$(2.2) \quad \mathcal{D}_a(x; q, t) P_\lambda(x) = t^{-\binom{a}{2}} e_a(s) P_\lambda(x) \quad (a = 1, 2, \dots, N),$$

where, $e_a(s)$ is the a th elementary symmetric function in $s = (s_1, \dots, s_N)$, with $s_i = t^{N-i} q^{\lambda_i}$. Macdonald polynomials form a basis for the space of symmetric polynomials of x .

2.2. Duality. The Macdonald polynomials $P_\lambda(x)$ satisfy a remarkable duality property, which is a symmetry under the interchange of the variables x and s . Under the specialization $x = t^\rho q^\mu$ with μ an integer partition, Macdonald showed that [Mac95]

$$(2.3) \quad \frac{P_\lambda(t^\rho q^\mu)}{P_\lambda(t^\rho)} = \frac{P_\mu(t^\rho q^\lambda)}{P_\mu(t^\rho)}$$

for all integer partitions λ, μ .

Remark 2.1. *The duality property has a formal generalization to arbitrary values of μ, λ , see the discussion on the universal function in Section 2.13.*

The denominators in Equation (2.3) can be obtained as specializations of an infinite product expression. Let

$$(2.4) \quad \Delta(x) = \prod_{n \geq 1} \prod_{\alpha \in R_+} \frac{1 - q^n x^{-\alpha}}{1 - t^{-1} q^n x^{-\alpha}} = \prod_{\beta \in \hat{R}_{++}} \frac{1 - x^{-\beta}}{1 - t^{-1} x^{-\beta}},$$

where the second product extends over the set \hat{R}_{++} of strictly positive affine roots $\beta = n\delta + \sum_i \beta_i \alpha_i$, $n \geq 1, \beta_i \in \mathbb{Z}_+$, and we use the shorthand notation $x^\beta = q^{-n} \prod_i (x_i/x_{i+1})^{\beta_i}$. Then [Mac95]¹

$$(2.5) \quad P_\lambda(t^\rho) = t^{\lambda \cdot \rho} \frac{\Delta(t^\rho)}{\Delta(t^\rho q^\lambda)}.$$

Next we use Macdonald's evaluation homomorphism u_μ defined on functions of x by $u_\mu(f(x)) = f(q^\mu t^\rho)$. It maps the generators of quantum torus \mathbb{T}_x to $u_\mu(x_i) = q^{\mu_i} t^{N-i}$ and $u_\mu(\Gamma_i) = e^{\partial_{\mu_i}}$ which acts as the translation $\mu_i \mapsto \mu_i + 1$ while leaving $\mu_j, j \neq i$ unchanged. Applying u_μ for μ an integer partition on the eigenvalue equation (2.2), and using duality (2.3) gives

$$\begin{aligned} e_a(t^\rho q^\lambda) P_\mu(t^\rho q^\lambda) &= t^{\binom{a}{2}} P_\mu(t^\rho) u_\mu(\mathcal{D}_a(x; q, t)) P_\mu(t^\rho)^{-1} P_\mu(t^\rho q^\lambda) \\ &= \left\{ t^{\binom{a}{2}} t^{\rho \cdot \mu} \Delta(t^\rho q^\mu)^{-1} u_\mu(\mathcal{D}_a(x; q, t)) \Delta(t^\rho q^\mu) t^{-\rho \cdot \mu} \right\} P_\mu(t^\rho q^\lambda). \end{aligned}$$

Let us finally interchange the partition labels $\lambda \leftrightarrow \mu$ in the above equation, and note moreover that since the polynomial $P_\lambda(x)$ is entirely determined by its values at the discrete specializations $x = t^\rho q^\mu$ with μ taking values in integer partitions, we can conclude that $P_\lambda(x)$ satisfies the Pieri equations:

$$(2.6) \quad \mathcal{H}_a(\Lambda; q, t) P_\lambda(x) = e_a(x) P_\lambda(x) \quad (a = 1, 2, \dots, N),$$

with

$$(2.7) \quad \mathcal{H}_a(\Lambda; q, t) := t^{\binom{a}{2}} t^{\rho \cdot \lambda} \Delta(s)^{-1} u_\lambda(\mathcal{D}_a(x; q, t)) \Delta(s) t^{-\rho \cdot \lambda}.$$

¹Our definition of the infinite product Δ differs slightly from that of Macdonald's Δ^+ , but is better suited for taking the limit $t \rightarrow \infty$ below.

The Pieri rules for the Macdonald polynomials express the multiplication of $P_\lambda(x)$ by an elementary symmetric function $e_a(x)$ as a linear combination of Macdonald polynomials with shifted partitions.

Remark 2.2. *Note that in (2.7) the evaluation u_λ maps $x_i \rightarrow q^{\lambda_i} t^{N-i} = s_i$ and $\Gamma_i \rightarrow e^{\partial \lambda_i}$. The same evaluation map sends the generators of the quantum torus \mathbb{T}_Λ to respectively $\Lambda_i \rightarrow q^{\lambda_i}$ and $T_i \rightarrow e^{\partial \lambda_i}$. By a slight abuse of notation, we simply express the operator $\mathcal{H}_a(\Lambda; q, t)$ in terms of the generators Λ_i and T_i and omit the mention of the evaluation map from now on. This amounts to evaluating the left action of the quantum torus \mathbb{T}_Λ on functions of the variables $(\Lambda_1, \dots, \Lambda_N)$. There is also a right action of the evaluation of \mathbb{T}_Λ on the basis of Macdonald polynomials P_λ in which Λ_i acts diagonally $P_\lambda \Lambda_i = P_\lambda q^{\lambda_i}$ and T_i by a shift $P_\lambda T_i = P_{\lambda+e_i}$. We adopt a single notation using the quantum torus generators, to denote either left or right action.*

Explicitly the commuting operators $H_a(\Lambda; q, t)$ acting on functions of Λ or s are

$$(2.8) \quad \mathcal{H}_a(\Lambda; q, t) = \sum_{\substack{I \subset [1, N] \\ |I|=a}} \prod_{\substack{i \in I, j \notin I \\ j < i}} \frac{t^{i-j-1} \Lambda_j - \Lambda_i}{t^{i-j} \Lambda_j - \Lambda_i} \frac{t^{i-j+1} \Lambda_j - q \Lambda_i}{t^{i-j} \Lambda_j - q \Lambda_i} \prod_{i \in I} T_i, \quad T_i \Lambda_j = q^{\delta_{ij}} \Lambda_j T_i.$$

They can be thought of as Macdonald Hamiltonians of type $A_{N-1}^{(1)}$: In the q -Whittaker limit, they become relativistic q -difference Toda Hamiltonians (see Sect. 2.12 below).

2.3. The q -Whittaker limit. By a slight abuse of language, we call the limit $t \rightarrow \infty$ of the difference operators and eigenfunctions above the q -Whittaker limit (as opposed to the standard $t \rightarrow 0$). The q -Whittaker functions² are the limits of the Macdonald polynomials [Che09]:

$$\Pi_\lambda(x) = \lim_{t \rightarrow \infty} P_\lambda(x).$$

In this limit we use the t -independent variables $\Lambda_i = q^{\lambda_i}$ instead of s_i . The symmetry between the eigenvalue equation and the Pieri rules under exchange of x and s is lost in this limit. However, the limit of the difference equations still makes sense. The q -Whittaker functions satisfy the difference equations

$$(2.9) \quad D_a(x; q) \Pi_\lambda(x) = \Lambda^{\omega_a} \Pi_\lambda(x), \quad \Lambda^{\omega_a} = \Lambda_1 \cdots \Lambda_a;$$

$$(2.10) \quad H_a(\Lambda; q) \Pi_\lambda(x) = e_a(x) \Pi_\lambda(x).$$

Here, the difference operators are

$$(2.11) \quad D_a(x; q) = \lim_{t \rightarrow \infty} t^{a(a-N)} \mathcal{D}_a(x; q, t) = \sum_{\substack{I \subset \{1, \dots, N\} \\ |I|=a}} \prod_{\substack{j \notin I \\ i \in I}} \frac{x_i}{x_i - x_j} \prod_{i \in I} \Gamma_i,$$

²This is a slight abuse of language, as these are strictly speaking q^{-1} -Whittaker functions. $\Pi_\lambda(x)$ is interpreted as a (class 1) q -Whittaker function where x is the representation index, and λ the argument.

and the Pieri operators are

$$(2.12) \quad H_a(\Lambda; q) = \lim_{t \rightarrow \infty} \mathcal{H}_a(\Lambda; q, t) = \sum_{\substack{I \subset [1, N] \\ |I|=a}} \prod_{\substack{i \in I \\ i-1 \notin I}} \left(1 - \frac{\Lambda_i}{\Lambda_{i-1}}\right) \prod_{i \in I} T_i$$

with the convention that $\Lambda_0 := \infty$. Note the simplified form of these equations: the Pieri operators are polynomials in $\Lambda^{-\alpha_i}$ where α_i are the simple roots, rather than rational functions. Moreover, the eigenvalues of operators $D_a(x; q)$ are monomials in Λ , because $e_a(s)$ is replaced by its leading term in t , Λ^{ω_a} , where ω_a are the fundamental weights.

2.4. Type $A_{N-1}^{(1)}$ quantum Q-system. The main result of [DFK18] is that the set of all τ_+ -translations of the operators $D_a(x; q)$ satisfy the quantum Q-system equations³ of type $A_{N-1}^{(1)}$. To define this system, let $\{\mathcal{Q}_{a;k} : a \in \{1, \dots, N\}, k \in \mathbb{Z}\}$ be invertible elements in some non-commutative algebra over $\mathbb{C}(q)$, subject to two types of relations: A q -commutation relation

$$(2.13) \quad \mathcal{Q}_{a;k} \mathcal{Q}_{b;k+i} = q^{i \min(a,b)} \mathcal{Q}_{b;k+i} \mathcal{Q}_{a;k}, \quad k \in \mathbb{Z}, a, b \in [1, N], |i| \leq 1,$$

and a cluster exchange-type relation, which can be thought of as a discrete time evolution in the variable k :

$$(2.14) \quad q^a \mathcal{Q}_{a;k+1} \mathcal{Q}_{a;k-1} = \mathcal{Q}_{a;k}^2 - \mathcal{Q}_{a-1;k} \mathcal{Q}_{a+1;k}, \quad a \in [1, N], k \in \mathbb{Z},$$

subject to the boundary conditions $\mathcal{Q}_{0;k} = 1, \mathcal{Q}_{N+1;k} = 0$. The algebra generated by the set $\{\mathcal{Q}_{a;k} : a \in [1, N], k \in \mathbb{Z}\}$ and their inverses is a subalgebra in a certain quantum cluster algebra [DFK11].

Define the generalized q -Whittaker difference operators, acting on the space of functions in the variables $\{x_1, \dots, x_N\}$ with coefficients in $\mathbb{C}(x_1, \dots, x_N, q)$:

$$(2.15) \quad D_{a;k}(x; q) = \sum_{\substack{I \subset \{1, \dots, N\} \\ |I|=a}} \prod_{i \in I} x_i^k \prod_{j \notin I} \left(\frac{x_i}{x_i - x_j} \right) \prod_{i \in I} \Gamma_i, \quad a \in [1, N], k \in \mathbb{Z}.$$

When $k = 0$ these are just the q -Whittaker difference operators (2.9). The main result of [DFK18] is that these operators provide a functional representation of the quantum Q-system:

Theorem 2.3 ([DFK18]). *The q -difference operators $D_{a;k}(x; q)$ of (2.15) satisfy the quantum Q-system relations (2.13) and (2.14).*

The original proof of the theorem consisted of working directly with the difference operators $D_{a;k}(x; q)$. Here, we present a simplification of the proof which uses the duality of Section 2.2. This method has the advantage that it can be generalized to other root systems.

³This was referred to as the type A_{N-1} Q-system in e.g. [DFK18].

2.5. The action of τ_+ on the difference operators. The difference operators $D_a(x; q, t)$ are representations of elements of the spherical DAHA acting on the space of functions in N variables. There is an action of $SL_2(\mathbb{Z})$ on DAHA, and the $SL_2(\mathbb{Z})$ -generator τ_+ acts on the functional representation by the adjoint action of Cherednik's Gaussian operator [Che05]

$$(2.16) \quad \gamma(x) = e^{\sum_{i=1}^N \frac{\log(x_i)^2}{2 \log(q)}}.$$

In the q -Whittaker limit, the adjoint action of $\gamma(x)$ is well-defined. In particular, the difference operators (2.11) are the τ_+ -translates of $D_a(x; q)$:

Lemma 2.4 ([DFK19]).

$$(2.17) \quad D_{a;k}(x; q) = q^{-ak/2} \gamma(x)^{-k} D_a(x; q) \gamma(x)^k.$$

The proof follows from

$$\gamma(x)^{-1} \Gamma_i \gamma(x) = q^{\frac{1}{2}} x_i \Gamma_i.$$

2.6. Fourier transform. The q -Whittaker functions $\{\Pi_\lambda(x)\}$, with λ ranging over integer partitions, form a complete basis of the space of symmetric polynomials. Suppose that a set of difference operators $\{D_a(x)\}$ in the variables of \mathbb{T}_x satisfies

$$D_a(x) \Pi_\lambda(x) = \bar{D}_a(\Lambda) \Pi_\lambda(x)$$

for all λ , where the difference operators $\bar{D}_a(\Lambda)$ act in the variables of $\mathbb{T}_{\Lambda=q^\lambda}$. If the operators $\{\bar{D}_a(\lambda)\}$ satisfy certain relations \mathcal{R} , then the set of operators $\{D_a(x)\}$ satisfy the relations \mathcal{R}^{op} with the opposite multiplication. The operators $\bar{D}_a(\Lambda)$ are the ‘‘Fourier transforms’’ of the operators $D_a(x)$.

The strategy is to define the operators $\bar{D}_{a;k}(\Lambda; q)$, starting with initial data $\{\bar{D}_{a;0}, \bar{D}_{a;1} : a \in [1, N]\}$, by using the quantum Q-system relations with the opposite multiplication. We will then show that the corresponding Fourier-dual operators $D_{a;k}(x; q)$ are the difference operators (2.15), which therefore satisfy the quantum Q-system. The simplification of the proof in the Fourier transformed picture is due to the simple form of the initial data in \mathbb{T}_Λ , which allows us to compute the τ_+ action on the Fourier transformed operators directly.

2.6.1. Initial data. To find appropriate initial data for the opposite quantum Q-system, we start with the eigenvalue equation (2.9), which we write as

$$(2.18) \quad D_{a;0}(x; q) \Pi_\lambda(x) = \bar{D}_{a;0}(\Lambda; q) \Pi_\lambda(x), \quad \bar{D}_{a;0}(\Lambda; q) = \Lambda^{\omega_a}, \quad a \in [1, N].$$

For the operators $\bar{D}_{a;k}(\Lambda; q) \in \mathbb{C}_q(\Lambda)[T]$ to be well-defined from the opposite quantum Q-system for all k , we need another set of initial data, $\{\bar{D}_{a;1}(\Lambda; q), a \in [1, N]\}$, such that (we drop the arguments Λ, q from now on):

$$(2.19) \quad \bar{D}_{a;0} \bar{D}_{b;1} = q^{-\min(a,b)} \bar{D}_{b;1} \bar{D}_{a;0}, \quad [\bar{D}_{a;1}, \bar{D}_{b;1}] = 0, \quad a, b \in [1, N].$$

These are $2N$ relations on the quantum torus \mathbb{T}_Λ of dimension $2N$, so they determine $\{\bar{D}_{a;1}\}$ up to scalar multiple, which commutes with \mathbb{T}_Λ , i.e. an element in $\mathbb{C}(q)$. We choose this constant to be 1:

$$(2.20) \quad \bar{D}_{a;1} = \Lambda^{\omega_a} T^{\omega_a} = \Lambda_1 \cdots \Lambda_a T_1 \cdots T_a.$$

2.6.2. *The set $\{\bar{D}_{a;k}\}$.* Given the set of $2N$ initial data $\{\bar{D}_{a;k}, k \in \{0, 1\}, a \in [1, N]\}$, which form an alternate basis for \mathbb{T}_Λ , all $\bar{D}_{a;k}$ are uniquely defined by the requirement that they satisfy the opposite quantum Q-system relation:

$$(2.21) \quad q^a \bar{D}_{a;k-1} \bar{D}_{a;k+1} = \bar{D}_{a;k}^2 - \bar{D}_{a+1;k} \bar{D}_{a-1;k}, \quad \bar{D}_{0;k} = 1, \quad \bar{D}_{N+1;k} = 0.$$

Remark 2.5. *The quantum Q-system variables are cluster variables in a quantum cluster algebra, and therefore, due to the Laurent property of these algebras, all $\bar{D}_{a;k}$ are in fact Laurent polynomials in the initial data (2.18) and (2.20). As the latter are monomial in the variables of the quantum torus \mathbb{T}_Λ , so are the $\bar{D}_{a;k} \in \mathbb{C}_q[\Lambda^{\pm 1}, T^{\pm 1}]$.*

The main theorem to be proven is

Theorem 2.6.

$$\bar{D}_{a;k}(\Lambda; q) \Pi_\lambda(x) = D_{a;k}(x; q) \Pi_\lambda(x)$$

for all $a \in [1, N]$ and $k \in \mathbb{Z}$, with $D_{a;k}(x; q)$ the difference operators of Equation (2.15)

Since $\bar{D}_{a;k}(\Lambda; q)$ satisfy the opposite quantum Q-system relations, then the operators $D_{a;k}(x; q)$ satisfy the quantum Q-system. Theorem 2.6 therefore implies Theorem 2.3.

The proof of this theorem performed over the next several subsections as follows. First, in 2.7 we compute a ‘‘time translation’’ operator $g(\Lambda)$, whose adjoint action on $\bar{D}_{a;k}(\Lambda; q)$ gives $\bar{D}_{a;k+1}(\Lambda; q)$ in accordance with the opposite Q-system recursion evolution. We then show in 2.8 that this time-translation commutes with the Pieri operator or Hamiltonian $H_1(\Lambda; q)$. In 2.9 we prove the uniqueness of the solution of the Pieri equation $H_1(\Lambda, q) f = \hat{e}(x) f$. This implies, as shown in 2.10, that $g(\Lambda)$ is the Fourier transform of the Gaussian $\gamma(x)$. The proof of Theorem 2.6 follows (Section 2.11) from a simple argument using the Fourier transform.

2.7. **The action of τ_+ on $\bar{D}_{a;k}(\Lambda; q)$.** The key to proving theorem 2.6 is to find an element $g(\Lambda)$ in the completion of the space of rational functions in \mathbb{T}_Λ which realizes the action of $\gamma(x)$ in the Fourier transformed picture⁴:

Theorem 2.7. *There is a unique, up to scalar multiple, function $g \equiv g(\Lambda)$ acting on the variables of \mathbb{T}_Λ , such that*

$$(2.22) \quad \bar{D}_{a;k} = q^{-ak/2} g^k \bar{D}_{a;0} g^{-k}$$

⁴The element $g(\Lambda)$ is referred to as the ‘‘Dehn twist’’ generator in the geometric formulation of Ref. [SS18], which uses a different but related definition of the Fourier transform, under the name of Whittaker transform.

is a solution of the opposite quantum Q-system for all a, k .

Proof. The element g is determined by the commutation relation (2.19) and the subset of Q-system relations (2.21) corresponding to $k = 1$. Indeed, assuming such an element g exists, we first note that $g^m[\bar{D}_{a,0}, \bar{D}_{b,0}]g^{-m} = q^{(a+b)m/2}[\bar{D}_{a,m}, \bar{D}_{b,m}] = 0$ for all $m \in \mathbb{Z}$. Similarly, conjugating (2.19) and (2.21) for $k = 1$ with g^m gives

$$\begin{aligned} g^m(\bar{D}_{a,0} \bar{D}_{b,1} - q^{\text{Min}(a,b)} \bar{D}_{b,1} \bar{D}_{a,0})g^{-m} &= q^{(a+b)m/2}(\bar{D}_{a,m} \bar{D}_{a,m+1} - q^{\text{Min}(a,b)} \bar{D}_{a,m+1} \bar{D}_{a,m}) = 0 \\ g^m(q^a \bar{D}_{a,-1} \bar{D}_{a,1} - \bar{D}_{a,0}^2 + \bar{D}_{a-1,0} \bar{D}_{a+1,0})g^{-m} &= q^{am}(q^a \bar{D}_{a,m-1} \bar{D}_{a,m+1} - \bar{D}_{a,m}^2 + \bar{D}_{a-1,m} \bar{D}_{a+1,m}) = 0 \end{aligned}$$

We therefore obtain the opposite of (2.13) and (2.14) for all $m \in \mathbb{Z}$.

We now prove the existence of g by construction. Assume $g = g_T g_\Lambda$, where g_T commutes with all T_a and g_Λ commutes with all Λ_a . Up to a scalar multiple, g_T is determined by the N equations (2.22) with $k = 1$:

$$q^{-a/2} g \bar{D}_{a;0} g^{-1} = q^{-a/2} g_T \Lambda^{\omega_a} g_T^{-1} = \bar{D}_{a;1} = \Lambda^{\omega_a} T^{\omega_a}, \quad a \in [1, N],$$

because g_Λ commutes with $\bar{D}_{a;0} = \Lambda^{\omega_a}$ by assumption. This has the solution

$$(2.23) \quad g_T = \prod_{i=a}^N e^{\frac{(\log T_a)^2}{2 \log q}},$$

where we used the identity

$$e^{\frac{(\log T_a)^2}{2 \log q}} \Lambda_a e^{-\frac{(\log T_a)^2}{2 \log q}} = \Lambda_a e^{\frac{(\log q T_a)^2 - (\log T_a)^2}{2 \log q}} = q^{1/2} \Lambda_a T_a.$$

To find g_Λ , we use the Q-system (2.21) with $k = 1$:

$$(2.24) \quad \bar{D}_{a;2} = \Lambda^{\omega_a} T^{2\omega_a} (1 - q \Lambda^{-\alpha_a} T^{-\alpha_a}), \quad a \in [1, N-1], \quad \bar{D}_{N;2} = \Lambda^{\omega_N} T^{2\omega_N}.$$

To simplify the equations below, we set $\Lambda_{N+1} = 0$ and $\Lambda_0 = \infty$, i.e. we define α_0, α_N by $\Lambda^{-\alpha_0} = \Lambda^{-\alpha_N} = 0$. The function g_Λ is defined from

$$(2.25) \quad \bar{D}_{a;1} = q^{a/2} g^{-1} \bar{D}_{a;2} g = q^{a/2} g_\Lambda^{-1} g_T^{-1} \bar{D}_{a;2} g_T g_\Lambda = \Lambda^{\omega_a} T^{\omega_a}.$$

Acting on (2.24) by the adjoint action of g_T^{-1} , we have

$$q^{a/2} g_T^{-1} \bar{D}_{a;2} g_T = \Lambda^{\omega_a} T^{\omega_a} (1 - \Lambda^{-\alpha_a}).$$

Using

$$\prod_{n=0}^{\infty} (1 - q^n \Lambda^{-\alpha_a}) T^{\omega_b} = T^{\omega_b} \prod_{n=0}^{\infty} (1 - q^{n+\delta_{ab}} \Lambda^{-\alpha_a}) = T^{\omega_b} (1 - \Lambda^{-\alpha_a})^{-1} \prod_{n=0}^{\infty} (1 - q^n \Lambda^{-\alpha_a})$$

we can choose

$$(2.26) \quad g_\Lambda = \prod_{a=1}^{N-1} (\Lambda^{-\alpha_a}; q)_\infty^{-1}, \quad (a; b)_\infty = \prod_{n=0}^{\infty} (1 - b^n a).$$

We conclude that the function

$$(2.27) \quad g = \left(\prod_{a=1}^N e^{\frac{(\log T_a)^2}{2 \log q}} \right) \prod_{a=1}^{N-1} \left(\frac{\Lambda_{a+1}}{\Lambda_a}; q \right)_{\infty}^{-1}$$

satisfies the conditions of Theorem 2.7. \square

2.8. Commutation with the first Pieri operator. In general, the translation operator g commutes with each of the Pieri operators $H_a(\Lambda; q)$. It is sufficient for our purposes to show that it commutes with the first, $H_1(\Lambda; q)$:

Theorem 2.8. *The function $g(\Lambda)$ of Theorem 2.7 commutes with the Pieri operator $H_1(\Lambda; q)$ of Equation (2.12).*

Proof. From (2.12),

$$H_1(\Lambda; q) = \sum_{a=1}^N (1 - \Lambda^{-\alpha_{a-1}}) T_a = \sum_{a=1}^N T_a - \sum_{a=1}^{N-1} \frac{\Lambda_{a+1}}{\Lambda_a} T_{a+1}.$$

For all $a \in [1, N]$,

$$g(1 - \Lambda^{-\alpha_{a-1}}) T_a g^{-1} = g_T (1 - q^{-1} \Lambda^{-\alpha_a}) T_a g_T^{-1} = (T_a - \Lambda^{-\alpha_a} T_{a+1}),$$

again with the convention that $\Lambda^{-\alpha_0} = \Lambda^{-\alpha_N} = 0$. Summing over all $a = 1, 2, \dots, N$ results in $gH_1(\Lambda; q)g^{-1} = H_1(\Lambda; q)$. \square

2.9. Uniqueness of the Pieri solution. To prove the result of next section, we resort to a uniqueness argument regarding the solutions to the first Pieri equation (2.30).

Recall that $\Pi_\lambda(x)$ is eigenfunction of the Macdonald operator: $D_1 \Pi_\lambda = \Lambda_1 \Pi_\lambda$. Writing $\Pi_\lambda(x) = x^\lambda p_\lambda(x)$, and $D_1 = \sum_i \phi_i(x) \Gamma_i$, the eigenvalue equation turns into

$$(2.28) \quad \left(1 - \sum_i \phi_i(x) \frac{\Lambda_i}{\Lambda_1} \Gamma_i \right) p_\lambda(x) = 0.$$

This equation makes it easy to analytically continue $p_\lambda(x)$ to a function $\hat{p}_\lambda(x)$ with $\lambda \in \mathbb{C}^N$, as the dependence on Λ is explicit (in fact polynomial of the variables $\Lambda^{-\alpha_i}$, α_i the simple roots of A_{N-1}). Writing $\hat{p}_\lambda(x)$ as a series of the variables $x^{-\alpha_i}$: $\hat{p}_\lambda(x) = \sum_{\beta \in Q_+} c_\beta(\Lambda) x^{-\beta}$, Q_+ the positive cone of the root lattice of A_{N-1} , (2.28) turns into a linear triangular system for the coefficients $c_\beta(\Lambda)$, which are uniquely determined (with $c_0(\Lambda) = 1$), and rational functions of the $\Lambda^{-\alpha_i}$. Moreover, specializing λ to an integer partition in $\hat{p}_\lambda(x)$ recovers $p_\lambda(x)$. Expanding $c_\beta(\Lambda) = \sum_{\delta \in Q_+} c_{\beta, \delta} \Lambda^{-\delta}$ allows to view $\Pi_\lambda(x)$ as the specialization of a series in Λ :

$$x^\lambda \hat{p}_\lambda(x) = x^\lambda \sum_{\delta \in Q_+} \hat{c}_\delta(x) \Lambda^{-\delta}, \quad \hat{c}_\delta(x) = \sum_{\beta \in Q_+} c_{\beta, \delta} x^{-\beta}.$$

Lemma 2.9. *Assume we have a (non necessarily polynomial) solution $\Theta_\lambda(x)$ of the first Pieri rule (2.30), which admits a series expansion of the form $\Theta_\lambda(x) = x^\lambda \sum_{\beta \in Q_+} \tau_\beta(x) \Lambda^{-\beta}$ for $\lambda \in \mathbb{C}^N$. Then we have, for λ an integer partition:*

$$\Pi_\lambda(x) = \frac{\hat{c}_0(x)}{\tau_0(x)} \Theta_\lambda(x),$$

where $\hat{c}_0(x)$ is the leading coefficient in the series $\hat{p}_\lambda(x)$ that specializes to $\Pi_\lambda(x)$ for integer partitions λ .

Proof. Write $\Theta_\lambda(x) = x^\lambda \theta_\lambda(x)$. By use of (2.12) for $a = 1$, the Pieri equation turns into

$$\left(\sum_{i=1}^N x_i \{ (1 - \Lambda^{-\alpha_i}) T_i - 1 \} \right) \theta_\lambda(x) = 0.$$

This is a linear triangular system for the coefficients $\tau_\beta(x)$, which are uniquely determined for $\beta \neq 0$, proportional to $\tau_0(x)$. The same holds for the coefficients $\hat{c}_\beta(x)$ of $\hat{p}_\lambda(x)$ in terms of $\hat{c}_0(x)$. We deduce that $x^\lambda \hat{p}_\lambda(x) = \frac{\hat{c}_0(x)}{\tau_0(x)} \Theta_\lambda(x)$, and the Lemma follows by specialization. \square

Remark 2.10. *The argument used in this section prefigures the reformulation in terms of universal solutions performed in Sect. 2.13 below, and extends to all types.*

2.10. Fourier duality of g and γ . The final piece of information we need to prove Theorem 2.6 is that g acting on the q -Whittaker functions is equal to the Gaussian $\gamma(x)$ acting on the same functions:

Theorem 2.11.

$$(2.29) \quad g(\Lambda) \Pi_\lambda(x) = \gamma(x) \Pi_\lambda(x).$$

Proof. Acting with $g(\Lambda)$ on both sides of the first Pieri rule

$$(2.30) \quad H_1(\Lambda; q) \Pi_\lambda(x) = e_1(x) \Pi_\lambda(x)$$

and noting that g and γ commute with both $H_1(\Lambda; q)$ and $e_1(x)$, we see that $\gamma^{-1} g \Pi_\lambda(x)$ obeys the *same* first Pieri rule. We use the uniqueness argument of the previous section for the solutions of (2.30). Applying Lemma 2.9 to $\Theta_\lambda = \gamma^{-1} g x^\lambda \hat{p}_\lambda(x)$, the latter must be proportional to $\Pi_\lambda(x)$ when specialized to an integer partition λ . To determine the proportionality constant, we compute the leading coefficient of the series expansion $x^{-\lambda} \gamma^{-1} g x^\lambda \hat{p}_\lambda(x)$. Since g_Λ is a power series in $\{\Lambda^{-\alpha_i}\}$ with leading term 1, the leading term is determined by the action of g_T on $x^\lambda \hat{c}_0(x)$.

We claim that

$$g_T x^\lambda = \gamma x^\lambda g'_T,$$

where g'_T acts only by shifting Λ by powers of q , leaving the leading term $\hat{c}_0(x)$ unchanged.

To see this, let a, b be elements in an algebra such that $[a, [a, b]] = 0$ and $[b, [a, b]] = c$ commutes with both a and b . Then the Campbell-Hausdorff formula implies:

$$(2.31) \quad e^a e^b e^{-a} e^{-b} = e^{a+b+\frac{1}{2}[a,b]-\frac{1}{12}[b,[a,b]]} e^{-a-b+\frac{1}{2}[a,b]+\frac{1}{12}[b,[a,b]]} = e^{[a,b]+\frac{c}{2}}.$$

Choosing $e^a = g_T$ and $e^b = x^\lambda$, i.e. $a = \sum_i \frac{\partial_{\lambda_i}^2}{2 \log q}$ and $b = \sum_i \lambda_i \log x_i$ leads to $[a, b] = \sum_i \frac{\log x_i \partial_{\lambda_i}}{\log q}$ and $e^{\frac{1}{2}c} = \gamma(x)^{-1}$, which commutes with $e^{[a,b]}$. We finally write

$$e^a e^b = g_T x^\lambda = e^{\frac{c}{2}} e^{[a,b]} e^b e^a = \gamma^{-1} e^{\sum_i \frac{\log x_i \partial_{\lambda_i}}{\log q}} e^{\sum_i \lambda_i \log x_i} g_T = \gamma x^\lambda \prod_i T_i^{\frac{\log x_i}{\log q}} g_T = \gamma x^\lambda g'_T$$

where the last step uses again (2.31), with $a = \sum_i \frac{\log x_i \partial_{\lambda_i}}{\log q}$, $b = \sum_i \lambda_i \log x_i$ hence $e^{[a,b]} = \gamma^2$ and $c = 0$.

We conclude that $\gamma^{-1} g \Pi_\lambda(x) = \Pi_\lambda(x)$, and the theorem follows. \square

Remark 2.12. *The relation $g(\Lambda) \Pi_\lambda(x) = \gamma(x) \Pi_\lambda(x)$ is equivalent to the recursion relation for the quantities denoted J_β^μ in [FJMM09] (Theorem 3.1), which we denote as $J_\beta(x = q^\mu)$ below. These can be identified as the coefficients in the formal expansion of the series $\tilde{\Pi}_\lambda(x) := g_\Lambda \Pi_\lambda(x) = x^\lambda \sum_{\beta \in Q_+} J_\beta(x) \Lambda^{-\beta}$. Theorem 2.11 says that $g_T g_\Lambda \Pi_\lambda(x) = \gamma(x) \Pi_\lambda(x)$, which implies $(g_\Lambda g_T) \tilde{\Pi}_\lambda(x) = \gamma(x) \tilde{\Pi}_\lambda(x)$. As a recursion relation for coefficients, this means*

$$J_\beta^\mu = \sum_\delta \frac{\gamma(q^\delta)}{(q, q)_{\beta-\delta}} x^{-\delta} J_\delta^\mu$$

with the notation $(q, q)_\alpha = \prod_i (q, q)_{\alpha_i}$, where we have used the series expansion $g_\Lambda = \sum_{\alpha \in Q_+} \frac{\Lambda^{-\alpha}}{(q; q)_\alpha}$, as well as the relation $\gamma(x)^{-1} x^{-\lambda} g_T x^\lambda \Lambda^{-\delta} = \gamma(q^\delta) (\Lambda x)^{-\delta}$. A similar connection holds for certain other root systems, see Remark 4.13.

Corollary 2.13. *The (higher) q -difference Toda Hamiltonians $H_a(\Lambda)$ are algebraically independent conserved quantities of the opposite quantum Q -system (2.21).*

Proof. Multiply the Pieri equation $e_a(x) \Pi_\lambda(x) = H_a(\Lambda) \Pi_\lambda(x)$ on the left by $g\gamma^{-1}$, and use Theorem 2.11 to rewrite:

$$g\gamma^{-1} e_a(x) \Pi_\lambda(x) = e_a(x) \Pi_\lambda(x) = g H_a(\Lambda) g^{-1} \Pi_\lambda(x) = H_a(\Lambda) \Pi_\lambda(x),$$

hence g commutes with all $H_a(\Lambda)$, $a = 1, 2, \dots, N$. The latter are Laurent polynomials of the elements of the quantum torus \mathbb{T}_Λ , hence of the initial data $\bar{D}_{i;0}, \bar{D}_{i;1}$ as well, which are invariant under any time translation $(\bar{D}_{i;0}, \bar{D}_{i;1}) \mapsto (\bar{D}_{i;n}, \bar{D}_{i;n+1})$, $n \in \mathbb{Z}$. These are independent conserved quantities of the opposite quantum Q -system that governs the $\bar{D}_{a;n}$, as any dependence between $H_a(\Lambda)$ would imply a dependence between the $e_a(x)$ by inverse Fourier transform. \square

2.11. **Proof of Theorem 2.6.** The proof of the main Theorem 2.6 now follows.

Proof. Acting on $D_{a;0}(x; q)\Pi_\lambda(x)$ by $\gamma(x)^{-k}g^k(\Lambda)$, we have

$$\gamma^{-k}g^k D_{a;0}(x; q)\Pi_\lambda(x) = \gamma^{-k} D_{a;0}(x; q)g^k \Pi_\lambda(x) = \gamma^{-k} D_{a;0}(x; q)\gamma^k \Pi_\lambda(x) = q^{ak/2} D_{a;k}(x; q)\Pi_\lambda(x).$$

Using the eigenvalue equation (2.9), this is equal to

$$\gamma^{-k}g^k \bar{D}_{a;0}\Pi_\lambda(x) = g^k \bar{D}_{a;0}\gamma^{-k}\Pi_\lambda(x) = g^k \bar{D}_{a;0}g^{-k}\Pi_\lambda(x) = q^{ak/2} \bar{D}_{a;k}\Pi_\lambda(x),$$

which is the statement of Theorem 2.6. \square

2.12. **Pieri operators, conserved quantities and Toda Hamiltonians.** Let us compare the result of Corollary 2.13 with the conserved quantities of the (opposite) quantum Q-system of [DFK11, DFK10, DFK18]. Those references use a different normalization of the Q-system variables, resulting in the system

$$(2.32) \quad \begin{aligned} \bar{Q}_{a;n}\bar{Q}_{b;n+1} &= v^{-\Lambda_{a,b}}\bar{Q}_{b;n+1}\bar{Q}_{a;n} \\ v^{\Lambda_{a,a}}\bar{Q}_{a;n-1}\bar{Q}_{a;n+1} &= (\bar{Q}_{a;n})^2 - \bar{Q}_{a-1;n}\bar{Q}_{a+1;n} \end{aligned}$$

with $\Lambda_{a,b} = \min(a, b)(N - \max(a, b))$. The precise relation with the operators $\bar{D}_{a;k}(\Lambda; q)$ is

$$(2.33) \quad \bar{Q}_{a;n} = q^{-\frac{na^2}{2N}}(\bar{D}_{N;0})^{-\frac{a}{N}}\bar{D}_{a;n}$$

with $v = q^{1/N}$. The conserved quantities \bar{C}_m of the quantum system (2.32) are expressed in terms of the initial cluster $\{\bar{Q}_{a;0}, \bar{Q}_{a;1}\}$ as hard particle partition functions (i.e. generating polynomials of independent sets of vertices) on a graph (Figure 3 of [DFK10]) with ordered vertices labeled $1, 2, \dots, 2N - 1$, with a weight \bar{y}_i per occupied vertex labeled i , where:

$$\begin{aligned} \bar{y}_{2a-1} &= \bar{Q}_{a-1;0}(\bar{Q}_{a;0})^{-1}(\bar{Q}_{a-1;1})^{-1}\bar{Q}_{a;1} & (a = 1, 2, \dots, N), \\ \bar{y}_{2a} &= -\bar{Q}_{a-1;0}(\bar{Q}_{a;0})^{-1}(\bar{Q}_{a;1})^{-1}\bar{Q}_{a+1;1} & (a = 1, 2, \dots, N - 1). \end{aligned}$$

Using (2.33),

$$\bar{y}_{2a-1} = v^{-\frac{1}{2}}T_a, \quad \bar{y}_{2a} = -v^{-\frac{1}{2}}\frac{\Lambda_{a+1}}{\Lambda_a}T_{a+1},$$

hence the resulting conserved quantities are related to the Hamiltonians (2.12) by $H_m(\Lambda) = v^{m/2}\bar{C}_m$.

In Ref. [DFK18], it was shown that the conserved quantities \bar{C}_m can be interpreted as type A (relativistic) q -difference Toda Hamiltonians [Eti99]. This justifies the identification of the Pieri operators (2.12) as q -difference Toda Hamiltonians.

2.13. **Universal solutions and duality.** The results of the previous sections may be rephrased in the more uniform context of universal solutions, obtained explicitly in the case of type A in [NS12]. The following discussion treats the two sets x and s on equal footing, as formal variables, and implies that the duality property is more general.

2.13.1. *Universal Macdonald and Pieri solutions and duality.* We now consider $s = t^\rho q^\lambda$ as a *formal* variable which we may specialize to integer partitions λ , and similarly write $x = q^\mu t^\rho$. If we consider the difference operators (2.1) in the space $\mathbb{C}(q, t)[[\{x^{-\alpha_i}\}]][\Gamma_1, \dots, \Gamma_N]$, i.e. expanded as power series of the variables x_{i+1}/x_i with $1 \leq i \leq N-1$, then there is a unique series solution $P(x; s)$ to the Macdonald eigenvalue equation

$$(2.34) \quad \mathcal{D}_1(x; q, t) P(x; s) = e_1(x) P(x; s)$$

of the form

$$(2.35) \quad P(x; s) = q^{\lambda \cdot \mu} \sum_{\beta \in Q_+} c_\beta(s; q, t) x^{-\beta}, \quad c_0(s; q, t) = 1.$$

Note that $q^{\lambda \cdot \mu} = x^\lambda t^{-\rho \cdot \lambda} = s^\mu t^{-\rho \cdot \mu}$ is symmetric with respect to the interchange of x and s . The uniqueness of the solution follows from the fact that (2.34) is a nonsingular triangular system for the coefficients $c_\beta(s; q, t)$.

The series (2.35) is called the universal Macdonald solution because, under the specialization of the variables $s = q^\lambda t^\rho$ with λ an integer partition, the function $t^{\rho \cdot \lambda} P(x; s)$ specializes to the symmetric Macdonald polynomial $P_\lambda(x)$:

$$(2.36) \quad P_\lambda(x) = t^{\rho \cdot \lambda} P(x; q^\lambda t^\rho), \quad \lambda = (\lambda_1 \geq \dots \geq \lambda_N), \quad \lambda_i \in \mathbb{Z}_{\geq 0}.$$

That is, the infinite series (2.35) truncates to a finite number of terms, and the prefactor $t^{\rho \cdot \lambda} q^{\lambda \cdot \mu} = x^\lambda$ ensures that the function is a symmetric polynomial of x_1, \dots, x_N . All Macdonald polynomials $P_\lambda(x)$ are obtained as specializations of $P(x; s)$. On the other hand, the series $P(x; s)$ also specializes to the Baker-Akhiezer quasi-polynomials introduced by Chalykh [Cha02]. In this case, the relevant specialization consists in taking $t = q^{-k}$, k some positive integer. One can check that another truncation occurs leaving us with a finite sum.

Example 2.14. *Let us illustrate the phenomenon of truncation in the simplest case of $A_1^{(1)}$ of rank $N = 2$. The universal Macdonald series is expressed in terms of the variables $u = x_2/x_1$ and $v = \Lambda_2/\Lambda_1$ as*

$$P(x; s) = x_1^{\lambda_1} x_2^{\lambda_2} t^{-\lambda_1} \sum_{n=0}^{\infty} u^n \prod_{i=0}^{n-1} \frac{1 - vq^i}{1 - t^{-1}vq^i} \frac{1 - t^{-1}q^{-i}}{1 - q^{-i-1}}.$$

The first type of truncation occurs for $v = q^{\lambda_2 - \lambda_1} = q^{-k}$ for some integer $k \geq 0$. We see that the first factor in the numerator of the coefficient in the series vanishes as soon as $n > k$ (for $i = k$). The second type of truncation occurs when $t = q^{-k}$ for some integer $k \geq 0$ and arbitrary λ . In that case the second factor vanishes as soon as $n > k$.

Similarly, if we consider $\mathcal{H}_1(\Lambda; q, t)$ as a power series in $\{s_{i+1}/s_i\}$, then the equation

$$(2.37) \quad t^{-\rho \cdot \lambda} \mathcal{H}_1(\Lambda; q, t) t^{\rho \cdot \lambda} Q(s; x) = e_1(x) Q(s; x)$$

has a unique ‘‘universal Pieri’’ solution of the form

$$(2.38) \quad Q(s; x) = q^{\lambda\mu} \sum_{\beta \in Q_+} \bar{c}_\beta(x; q, t) s^{-\beta}, \quad \bar{c}_0(x; q, t) = 1.$$

An outcome of the study of [NS12] is a relation between the universal Macdonald and Pieri solutions, which we re-prove below.

Theorem 2.15. *The universal Macdonald and Pieri solutions are related via*

$$P(x; s) = \Delta(x) Q(s; x),$$

with $\Delta(x)$ as in (2.4).

Proof. First note that $P(x; s)$ obey both Macdonald eigenvalue and Pieri equations [Che95b, Cha02]. Expanding the coefficients of $P(x; s)$ as series of $s^{-\alpha_i}$: $c_\beta(s) = \sum_{\delta \in Q_+} c_{\beta, \delta} s^{-\delta}$ allows to rewrite $P(x; s) = q^{\lambda\mu} \sum_{\delta \in Q_+} \hat{c}_\delta(x) s^{-\delta}$, where $\hat{c}_\delta(x) = \sum_{\beta \in Q_+} c_{\beta, \delta} x^{-\beta}$. Now both $P(x; s)$ and $Q(s; x)$ can be viewed as two different solutions of the Pieri equation. By uniqueness, they must be proportional up to an s -independent factor. We deduce that $P(x; s) = \hat{c}_0(x) Q(s; x)$. To compute $\hat{c}_0(x)$, note that it is the limit of $q^{-\lambda\mu} P(x; s) = \hat{c}_0(x) + O(s^{-\alpha_i})$, when we take $|s_1| \gg |s_2| \gg \dots |s_N| \gg 1$, so that all $s^{-\alpha_i} \rightarrow 0$. Using the a -th Macdonald eigenvalue equation (2.2) and the explicit formula (2.1), and dividing by $\Lambda_1 \Lambda_2 \dots \Lambda_a$, we have for $a = 1, 2, \dots, N$:

$$\left\{ t^{-\binom{a}{2}} \frac{e_a(s)}{\Lambda_1 \Lambda_2 \dots \Lambda_a} - \sum_{\substack{I \subset [1, N] \\ |I|=a}} \prod_{\substack{j \notin I \\ i \in I}} \frac{tx_i - x_j}{x_i - x_j} \frac{1}{\Lambda_1 \Lambda_2 \dots \Lambda_a} \prod_{i \in I} \Lambda_i \Gamma_i \right\} (\hat{c}_0(x) + O(x^{\alpha_i})) = 0.$$

In the limit $\Lambda_{i+1}/\Lambda_i \rightarrow 0$, all the terms tend to zero except for the leading term in the eigenvalue, and the term corresponding to $I = \{1, 2, \dots, a\}$ in the sum:

$$(2.39) \quad \left\{ 1 - \prod_{1 \leq i < a < j \leq N} \frac{1 - \frac{x_j}{tx_i}}{1 - \frac{x_j}{x_i}} \Gamma_1 \Gamma_2 \dots \Gamma_a \right\} \hat{c}_0(x) = 0.$$

We verify that $\hat{c}_0(x)$ and $\Delta(x)$ of (2.4) both obey (2.39) for $a = 1, 2, \dots, N$. Their ratio must therefore be independent of x as it is invariant under each Γ_i , and is easily identified to 1 by taking the limit when all $x^{-\alpha_i} \rightarrow 0$ and using $c_{0,0} = 1$. The Theorem follows. \square

Repeating the argument of Section 2.2 starting with (2.34), including the interchange of the labels x and s , we see that

$$t^{-\rho\lambda} \mathcal{H}_1(\Lambda; q, t) t^{\rho\lambda} Q(x; s) = e_1(x) Q(x; s),$$

where the expansion in s of $Q(x; s) = P(s; x)/\Delta(s)$ has the form $q^{\lambda\mu}(1 + \mathcal{O}(s^{-\alpha_i}))$. By uniqueness of the solution,

$$(2.40) \quad Q(x; s) = Q(s; x) \quad \Leftrightarrow \quad \Delta(s) P(x; s) = \Delta(x) P(s; x).$$

The duality (2.40) states that the series $Q(s; x)$ in s is equal to the series $Q(x; s)$ in x . The specialization of λ and μ to integer partitions reduces to the Macdonald polynomial duality relation (2.3). We may therefore interpret (2.40) as a universal extension of the duality of Macdonald theory.

2.13.2. *q-Whittaker limit.* The symmetry (2.40) is replaced in the $t \rightarrow \infty$ limit by

$$(2.41) \quad \Pi(x; \Lambda) = \overline{\Delta}(x)K(\Lambda; x), \quad \overline{\Delta}(x) = \prod_{\alpha \in R_+} (qx^{-\alpha}; q)_\infty,$$

where the universal q -Whittaker function $\Pi(x; \Lambda)$ and Pieri solution $K(\Lambda; x)$ read respectively

$$\Pi(x; \Lambda) = \lim_{t \rightarrow \infty} t^{\rho\lambda} P(x; s), \quad K(\Lambda; x) = \lim_{t \rightarrow \infty} t^{\rho\lambda} Q(s; x)$$

and satisfy the limiting Macdonald eigenvalue equations and Pieri rules:

$$(2.42) \quad D_a(x; q) \Pi(x; \Lambda) = \Lambda^{\omega_a} \Pi(x; \Lambda)$$

$$(2.43) \quad H_a(\Lambda; q) K(\Lambda; x) = e_a(x) K(\Lambda; x)$$

As in the case of the Macdonald function, the universal functions $\Pi(x; \Lambda)$ and $K(\Lambda; x)$ are uniquely determined from the difference equations (2.42) and (2.43) as series of the form

$$(2.44) \quad \Pi(x; \Lambda) = x^\lambda \sum_{\beta \in Q_+} c_\beta(\Lambda; q) x^{-\beta}, \quad c_0(\Lambda; q) = 1,$$

$$(2.45) \quad K(\Lambda; x) = x^\lambda \sum_{\beta \in Q_+} \bar{c}_\beta(x; q) \Lambda^{-\beta}, \quad \bar{c}_0(x; q) = 1.$$

Example 2.16. *Let us compute $K(\Lambda; x)$ explicitly as a solution of the first Pieri equation $H_1(\Lambda; q) K(\Lambda; x) = e_1(x) K(\Lambda; x)$ with $H_1(\Lambda; q)$ as in (2.12) for $a = 1$, by use of a path model. The equation reduces to the following triangular system for the coefficients $\bar{c}_\beta(x; q)$ from (2.45):*

$$\left\{ \sum_{i=1}^N x_i (q^{\beta_i-1-\beta_i} - 1) \right\} \bar{c}_\beta(x; q) = \sum_{i=1}^{N-1} x_{i+1} q^{\beta_i-\beta_{i+1}-1} \bar{c}_{\beta-e_i}(x; q),$$

with $\beta = \sum_{i=1}^{N-1} \beta_i e_i$ and $\beta_0 = \beta_N = 0$ by convention. The system is nonsingular for x generic, and we may express $\bar{c}_\beta(x; q)$ as a path model partition function, namely as the sum over all paths p from 0 to β in the positive quadrant \mathbb{Z}_+^{N-1} of path weights $w(p)$. The path weight is defined as a product along the path of its vertex weights $w_v(a) = (\sum_{i=1}^n x_i (q^{a_i-1-a_i} - 1))^{-1}$ per vertex a visited and edge step weights $w_e(s) = x_{i+1} q^{b_i-b_{i+1}-1}$ per edge step $s = b - e_i \rightarrow b$:

$$\bar{c}_\beta(x; q) = \sum_{\text{paths } p: 0 \rightarrow \beta} \prod_{\text{vertex } a \text{ of } p} w_v(a) \prod_{\text{edge step } s \text{ of } p} w_e(s)$$

This construction parallels that of Whittaker vectors performed in [DFKT17].

The Fourier transform may be restated in terms of the universal q -Whittaker function as follows: $f(x) \Pi(x; \Lambda) = \bar{f}(\Lambda) \Pi(x; \Lambda)$, and our main theorem 2.6 as $D_{a;k}(x; q) \Pi(x; \Lambda) = \bar{D}_{a;k}(\Lambda; q) \Pi(x; \Lambda)$, also a consequence of $g(\Lambda) \Pi(x; \Lambda) = \gamma(x) \Pi(x; \Lambda)$.

3. DUALITY PROPERTY FOR OTHER CLASSICAL ROOT SYSTEMS

In this section we present a generalization of the methods of Section 2 for the root systems corresponding to the affine algebras in the classical series, listed in Table 1. In these cases, the underlying double affine Hecke algebra is of BC -type, corresponding to finite-type Weyl groups of types C_N or D_N , and the commuting q -difference operators and their eigenfunctions are Koornwinder/Macdonald operators and polynomials. A key ingredient is the duality property, which implies a relation between the Koornwinder/Macdonald eigenvalue equations and the Pieri rules [vD96, Sah99, Che97].

Our goal is to choose a set of N commuting difference operators for each algebra \mathfrak{g} such that, in the q -Whittaker limit, these difference operators and their $SL(2, \mathbb{Z})$ -translates satisfy the \mathfrak{g} -type quantum Q-system. There are several known constructions of the difference operators. Koornwinder introduced a first order operator [Koo92] depending on generic parameters (q, t, a, b, c, d) , which was extended by van Diejen to a complete set of N commuting difference operators [vD95]. Upon specialization of the parameters (a, b, c, d) , these correspond to the root systems of \mathfrak{g} . In some cases, van Diejen's operators must be supplemented by operators introduced by Macdonald [Mac01], who used the structure of specific root systems to produce difference operators for miniscule co-weights. We combine these constructions and define a set of N commuting \mathfrak{g} -Macdonald operators in Appendix A, whose eigenvalues form a basis for the space of fundamental characters of the finite algebra $R^* = R(\mathfrak{g}^*)$ (see Table 1).

This section is organized as follows. We introduce the Koornwinder operators for generic values of (a, b, c, d) in (Section 3.1). The specialization of the parameters corresponding to \mathfrak{g} of Table 1 are explained in Section 3.2. Finally, the q -Whittaker limit of the \mathfrak{g} -Macdonald operators is described in Section 3.3, together with their τ_+ -translates. We will prove that the τ_+ translation is the discrete time evolution of the associated quantum Q-systems in Section 4.

3.1. Koornwinder Operators, Polynomials and Pieri rules.

3.1.1. *The Koornwinder operators.* Let \mathbb{F} be the field of rational functions in the indeterminates $a, b, c, d, q^{\frac{1}{2}}, t^{\frac{1}{2}}$ and W the Weyl group of type C_N . It acts on functions in $\mathbb{F}(x_1, \dots, x_N)$ by permutations and inversions of the variables. The W -invariant space of Laurent polynomials in x is generated by the elementary symmetric functions $\hat{e}(x)$, defined via the

generating function

$$(3.1) \quad \hat{E}(z; x) := \prod_{i=1}^N \prod_{\epsilon=\pm 1} (1 + zx_i^\epsilon) = \sum_{k=0}^{2N} z^k \hat{e}_k(x),$$

so that $\hat{e}_1(x) = \sum_{i=1}^n (x_i + x_i^{-1})$. The space $\mathbb{F}[x, x^{-1}]^W$ is preserved by the action of the Koornwinder operator, defined as follows.

Definition 3.1. *The Koornwinder operator $\mathcal{K}_1^{(a,b,c,d)}(x; q, t)$ is the q -difference operator acting on $\mathbb{F}[x, x^{-1}]$:*

$$(3.2) \quad \mathcal{K}_1^{(a,b,c,d)}(x; q, t) = \sum_{i=1}^N \sum_{\epsilon=\pm 1} \Phi_{i,\epsilon}^{(a,b,c,d)}(x) (\Gamma_i^\epsilon - 1), \quad \Gamma_i x_j = q^{\delta_{ij}} x_j \Gamma_i,$$

where

$$\Phi_{i,\epsilon}^{(a,b,c,d)}(x) = \frac{(1 - ax_i^\epsilon)(1 - bx_i^\epsilon)(1 - cx_i^\epsilon)(1 - dx_i^\epsilon)}{(1 - x_i^{2\epsilon})(1 - qx_i^{2\epsilon})} \prod_{j \neq i} \frac{tx_i^\epsilon - x_j}{x_i^\epsilon - x_j} \frac{tx_i^\epsilon x_j - 1}{x_i^\epsilon x_j - 1}.$$

Koornwinder polynomials are the unique monic, symmetric, Laurent polynomial eigenfunctions of the eigenvalue equation

$$\mathcal{K}_1^{(a,b,c,d)}(x; q, t) P_\lambda^{(a,b,c,d)}(x) = \sigma t^{N-1} (\hat{e}_1(s) - \hat{e}_1(\sigma t^\rho)) P_\lambda^{(a,b,c,d)}(x)$$

where λ is any integer partition coding the leading term x^λ of $P_\lambda^{(a,b,c,d)}(x)$. We make use of the notations

$$(3.3) \quad s_i = q^{\lambda_i} t^{\rho_i} \sigma, \quad \sigma = \sqrt{abcd/q}, \quad \rho_i = N - i \quad (i = 1, 2, \dots, N).$$

As an example, the first two Koornwinder polynomials are $P_\emptyset^{(a,b,c,d)}(x) = 1$, with eigenvalue 0, and

$$P_{1,0,0,\dots,0}^{(a,b,c,d)}(x) = \hat{e}_1(x) + \frac{1 - t^N}{1 - t} \frac{abcd(a^{-1} + b^{-1} + c^{-1} + d^{-1})t^{N-1} - (a + b + c + d)}{1 - abcd t^{2N-2}}.$$

3.1.2. Koornwinder-Macdonald operators. We define the Koornwinder-Macdonald operators to be the set of mutually commuting difference operators, which commute with the Koornwinder operator, chosen so that their eigenvalues are proportional to the elementary symmetric functions $\hat{e}_m(s)$. The first order Koornwinder-Macdonald operator (A.23) is

$$(3.4) \quad \mathcal{D}_1^{(a,b,c,d)}(x; q, t) = \mathcal{K}_1^{(a,b,c,d)}(x; q, t) + \frac{1 - t^N}{1 - t} \left(1 + \frac{abcd}{q} t^{N-1} \right).$$

The additive constant in (3.4) is $\sigma t^{N-1} \hat{e}_1(\sigma t^\rho)$, so that Koornwinder polynomials satisfy the eigenvalue equation

$$(3.5) \quad \mathcal{D}_1^{(a,b,c,d)}(x; q, t) P_\lambda^{(a,b,c,d)}(x) = \sigma t^{N-1} \hat{e}_1(s) P_\lambda^{(a,b,c,d)}(x).$$

Definition 3.2. *The Koornwinder-Macdonald operators are a commuting family of difference operators $\{\mathcal{D}_m^{(a,b,c,d)}(x; q, t) : m \in [1, N]\}$, which are linear combinations of van Diejen's commuting difference operators [vD96], uniquely defined by their eigenvalues:*

$$(3.6) \quad \mathcal{D}_m^{(a,b,c,d)}(x; q, t) P_\lambda^{(a,b,c,d)}(x) = d_{\lambda;m}^{(a,b,c,d)} P_\lambda^{(a,b,c,d)}(x), \quad d_{\lambda;m}^{(a,b,c,d)} = \sigma^m t^{m(N-\frac{m+1}{2})} \hat{e}_m(s).$$

These operators are explicitly constructed in Appendix A, see Definitions A.2, A.8 and Theorem A.9.

3.1.3. Rains operators. A useful alternative “ N -th order” difference operator was constructed by Rains [Rai05]⁵ in his study of BC-symmetric polynomials. Rains shows that the operators

$$(3.7) \quad \mathcal{R}_N^{(u,v)}(x; q, t) = \sum_{\epsilon_1, \epsilon_2, \dots, \epsilon_N = \pm 1} \prod_{i=1}^N \frac{(1 - ux_i^{\epsilon_i})(1 - vx_i^{\epsilon_i})}{1 - x_i^{2\epsilon_i}} \prod_{1 \leq i < j \leq n} \frac{1 - tx_i^{\epsilon_i} x_j^{\epsilon_j}}{1 - x_i^{\epsilon_i} x_j^{\epsilon_j}} \prod_{i=1}^N \Gamma_i^{\epsilon_i/2}$$

are maps between W -invariant spaces with different values of the parameters (a, b, c, d) . They act between bases of Koornwinder polynomials as follows:

$$(3.8) \quad \begin{aligned} \mathcal{R}_N^{(q^{-\frac{1}{2}a, q^{-\frac{1}{2}b})}(x; q, t) P_\lambda^{(a,b,c,d)}(x) &= q^{-|\lambda|/2} \prod_{i=1}^N (1 - abq^{\lambda_i-1} t^{N-i}) P_\lambda^{(q^{-\frac{1}{2}a, q^{-\frac{1}{2}b, q^{\frac{1}{2}c, q^{\frac{1}{2}d})}(x), \\ \mathcal{R}_N^{(q^{-\frac{1}{2}c, q^{-\frac{1}{2}d})}(x; q, t) P_\lambda^{(a,b,c,d)}(x) &= q^{-|\lambda|/2} \prod_{i=1}^N (1 - cdq^{\lambda_i-1} t^{N-i}) P_\lambda^{(q^{\frac{1}{2}a, q^{\frac{1}{2}b, q^{-\frac{1}{2}c, q^{-\frac{1}{2}d})}(x), \end{aligned}$$

where $|\lambda| = \sum_{i=1}^N \lambda_i$. The product

$$(3.9) \quad \widehat{\mathcal{D}}_N^{(a,b,c,d)}(x; q, t) = \mathcal{R}_N^{(a,b)}(x; q, t) \mathcal{R}_N^{(q^{-\frac{1}{2}c, q^{-\frac{1}{2}d})}(x; q, t)$$

commutes with the Koornwinder operators, since Koornwinder polynomials satisfy the eigenvalue equation

$$\widehat{\mathcal{D}}_N^{(a,b,c,d)}(x; q, t) P_\lambda^{(a,b,c,d)}(x) = \hat{d}_{\lambda;N}^{(a,b,c,d)} P_\lambda^{(a,b,c,d)}(x),$$

with eigenvalues

$$(3.10) \quad \hat{d}_{\lambda;N}^{(a,b,c,d)} = q^{-|\lambda|} \prod_{i=1}^N (1 - abq^{\lambda_i} t^{N-i})(1 - cdq^{\lambda_i-1} t^{N-i}).$$

The operator (3.9) is not linearly independent of the set of Koornwinder-Macdonald operators: See Section A.2.3, Lemma A.10 for the explicit expression. However, its factored form will play a crucial role in the proof of our main theorems below.

⁵Some of these operators actually appeared in earlier works of van Diejen, but Rains' construction is more systematic.

3.1.4. *Duality.* Due to the existence of an anti-involution $*$ of the DAHA, Koornwinder polynomials obey a remarkable duality property [vD96, Sah99]. The involution acts on the parameters (a, b, c, d) as

$$(3.11) \quad a^* = \left(\frac{abcd}{q}\right)^{1/2}, \quad b^* = -\left(q\frac{ab}{cd}\right)^{1/2}, \quad c^* = \left(q\frac{ac}{bd}\right)^{1/2}, \quad d^* = -\left(q\frac{ad}{bc}\right)^{1/2},$$

so that $\sigma^* = a$. The duality property for Koornwinder polynomials is

$$(3.12) \quad \frac{P_\lambda^{(a,b,c,d)}(q^\mu t^{\rho^*})}{P_\lambda^{(a,b,c,d)}(t^{\rho^*})} = \frac{P_\mu^{(a^*,b^*,c^*,d^*)}(q^\lambda t^\rho)}{P_\mu^{(a^*,b^*,c^*,d^*)}(t^\rho)}.$$

where μ, λ are both integer partitions. ρ, ρ^* are defined by

$$(3.13) \quad t^{\rho_i} = t^{\rho_i^{(a,b,c,d)}} = \sigma t^{N-i}, \quad t^{\rho_i^*} = t^{\rho_i^{(a^*,b^*,c^*,d^*)}} = at^{N-i}.$$

3.1.5. *Pieri rules.* Using the duality relation (3.12), one obtains the Pieri rules for Koornwinder polynomials with parameters (a, b, c, d) from the eigenvalue equation with parameters (a^*, b^*, c^*, d^*) . It is useful to define the function

$$(3.14) \quad \Delta^{(a,b,c,d)}(x) := \prod_{i=1}^N \frac{\left(\frac{q}{x_i^2}; q\right)_\infty}{\left(\frac{q}{ax_i}; q\right)_\infty \left(\frac{q}{bx_i}; q\right)_\infty \left(\frac{q}{cx_i}; q\right)_\infty \left(\frac{q}{dx_i}; q\right)_\infty} \prod_{1 \leq i < j \leq N} \prod_{\epsilon = \pm 1} \frac{\left(\frac{qx_j^\epsilon}{x_i}; q\right)_\infty}{\left(\frac{qx_i^\epsilon}{tx_j}; q\right)_\infty}.$$

The normalization factor in the duality (3.12) is given by (See Theorem 5.1 of [vD96]):

$$(3.15) \quad P_\lambda^{(a,b,c,d)}(t^{\rho^*}) = t^{\rho^* \cdot \lambda} \frac{\Delta^{(a^*,b^*,c^*,d^*)}(t^\rho)}{\Delta^{(a^*,b^*,c^*,d^*)}(q^\lambda t^\rho)}, \quad P_\mu^{(a^*,b^*,c^*,d^*)}(t^\rho) = t^{\rho \cdot \mu} \frac{\Delta^{(a,b,c,d)}(t^{\rho^*})}{\Delta^{(a,b,c,d)}(q^\mu t^{\rho^*})},$$

Using the parametrization $x = q^\mu t^{\rho^*}$ in the eigenvalue equation (3.6),

$$(3.16) \quad \mathcal{D}_m(q^\mu t^{\rho^*}; q, t) P_\lambda(q^\mu t^{\rho^*}) = \theta_m \hat{e}_m(q^\lambda t^\rho) P_\lambda(q^\mu t^{\rho^*}), \quad m \in [1, N],$$

where $\theta_m = \sigma^m t^{m(N - \frac{m+1}{2})}$ and $\Gamma_i : \mu_j \mapsto \mu_j + \delta_{i,j}$ in the difference operator $\mathcal{D}_m = \mathcal{D}_m^{(a,b,c,d)}$. Upon specializing μ to integer partitions, we use the duality relation (3.12), where we replace the normalization factor by (3.15). Equation (3.16) becomes

$$\begin{aligned} \theta_m \hat{e}_m(q^\lambda t^\rho) P_\mu^*(q^\lambda t^\rho) &= P_\mu^*(t^\rho) \mathcal{D}_m(q^\mu t^{\rho^*}; q, t) P_\mu^*(t^\rho)^{-1} P_\mu^*(q^\lambda t^\rho) \\ &= (t^{\rho \cdot \mu} \Delta(q^\mu t^{\rho^*})^{-1} \mathcal{D}_m(q^\mu t^{\rho^*}; q, t) \Delta(q^\mu t^{\rho^*}) t^{-\rho \cdot \mu}) P_\mu^*(q^\lambda t^\rho). \end{aligned}$$

Acting with the involution $*$ on the parameters and interchanging the roles of λ and μ (i.e. x and s) above, we arrive at the following Pieri rules.

Theorem 3.3. *The Pieri rules for the Koornwinder polynomials are*

$$(3.17) \quad \mathcal{H}_m^{(a,b,c,d)}(s; q, t) P_\lambda^{(a,b,c,d)}(x) = \hat{e}_m(x) P_\lambda^{(a,b,c,d)}(x),$$

| \mathfrak{g} | \mathfrak{g}^* | a | b | c | d | R | S | R^* | $\xi_{\mathfrak{g}}$ |
|------------------|------------------|-------------------|--------------------|-----------------------------------|------------------------------------|--------|-------|--------|----------------------|
| $D_N^{(1)}$ | $D_N^{(1)}$ | 1 | -1 | $q^{\frac{1}{2}}$ | $-q^{\frac{1}{2}}$ | D_N | D_N | D_N | 0 |
| $B_N^{(1)}$ | $C_N^{(1)}$ | t | -1 | $q^{\frac{1}{2}}$ | $-q^{\frac{1}{2}}$ | B_N | B_N | C_N | $\frac{1}{2}$ |
| $C_N^{(1)}$ | $B_N^{(1)}$ | $t^{\frac{1}{2}}$ | $-t^{\frac{1}{2}}$ | $t^{\frac{1}{2}} q^{\frac{1}{2}}$ | $-t^{\frac{1}{2}} q^{\frac{1}{2}}$ | C_N | C_N | B_N | 1 |
| $A_{2N-1}^{(2)}$ | $A_{2N-1}^{(2)}$ | $t^{\frac{1}{2}}$ | $-t^{\frac{1}{2}}$ | $q^{\frac{1}{2}}$ | $-q^{\frac{1}{2}}$ | C_N | B_N | C_N | $\frac{1}{2}$ |
| $D_{N+1}^{(2)}$ | $D_{N+1}^{(2)}$ | t | -1 | $t q^{\frac{1}{2}}$ | $-q^{\frac{1}{2}}$ | B_N | C_N | B_N | 1 |
| $A_{2N}^{(2)}$ | $A_{2N}^{(2)}$ | t | -1 | $t^{\frac{1}{2}} q^{\frac{1}{2}}$ | $-t^{\frac{1}{2}} q^{\frac{1}{2}}$ | BC_N | - | BC_N | 1 |

TABLE 1. Specialization of the Koornwinder parameters a, b, c, d corresponding to the affine algebra \mathfrak{g} . The pair (R, S) refer to a pair of classical root systems corresponding to Macdonald's notation [Mac01], used in Appendix A, except for $A_{2n}^{(2)}$.

where the Pieri operators $\mathcal{H}_m^{(a,b,c,d)}(s; q, t)$ are

$$(3.18) \quad \mathcal{H}_m^{(a,b,c,d)}(s; q, t) = \frac{1}{\theta_m^*} t^{\rho^* \cdot \lambda} \Delta^{(a^*, b^*, c^*, d^*)}(s)^{-1} \mathcal{D}_m^{(a^*, b^*, c^*, d^*)}(s; q, t) \Delta^{(a^*, b^*, c^*, d^*)}(s) t^{-\rho^* \cdot \lambda}.$$

In the difference operators \mathcal{H}_m , s is specialized to $q^\lambda t^\rho$ with λ an integer partition, $\theta_m^* = a^m t^{m(N - \frac{m+1}{2})}$, and $T_i : \lambda_j \mapsto \lambda_j + \delta_{i,j}$.

The explicit first Pieri operator $\mathcal{H}_1^{(a,b,c,d)}(s; q, t)$ is derived in Theorem B.1 of Appendix B.

3.2. Specialization of the parameters (a, b, c, d) .

3.2.1. *\mathfrak{g} -Macdonald operators and polynomials.* The Macdonald operators for type \mathfrak{g} are a set of N commuting operators which commute with the specialization of the Koornwinder operators at values of (a, b, c, d) indicated in Table 1. The list of operators is given in Definition A.8. In most cases, these are just the specialized Koornwinder-Macdonald operators, but for a few exceptions, where the operators are chosen so that their eigenvalues be fundamental characters of the finite-dimensional algebra $R^* = R(\mathfrak{g}^*)$. This occurs when $R^* = B_N$ or D_N .

| R | $\hat{e}_a^{(R)}(x)$ |
|-------------|---|
| D_N | $\hat{e}_1, \dots, \hat{e}_{N-2}, \hat{e}_{N-1}^{(D_N)}, \hat{e}_N^{(D_N)}$ |
| B_N | $\hat{e}_1, \hat{e}_2, \dots, \hat{e}_{N-1}, \hat{e}_N^{(B_N)}$ |
| C_N, BC_N | $\hat{e}_1, \hat{e}_2, \dots, \hat{e}_N$ |

TABLE 2. The list of chosen symmetric functions forming a basis for the fundamental characters of the finite Lie algebra R . The exceptional symmetric functions in the table are given in (A.32).

Remark 3.4. *The case of $A_{2N}^{(2)}$ is special, as $R = BC_N = R^*$ is non-reduced. In this case, we must use the set of simple roots $\alpha_a = \alpha_a^*$ for B_N , and the set of fundamental weights $\omega_a = \omega_a^*$ for C_N .*

The principle for the choice of \mathfrak{g} -Macdonald operators is that their spectrum generates the Grothendieck ring of the algebra R^* associated to each \mathfrak{g} . The choice is not unique. Define

$$(3.19) \quad N_{\mathfrak{g}} := \begin{cases} N, & \text{for } \mathfrak{g} = A_{N-1}^{(1)}, B_N^{(1)}, A_{2N-1}^{(2)}, A_{2N}^{(2)}, \\ N-1, & \text{for } \mathfrak{g} = C_N^{(1)}, D_{N+1}^{(2)}, \\ N-2, & \text{for } \mathfrak{g} = D_N^{(1)}. \end{cases}$$

Definition 3.5. *For $m \leq N_{\mathfrak{g}}$, the \mathfrak{g} -Macdonald operators $\mathcal{D}_m^{(\mathfrak{g})} = \mathcal{D}_m^{(\mathfrak{g})}(x)$ are*

$$\mathcal{D}_m^{(\mathfrak{g})} := \mathcal{D}_m^{(a,b,c,d)}, \quad m \in [1, N_{\mathfrak{g}}],$$

where $\mathcal{D}_m^{(a,b,c,d)}$ are as in Definition 3.2 with (a, b, c, d) specialized according Table 1. For $N_{\mathfrak{g}} < m \leq N$, we use the operators constructed by Macdonald, described in Section A.1:

$$(3.20) \quad D_N^{(1)} : \quad \mathcal{D}_a^{(D_N^{(1)})} := \mathcal{E}_{\omega_a}^{(D_N^{(1)})}, \quad a = N-1, N;$$

$$(3.21) \quad C_N^{(1)} : \quad \mathcal{D}_N^{(C_N^{(1)})} := \mathcal{E}_{\omega_N}^{(C_N^{(1)})},$$

$$(3.22) \quad D_{N+1}^{(2)} : \quad \mathcal{D}_N^{(D_{N+1}^{(2)})} := \mathcal{E}_{\omega_N}^{(D_{N+1}^{(2)})}.$$

where $\mathcal{E}_{\omega_a}^{(D_N^{(1)})}$, $\mathcal{E}_{\omega_N}^{(C_N^{(1)})}$, $\mathcal{E}_{\omega_N}^{(D_{N+1}^{(2)})}$ are as in (A.4-A.5), (A.6), (A.7) respectively.

The choice of higher commuting difference operators (3.5) is such that the eigenvalue equation for the m th \mathfrak{g} -Macdonald operator is (see Theorem A.15):

$$(3.23) \quad \mathcal{D}_m^{(\mathfrak{g})}(x; q, t) P_\lambda^{(\mathfrak{g})}(x) = \theta_m^{(\mathfrak{g})} \hat{e}_m^{(R^*)}(s) P_\lambda^{(\mathfrak{g})}(x),$$

where $\theta_m^{(\mathfrak{g})}$ are as in (A.34-A.35) and $\hat{e}_m^{(R)}(x)$ are listed in Table 2 (as these involve the fundamental weights we choose the same eigenvalues for type BC_N and C_N , see Remark 3.4).

In particular, the first \mathfrak{g} -Macdonald operators are simply the specialization of the Koornwinder-Macdonald operator (3.4).

$$(3.24) \quad \mathcal{D}_1^{(\mathfrak{g})}(x; q, t) = t^{\rho_1^{\mathfrak{g}}} \hat{e}_1(t^{\rho_1^{\mathfrak{g}}}) + \sum_{i=1}^N \sum_{\epsilon=\pm 1} \phi_{i,\epsilon}^{(\mathfrak{g})}(x; q, t) (\Gamma_i^\epsilon - 1).$$

The factor σ under the specialization is $t^{\xi_{\mathfrak{g}}}$ (see Table 1), so that

$$(3.25) \quad s_i = q^{\lambda_i} t^{N+\xi_{\mathfrak{g}}-i} = q^{\lambda_i} t^{\rho_i^{(\mathfrak{g})}}, \quad (i = 1, 2, \dots, N),$$

where $\rho^{(\mathfrak{g})} = \rho(S)$ is the half-sum of positive roots of S (except when $\mathfrak{g} = A_{2N}^{(2)}$). The functions $\phi_{i,\epsilon}^{(\mathfrak{g})}(x; q, t)$ are

$$\phi_{i,\epsilon}^{(\mathfrak{g})}(x; q, t) = \prod_{j \neq i} \prod_{\epsilon'=\pm 1} \frac{tx_i^\epsilon x_j^{\epsilon'} - 1}{x_i^\epsilon x_j^{\epsilon'} - 1} \times \begin{cases} 1, & \mathfrak{g} = D_N^{(1)}; \\ \frac{tx_i^\epsilon - 1}{x_i^\epsilon - 1}, & \mathfrak{g} = B_N^{(1)}; \\ \frac{tx_i^{2\epsilon} - 1}{x_i^{2\epsilon} - 1} \frac{tqx_i^{2\epsilon} - 1}{qx_i^{2\epsilon} - 1}, & \mathfrak{g} = C_N^{(1)}; \\ \frac{tx_i^{2\epsilon} - 1}{x_i^{2\epsilon} - 1}, & \mathfrak{g} = A_{2N-1}^{(2)}; \\ \frac{tx_i^\epsilon - 1}{x_i^\epsilon - 1} \frac{tq^{1/2}x_i^\epsilon - 1}{q^{1/2}x_i^\epsilon - 1}, & \mathfrak{g} = D_{N+1}^{(2)}; \\ \frac{tx_i^\epsilon - 1}{x_i^\epsilon - 1} \frac{tqx_i^{2\epsilon} - 1}{qx_i^{2\epsilon} - 1}, & \mathfrak{g} = A_{2N}^{(2)}. \end{cases}$$

The unique monic eigenfunction $P_\lambda^{(\mathfrak{g})}(x)$ of $\mathcal{D}_1^{(\mathfrak{g})}(x; q, t)$ with eigenvalue $t^{\rho_1^{\mathfrak{g}}} \hat{e}_1(s)$, where λ is \mathfrak{g} -partition, is the \mathfrak{g} -type Macdonald (Laurent) polynomial.

Definition 3.6. A \mathfrak{g} -partition is a set $\lambda = (\lambda_1, \dots, \lambda_N)$ such that $\sum_i \lambda_i e_i$ is a dominant integral weight of $R = R(\mathfrak{g})$.

| G | affine roots in \widehat{R}_{++} |
|------------------|---|
| $D_N^{(1)}$ | $n\delta + (e_i \pm e_j) \quad (1 \leq i < j \leq N; n \geq 1)$ |
| $B_N^{(1)}$ | $n\delta + e_i \quad (1 \leq i \leq N), n\delta + (e_i \pm e_j) \quad (1 \leq i < j \leq N; n \geq 1)$ |
| $C_N^{(1)}$ | $n\delta + 2e_i \quad (1 \leq i \leq N), n\delta + (e_i \pm e_j) \quad (1 \leq i < j \leq N; n \geq 1)$ |
| $A_{2N-1}^{(2)}$ | $2n\delta + 2e_i \quad (1 \leq i \leq N), n\delta + (e_i \pm e_j) \quad (1 \leq i < j \leq N; n \geq 1)$ |
| $D_{N+1}^{(2)}$ | $\frac{n}{2}\delta + e_i \quad (1 \leq i \leq N), n\delta + (e_i \pm e_j) \quad (1 \leq i < j \leq N; n \geq 1)$ |
| $A_{2N}^{(2)}$ | $n\delta + e_i, (2n-1)\delta + 2e_i \quad (1 \leq i \leq N), n\delta + (e_i \pm e_j) \quad (1 \leq i < j \leq N; n \geq 1)$ |

TABLE 3. The subset \widehat{R}_{++} of the affine roots for each affine algebra \mathfrak{g} .

The \mathfrak{g} -partitions are simply integer partitions except in the cases $R = B_N, D_N$, where

$$\lambda \in \begin{cases} \{(\mathbb{Z}_+)^{N-1} \times \mathbb{Z}\} \cup \{(\mathbb{Z}_{+\frac{1}{2}})^{N-1} \times (\mathbb{Z}_{+\frac{1}{2}})\}, & \lambda_1 \geq \cdots \geq \lambda_{N-1} \geq |\lambda_N|, \quad R = D_N; \\ (\mathbb{Z}_+)^N \cup (\mathbb{Z}_{+\frac{1}{2}})^N, & \lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_N \geq 0, \quad R = B_N. \end{cases}$$

Remark 3.7. *The \mathfrak{g} -Macdonald polynomial $P_\lambda^{(\mathfrak{g})}(x)$ with λ an integer partition is the specialization of the Koornwinder polynomial, indexed by the same partition, to the parameters of Table 1. For non-integer partitions, \mathfrak{g} -Macdonald polynomials can be obtained from a different specialization of the parameters [vD95], see Section 4.3.3 in type B. Alternatively, we can use the specialization of the universal functions of Section 5, see remark 5.2.*

3.2.2. Duality. The involution $*$ acting on the parameters (a, b, c, d) implies an involution $\mathfrak{g} \mapsto \mathfrak{g}^*$ and $R \mapsto R^*$, as listed in Table 1. In particular, $\mathfrak{g} = \mathfrak{g}^*$ except in the cases $(C_N^{(1)})^* = B_N^{(1)}$ and $(B_N^{(1)})^* = C_N^{(1)}$.

The duality relation (3.12), specialized to \mathfrak{g} -Macdonald polynomials is

$$(3.26) \quad \frac{P_\lambda^{(\mathfrak{g})}(q^\mu t^\rho)}{P_\lambda^{(\mathfrak{g})}(t^\rho)} = \frac{P_\mu^{(\mathfrak{g}^*)}(q^\lambda t^\rho)}{P_\mu^{(\mathfrak{g}^*)}(t^\rho)}.$$

Note, however, that the range of validity of (3.26) is wider, as λ can be any \mathfrak{g} -partition, while μ is any \mathfrak{g}^* -partition, and these are not necessarily integer partitions as in (3.12). First conjectured by Macdonald, the duality (3.26) was successively proved for all types in the case of integer partitions: for type A, it appears in [Mac95], for other types the main proof is in [Che95b], supplemented by [vD96] and [Sah99] upon specialization of (3.12). See Section 5 for a more general duality statement.

For $\alpha = n\delta + \alpha_1 + \cdots + \alpha_N \in \widehat{R}$, define $x^\alpha = q^{-n}x_1^{\alpha_1} \cdots x_N^{\alpha_N}$. The function Δ in Equation (3.14) has the following specializations:

$$(3.27) \quad \Delta^{(\mathfrak{g})}(x) = \prod_{\alpha \in \widehat{R}_{++}} \frac{1 - x^{-\alpha}}{1 - t^{-1}x^{-\alpha}},$$

where $\widehat{R}_{++} = \{\alpha + n\delta \in \widehat{R} : \alpha \in R_+, n > 0\}$ is a subset of the affine roots of \mathfrak{g} , see Table 3. In all cases but $\mathfrak{g} = A_{2N}^{(2)}$, Δ is closely related to a simplified version of Macdonald's function Δ^+ [Mac01], suitable for taking the dual q -Whittaker limit $t \rightarrow \infty$. The function Δ enters the duality relation (3.26) via the evaluation formulas (see (0.6) in [Che95b])

$$(3.28) \quad P_\lambda^{(\mathfrak{g})}(t^{\rho^*}) = t^{\rho^* \cdot \lambda} \frac{\Delta^{(\mathfrak{g}^*)}(t^\rho)}{\Delta^{(\mathfrak{g}^*)}(q^\lambda t^\rho)}, \quad P_\mu^{(\mathfrak{g}^*)}(t^\rho) = t^{\rho \cdot \mu} \frac{\Delta^{(\mathfrak{g})}(t^{\rho^*})}{\Delta^{(\mathfrak{g})}(q^\mu t^{\rho^*})}.$$

3.2.3. Pieri rules. As in the case of generic (a, b, c, d) , the Pieri rules follow from the eigenvalue equations (3.23) and the duality (3.26). Start with (3.23) with $x = q^\mu t^{\rho^*}$, and specialize μ to a \mathfrak{g}^* -partition:

$$(3.29) \quad \mathcal{D}_m^{(\mathfrak{g})}(q^\mu t^{\rho^*}; q, t) P_\lambda^{(\mathfrak{g})}(q^\mu t^{\rho^*}) = \theta_m^{(\mathfrak{g})} \hat{e}_m^{(R^*)}(q^\lambda t^\rho) P_\lambda^{(\mathfrak{g})}(q^\mu t^{\rho^*}),$$

where (x, Γ) is specialized to $(q^\mu t^{\rho^*}, e^{\partial_\mu})$. Following the same steps as in Section 3.1.5, using the duality relation (3.26), one obtains the \mathfrak{g} -Pieri formulas:

Theorem 3.8. *The Pieri rules for \mathfrak{g} -Macdonald polynomials are*

$$(3.30) \quad \mathcal{H}_m^{(\mathfrak{g})}(s; q, t) P_\lambda^{(\mathfrak{g})}(x) = \hat{e}_m^{(R)}(x) P_\lambda^{(\mathfrak{g})}(x),$$

where the difference operators $\mathcal{H}_m^{(\mathfrak{g})}(s; q, t)$ are given in terms of $\mathcal{D}_m^{(\mathfrak{g}^*)}(s; q, t)$:

$$(3.31) \quad \mathcal{H}_m^{(\mathfrak{g})}(s; q, t) = \frac{1}{\theta_m^{(\mathfrak{g}^*)}} t^{\rho^* \cdot \lambda} \Delta^{(\mathfrak{g}^*)}(s)^{-1} \mathcal{D}_m^{(\mathfrak{g}^*)}(s; q, t) \Delta^{(\mathfrak{g}^*)}(s) t^{-\rho^* \cdot \lambda}.$$

The explicit Pieri operators $\mathcal{H}_1^{(\mathfrak{g})}(s; q, t)$ are listed in Section B.2.

3.3. The q -Whittaker limit. The Macdonald polynomials have a symmetry $(t, q) \mapsto (t^{-1}, q^{-1})$. For certain root systems, the q -Whittaker polynomials are the $t \rightarrow 0$ limit of the Macdonald polynomials, and therefore the q^{-1} -Whittaker polynomials are the $t \rightarrow \infty$ limit of these polynomials. In this paper, we define the various functions and operators so that they have well-defined limits as $t \rightarrow \infty$. We refer to this as the q -Whittaker limit by slight abuse of terminology. We also call q -Whittaker polynomials what technically should be called q^{-1} -Whittaker polynomials.

3.3.1. *Whittaker difference operators.* The eigenvalue equations and Pieri rules of the previous section have well-defined q -Whittaker limits. Define

$$(3.32) \quad \Pi_\lambda^{(\mathfrak{g})}(x) = \lim_{t \rightarrow \infty} P_\lambda^{(\mathfrak{g})}(x)$$

and

$$(3.33) \quad D_a^{(\mathfrak{g})}(x; q) = \lim_{t \rightarrow \infty} (\theta_a^{(\mathfrak{g})})^{-2} \mathcal{D}_a^{(\mathfrak{g})}(x; q, t), \quad (a = 1, 2, \dots, N).$$

The first q -Whittaker difference operators are obtained from Equation (3.2.1):

$$(3.34) \quad D_1^{(\mathfrak{g})}(x; q) = 1 + \sum_{\substack{i=1 \\ \epsilon=\pm 1}}^N \phi_{i,\epsilon}^{(\mathfrak{g})}(x) (\Gamma_i^\epsilon - 1),$$

with

$$(3.35) \quad \phi_{i,\epsilon}^{(\mathfrak{g})}(x) = \prod_{j \neq i} \frac{x_i^\epsilon}{x_i^\epsilon - x_j} \frac{x_i^\epsilon x_j}{x_i^\epsilon x_j - 1} \times \begin{cases} 1 & (D_N^{(1)}); \\ \frac{x_i^\epsilon}{x_i^\epsilon - 1}, & (B_N^{(1)}), \\ \frac{x_i^{2\epsilon}}{x_i^{2\epsilon} - 1} \frac{qx_i^{2\epsilon}}{qx_i^{2\epsilon} - 1}, & (C_N^{(1)}), \\ \frac{x_i^{2\epsilon}}{x_i^{2\epsilon} - 1}, & (A_{2N-1}^{(2)}), \\ \frac{x_i^\epsilon}{x_i^\epsilon - 1} \frac{q^{\frac{1}{2}} x_i^\epsilon}{q^{\frac{1}{2}} x_i^\epsilon - 1}, & (D_{N+1}^{(2)}), \\ \frac{x_i^\epsilon}{x_i^\epsilon - 1} \frac{qx_i^{2\epsilon}}{qx_i^{2\epsilon} - 1}, & (A_{2N}^{(2)}). \end{cases}$$

The limiting eigenvalues of $D_m^{(\mathfrak{g})}(x; q)$ are not symmetric functions because of the dependence of s on t . Instead, they are given by the dominant term in t in the functions $\hat{e}_a^{(R^*)}(s)$, $\Lambda^{\omega_a^*}$. The eigenvalue equations are (see Section A.4, Theorem A.18)

$$(3.36) \quad D_a^{(\mathfrak{g})}(x; q) \Pi_\lambda^{(\mathfrak{g})}(x) = \Lambda^{\omega_a^*} \Pi_\lambda^{(\mathfrak{g})}(x).$$

3.3.2. *Pieri rules and Toda Hamiltonians.* The Pieri operators (3.31) have well-defined limits as $t \rightarrow \infty$. Let

$$(3.37) \quad H_m^{(\mathfrak{g})}(\Lambda; q) := \lim_{t \rightarrow \infty} \mathcal{H}_m^{(\mathfrak{g})}(s = \Lambda t^{\rho^{(\mathfrak{g})}}; q, t).$$

Using the Pieri operators from Section B.2 (see also Remark B.3), and taking the $t \rightarrow \infty$ limit, we obtain the following list of first Pieri operators.

Theorem 3.9. *For all \mathfrak{g} , the first Pieri operators take the form*

$$(3.38) \quad H_1^{(\mathfrak{g})}(\Lambda; q) = \sum_{i=1}^N (1 - \Lambda^{-\alpha_{i-1}^*}) T_i + \sum_{i=1}^{N_{R^*}} (1 - \Lambda^{-\alpha_i^*}) T_i^{-1} + M^{(\mathfrak{g})}(\Lambda; q),$$

where the roots α_i^* are the simple roots of R^* and $\alpha_0^* = 0$ by convention. Here, $N_{R^*} = N - 1$ if $R^* = B_N$, $N - 2$ if $R^* = D_N$, and is equal to N otherwise. The boundary terms $M^{(\mathfrak{g})}(\Lambda; q)$ are as follows:

$$(3.39) \quad M^{(\mathfrak{g})}(\Lambda; q) = \begin{cases} (1 - \Lambda^{-\alpha_N^*})(T_N^{-1} + (1 - \Lambda^{-\alpha_{N-1}^*})T_{N-1}^{-1}), & \mathfrak{g} = D_N^{(1)}, \\ (1 - \Lambda^{-\alpha_N^*})(1 - q\Lambda^{-\alpha_N^*})T_N^{-1} + \Lambda^{-\alpha_N^*}(q^{-1}\Lambda^{-\alpha_{N-1}^*} - (1 + q^{-1})), & \mathfrak{g} = B_N^{(1)}, \\ 0, & \mathfrak{g} = C_N^{(1)}, A_{2N-1}^{(2)}, \\ (1 - \Lambda^{-\alpha_N^*})(1 - q^{\frac{1}{2}}\Lambda^{-\alpha_N^*})T_N^{-1}, & \mathfrak{g} = D_{N+1}^{(2)}, \\ -\Lambda^{-\alpha_N^*}, & \mathfrak{g} = A_{2N}^{(2)}. \end{cases}$$

Explicitly, the first Pieri operators are:

$$\begin{aligned} H_1^{(D_N^{(1)})}(\Lambda) &= T_1 + \sum_{i=2}^N \left(1 - \frac{\Lambda_i}{\Lambda_{i-1}}\right) T_i + \sum_{i=1}^{N-2} \left(1 - \frac{\Lambda_{i+1}}{\Lambda_i}\right) T_i^{-1} \\ &\quad + \left(1 - \frac{\Lambda_N}{\Lambda_{N-1}}\right) \left(1 - \frac{1}{\Lambda_{N-1}\Lambda_N}\right) T_{N-1}^{-1} + \left(1 - \frac{1}{\Lambda_{N-1}\Lambda_N}\right) T_N^{-1}, \\ H_1^{(B_N^{(1)})}(\Lambda) &= T_1 + \sum_{i=2}^N \left(1 - \frac{\Lambda_i}{\Lambda_{i-1}}\right) T_i + \sum_{i=1}^{N-1} \left(1 - \frac{\Lambda_{i+1}}{\Lambda_i}\right) T_i^{-1} \\ &\quad + \left(1 - \frac{1}{\Lambda_N^2}\right) \left(1 - \frac{q}{\Lambda_N^2}\right) T_N^{-1} + \frac{q^{-1}}{\Lambda_{N-1}\Lambda_N} - \frac{1 + q^{-1}}{\Lambda_N^2}, \\ H_1^{(C_N^{(1)})}(\Lambda) &= T_1 + \sum_{i=2}^N \left(1 - \frac{\Lambda_i}{\Lambda_{i-1}}\right) T_i + \sum_{i=1}^{N-1} \left(1 - \frac{\Lambda_{i+1}}{\Lambda_i}\right) T_i^{-1} + \left(1 - \frac{1}{\Lambda_N}\right) T_N^{-1}, \\ H_1^{(A_{2N-1}^{(2)})}(\Lambda) &= T_1 + \sum_{i=2}^N \left(1 - \frac{\Lambda_i}{\Lambda_{i-1}}\right) T_i + \sum_{i=1}^{N-1} \left(1 - \frac{\Lambda_{i+1}}{\Lambda_i}\right) T_i^{-1} + \left(1 - \frac{1}{\Lambda_N^2}\right) T_N^{-1}, \\ H_1^{(D_{N+1}^{(2)})}(\Lambda) &= T_1 + \sum_{i=2}^N \left(1 - \frac{\Lambda_i}{\Lambda_{i-1}}\right) T_i + \sum_{i=1}^{N-1} \left(1 - \frac{\Lambda_{i+1}}{\Lambda_i}\right) T_i^{-1} \\ &\quad + \left(1 - \frac{1}{\Lambda_N}\right) \left(1 - \frac{q^{\frac{1}{2}}}{\Lambda_N}\right) T_N^{-1} - \frac{1 + q^{-\frac{1}{2}}}{\Lambda_N}, \\ H_1^{(A_{2N}^{(2)})}(\Lambda) &= T_1 + \sum_{i=2}^N \left(1 - \frac{\Lambda_i}{\Lambda_{i-1}}\right) T_i + \sum_{i=1}^{N-1} \left(1 - \frac{\Lambda_{i+1}}{\Lambda_i}\right) T_i^{-1} + \left(1 - \frac{1}{\Lambda_N}\right) T_N^{-1} - \frac{1}{\Lambda_N}. \end{aligned}$$

The first q -Whittaker Pieri rule is

$$(3.40) \quad H_1^{(\mathfrak{g})}(\Lambda) \Pi_\lambda^{(\mathfrak{g})}(x) = \hat{e}_1(x) \Pi_\lambda^{(\mathfrak{g})}(x).$$

Alternatively this equation can be interpreted as the eigenvector equation for q -Whittaker functions in which the roles of variables Λ and x are interchanged.

Remark 3.10. *The Pieri operators $H_1^{(\mathfrak{g})}(\Lambda; q)$ match the relativistic $U_q(R)$ -Toda Hamiltonians acting on functions of Λ [vDE15, GT21] upon the following correspondence:*

$$D_N^{(1)} \rightarrow D_N, \quad D_{N+1}^{(2)} \rightarrow B_N, \quad A_{2N-1}^{(2)} \rightarrow C_N.$$

The equivalence uses the Etingof automorphism [Eti99] $T_i \mapsto T_{2\varpi_i}$, $\Lambda_i \mapsto \Lambda_i T_{-\varpi_i}$ and $v = q^{\frac{1}{2}}$ in the notations of [GT21]. To our knowledge, the cases $\mathfrak{g} = B_N^{(1)}, C_N^{(1)}, A_{2N}^{(2)}$ do not appear in the literature in relation to standard constructions of q -Whittaker functions for quantum groups. By a slight abuse of terminology, we still call the corresponding limits of Macdonald polynomials q -Whittaker functions, and the limiting Pieri operators Toda Hamiltonians, and keep our labeling with (twisted) affine algebras to avoid confusion. The same correspondence in the $q = 1$ limit occurs in relation to factorization dynamics [Wil15], where the B_N, C_N cases match classical Q -system evolutions for the twisted algebras $\mathfrak{g} = D_{N+1}^{(2)}, A_{2N-1}^{(2)}$ respectively.

3.3.3. Generalized q -Whittaker difference operators. In this section, we consider the $\tau_+ \in SL(2, \mathbb{Z})$ -action on the q -Whittaker difference operators of Section 3.3.1. Let $\gamma(x)$ be as in Equation (2.16). We will define “translated” operators $D_{a;n}^{(\mathfrak{g})}$ for all $a = 1, 2, \dots, N$ and $n \in \mathbb{Z}$, by suitable use of iterated conjugations with γ^{-1} . There is a subtlety arising from a distinction according to whether a is a long or short label. These do not necessarily correspond to long and short roots of R , but are determined instead by the Q -system evolutions described in the next section.

Definition 3.11. *All labels $a \in [1, N]$ are long except for the following cases: $a = N$ for $B_N^{(1)}$, $a \in [1, N-1]$ for $C_N^{(1)}$, and $a \in [1, N]$ for $A_{2N}^{(2)}$.*

In general, we think of the integer n as a discrete time, and as $\text{Ad}_{\gamma^{-1}}$ as a “time translation” operator.

Definition of $D_{1;n}^{(\mathfrak{g})}(x)$ and properties. We start with the definition of the operators $D_{1;n}^{(\mathfrak{g})}(x)$, which depends on whether $a = 1$ is a long or short label.

Definition 3.12. *When the label $a = 1$ is long ($\mathfrak{g} = D_N^{(1)}, B_N^{(1)}, A_{2N-1}^{(2)}, D_{N+1}^{(2)}$), define, for any $n \in \mathbb{Z}$,*

$$(3.41) \quad D_{1;n}^{(\mathfrak{g})}(x) := q^{-\frac{n}{2}} \gamma^{-n} D_1^{(\mathfrak{g})}(x; q) \gamma^n = q^{-\frac{n}{2}} + \sum_{i, \epsilon} \phi_{i, \epsilon}^{(\mathfrak{g})}(x) (x_i^{n\epsilon} \Gamma_i^\epsilon - q^{-\frac{n}{2}}),$$

with $\phi_{i, \epsilon}^{(\mathfrak{g})}(x)$ as in (3.35).

Definition 3.13. When the label $a = 1$ is short ($\mathfrak{g} = C_N^{(1)}, A_{2N}^{(2)}$) define

$$(3.42) \quad D_{1;-1}^{(\mathfrak{g})}(x) = \sum_{i,\epsilon} \phi_{i,\epsilon}^{(\mathfrak{g})}(x) x_i^{-\epsilon} (\Gamma_i^\epsilon - 1),$$

with $\phi_{i,\epsilon}^{(\mathfrak{g})}(x)$ as in (3.35). Then for all $n \in \mathbb{Z}$ and for $i = 0, -1$, define

$$(3.43) \quad D_{1;2n+i}^{(\mathfrak{g})}(x) = q^{-n} \gamma^{-2n} D_{1,i}^{(\mathfrak{g})}(x; q) \gamma^{2n},$$

where $D_{1,0}^{(\mathfrak{g})}(x; q) = D_1^{(\mathfrak{g})}(x; q)$.

Conjugating (3.34) and (3.42) by γ^{-2n} , we have, when $\mathfrak{g} = C_N^{(1)}, A_{2N}^{(2)}$,

$$(3.44) \quad D_{1;2n}^{(\mathfrak{g})}(x) = q^{-n} + \sum_{i,\epsilon} \phi_{i,\epsilon}^{(\mathfrak{g})}(x) (x_i^{2n\epsilon} \Gamma_i^\epsilon - q^{-n}),$$

$$(3.45) \quad D_{1;2n-1}^{(\mathfrak{g})}(x) = \sum_{i,\epsilon} \phi_{i,\epsilon}^{(\mathfrak{g})}(x) x_i^{-\epsilon} (x_i^{2n\epsilon} \Gamma_i^\epsilon - q^{-n}).$$

The main difference between the cases of Definitions 3.12 and 3.13 is that in the latter case the translation splits into distinct even and odd time evolution equations (3.44-3.45).

Lemma 3.14. We have the commutation relation

$$(3.46) \quad [D_{1;n}^{(\mathfrak{g})}(x), \hat{e}_1(x)] = (q-1) D_{1;n+1}^{(\mathfrak{g})}(x) + (q^{-1}-1) D_{1;n-1}^{(\mathfrak{g})}(x).$$

Proof. Using

$$[\Gamma_i^\epsilon, \hat{e}_1(x)] = (q^\epsilon - 1) x_i^\epsilon \Gamma_i^\epsilon + (q^{-\epsilon} - 1) x_i^{-\epsilon} \Gamma_i^\epsilon = \sum_{\epsilon'} (q^{\epsilon'} - 1) x_i^{\epsilon\epsilon'} \Gamma_i^\epsilon.$$

we have

$$[D_{1,n}^{(\mathfrak{g})}, e_1^{(x)}] = \sum_{i=1}^N \sum_{\epsilon=\pm 1} \phi_{i,\epsilon} x^{n\epsilon} [\Gamma_i^\epsilon, \hat{e}_1(x)] = \sum_{i,\epsilon,\epsilon'} \phi_{i,\epsilon} x^{(n+\epsilon')\epsilon} (q^{\epsilon'} - 1) \Gamma_i^\epsilon.$$

□

Definition of higher generalized q -Whittaker operators.

The higher, generalized q -Whittaker operators are denoted by $D_{a;n}^{(\mathfrak{g})}(x)$, with $a \in [1, N]$ and $n \in \mathbb{Z}$. They are defined as follows.

Definition 3.15. For all $a \in [1, N]$ and for all n ,

$$D_{a,t_a n}^{(\mathfrak{g})}(x) = q^{-nt_a \omega_a^* \cdot \omega_a / 2} \gamma^{-t_1 n} D_a^{(\mathfrak{g})}(x) \gamma^{t_1 n},$$

where $D_a^{(\mathfrak{g})}(x)$ are defined in (3.33) and ω_a (ω_a^*) are fundamental weights of R (R^*), $t_a = 2$ for short labels and $t_a = 1$ for long labels in Definition 3.11.

Definition 3.16. For the short labels $a \geq 2$ of $C_N^{(1)}$ and $A_{2N}^{(2)}$, define

$$(3.47) \quad D_{a;-1}^{(\mathfrak{g})} := \frac{(-1)^a}{q-1} [D_{1;-a}^{(\mathfrak{g})}, D_{a-1;0}^{(\mathfrak{g})}]_{q^a}, \quad (n \in \mathbb{Z}),$$

where $[A, B]_p = [A, B]_p = AB - pBA$. Then

$$D_{a;2n-1}^{(\mathfrak{g})} = q^{-na} \gamma^{-2n} D_{a;-1}^{(\mathfrak{g})} \gamma^{2n} = \frac{(-1)^a}{q-1} [D_{1;2n-a}^{(\mathfrak{g})}, D_{a-1;2n}^{(\mathfrak{g})}]_{q^a}.$$

Definition 3.16 anticipates on the desired connection to the quantum Q-systems of the next section and uses a recursive reformulation of the quantum Q-system solution obtained in [DFK17b].

Finally, to define the generalized difference operators with odd n corresponding to the label $a = N$, $\mathfrak{g} = B_N^{(1)}$, we use the Rains operators $\mathcal{R}_N^{(u,v)}$ and $\widehat{\mathcal{D}}_N^{(a,b,c,d)}$ of (3.7) and (3.9), with the specialization $(a, b, c, d) = (t, -1, q^{1/2}, -q^{1/2})$:

$$\widehat{\mathcal{D}}_N^{(B_N^{(1)})} = \widehat{\mathcal{D}}_N^{(t, -1, q^{1/2}, -q^{1/2})} = \mathcal{R}_N^{(1)} \mathcal{R}_N^{(0)}$$

with

$$\mathcal{R}_N^{(1)} = \mathcal{R}_N^{(t, -1)}, \quad \mathcal{R}_N^{(0)} = \mathcal{R}_N^{(1, -1)}.$$

In the q -Whittaker limit $t \rightarrow \infty$, these become

$$(3.48) \quad \widehat{\mathcal{D}}_N^{(B_N^{(1)})} = \lim_{t \rightarrow \infty} t^{-N^2} \widehat{\mathcal{D}}_N^{(B_N^{(1)})} = R_N^{(1)} R_N^{(0)},$$

with

$$(3.49) \quad R_N^{(0)} = \lim_{t \rightarrow \infty} t^{-N(N-1)/2} \mathcal{R}_N^{(1, -1)} = \sum_{\epsilon_1, \dots, \epsilon_N = \pm 1} \prod_{1 \leq i < j \leq N} \frac{x_i^{\epsilon_i} x_j^{\epsilon_j}}{x_i^{\epsilon_i} x_j^{\epsilon_j} - 1} \prod_{i=1}^N \Gamma_i^{\epsilon_i/2}$$

$$(3.50) \quad R_N^{(1)} = \lim_{t \rightarrow \infty} t^{-N(N+1)/2} \mathcal{R}_N^{(t, -1)} = \sum_{\epsilon_1, \dots, \epsilon_N = \pm 1} \prod_{i=1}^N \frac{x_i^{\epsilon_i}}{x_i^{\epsilon_i} - 1} \prod_{1 \leq i < j \leq N} \frac{x_i^{\epsilon_i} x_j^{\epsilon_j}}{x_i^{\epsilon_i} x_j^{\epsilon_j} - 1} \prod_{i=1}^N \Gamma_i^{\epsilon_i/2}.$$

The factorized form of $\widehat{\mathcal{D}}_N^{(B_N^{(1)})}$ can be used to define the discrete time evolution as follows:

Definition 3.17. Let $n \in \mathbb{Z}$. Define

$$(3.51) \quad \begin{aligned} R_{N;n}^{(i)} &= q^{-\frac{Nn}{8}} \gamma^{-n} R_N^{(i)} \gamma^n, \quad (i = 0, 1), \\ D_{N;2n-1}^{(B_N^{(1)})} &= q^{-\frac{Nn}{4}} R_{N;n-1}^{(1)} R_{N;n}^{(0)}. \end{aligned}$$

Note in particular that $D_{N;2n-1}^{(B_N^{(1)})} = q^{-\frac{nN}{2}} \gamma^{-n} D_{N;-1}^{(B_N^{(1)})} \gamma^n$, similarly to the even n case where $D_{N;2n}^{(B_N^{(1)})} = q^{-\frac{nN}{2}} \gamma^{-n} D_{N;0}^{(B_N^{(1)})} \gamma^n$.

4. QUANTUM Q-SYSTEMS AND MACDONALD OPERATOR CONJECTURE

For any affine algebra \mathfrak{g} , there is a corresponding Q-system [HKO⁺99, HKO⁺02], which is a recursion relation satisfied by the characters of the KR-modules [CM06] of the Yangian $Y(\mathfrak{g})$. Their deformation into recursion relations for non-commuting variables, called quantum Q-systems, were first defined [DFK14] using the identification of the Q-system recursion relations as mutations in a cluster algebra. In that case, one may use the canonical quantization of the cluster algebra [BZ05], and in [DFK14, Lin21, Lin] it is shown that this quantization is related to the graded fusion characters for \mathfrak{g} . In special cases, these are known to be related to q -Whittaker functions.

One of the main conjectures presented in [DFK21] is that for untwisted \mathfrak{g} in Table 1, some q -difference operators $D_{a;n}^{(\mathfrak{g})}(x)$ satisfy the quantum Q-system relations of type \mathfrak{g} , and that the operators $D_{a;1}^{(\mathfrak{g})}$ and $D_{a;-1}^{(\mathfrak{g})}$ act as raising and lowering operators when acting on the eigenfunctions $\Pi_\lambda^{(\mathfrak{g})}(\mathbf{x})$ of $D_{a;0}^{(\mathfrak{g})}$. The purpose of this section is to prove these statements for all \mathfrak{g} of Table 1.

4.1. Statement of the main theorems.

4.1.1. *Quantum Q-systems.* For each \mathfrak{g} , we consider an algebra generated by invertible, non-commuting elements $\{Q_{a;n}^{\pm 1} : a \in [1, N], n \in \mathbb{Z}\}$. The quantum Q-system of type \mathfrak{g} is a set of relations among these generators, which depend on the root data of \mathfrak{g} .

Definition 4.1. For each \mathfrak{g} , let $\Lambda^{(\mathfrak{g})}$ be the $N \times N$ matrix defined by

$$(4.1) \quad \Lambda_{a,b}^{(\mathfrak{g})} = \omega_a^* \cdot \omega_b, \quad a, b \in [1, N],$$

where ω_a and ω_a^* are the fundamental weights of type R and R^* respectively, and \cdot is the standard inner product.

In the case of untwisted \mathfrak{g} , $\Lambda^{(\mathfrak{g})}$ is the inverse of the Cartan matrix C of R , since $\omega_a^* = \omega_a^\vee$. For the case $\mathfrak{g} = D_{N+1}^{(2)}$, $\Lambda^{(\mathfrak{g})}$ is the inverse of the *symmetrized* Cartan matrix of type B_N , and for $\mathfrak{g} = A_M^{(2)}$, it is the inverse of the symmetrized Cartan matrix of type $C_{\lfloor \frac{M+1}{2} \rfloor}$:

$$\begin{aligned} \Lambda_{a,b}^{(D_{N+1}^{(2)})} &= (D_{a,a})^{-1} \Lambda_{a,b}^{(B_N^{(1)})} = \begin{cases} \min(a, b), & (a, b < N) \\ \frac{1}{2} \min(a, b), & (a = N, \text{ or } b = N) \\ \frac{1}{4} N, & (a = b = N), \end{cases} \\ \Lambda_{a,b}^{(A_{2N-1}^{(2)})} &= \Lambda_{a,b}^{(A_{2N}^{(2)})} = (D'_{a,a})^{-1} \Lambda_{a,b}^{(C_N^{(1)})} = \min(a, b). \end{aligned}$$

Here, $D = \text{diag}(1, 1, \dots, 1, 2)$ and $D' = D^{-1}$ are such that $C = C^{(B_N)} D$ and $C = C^{(C_N)} D'$ are symmetric. These are chosen so that $\Lambda_{1,1}^{(\mathfrak{g})} = 1$. Defining $t_a = 2$ for α_a a short simple root of R , $t_a = 1$ for α_a long, t_1/t_a is $D_{a,a}$ for type B and $D'_{a,a}$ for type C .

The quantum Q -system relations are of two types. The first are q -commutation relations:

$$(4.2) \quad \mathcal{Q}_{a;ta k+i} \mathcal{Q}_{b;tb k+j} = q^{\Lambda_{a,b}^{(\mathfrak{g})} j - \Lambda_{b,a}^{(\mathfrak{g})} i} \mathcal{Q}_{b;tb k+j} \mathcal{Q}_{a;ta k+i}, \quad (i, j = 0, 1, k \in \mathbb{Z}).$$

The second type of relations are evolution equations in the discrete time variable n , and have the form $q^{\Lambda_{aa}} \mathcal{Q}_{a;n+1} \mathcal{Q}_{a;n-1} = \mathcal{Q}_{a;n}^2 - \mathcal{T}_{a;n}$ for some monomials $\mathcal{T}_{a;n}$. Let $\bar{N}_{\mathfrak{g}}$ be the maximal integer such that $(\Lambda^{(\mathfrak{g})})_{a,a} = a$ for $a \leq \bar{N}_{\mathfrak{g}} + 1$:

$$\bar{N}_{D_N^{(1)}} = N - 3, \quad \bar{N}_{\mathfrak{g}} = N - 2, \quad (\mathfrak{g} = B_N^{(1)}, C_N^{(1)}, D_{N+1}^{(2)}), \quad \bar{N}_{\mathfrak{g}} = N - 1, \quad (\mathfrak{g} = A_{2N}^{(2)}, A_{2N-1}^{(2)}).$$

The evolution equations are

$$(4.3) \quad q^a \mathcal{Q}_{a;n+1} \mathcal{Q}_{a;n-1} = \mathcal{Q}_{a;n}^2 - \mathcal{Q}_{a+1;n} \mathcal{Q}_{a-1;n}, \quad a \in [1, \bar{N}_{\mathfrak{g}}], \text{ all } \mathfrak{g}.$$

$$(4.4) \quad \begin{aligned} D_N^{(1)} : \quad & q^{N-2} \mathcal{Q}_{N-2;n+1} \mathcal{Q}_{N-2;n-1} = \mathcal{Q}_{N-2;n}^2 - q^{-\frac{(N-2)n}{4}} \mathcal{Q}_{N-3;n} \mathcal{Q}_{N-1;n} \mathcal{Q}_{N;n}, \\ & q^{\frac{N}{4}} \mathcal{Q}_{N-1;n+1} \mathcal{Q}_{N-1;n-1} = \mathcal{Q}_{N-1;n}^2 - q^{\frac{(N-4)n}{4}} \mathcal{Q}_{N-2;n}, \\ & q^{\frac{N}{4}} \mathcal{Q}_{N;n+1} \mathcal{Q}_{N;n-1} = \mathcal{Q}_{N;n}^2 - q^{\frac{(N-4)n}{4}} \mathcal{Q}_{N-2;n}, \\ B_N^{(1)} : \quad & q^{N-1} \mathcal{Q}_{N-1;n+1} \mathcal{Q}_{N-1;n-1} = \mathcal{Q}_{N-1;n}^2 - \mathcal{Q}_{N-2;n} \mathcal{Q}_{N;2n}, \\ & q^{\frac{N}{2}} \mathcal{Q}_{N;2n+1} \mathcal{Q}_{N;2n-1} = \mathcal{Q}_{N;2n}^2 - q^{-n} \mathcal{Q}_{N-1;n}^2, \\ & q^{\frac{N}{2}} \mathcal{Q}_{N;2n+2} \mathcal{Q}_{N;2n} = \mathcal{Q}_{N;2n+1}^2 - q^{\frac{N}{2}-n-1} \mathcal{Q}_{N-1;n+1} \mathcal{Q}_{N-1;n}, \\ C_N^{(1)} : \quad & q^{N-1} \mathcal{Q}_{N-1;2n+1} \mathcal{Q}_{N-1;2n-1} = \mathcal{Q}_{N-1;2n}^2 - q^{-\frac{Nn}{2}} \mathcal{Q}_{N-2;2n} \mathcal{Q}_{N;n}^2, \\ & q^{N-1} \mathcal{Q}_{N-1;2n+2} \mathcal{Q}_{N-1;2n} = \mathcal{Q}_{N-1;2n+1}^2 - q^{-\frac{Nn}{2}} \mathcal{Q}_{N-2;2n+1} \mathcal{Q}_{N;n+1} \mathcal{Q}_{N;n}, \\ & q^{\frac{N}{2}} \mathcal{Q}_{N;n+1} \mathcal{Q}_{N;n-1} = \mathcal{Q}_{N;n}^2 - q^{\frac{(N-2)n}{2}} \mathcal{Q}_{N-1;2n}, \\ D_{N+1}^{(2)} : \quad & q^{N-1} \mathcal{Q}_{N-1;n+1} \mathcal{Q}_{N-1;n-1} = \mathcal{Q}_{N-1;n}^2 - q^{-\frac{Nn}{4}} \mathcal{Q}_{N-2;n} \mathcal{Q}_{N;n}^2, \\ & q^{\frac{N}{4}} \mathcal{Q}_{N;n+1} \mathcal{Q}_{N;n-1} = \mathcal{Q}_{N;n}^2 - q^{\frac{(N-2)n}{4}} \mathcal{Q}_{N-1;n}, \\ A_{2N-1}^{(2)} : \quad & q^N \mathcal{Q}_{N;n+1} \mathcal{Q}_{N;n-1} = \mathcal{Q}_{N;n}^2 - q^{-n} \mathcal{Q}_{N-1;n}^2, \\ A_{2N}^{(2)} : \quad & q^N \mathcal{Q}_{N;2n+1} \mathcal{Q}_{N;2n-1} = \mathcal{Q}_{N;2n}^2 - q^{-n} \mathcal{Q}_{N-1;2n} \mathcal{Q}_{N;2n}, \\ & q^N \mathcal{Q}_{N;2n+2} \mathcal{Q}_{N;2n} = \mathcal{Q}_{N;2n+1}^2 - q^{-n} \mathcal{Q}_{N-1;2n+1} \mathcal{Q}_{N;2n+1}. \end{aligned}$$

Remark 4.2. If $\mathfrak{g} \neq A_{2N}^{(2)}$, the evolution equations above are equivalent to the quantization of the Q -system cluster algebras. These correspond to exchange matrices [Ked08, DFK09, Wil15]: $B = \begin{pmatrix} C^t & -C & -C^t \\ C & & 0 \end{pmatrix}$ for untwisted \mathfrak{g} and $B = \begin{pmatrix} 0 & -C \\ C & 0 \end{pmatrix}$ for twisted \mathfrak{g} , where C is the Cartan matrix of R . These correspond to the initial cluster data $(Q_{a;i} : a \in [1, N], i = 0, 1)$. We choose a skew-symmetric q -commutation matrix to be the inverse of

the skew-symmetrized matrix B , of the form $\begin{pmatrix} 0 & \Lambda \\ -\Lambda^t & \Lambda^t - \Lambda \end{pmatrix}$ with $\Lambda = \Lambda^{(\mathfrak{g})}$ as in Definition 4.1. The quantized cluster variables obey the commutation relations (4.2), and the relevant quantum mutations [DFK09] are

$$(4.5) \quad q^{\Lambda_{a,a}^{(\mathfrak{g})}} \mathcal{Q}_{a;n+1} \mathcal{Q}_{a;n-1} = (\mathcal{Q}_{a;n})^2 - q^{\frac{1}{2} D_{a,a}^{-1}} : T_{a;n} : \quad (a = 1, 2, \dots, N),$$

with $D_{a,a} = 1$ except for $a = N$ in types $A_{2N-1}^{(2)}, D_{N+1}^{(2)}$, where it is equal to $\frac{1}{2}, 2$, respectively. The monomial $T_{a;k}$ is the product of $\mathcal{Q}_{b;k'}$ appearing as the second term in the right hand side of each Q-system relation, not including any factors of q , and the normal ordering $:\cdot:$ is defined as $:\prod X_i^{\mu_i}:$ $= q^{-\frac{1}{2} \sum_{i < j} a_{i,j} \mu_i \mu_j} X_1^{\mu_1} \dots X_k^{\mu_k}$ if $X_i X_j = q^{a_{i,j}} X_j X_i$. The quantum Q-system relations (4.3-4.4) are equivalent to the quantum mutations (4.5) after a renormalization of the cluster variables (see [DFK21] Lemma 4.4).

In this section, we prove the following Theorem, which is one of the main results of this paper:

Theorem 4.3. *For each \mathfrak{g} in Table 1, the limit $t \rightarrow \infty$ of the generalized \mathfrak{g} -Macdonald operators $D_{a;n}^{(\mathfrak{g})}(x, q)$ of Section 3.3.3 satisfy the corresponding quantum Q-system relations (4.2-4.4).*

Therefore the algebra generated by the elements $\mathcal{Q}_{a;k}$ subject to the quantum Q-system relations has a functional representation given by the difference operators of Section 3.3.3.

4.1.2. *Raising/lowering operator conditions.* We refer to the polynomials $\Pi_\lambda^{(\mathfrak{g})}(x)$ (3.32), which are the common eigenfunctions of $D_{a;0}^{(\mathfrak{g})}(x; q) = D_a^{(\mathfrak{g})}(x; q)$, as q -Whittaker polynomials. For non-twisted \mathfrak{g} , we conjectured in [DFK21] that $D_{a;1}^{(\mathfrak{g})}, D_{a;-1}^{(\mathfrak{g})}$ are raising and lowering operators acting on $\Pi_\lambda^{(\mathfrak{g})}(x)$. The following are the statement of this result for all \mathfrak{g} in Table 1, and will be proven in this Section:

Theorem 4.4. *For any \mathfrak{g} in Table 1,*

$$(4.6) \quad D_{a;0}^{(\mathfrak{g})}(x) \Pi_\lambda^{(\mathfrak{g})}(x) = \Lambda^{\omega_a^*} \Pi_\lambda^{(\mathfrak{g})}(x),$$

$$(4.7) \quad D_{a;1}^{(\mathfrak{g})}(x) \Pi_\lambda^{(\mathfrak{g})}(x) = \Lambda^{\omega_a^*} \Pi_{\lambda+\omega_a}^{(\mathfrak{g})}(x),$$

where ω_a and ω_a^* are fundamental weights of R or R^* , respectively.

Combining Theorem 4.4 and Theorem 4.3 with the relevant quantum Q-system relations, it follows that $D_{a;-1}^{(\mathfrak{g})}(x)$ is a lowering operator:

Corollary 4.5. *For all \mathfrak{g} , we have the following lowering operator conditions*

$$(4.8) \quad D_{a;-1}^{(\mathfrak{g})}(x) \Pi_\lambda^{(\mathfrak{g})}(x) = \Lambda^{\omega_a^*} (1 - \Lambda^{-\alpha_a^*}) \Pi_{\lambda-\omega_a}^{(\mathfrak{g})}(x),$$

where α_a^* are the simple roots of R^* .

Note that the prefactor guarantees that the result is 0 whenever $\lambda - \omega_a$ is not a dominant integral R -weight, i.e. not a \mathfrak{g} -partition.

The proof follows the steps of Section 2, using the Fourier transformed operators $\bar{D}_{a;n}^{(\mathfrak{g})}(\Lambda)$.

4.2. Opposite quantum Q-systems and integrability. For each \mathfrak{g} , we consider the solutions $\bar{Q}_{a;n}$ which satisfy the quantum Q-system with the opposite multiplication, referred to as the quantum \bar{Q} -system. We construct a representation $\bar{D}_{a;n}(\Lambda)$ of the solutions $\bar{Q}_{a;n}$ of the quantum \bar{Q} -system, subject to appropriate initial data, in terms of q -difference operators written in terms of the quantum torus $\mathbb{T}_\Lambda = \langle \Lambda_i, T_i \rangle_{i=1}^N$. Elements in this algebra act on functions of Λ as q -difference operators in Λ . The initial data $\bar{D}_{a;0}(\Lambda)$ are deduced from the eigenvalue equation (4.6). These are supplemented with a choice of $\bar{D}_{a;1}(\Lambda)$ ensuring that the quantum commutation relations opposite to those of Eq. (4.2) are satisfied:

$$(4.9) \quad \bar{D}_{a;0} \bar{D}_{b;1} = q^{-\Lambda_{a,b}^{(\mathfrak{g})}} \bar{D}_{b;1} \bar{D}_{a;0},$$

with $\Lambda^{(\mathfrak{g})}$ as in (4.1).

Definition 4.6. For all \mathfrak{g} , define

$$\bar{D}_{a;0} := \bar{D}_{a;0}^{(\mathfrak{g})}(\Lambda) = \Lambda^{\omega_a^*}, \quad \bar{D}_{a;1} := \bar{D}_{a;1}^{(\mathfrak{g})}(\Lambda) = \Lambda^{\omega_a^*} T^{\omega_a},$$

where ω_a, ω_a^* are fundamental weights of R, R^* .

These obey the relations (4.9), since $\Lambda^{\omega_a^*} T^{\omega_b} = q^{-\omega_a^* \cdot \omega_b} T^{\omega_b} \Lambda^{\omega_a^*}$. Since all Q-system evolutions are two-step recursion relations, the following are uniquely defined:

Definition 4.7. Define $\bar{D}_{a;n}^{(\mathfrak{g})}(\Lambda) = \bar{Q}_{a;n}$ for all $n \neq 0, 1$ to be the solutions of the \mathfrak{g} -type quantum \bar{Q} -system relations subject to the initial data in Definition 4.6.

Remark 4.8. Due to the Laurent property of quantum cluster algebras [BZ05], the solutions $\bar{D}_{a;n}$ are Laurent in the initial data $\{\bar{D}_{a;i} : i = 0, 1, a \in [1, N]\}$. Since these are monomials in $\{T_a, \Lambda_a, a \in [1, N]\}$, all quantum cluster variables are Laurent in the quantum torus generators, therefore they are q -difference operators. Although this argument doesn't apply to $\mathfrak{g} = A_{2N}^{(2)}$, we will show that all solutions of the \bar{Q} -system are Laurent in this case also.

4.2.1. Time translation operator g .

Theorem 4.9. Let $\bar{D}_{a;n}^{(\mathfrak{g})}$ be as in Definition 4.7. For each \mathfrak{g} there exists an element $g = g^{(\mathfrak{g})}$ in a completion of the quantum torus \mathbb{T}_Λ , such that for all $n \in \mathbb{Z}$:

$$(4.10) \quad \bar{D}_{a,n} = q^{-\frac{n}{2} \Lambda_{a,a}^{(\mathfrak{g})}} g^n \bar{D}_{a,0} g^{-n}, \quad a \text{ long},$$

$$(4.11) \quad \bar{D}_{a,2n+i} = q^{-n \Lambda_{a,a}^{(\mathfrak{g})}} g^n \bar{D}_{a,i} g^{-n}, \quad i = 0, 1, a \text{ short}.$$

These elements are

$$(4.12) \quad g^{(\mathfrak{g})} = \begin{cases} g_T g_\Lambda \left(\frac{1}{\Lambda_{N-1}\Lambda_N}; q \right)_\infty^{-1}, & \mathfrak{g} = D_N^{(1)}, \\ \left(g_T^{1/2} \left(\frac{1}{\Lambda_N^2}; q \right)_\infty^{-1} \right)^2 g_\Lambda, & \mathfrak{g} = B_N^{(1)}, \\ (g_T g_\Lambda)^2 \left(\frac{1}{\Lambda_N}; q \right)_\infty^{-1}, & \mathfrak{g} = C_N^{(1)}, \\ g_T g_\Lambda \left(\frac{1}{\Lambda_N^2}; q^2 \right)_\infty^{-1}, & \mathfrak{g} = A_{2N-1}^{(2)}, \\ g_T g_\Lambda \prod_{n=0}^{\infty} \left(\frac{1}{\Lambda_N}; q^{1/2} \right)_\infty^{-1}, & \mathfrak{g} = D_{N+1}^{(2)}, \\ g_T g_\Lambda \left(q^{1/2} \frac{1}{\Lambda_N}; q \right)_\infty^{-1} g_T g_\Lambda \left(\frac{1}{\Lambda_N}; q \right)_\infty^{-1}, & \mathfrak{g} = A_{2N}^{(2)}, \end{cases}$$

where we use the shorthand g_T, g_Λ of Equations (2.23) and (2.26).

Note the Λ -dependence of g is only via $\Lambda^{-\alpha_a^*}$ where α_a^* are the simple roots of R^* .

Proof. The proof is by induction on n . The inductive step relies on the fact that Equations (4.10) and (4.11), which can be written as

$$(4.13) \quad \bar{D}_{a;m+t_a} = q^{-\Lambda_{a,a}^{(\mathfrak{g})} t_a / 2} g \bar{D}_{a;m} g^{-1}, \quad t_a = \begin{cases} 1 & \text{if } a \text{ long} \\ 2 & \text{if } a \text{ short} \end{cases},$$

are compatible with the \bar{Q} -system evolution, from which $\bar{D}_{a;n}$ are defined. Define for all labels a : $\bar{\mathcal{T}}_{a;n} := \bar{Q}_{a;n}^2 - q^{\Lambda_{a,a}} \bar{Q}_{a;n-1} \bar{Q}_{a;n+1}$. If (4.13) hold for all $m \leq n$, then they hold for $m = n + 1$ iff $g \bar{\mathcal{T}}_{a;n} g^{-1} = q^{\Lambda_{a,a}^{(\mathfrak{g})} t_a} \bar{\mathcal{T}}_{a;n+t_a}$. This is easily checked case-by-case. To complete the proof, we must check (4.10-4.11) for some initial values of n . Using the form of $\bar{\mathcal{T}}_{a;n}$, we see that for long roots, it is enough to show that (4.10) holds for $n = -1$ and $n = 0$, so we need the explicit expressions for $\bar{D}_{a,n}$ with $n = -1, 0, 1$ to start the induction, and for short roots, we also need the expression for $n = 2$:

$$(4.14) \quad \bar{D}_{a,-1} = \bar{D}_{a,0} (1 - \Lambda^{-\alpha_a^*}) T^{-\omega_a}, \quad a \in [1, N],$$

$$(4.15) \quad \bar{D}_{a,2} = \bar{D}_{a,1} (1 - \Lambda^{-\alpha_a^*} T^{-\alpha_a}) T^{\omega_a}, \quad a \text{ short}.$$

Let $h_\alpha(u) = (\Lambda^{-\alpha}; q^u)_\infty^{-1}$, $h_\alpha = h_\alpha(1)$ and $h_\alpha^+ = (q^{\frac{1}{2}}\Lambda^{-\alpha}; q)_\infty^{-1}$. Then $g_\Lambda = \prod_{a=1}^{N-1} h_{\alpha_a^*}$. It is useful to rewrite the g operators of (4.12) as

$$g = g_T g_\Lambda h_{\alpha_N^*}(u), \quad \mathfrak{g} = D_N^{(1)}, A_{2N-1}^{(2)}, D_{N+1}^{(2)}, \quad u = 1, 2, \frac{1}{2} \text{ resp.},$$

$$g = g_1 g_2 \text{ where } g_1 = \begin{cases} g_T^{\frac{1}{2}} h_{\alpha_N^*}, \\ g_T g_\Lambda, \\ g_T g_\Lambda h_{\alpha_N^+}, \end{cases} \quad \text{and } g_2 = \begin{cases} g_1 g_\Lambda, & \mathfrak{g} = B_N^{(1)}, \\ g_1 h_{\alpha_N^*}, & \mathfrak{g} = C_N^{(1)}, \\ g_T g_\Lambda h_{\alpha_N^*}, & \mathfrak{g} = A_{2N}^{(2)}, \end{cases}$$

We use the exchange relations, valid for any $x, y \in \mathbb{Q}$,

$$(4.16) \quad (g_T)^x (\Lambda_a)^y = q^{\frac{xy^2}{2}} (\Lambda_a)^y (T_a)^{xy} (g_T)^x,$$

$$(4.17) \quad h_{\alpha_b^*} T^{\omega_a} = T^{\omega_a} (1 - \Lambda^{-\alpha_a^*}) h_{\alpha_b^*}, \quad h_{\alpha_b^*}^{-1} T^{-\omega_a} = (1 - \Lambda^{-\alpha_a^*}) T^{-\omega_a} h_{\alpha_b^*}^{-1}.$$

Long labels. In this case, we need only verify that (4.10) hold for $n = \pm 1$, Note that $\bar{D}_{a;0} = \Lambda^{\omega_a^*}$ commutes with g_Λ . In the self-dual cases $\mathfrak{g} = D_N^{(1)}, A_{2N-1}^{(2)}, D_{N+1}^{(2)}$, all labels are long and Equation (4.10) with $n = 1$, $g\bar{D}_{a;0}g^{-1} = q^{\Lambda_{a,a}/2} \bar{D}_{a;1}$, follows from

$$g_T \bar{D}_{a;0} = q^{\frac{\Lambda_{a,a}}{2}} \Lambda^{\omega_a^*} T^{\omega_a} g_T = q^{\frac{\Lambda_{a,a}}{2}} \bar{D}_{a;1} g_T.$$

For $n = -1$ it is equivalent to

$$\begin{aligned} g^{-1} \bar{D}_{a;0} &= h_{\alpha_N^*}^{-1} g_\Lambda^{-1} g_T^{-1} \Lambda^{\omega_a^*} = q^{-\frac{\Lambda_{a,a}}{2}} \Lambda^{\omega_a^*} g_\Lambda^{-1} T^{-\omega_a} h_{\alpha_N^*}^{-1} g_\Lambda^{-1} \\ &= q^{-\frac{\Lambda_{a,a}}{2}} \bar{D}_{a;0} (1 - \Lambda^{-\alpha_a^*}) T^{-\omega_a} g^{-1} = q^{-\frac{\Lambda_{a,a}}{2}} \bar{D}_{a;-1} g^{-1}. \end{aligned}$$

For the long labels $a < N$ of $B_N^{(1)}$, $\omega_a^* = \omega_a$, $\Lambda_{a,a} = a$ and (4.10) with $n = \pm 1$ follow from $g_T^{1/2} h_{\alpha_N^*} g_T^{1/2} \bar{D}_{a;0} = q^{\frac{a}{4}} g_T^{1/2} \Lambda^{\omega_a^*} T^{\omega_a/2} h_{\alpha_N^*} g_T^{1/2} = q^{\frac{a}{2}} \Lambda^{\omega_a^*} T^{\omega_a} g_T^{1/2} h_{\alpha_N^*} g_T^{1/2} = q^{\frac{a}{2}} \bar{D}_{a;1} g_T^{1/2} h_{\alpha_N^*} g_T^{1/2}$ and

$$\begin{aligned} g^{-1} \bar{D}_{a;0} &= g_\Lambda^{-1} (h_{\alpha_N^*}^{-1} g_T^{-1/2})^2 \Lambda^{\omega_a^*} = q^{-\frac{a}{2}} \Lambda^{\omega_a^*} g_\Lambda^{-1} T^{-\omega_a^*} h_{\alpha_N^*}^{-1} g_T^{-1/2} h_{\alpha_N^*}^{-1} g_T^{-1/2} \\ &= q^{-\frac{a}{2}} \bar{D}_{a;0} (1 - \Lambda^{-\alpha_a^*}) T^{-\omega_a^*} g^{-1} = q^{-\frac{\Lambda_{a,a}}{2}} \bar{D}_{a;-1} g^{-1}. \end{aligned}$$

For the long label $a = N$ of $\mathfrak{g} = C_N^{(1)}$, $2\omega_N^* = \omega_N$, (4.10) with $n = \pm 1$ follows from

$$g_T g_\Lambda g_T \bar{D}_{N;0} = q^{\frac{N}{2}} g_T \Lambda^{\omega_N^*} g_\Lambda T^{\omega_N^*} g_T = q^N \Lambda^{\omega_N^*} T^{2\omega_N^*} g_T g_\Lambda g_T = q^N \bar{D}_{N;1} g_T g_\Lambda g_T$$

and

$$\begin{aligned} g^{-1} \bar{D}_{N;0} &= h_{\alpha_N^*}^{-1} g_\Lambda^{-1} g_T^{-1} g_\Lambda^{-1} g_T^{-1} \Lambda^{\omega_N^*} = q^{-\frac{N}{4}} h_{\alpha_N^*}^{-1} g_\Lambda^{-1} g_T^{-1} \Lambda^{\omega_N^*} T^{-\omega_N^*} g_\Lambda^{-1} g_T^{-1} \\ &= q^{-\frac{N}{2}} \Lambda^{\omega_N^*} h_{\alpha_N^*}^{-1} T^{-2\omega_N^*} (g_\Lambda^{-1} g_T^{-1})^2 = q^{-\frac{N}{2}} \bar{D}_{N;0} (1 - \Lambda^{-\alpha_N^*}) T^{-2\omega_N^*} g^{-1} = q^{-\frac{N}{2}} \bar{D}_{N;-1} g^{-1}. \end{aligned}$$

Therefore (4.10) holds when $n = \pm 1$ and a long.

Short labels. Next we verify (4.11) for short labels a . We claim the following half-evolution equations hold in types $B_N^{(1)}, C_N^{(1)}, A_{2N}^{(2)}$:

$$(4.18) \quad q^{-\frac{1}{2}\Lambda_{a,a}} g_2 \bar{D}_{a;0} g_2^{-1} = \bar{D}_{a;1},$$

$$(4.19) \quad q^{-\frac{1}{2}\Lambda_{a,a}} g_1 \bar{D}_{a;1} g_1^{-1} = \bar{D}_{a;2}.$$

To see this, Equation (4.18) for $B_N^{(1)}$ with $a = N$, where $\omega_N^* = 2\omega_N$, follows from

$$g_2 \bar{D}_{N;0} g_2^{-1} = g_T^{-\frac{1}{2}} \Lambda^{\omega_N^*} g_T^{-\frac{1}{2}} = q^{\frac{N}{4}} \Lambda^{\omega_N^*} T^{\omega_N} = q^{\frac{\Lambda_{N,N}}{2}} \bar{D}_{N;1}.$$

For the short labels of $C_N^{(1)}, A_{2N}^{(2)}$, where $\omega_a = \omega_a^*$, it follows from

$$g_2 \bar{D}_{a;0} g_2^{-1} = g_T \Lambda^{\omega_a^*} g_T^{-1} = q^{\frac{a}{2}} \Lambda^{\omega_a^*} T^{\omega_a^*} = q^{\frac{\Lambda_{a,a}}{2}} \bar{D}_{a;1}.$$

Equation (4.19) $\mathfrak{g} = B_N^{(1)}$ with $a = N$, where $\alpha_N^* = 2\alpha_N$, follows from

$$\begin{aligned} g_1 \bar{D}_{N;1} &= g_T^{-\frac{1}{2}} \Lambda^{\omega_N^*} h_{\alpha_N^*} T^{\omega_N} = g_T^{-\frac{1}{2}} \Lambda^{\omega_N^*} T^{\omega_N} (1 - \Lambda^{-\alpha_N^*}) h_{\alpha_N^*} \\ &= q^{\frac{N}{4}} \Lambda^{\omega_N^*} T^{2\omega_N} (1 - q \Lambda^{-\alpha_N^*} T^{-\alpha_N^*/2}) g_T^{-\frac{1}{2}} h_{\alpha_N^*} = q^{\frac{\Lambda_{N,N}}{2}} \bar{D}_{N;2} g_1. \end{aligned}$$

When $a \leq N - 1$ for $C_N^{(1)}, A_{2N}^{(2)}$, for $a \leq N - 1$, $\alpha_a^* = \alpha_a$ and

$$\begin{aligned} g_1 \bar{D}_{a;1} &= g_T \Lambda^{\omega_a^*} g_T T^{\omega_a} = g_T \Lambda^{\omega_a^*} T^{\omega_a} (1 - \Lambda^{-\alpha_a^*}) g_T \\ &= q^{\frac{a}{2}} \bar{D}_{a;1} (1 - \Lambda^{-\alpha_a^*} T^{-\alpha_a^*}) T^{\omega_a} g_T g_T = q^{\frac{a}{2}} \bar{D}_{a;2} g_1. \end{aligned}$$

For $A_{2N}^{(2)}$, $a = N$,

$$\begin{aligned} g_1 \bar{D}_{N;1} &= g_T \Lambda^{\omega_N^*} g_T h_{\alpha_N^*}^+ T^{\omega_N} = g_T \Lambda^{\omega_N^*} (1 - q^{-\frac{1}{2}} \Lambda^{-\alpha_N^*}) T^{\omega_N} g_T h_{\alpha_N^*}^+ \\ &= q^{\frac{N}{2}} \bar{D}_{N;1} (1 - \Lambda^{-\alpha_N^*} T^{-\alpha_N^*}) g_T g_T h_{\alpha_N^*}^+ = q^{\frac{N}{2}} \bar{D}_{N;2} g_1. \end{aligned}$$

Equations (4.18),(4.19) imply (4.11) follows for $i = 1, n = 1$.

We also have the half-evolution equations for short labels:

$$(4.20) \quad q^{\frac{1}{2}\Lambda_{a,a}} g_1^{-1} \bar{D}_{a;1} g_1 = \bar{D}_{a;0},$$

$$(4.21) \quad q^{\frac{1}{2}\Lambda_{a,a}} g_2^{-1} \bar{D}_{a;0} g_2 = \bar{D}_{a;-1}.$$

Equation (4.20) is equivalent to $\bar{D}_{a;1} = q^{-\frac{1}{2}\Lambda_{a,a}} g_1 \bar{D}_{a;0} g_1^{-1}$, which is unchanged if we replace g_1 by g_2 , as only the T -dependent part g_T acts on $\bar{D}_{a;0}$, and the equation is therefore equivalent to (4.18). To show (4.21) for $B_N^{(1)}$,

$$\begin{aligned} g_2^{-1} \bar{D}_{N;0} &= g_\Lambda^{-1} h_{\alpha_N^*}^{-1} g_T^{-\frac{1}{2}} \Lambda^{\omega_N^*} = q^{\frac{N}{2}} \Lambda^{\omega_N^*} g_\Lambda^{-1} h_{\alpha_N^*}^{-1} T^{-\omega_N} g_T^{-\frac{1}{2}} \\ &= q^{\frac{N}{2}} \Lambda^{\omega_N^*} (1 - \Lambda^{-\alpha_N^*}) T^{-\omega_N} g_\Lambda^{-1} h_{\alpha_N^*}^{-1} g_T^{-\frac{1}{2}} = q^{\frac{N}{2}} \bar{D}_{N;-1} g_2^{-1}, \end{aligned}$$

and for a short in $C_N^{(1)}, A_{2N}^{(2)}$,

$$g_2^{-1} \bar{D}_{a;0} = q^{\frac{a}{2}} \Lambda^{\omega_a^*} h_{\alpha_N^*}^{-1} g_\Lambda^{-1} T^{-\omega_a} g_T^{-1} = q^{\frac{a}{2}} \Lambda^{\omega_a^*} (1 - \Lambda^{-\alpha_a^*}) T^{-\omega_a} h_{\alpha_N^*}^{-1} g_\Lambda^{-1} g_T^{-1} = q^{\frac{a}{2}} \bar{D}_{a;-1} g_2^{-1}.$$

These half-evolutions imply Equation (4.11) with $i = 1, n = -1$.

The Theorem follows by induction. \square

4.2.2. Integrability and conserved quantities. We claim that the time-translation operators g commute with the Pieri operators for each \mathfrak{g} , which therefore have an interpretation as the conserved quantities of the quantum \bar{Q} -system.

Theorem 4.10. *For all \mathfrak{g} , the operator $g^{(\mathfrak{g})}(\Lambda)$ commutes with the first Pieri operator $H_1^{(\mathfrak{g})}(\Lambda)$.*

The proof is by explicit calculation for each \mathfrak{g} . It is given in Appendix C.

In what follows, we need to generalize to type \mathfrak{g} the statement of Section 2.9 about the uniqueness of the solutions to the first Pieri equation. Starting with $D_1^{(\mathfrak{g})}(x) \Pi_\lambda^{(\mathfrak{g})}(x) = \Lambda_1 \Pi_\lambda^{(\mathfrak{g})}(x)$, writing $\Pi_\lambda^{(\mathfrak{g})}(x) = x^\lambda p_\lambda^{(\mathfrak{g})}(x)$, and conjugating (3.34) with $x^{-\lambda}$ results in the equation

$$\left(\frac{1}{\Lambda_1} + \sum_{\substack{i=1 \\ \epsilon=\pm 1}}^N \phi_{i,\epsilon}^{(\mathfrak{g})}(x) \left(\frac{\Lambda_i^\epsilon}{\Lambda_1} \Gamma_i^\epsilon - \frac{1}{\Lambda_1} \right) - 1 \right) p_\lambda^{(\mathfrak{g})}(x) = 0.$$

The difference operator is polynomial in $\{\Lambda^{-\alpha_i^*}\}$, α_i^* the simple roots of R^* , so we may analytically continue the solution $p_\lambda^{(\mathfrak{g})}(x)$ to $\hat{p}_\lambda^{(\mathfrak{g})}(x)$, $\lambda \in \mathbb{C}^N$, and then consider Λ to be a formal parameter. The coefficients $\phi_{i,\epsilon}^{(\mathfrak{g})}(x)$ have series expansions in the variables $x^{-\alpha_i}$, α_i the simple roots of R , hence we may expand $\hat{p}_\lambda^{(\mathfrak{g})}(x) = \sum_{\beta \in Q_+(R)} c_\beta^{(\mathfrak{g})}(\Lambda) x^{-\beta}$, as well as $c_\beta^{(\mathfrak{g})}(\Lambda) = \sum_{\delta \in Q_+(R^*)} c_{\beta,\delta}^{(\mathfrak{g})} \Lambda^{-\delta}$. When dealing with formal variables, we may exchange the summations and write a new series expansion,

$$\hat{p}_\lambda^{(\mathfrak{g})}(x) = \sum_{\delta \in Q_+(R^*)} \hat{c}_\delta^{(\mathfrak{g})}(x) \Lambda^{-\delta}, \quad \hat{c}_\delta^{(\mathfrak{g})}(x) = \sum_{\beta \in Q_+(R)} c_{\beta,\delta}^{(\mathfrak{g})} x^{-\beta}.$$

The first Pieri equation (3.40) is also easily extended to formal λ , as the dependence on $\Lambda^{-\alpha_a^*}$ is polynomial, while $T_i : \Lambda_j \mapsto q^{\delta_{ij}} \Lambda_j$.

Lemma 4.11. *Let $\Theta_\lambda(x)$ (λ arbitrary) be a solution of the first Pieri equation (3.40), such that $\Theta_\lambda(x) = x^\lambda \sum_{\beta \in Q_+(R^*)} \tau_\beta(x) \Lambda^{-\beta}$. Then when λ is evaluated as a \mathfrak{g} -partition,*

$$\Pi_\lambda^{(\mathfrak{g})} = \frac{\hat{c}_0^{(\mathfrak{g})}(x)}{\tau_0(x)} \Theta_\lambda(x).$$

Proof. The Pieri equation is a triangular linear system for the coefficients $\tau_\beta(x)$, determined uniquely for $\beta \neq 0$ up to the factor $\tau_0(x)$. The same holds for the coefficients $\hat{c}_\beta^{(\mathfrak{g})}(x)$ in the expansion of $\hat{p}_\lambda^{(\mathfrak{g})}(x)$, up to $\hat{c}_0^{(\mathfrak{g})}(x)$. We deduce that $x^\lambda \hat{p}_\lambda^{(\mathfrak{g})}(x) = \frac{\hat{c}_0^{(\mathfrak{g})}(x)}{\tau_0(x)} \Theta_\lambda(x)$. The Lemma follows by specialization to \mathfrak{g} -partitions. \square

Theorem 4.12. *The action of $g(\Lambda)$ on $\Pi_\lambda^{(\mathfrak{g})}(x)$ is equivalent to acting by a Gaussian in x :*

$$(4.22) \quad g^{(\mathfrak{g})}(\Lambda) \Pi_\lambda^{(\mathfrak{g})}(x) = \gamma^{(\mathfrak{g})}(x) \Pi_\lambda^{(\mathfrak{g})}(x), \quad \gamma^{(\mathfrak{g})}(x) = \gamma(x)^{t_1},$$

where $t_1 = 2$ if 1 is a short label (i.e. $\mathfrak{g} = C_N^{(1)}, A_{2N}^{(2)}$), and is equal to 1 otherwise.

In other words, $g^{(\mathfrak{g})}(\Lambda)$ is the Fourier transform of $\gamma(x)^{t_1}$.

Proof. Like $\Pi_\lambda^{(\mathfrak{g})}(x)$, the quantity $g^{(\mathfrak{g})}(\Lambda) \Pi_\lambda^{(\mathfrak{g})}(x)$ can be continued to arbitrary λ , and we write it as $g^{(\mathfrak{g})}(\Lambda) x^\lambda \hat{p}_\lambda^{(\mathfrak{g})}(x)$. Multiplying the first Pieri equation by $g^{(\mathfrak{g})}$ on the left, we get:

$$g^{(\mathfrak{g})} H_1^{(\mathfrak{g})}(\Lambda) x^\lambda \hat{p}_\lambda^{(\mathfrak{g})}(x) = H_1^{(\mathfrak{g})}(\Lambda) \left(g^{(\mathfrak{g})} x^\lambda \hat{p}_\lambda^{(\mathfrak{g})}(x) \right) = \hat{e}_1(x) \left(g^{(\mathfrak{g})} x^\lambda \hat{p}_\lambda^{(\mathfrak{g})}(x) \right),$$

hence both $g^{(\mathfrak{g})} x^\lambda \hat{p}_\lambda^{(\mathfrak{g})}(x)$ and $x^\lambda \hat{p}_\lambda^{(\mathfrak{g})}(x)$ obey the same first Pieri rule. Apply Lemma 4.11 to the function $\Theta_\lambda(x) = g^{(\mathfrak{g})} x^\lambda \hat{p}_\lambda^{(\mathfrak{g})}(x) = g^{(\mathfrak{g})} x^\lambda (\hat{c}_0^{(\mathfrak{g})}(x) + O(\Lambda^{-\alpha_i^*}))$. Expanding $g^{(\mathfrak{g})} = 1 + O(\Lambda^{-\alpha_i^*})$, we see that the leading order term in $g^{(\mathfrak{g})} x^\lambda \hat{p}_\lambda^{(\mathfrak{g})}(x)$ has only contributions from the action of the g_T parts of $g^{(\mathfrak{g})}$ on the leading term x^λ . The total contribution of the g_T terms is $g_T^{t_1}$, as directly read off (4.12). Finally noting that $(g_T)^{t_1} x^\lambda = \gamma(x)^{t_1} x^\lambda T^{t_1 \xi} (g_T)^{t_1}$, where $x = q^\xi$ (using (4.16)), we find that the leading term is $\tau_0(x) = \gamma(x)^{t_1} \hat{c}_0^{(\mathfrak{g})}(x)$, and the Theorem follows from the relation $\tau_0(x)/\hat{c}_0^{(\mathfrak{g})}(x) = \gamma(x)^{t_1}$. \square

Remark 4.13. *As noted in Remark 2.12, in the cases $\mathfrak{g} = D_{N+1}^{(2)}, A_{2N-1}^{(2)}, D_N^{(1)}$, the relation (4.22) boils down to the recursion relation of [FJMM09] (Theorem 3.1) for the coefficients $J_\beta^\mu = J_\beta(x = q^\mu)$ for the root systems $R = B_N, C_N, D_N$ respectively (see Table 1). These are the coefficients in the series expansion $\tilde{\Pi}_\lambda^{(\mathfrak{g})}(x) = x^\lambda g_\Lambda^{(\mathfrak{g})} \Pi_\lambda^{(\mathfrak{g})}(x) = x^\lambda \sum_{\beta \in Q_+} J_\beta(x) \Lambda^{-\beta}$, up to a rescaling $q \rightarrow q^2$ in the cases $\mathfrak{g} = D_N^{(1)}, D_{N+1}^{(2)}$. Here we use the notation*

$$g_\Lambda^{(\mathfrak{g})} = g_\Lambda \times \begin{cases} 1 / \left(\frac{1}{\Lambda_N}; q^{1/2} \right)_\infty & \text{for } \mathfrak{g} = D_{N+1}^{(2)}, R = B_N \\ 1 / \left(\frac{1}{\Lambda_N^2}; q^2 \right)_\infty & \text{for } \mathfrak{g} = A_{2N-1}^{(2)}, R = C_N \\ 1 / \left(\frac{1}{\Lambda_{N-1} \Lambda_N}; q \right)_\infty & \text{for } \mathfrak{g} = D_N^{(1)}, R = D_N \end{cases}$$

Corollary 4.14. *The Pieri operators $H_a^{(\mathfrak{g})}(\Lambda)$, $a = 1, 2, \dots, N$ of Eq. (3.37) are algebraically independent conserved quantities of the \mathfrak{g} -quantum \bar{Q} -systems.*

Proof. To see that the time translation operator $g^{(\mathfrak{g})}(\Lambda)$ commutes with all higher Pieri operators $H_a^{(\mathfrak{g})}(\Lambda)$, $a \in [1, N]$, act with $(g^{(\mathfrak{g})}(\Lambda))^{-1}\gamma(x)^{t_1}$ on the Pieri equations, and use Theorem 4.12

$$g^{(\mathfrak{g})}(\Lambda)^{-1}\gamma(x)^{t_1} H_a^{(\mathfrak{g})}(\Lambda) \Pi_\lambda^{(\mathfrak{g})}(x) = g^{(\mathfrak{g})}(\Lambda)^{-1} H_a^{(\mathfrak{g})}(\Lambda) g^{(\mathfrak{g})}(\Lambda) \Pi_\lambda^{(\mathfrak{g})}(x) = \hat{e}_a^{(\mathfrak{g})}(x) \Pi_\lambda^{(\mathfrak{g})}(x),$$

which implies that $g^{(\mathfrak{g})}(\Lambda)^{-1} H_a^{(\mathfrak{g})}(\Lambda) g^{(\mathfrak{g})}(\Lambda) = H_a^{(\mathfrak{g})}(\Lambda)$ by the definition of the Pieri operators. The statement follows by noting that any dependence between $H_a^{(\mathfrak{g})}(\Lambda)$ would imply a dependence between $\hat{e}_a^{(\mathfrak{g})}(x)$, which are independent. \square

4.3. Proof of Theorems 4.3 and 4.4. The proof of both Theorems relies on the following.

Theorem 4.15. *For all $\mathfrak{g} = D_N^{(1)}, B_N^{(1)}, C_N^{(1)}, D_{N+1}^{(2)}, A_{2N-1}^{(2)}, A_{2N}^{(2)}$ we have the relation*

$$(4.23) \quad D_{a;n}^{(\mathfrak{g})}(x) \Pi_\lambda^{(\mathfrak{g})}(x) = \bar{D}_{a;n}^{(\mathfrak{g})}(\Lambda) \Pi_\lambda^{(\mathfrak{g})}(x), \quad a \in [1, N], n \in \mathbb{Z}$$

valid for any \mathfrak{g} -partition λ .

Theorem 4.3 follows from Theorem 4.15, as any relation satisfied by the difference operators $\{\bar{D}_{a;n}\}$ implies the opposite relation for the difference operators $\{D_{a;n}\}$. Theorem 4.4 is the particular case of (4.23) with $n = 0, 1$. We provide a proof of (4.23) for long labels, short labels except for $B_N^{(1)}$, and for the short label of $B_N^{(1)}$ separately.

4.3.1. Long labels and even time short labels. In this case, the proof is similar to that in type $A_{N-1}^{(1)}$. Acting on both sides of the eigenvalue equation (4.6), which we write as (4.23) with $n = 0$, with $\gamma(x)^{-t_1 n} (g^{(\mathfrak{g})}(\Lambda))^n$ and using Theorem 4.12,

$$\begin{aligned} \gamma^{-t_1 n} D_{a;0}^{(\mathfrak{g})}(x) (g^{(\mathfrak{g})})^n \Pi_\lambda^{(\mathfrak{g})}(x) &= \gamma^{-t_1 n} D_{a;0}^{(\mathfrak{g})}(x) \gamma^{t_1 n} \Pi_\lambda^{(\mathfrak{g})}(x) = q^{\frac{\Lambda_{a,a} t_a n}{2}} D_{a;nt_a}^{(\mathfrak{g})}(x) \Pi_\lambda^{(\mathfrak{g})}(x) \\ &= (g^{(\mathfrak{g})})^n \bar{D}_{a;0}^{(\mathfrak{g})}(\Lambda) \gamma^{-t_1 n} \Pi_\lambda^{(\mathfrak{g})}(x) = (g^{(\mathfrak{g})})^n \bar{D}_{a;0}^{(\mathfrak{g})}(\Lambda) (g^{(\mathfrak{g})})^{-n} \Pi_\lambda^{(\mathfrak{g})}(x) = q^{\frac{\Lambda_{a,a} t_a n}{2}} \bar{D}_{a;nt_a}^{(\mathfrak{g})}(\Lambda) \Pi_\lambda^{(\mathfrak{g})}(x), \end{aligned}$$

where we have used Definition 3.15. Therefore, Eq. (4.23) holds for n a multiple of t_a .

4.3.2. Short labels, odd n for $\mathfrak{g} = C_N^{(1)}, A_{2N}^{(2)}$. In both cases, $a = 1$ is a short label, $t_1 = 2$.

Lemma 4.16. *For $\mathfrak{g} = C_N^{(1)}, A_{2N}^{(2)}$ and for any \mathfrak{g} -partition λ ,*

$$\left(D_{1,1}^{(\mathfrak{g})}(x) - \bar{D}_{1,1}^{(\mathfrak{g})}(\Lambda) \right) \Pi_\lambda^{(\mathfrak{g})}(x) = q^{-1} \left(D_{1,-1}^{(\mathfrak{g})}(x) - \bar{D}_{1,-1}^{(\mathfrak{g})}(\Lambda) \right) \Pi_\lambda^{(\mathfrak{g})}(x).$$

Proof. Applying Equation 3.46 with $n = 0$, acting on $\Pi_\lambda^{(\mathfrak{g})}(x)$,

$$(4.24) \quad \begin{aligned} [D_{1,0}^{(\mathfrak{g})}(x), \hat{e}_1(x)] \Pi_\lambda^{(\mathfrak{g})}(x) &= \left(D_{1,0}^{(\mathfrak{g})}(x) H_1^{(\mathfrak{g})}(\Lambda) - \hat{e}_1(x) \bar{D}_{1,0}^{(\mathfrak{g})}(\Lambda) \right) \Pi_\lambda^{(\mathfrak{g})}(x) \\ &= \left[H_1^{(\mathfrak{g})}(\Lambda), \bar{D}_{1,0}^{(\mathfrak{g})}(\Lambda) \right] \Pi_\lambda^{(\mathfrak{g})}(x) = (q-1) \left(D_{1,1}^{(\mathfrak{g})}(x) - q^{-1} D_{1,-1}^{(\mathfrak{g})}(x) \right) \Pi_\lambda^{(\mathfrak{g})}(x), \end{aligned}$$

where we used the Pieri rule (3.40), and the eigenvalue equation (4.23) with $n = 0, a = 1$. The only terms in $H_1^{(\mathfrak{g})}(\Lambda)$ of Equations (3.38-3.39) which fail to commute with $\bar{D}_{1,0}^{(\mathfrak{g})}(\Lambda) = \Lambda_1$ are $T_1 + \left(1 - \frac{\Lambda_2}{\Lambda_1}\right) T_1^{-1}$ if $N \geq 2$. Therefore,

$$\begin{aligned} [H_1^{(\mathfrak{g})}(\Lambda), \bar{D}_{1,0}^{(\mathfrak{g})}(\Lambda)] &= [T_1 + \left(1 - \frac{\Lambda_2}{\Lambda_1}\right) T_1^{-1}, \Lambda_1] = (q-1)\Lambda_1 T_1 + (q^{-1}-1)(\Lambda_1 - \Lambda_2)T_1^{-1} \\ &= (q-1)\bar{D}_{1,1}^{(\mathfrak{g})}(\Lambda) + (q^{-1}-1)\bar{D}_{1,-1}^{(\mathfrak{g})}(\Lambda), \end{aligned}$$

where we used $\bar{D}_{1,-1}^{(\mathfrak{g})}(\Lambda) = (\Lambda_1 - \Lambda_2)T_1^{-1}$. The lemma follows. \square

Defining $\Sigma_n^{(\mathfrak{g})} = (D_{1,n}^{(\mathfrak{g})}(x) - \bar{D}_{1,n}^{(\mathfrak{g})}(\Lambda)) \Pi_\lambda^{(\mathfrak{g})}(x)$, Lemma 4.16 says that $\Sigma_1^{(\mathfrak{g})} = q^{-1} \Sigma_{-1}^{(\mathfrak{g})}$.

Lemma 4.17. For $\mathfrak{g} = C_N^{(1)}, A_{2N}^{(2)}$, we have:

$$H_1^{(\mathfrak{g})}(\Lambda) \Sigma_i^{(\mathfrak{g})} = \hat{e}_1(x) \Sigma_i^{(\mathfrak{g})}, \quad (i = \pm 1).$$

Proof. As $\Sigma_{-1}^{(\mathfrak{g})} = q \Sigma_1^{(\mathfrak{g})}$, we may restrict ourselves to $i = 1$.

$$\begin{aligned} (4.25) \quad H_1^{(\mathfrak{g})}(\Lambda) \Sigma_1^{(\mathfrak{g})} &= \left(D_{1,1}^{(\mathfrak{g})}(x) H_1^{(\mathfrak{g})}(\Lambda) - H_1^{(\mathfrak{g})}(\Lambda) \bar{D}_{1,1}^{(\mathfrak{g})}(\Lambda) \right) \Pi_\lambda^{(\mathfrak{g})}(x) \\ &= \left\{ \hat{e}_1(x) D_{1,1}^{(\mathfrak{g})}(x) + (q-1)(D_{1,2}^{(\mathfrak{g})}(x) - q^{-1}D_{1,0}^{(\mathfrak{g})}(x)) \right. \\ &\quad \left. - \bar{D}_{1,1}^{(\mathfrak{g})}(\Lambda) H_1^{(\mathfrak{g})}(\Lambda) - (q-1)(\bar{D}_{1,2}^{(\mathfrak{g})}(\Lambda) - q^{-1}\bar{D}_{1,0}^{(\mathfrak{g})}(\Lambda)) \right\} \Pi_\lambda^{(\mathfrak{g})}(x) \\ &= \left\{ \hat{e}_1(x) D_{1,1}^{(\mathfrak{g})}(x) - \bar{D}_{1,1}^{(\mathfrak{g})}(\Lambda) H_1^{(\mathfrak{g})}(\Lambda) \right\} \Pi_\lambda^{(\mathfrak{g})}(x) = \hat{e}_1(x) \Sigma_1^{(\mathfrak{g})}, \end{aligned}$$

where we used Lemma 3.14, as well as the $a = 1, n = 0, 2$ cases of (4.23) proven above. \square

Lemma 4.16, together with a uniqueness argument (Lemma 4.11), implies that there exists a function $\alpha^{(\mathfrak{g})}(x)$ such that $\Sigma_1^{(\mathfrak{g})} = \alpha^{(\mathfrak{g})}(x) \Pi_\lambda^{(\mathfrak{g})}(x)$, $\Sigma_{-1}^{(\mathfrak{g})} = q \alpha^{(\mathfrak{g})}(x) \Pi_\lambda^{(\mathfrak{g})}(x)$:

$$\{D_{1,1}^{(\mathfrak{g})}(x) - \bar{D}_{1,1}^{(\mathfrak{g})}(\Lambda) - \alpha^{(\mathfrak{g})}(x)\} \Pi_\lambda^{(\mathfrak{g})}(x) = \{D_{1,-1}^{(\mathfrak{g})}(x) - \bar{D}_{1,-1}^{(\mathfrak{g})}(\Lambda) - q \alpha^{(\mathfrak{g})}(x)\} \Pi_\lambda^{(\mathfrak{g})}(x) = 0.$$

Using this equation with $\lambda = 0$, $D_{1,-1}^{(\mathfrak{g})}(x) \Pi_0^{(\mathfrak{g})}(x) = D_{1,-1}^{(\mathfrak{g})}(x) 1 = 0$, using (3.42). Similarly, $\bar{D}_{1,-1}^{(\mathfrak{g})}(\Lambda) \Pi_0^{(\mathfrak{g})}(x) = (\Lambda_1 - \Lambda_2)T_1^{-1} \Pi_\lambda^{(\mathfrak{g})}(x)|_{\lambda=0} = 0$, since the prefactor vanishes. Therefore, $\alpha^{(\mathfrak{g})}(x) = 0$ and $\Sigma_1^{(\mathfrak{g})} = \Sigma_{-1}^{(\mathfrak{g})} = 0$. Multiplying $\Sigma_{-1}^{(\mathfrak{g})}$ by $(g^{(\mathfrak{g})})^n \gamma^{-2n}$, we conclude that for $\mathfrak{g} = C_N^{(1)}, A_{2N}^{(2)}$, $a = 1, n \in \mathbb{Z}$ and any integer partition λ , Equation (4.23) holds.

This result can be extended to all short labels for these algebras, as follows.

Lemma 4.18. Equation (4.23) holds for all short labels a and n odd in the case of $\mathfrak{g} = C_N^{(1)}, A_{2N}^{(2)}$.

Proof. Let $N_{\mathfrak{g}}$ be as in Equation (3.19). Then if $a \leq N_{\mathfrak{g}} - 1$, the relations of the quantum $\bar{\mathcal{Q}}$ -system are identical to those of type $A_{N_{\mathfrak{g}}-1}^{(1)}$, and hence we can apply Theorem 2.8 of [DFK17b], which states that any solution of the relations for $a = 1, 2, \dots, N_{\mathfrak{g}} - 1$

$$q^a \mathcal{Q}_{a;n+1} \mathcal{Q}_{a;n-1} = (\mathcal{Q}_{a;n})^2 - \mathcal{Q}_{a+1;n} \mathcal{Q}_{a-1;n}, \quad \mathcal{Q}_{a;n} \mathcal{Q}_{b;n+1} = q^{\min(a,b)} \mathcal{Q}_{b;n+1} \mathcal{Q}_{a;n},$$

satisfies

$$(-1)^a (q-1) \mathcal{Q}_{a;n} = [\mathcal{Q}_{1;n-a+1}, \mathcal{Q}_{a-1;n+1}]_{q^a}, \quad (a = 1, 2, \dots, N_{\mathfrak{g}}).$$

Therefore the \mathfrak{g} -quantum $\bar{\mathcal{Q}}$ -system solution $\bar{D}_{a;n}^{(\mathfrak{g})}$ obeys the opposite relations

$$(-1)^a (q-1) \bar{D}_{a;n}^{(\mathfrak{g})} = [\bar{D}_{a-1;n+1}^{(\mathfrak{g})}, \bar{D}_{1;n-a+1}^{(\mathfrak{g})}]_{q^a}.$$

Using this and Definition 3.16, we deduce that

$$\begin{aligned} & \{D_{a;2n-1}^{(\mathfrak{g})}(x) - \bar{D}_{a;2n-1}^{(\mathfrak{g})}(\Lambda)\} \Pi_{\lambda}^{(\mathfrak{g})}(x) \\ &= \frac{(-1)^a}{q-1} \{[D_{1;2n-a}^{(\mathfrak{g})}(x), D_{a-1;2n}^{(\mathfrak{g})}(x)]_{q^a} - [\bar{D}_{a-1;2n}^{(\mathfrak{g})}(\Lambda), \bar{D}_{1;2n-a}^{(\mathfrak{g})}(\Lambda)]_{q^a}\} \Pi_{\lambda}^{(\mathfrak{g})}(x) = 0, \end{aligned}$$

by use of Eq. (4.23) for $D_{a-1;2n}^{(\mathfrak{g})}$ and $D_{1;2n-a}^{(\mathfrak{g})}$ with a even and for $D_{1;2n-a}^{(\mathfrak{g})}$ when a is odd. \square

4.3.3. *The case of $B_N^{(1)}$ for odd n .* It remains to prove (4.23) for $\mathfrak{g} = B_N^{(1)}$, $a = N$, n odd. The methods above are inapplicable, and instead we use expression (3.51) for $D_{N;2n-1}^{(B_N^{(1)})}$ in terms of Rains operators.

Recall that $B_N^{(1)}$ -partitions are both integer and half-integer partitions, due to the spin representation with highest weight ω_N and character $s_{\omega_N}(x) = \prod_{i=1}^N \frac{1+x_i}{\sqrt{x_i}}$. Denote

$$(4.26) \quad P_{\lambda} = P_{\lambda}^{(B_N^{(1)})} = P_{\lambda}^{(t, -1, q^{\frac{1}{2}}, -q^{\frac{1}{2}})}, \quad \tilde{P}_{\lambda} = P_{\lambda}^{(t, -q, q^{\frac{1}{2}}, -q^{\frac{1}{2}})}.$$

There is a factorization of Macdonald polynomials [vD95],

$$(4.27) \quad P_{\lambda+\omega_N}(x) = s_{\omega_N}(x) \tilde{P}_{\lambda}(x), \quad \lambda \text{ integer partition.}$$

The parameter specialization (4.26) for \tilde{P}_{λ} is obtained by conjugating $\mathcal{D}_1^{(B_N^{(1)})}(x, q, t)$ by $s_{\omega_N}^{-1}$ and identifying the resulting parameters (a, b, c, d) . We must therefore consider the action of the difference operators on both functions $\Pi_{\lambda} = \lim_{t \rightarrow \infty} P_{\lambda}$ and $\tilde{\Pi}_{\lambda} = \lim_{t \rightarrow \infty} \tilde{P}_{\lambda}$ with λ integer partitions.

Moreover, the Rains operators map the eigenfunctions of the $B_N^{(1)}$ -type difference operators to those of $B_N^{(1)'}$ -type, corresponding to different parameters. We denote the corresponding q -Whittaker limit of the eigenfunctions functions as Π'_{λ} . To these parameters there corresponds a different quantization of the Q-system, a time-translation operator $g'(\Lambda)$, and Hamiltonians.

Specializing to the $B_N^{(1)}$ parameters $(a, b, c, d) = (t, -1, q^{\frac{1}{2}}, -q^{\frac{1}{2}})$ and taking the $t \rightarrow \infty$ limit as in (3.48), (3.8) become

$$(4.28) \quad \begin{aligned} R_N^{(0)}(x) \Pi_\lambda &= (\Lambda_1 \Lambda_2 \cdots \Lambda_N)^{\frac{1}{2}} \left(1 + \frac{1}{\Lambda_N}\right) \Pi'_\lambda, \\ R_N^{(1)}(x) \Pi'_\lambda &= (\Lambda_1 \Lambda_2 \cdots \Lambda_N)^{\frac{1}{2}} \Pi_\lambda, \end{aligned}$$

for integer partitions λ . We identify the $B_N^{(1)'}$ specialization to be $(a, b, c, d) = (tq^{\frac{1}{2}}, -q^{\frac{1}{2}}, 1, -1)$,

$$\Pi'_\lambda = \Pi_\lambda^{(B_N^{(1)'})} = \lim_{t \rightarrow \infty} P_\lambda^{(tq^{\frac{1}{2}}, -q^{\frac{1}{2}}, 1, -1)}.$$

Lemma 4.19. *Equations (4.28) also hold for half-integer partitions.*

Proof. Using the specialization $(a, b, c, d) = (t, -q, q^{\frac{1}{2}}, -q^{\frac{1}{2}})$ as in (4.26), in the limit $t \rightarrow \infty$ (3.8) becomes

$$\tilde{R}_N^{(0)} \tilde{\Pi}_\lambda = q^{\frac{N}{4}} (\Lambda_1 \Lambda_2 \cdots \Lambda_N)^{\frac{1}{2}} \left(1 + \frac{q^{-\frac{1}{2}}}{\Lambda_N}\right) \tilde{\Pi}'_\lambda, \quad \tilde{R}_N^{(1)} \tilde{\Pi}'_\lambda = q^{\frac{N}{4}} (\Lambda_1 \Lambda_2 \cdots \Lambda_N)^{\frac{1}{2}} \tilde{\Pi}_\lambda,$$

where

$$\begin{aligned} \tilde{R}_N^{(0)} &= \lim_{t \rightarrow \infty} t^{-N(N-1)/2} q^{-\frac{N}{4}} \mathcal{R}_N^{(1, -q^{\frac{1}{2}})} = s_{\omega_N}^{-1} R_N^{(0)} s_{\omega_N}, \\ \tilde{R}_N^{(1)} &= \lim_{t \rightarrow \infty} t^{-N(N+1)/2} q^{-\frac{N}{4}} \mathcal{R}_N^{(t, -q^{\frac{1}{2}})} = s_{\omega_N}^{-1} R_N^{(1)} s_{\omega_N}, \end{aligned}$$

by use of the limit of (4.27). Here, $R_N^{(0)}$ and $R_N^{(1)}$ are as in (3.48). Restoring the factors s_{ω_N} ,

$$\begin{aligned} R_N^{(0)}(x) s_{\omega_N} \tilde{\Pi}_\lambda &= (\tilde{\Lambda}_1 \tilde{\Lambda}_2 \cdots \tilde{\Lambda}'_N)^{\frac{1}{2}} \left(1 + \frac{1}{\tilde{\Lambda}_N}\right) s_{\omega_N} \tilde{\Pi}'_\lambda, \\ R_N^{(1)}(x) s_{\omega_N} \tilde{\Pi}'_\lambda &= (\tilde{\Lambda}_1 \tilde{\Lambda}_2 \cdots \tilde{\Lambda}'_N)^{\frac{1}{2}} s_{\omega_N} \tilde{\Pi}_\lambda, \end{aligned}$$

where $\tilde{\Lambda} = q^{\omega_N} \Lambda = q^{\omega_N + \lambda}$. Combining this with the limit of (4.27) implies that (4.28) is satisfied for half-integer partitions $\tilde{\lambda} = \omega_N + \lambda$ as well as integer partitions. \square

Combining equations (4.28) leads to the eigenvalue equation for $\hat{D}^{(B_N^{(1)})}(x)$ of (3.48)

$$\hat{D}_N^{(B_N^{(1)})}(x) \Pi_\lambda(x) = R_N^{(1)}(x) R_N^{(0)}(x) \Pi_\lambda = \Lambda_1 \Lambda_2 \cdots \Lambda_N \left(1 + \frac{1}{\Lambda_N}\right) \Pi_\lambda,$$

which is consistent with the relation $\hat{D}_N^{(B_N^{(1)})} = R_N^{(1)} R_N^{(0)} = D_{N,0} + D_{N-1,0}$ and the eigenvalues $\bar{D}_{a,0}$ in Definition (4.6).

The q -Whittaker limit of the difference operators for the $B_N^{(1)'}$ parameters are obtained as in (A.38), and are denoted by $D'_a(x) = D_a^{(B_N^{(1)'})}(x)$, $a = 1, 2, \dots, N$. The corresponding limit of the Rains operator (3.9) is $\hat{D}'_N(x) = R_N^{(0)}(x) R_N^{(1)}(x) = D'_N(x) + D'_{N-1}(x)$. This identification follows from the $t \rightarrow \infty$ limit of Lemma A.10, and from the eigenvalues, which are identical to those of the $B_N^{(1)}$ case, since $\xi_{\mathfrak{g}} = \xi_{\mathfrak{g}'}$. As in the $B_N^{(1)}$ case, We define the generalized Macdonald operators

$$(4.29) \quad \begin{aligned} D'_{a;n}(x) &= q^{-na/2} \gamma^{-n} D'_a(x) \gamma^n, & (a = 1, 2, \dots, N-1), \\ D'_{N;2n}(x) &= q^{-nN/2} \gamma^{-n} D'_N(x) \gamma^n, \\ D'_{N;2n+1}(x) &= q^{-nN/4} R_{N;n+1}^{(0)}(x) R_{N;n}^{(1)}(x) = q^{-nN/2} \gamma^{-n} D'_{N;1}(x) \gamma^n. \end{aligned}$$

Note the reversal of the order in the last product and the different time index compared to (3.51). The operators $D'_{a;n}$ will be shown to satisfy the set of recursion relations below.

Definition 4.20. *The type $B_N^{(1)'}$ -quantum Q-system relations⁶ are the same as the type $B_N^{(1)}$ relations in (4.2)–(4.4) except for the two equations with $a = N$:*

$$\begin{aligned} q^{N/2} \mathcal{Q}_{N;2n+2} \mathcal{Q}_{N;2n} &= \mathcal{Q}_{N;2n+1}^2 - q^{\frac{N-1}{2}-n} \mathcal{Q}_{N-1;n+1} \mathcal{Q}_{N-1;n}, \\ q^{N/2} \mathcal{Q}_{N;2n+1} \mathcal{Q}_{N;2n-1} &= (\mathcal{Q}_{N;2n} - q^{-\frac{n}{2}} \mathcal{Q}_{N-1;n})(\mathcal{Q}_{N;2n} + q^{\frac{1-n}{2}} \mathcal{Q}_{N-1;n}). \end{aligned}$$

In particular, the q -commutation relations are as in type $B_N^{(1)}$ (4.2).

The eigenvalue equation for D'_a corresponding to any λ is $D'_{a;0}(x) \Pi'_\lambda = \Lambda^{\omega_a^*} \Pi'_\lambda$, with ω_a^* a fundamental weight of type C_N . Therefore, the candidate Fourier transforms $\bar{D}'_{a,n}(\Lambda)$ are defined so that they satisfy the opposite quantum Q-system to that of Definition 4.20, subject to the same initial data as in type $B_N^{(1)}$, $\bar{D}'_{a;0} = \Lambda^{\omega_a^*}$ and $\bar{D}'_{a;1} = \Lambda^{\omega_a^*} T^{\omega_a}$, $\bar{D}'_{N;1} = \Lambda^{\omega_N^*} T^{\omega_N}$. Together with the recursions of Def. 4.20, this determines $\bar{D}'_{a;n}$ for all a, n . In particular,

$$\begin{aligned} \bar{D}'_{a;-1} &= \bar{D}'_{a;0} \left(1 - \frac{\Lambda_{a+1}}{\Lambda_a} \right) T^{-\omega_a} & (a = 1, 2, \dots, N-1), \\ \bar{D}'_{N;-1} &= \bar{D}'_{N;0} \left(1 - \frac{1}{\Lambda_N} \right) \left(1 + \frac{q^{\frac{1}{2}}}{\Lambda_N} \right) T^{-\omega_N}, & \bar{D}'_{N;2} = \bar{D}'_{N;1} \left(1 - q^{\frac{1}{2}} \frac{1}{\Lambda_N^2} \frac{1}{T_N} \right) T^{\omega_N}. \end{aligned}$$

As before, these are sufficient to determine the form of the time-translation operator $g'(\Lambda)$.

⁶This “quantum Q-system” is new, and we call it a Q-system by analogy with the other \mathfrak{g} cases.

Lemma 4.21. *The operator*

$$(4.30) \quad \begin{aligned} g' &= g_T^{\frac{1}{2}} g'_{\Lambda_N} g_T^{\frac{1}{2}} g''_{\Lambda_N} g_{\Lambda}, \\ g'_{\Lambda_N} &= \frac{1}{(q^{\frac{1}{2}} \Lambda_N^{-2}; q)_{\infty}}, \quad g''_{\Lambda_N} = \frac{1}{(\Lambda_N^{-1}; q^{\frac{1}{2}})_{\infty} (-q^{\frac{1}{2}} \Lambda_N^{-1}; q^{\frac{1}{2}})_{\infty}}, \end{aligned}$$

where g_T is as in (2.23) and g_{Λ} as in (2.26), is the time translation operator for the $B_N^{(1)'}$ opposite quantum Q -system. That is, for all $n \in \mathbb{Z}$,

$$\begin{aligned} \bar{D}'_{a;n} &= q^{-an/2} (g')^n \bar{D}'_a (g')^{-n} \quad (a = 1, 2, \dots, N-1), \\ \bar{D}'_{N;2n+i} &= q^{-Nn/2} (g')^n \bar{D}'_{N;i} (g')^{-n}, \quad i = 0, 1. \end{aligned}$$

The Pieri rules for $B_N^{(1)'}$ are obtained by duality. Using Theorem B.1 with $(a, b, c, d) = (1, -1, tq^{\frac{1}{2}}, -q^{\frac{1}{2}})$, $(a^*, b^*, c^*, d^*) = (t^{\frac{1}{2}}, -t^{-\frac{1}{2}}, t^{\frac{1}{2}}q^{\frac{1}{2}}, -t^{-\frac{1}{2}}q^{\frac{1}{2}})$, the q -Whittaker limit of the first Pieri operator is

$$\begin{aligned} H_1^{(B_N^{(1)'})}(\Lambda) &= T_1 + \sum_{a=2}^N \left(1 - \frac{\Lambda_a}{\Lambda_{a-1}}\right) T_a + \sum_{a=1}^{N-1} \left(1 - \frac{\Lambda_{a+1}}{\Lambda_a}\right) T_a^{-1} \\ &+ \left(1 - \frac{1}{\Lambda_N}\right) \left(1 + \frac{q}{\Lambda_N}\right) \left(1 - \frac{q}{\Lambda_N^2}\right) T_N^{-1} + \frac{q^{\frac{1}{2}} - q^{-\frac{1}{2}}}{\Lambda_N} - \frac{q^{\frac{1}{2}} + q^{-\frac{1}{2}}}{\Lambda_N^2} + \frac{q^{-\frac{1}{2}}}{\Lambda_{N-1}\Lambda_N}. \end{aligned}$$

By direct calculation, this operator commutes with $g'(\Lambda)$ and as a consequence

$$(4.31) \quad g'(\Lambda) \Pi'_{\lambda}(x) = \gamma(x) \Pi'_{\lambda}(x).$$

We want to show that $D_{N;1}(x)\Pi_{\lambda} = \bar{D}_{N;1}(\Lambda)\Pi_{\lambda}$ (case $a = N, n = 1$ of (4.23)). Using 4.12, Definition (3.51) and the relations (4.28), we find

$$(4.32) \quad \begin{aligned} D_{N;1}(x) \Pi_{\lambda} &= q^{-\frac{N}{4}} R_{N;0}^{(1)} R_{N;1}^{(0)} \Pi_{\lambda} \\ &= q^{-\frac{3N}{8}} g(\Lambda_1 \cdots \Lambda_N)^{\frac{1}{2}} \left(1 + \frac{1}{\Lambda_N}\right) (g')^{-1} (\Lambda_1 \cdots \Lambda_N)^{\frac{1}{2}} \Pi_{\lambda}. \end{aligned}$$

Recall the notation $h_{\alpha_N^*} = \prod_{n=0}^{\infty} \left(1 - \frac{q^n}{\Lambda_N^2}\right)^{-1}$, then we have the relations $\left(1 + \frac{1}{\Lambda_N}\right) (g''_{\Lambda_N})^{-1} = h_{\alpha_N^*}^{-1}$ and $T_N^{\frac{1}{4}}(g'_{\Lambda_N})^{-1} = h_{\alpha_N^*}^{-1} T_N^{\frac{1}{4}}$. These allow to compute

$$\begin{aligned}
& g(\Lambda_1 \cdots \Lambda_N)^{\frac{1}{2}} \left(1 + \frac{1}{\Lambda_N}\right) (g')^{-1} (\Lambda_1 \cdots \Lambda_N)^{\frac{1}{2}} \\
&= g_T^{\frac{1}{2}} h_{\alpha_N^*} g_T^{\frac{1}{2}} h_{\alpha_N^*} g_{\Lambda} (\Lambda_1 \cdots \Lambda_N)^{\frac{1}{2}} \left(1 + \frac{1}{\Lambda_N}\right) g_{\Lambda}^{-1} (g''_{\Lambda_N})^{-1} g_T^{-\frac{1}{2}} (g'_{\Lambda_N})^{-1} g_T^{-\frac{1}{2}} (\Lambda_1 \cdots \Lambda_N)^{\frac{1}{2}} \\
&= g_T^{\frac{1}{2}} h_{\alpha_N^*} g_T^{\frac{1}{2}} (\Lambda_1 \cdots \Lambda_N)^{\frac{1}{2}} g_T^{-\frac{1}{2}} (g'_{\Lambda_N})^{-1} g_T^{-\frac{1}{2}} (\Lambda_1 \cdots \Lambda_N)^{\frac{1}{2}} \\
&= q^{\frac{N}{16}} g_T^{\frac{1}{2}} h_{\alpha_N^*} (\Lambda_1 \cdots \Lambda_N)^{\frac{1}{2}} (T_1 \cdots T_N)^{\frac{1}{4}} (g'_{\Lambda_N})^{-1} g_T^{-\frac{1}{2}} (\Lambda_1 \cdots \Lambda_N)^{\frac{1}{2}} \\
&= q^{\frac{N}{16}} g_T^{\frac{1}{2}} (\Lambda_1 \cdots \Lambda_N)^{\frac{1}{2}} (T_1 \cdots T_N)^{\frac{1}{4}} g_T^{-\frac{1}{2}} (\Lambda_1 \cdots \Lambda_N)^{\frac{1}{2}} \\
&= q^{\frac{N}{8}} (\Lambda_1 \cdots \Lambda_N)^{\frac{1}{2}} (T_1 \cdots T_N)^{\frac{1}{2}} (\Lambda_1 \cdots \Lambda_N)^{\frac{1}{2}} \\
(4.33) \quad &= q^{\frac{3N}{8}} \Lambda_1 \cdots \Lambda_N (T_1 \cdots T_N)^{\frac{1}{2}} = q^{\frac{3N}{8}} \bar{D}_{N;1}.
\end{aligned}$$

Combining this with (4.32) results in the relation $D_{N;1}(x)\Pi_{\lambda} = \bar{D}_{N;1}(\Lambda)\Pi_{\lambda}$. Multiplying by $\gamma^{-n}g^n$ and using (4.31):

$$(4.34) \quad D_{N;2n+1}^{(B_N^{(1)})}(x) \Pi_{\lambda}^{(B_N^{(1)})}(x) = \bar{D}_{N;2n+1}^{(B_N^{(1)})}(\Lambda) \Pi_{\lambda}^{(B_N^{(1)})}(x), \quad (n \in \mathbb{Z}).$$

This completes the proof of (4.23), and Theorems 4.3 and 4.4 follow.

Corollary 4.22. *The D' -operators satisfy the quantum $B_N^{(1)'}$ -quantum Q -system of Definition 4.20. Moreover we have the raising operator conditions for any $B_N^{(1)}$ -partition λ :*

$$D'_{a;1}(x) \Pi'_{\lambda} = \Lambda^{\omega_a^*} \Pi'_{\lambda + \omega_a}, \quad (a \in [1, N]),$$

ω_a, ω_a^* the fundamental weights of B_N, C_N .

Proof. Starting from the eigenvalue equations $D'_a(x) \Pi'_{\lambda} = \bar{D}'_a \Pi'_{\lambda}$ for $a = 1, 2, \dots, N$ and multiplying with $\gamma^{-n}(g')^n$, and still denoting \bar{X} the prime Fourier transform of X ,

$$D'_{a;t_a n}(x) \Pi'_{\lambda} = q^{-na/2} \gamma^{-n} D'_a(x) \gamma^n \Pi'_{\lambda} = q^{-na/2} (g')^n \bar{D}'_a (g')^{-n} \Pi'_{\lambda} = \bar{D}'_{a;t_a n} \Pi'_{\lambda}.$$

For the short label $a = N$, using (4.29), (4.28), (4.33) and $\bar{D}_{N;1} = \bar{D}'_{N;1} = \Lambda^{\omega_N^*} T^{\omega_N}$:

$$\begin{aligned}
D'_{N;1}(x) \Pi'_{\lambda} &= R_{N;1}^{(0)} R_N^{(1)} \Pi'_{\lambda} = q^{-\frac{N}{8}} (\Lambda_1 \cdots \Lambda_N)^{\frac{1}{2}} g (\Lambda_1 \cdots \Lambda_N)^{\frac{1}{2}} \left(1 + \frac{1}{\Lambda_N}\right) (g')^{-1} \Pi_{\lambda} \\
&= q^{-\frac{N}{8}} (\Lambda_1 \cdots \Lambda_N)^{\frac{1}{2}} q^{\frac{3N}{8}} \bar{D}_{N;1} (\Lambda_1 \cdots \Lambda_N)^{-\frac{1}{2}} \Pi_{\lambda} = \bar{D}'_{N;1} \Pi'_{\lambda}.
\end{aligned}$$

The relation is easily extended by multiplication with $\gamma^{-n}(g')^n$, resulting in $D'_{N;2n+1}(x) \Pi'_{\lambda} = \bar{D}'_{N;2n+1} \Pi'_{\lambda}$ for all $n \in \mathbb{Z}$. The Corollary follows. \square

5. UNIVERSAL SOLUTIONS

The relations between Macdonald eigenvalue equations and Pieri rules are embodied by the duality property of Macdonald polynomials (3.26). We now discuss a reformulation of this duality in terms of universal solutions along the same lines as the case of type A , treated in Section 2.13. This allows to re-prove the main results of this paper in terms of universal solutions.

5.1. Universal solutions.

5.1.1. *Universal Koornwinder-Macdonald and \mathfrak{g} -Macdonald eigenvalue solutions.* As in the case of type A , it is extremely fruitful to think directly of x and s as dual variables, whose roles may be interchanged. To this end, we use changes of variables:

$$x_i = q^{\mu_i} t^{\rho_i^*} = q^{\mu_i} a t^{N-i}, \quad s_i = q^{\lambda_i} t^{\rho_i} = q^{\lambda_i} \sigma t^{N-i}, \quad \lambda, \mu \in \mathbb{C}^N,$$

where we use the notations of Section 3.1.4, see Equation (3.13). Note that if we write $P_\lambda^{(a,b,c,d)}(x) = x^\lambda p_\lambda^{(a,b,c,d)}(x)$ and substitute this into the Koornwinder-Macdonald eigenvalue equation (3.5), we obtain

$$\left(\frac{(1-t^N)(1+\sigma t^{N-1})}{1-t} - \sigma t^{N-1} \hat{e}_1(s) + \sum_{\substack{i=1 \\ \epsilon=\pm 1}}^N \Phi_{i,\epsilon}^{(a,b,c,d)}(x) (\Lambda_i^\epsilon \Gamma_i^\epsilon - 1) \right) p_\lambda^{(a,b,c,d)}(x; s) = 0,$$

where we have conjugated (3.4) with $x^{-\lambda}$. Noting that $\Phi_{i,\epsilon}^{(a,b,c,d)}(x)$ can be expanded in series of the variables $x^{-\alpha_i}$, α_i the B_N simple roots, this suggests what we call a universal solution $P^{(a,b,c,d)}(s; x)$ of the Koornwinder-Macdonald eigenvalue equation (3.5) in the form

$$(5.1) \quad P^{(a,b,c,d)}(x; s) = q^{\lambda \cdot \mu} \sum_{\beta \in Q_+} c_\beta^{(a,b,c,d)}(s) x^{-\beta}, \quad c_0^{(a,b,c,d)}(s) = 1,$$

where Q_+ denotes the positive cone of the root lattice of B_N . The normalizing prefactor for $P^{(a,b,c,d)}(x; s)$ is such that $q^{\lambda \cdot \mu} = x^\lambda t^{-\rho^* \cdot \lambda} = s^\mu t^{-\rho \cdot \mu}$, and is invariant under the interchange of $\lambda \leftrightarrow \mu$, therefore under $x \leftrightarrow s$. Equation (3.5) for $P^{(a,b,c,d)}(x; s)$ and generic s ,

$$(5.2) \quad \mathcal{D}_1^{(a,b,c,d)} P^{(a,b,c,d)}(x; s) = \sigma t^{N-1} \hat{e}_1(s) P^{(a,b,c,d)}(x; s),$$

is equivalent to a linear triangular, generically nonsingular, system for the coefficients $c_\beta^{(a,b,c,d)}(s)$, which uniquely fixes them for all $\beta \in Q_+$. The solution $P^{(a,b,c,d)}(x; s)$ is therefore unique, and we refer to it as *the universal Koornwinder-Macdonald solution*. The normalization $c_0^{(a,b,c,d)}(s)$ uniquely fixes the solution, which is otherwise determined up to the overall normalization given by this coefficient.

Specializing the Koornwinder parameters (a, b, c, d) according to Table 1, we obtain the universal solutions $P^{(\mathfrak{g})}(x; s)$ of the first \mathfrak{g} -Macdonald eigenvalue equations (3.23):

$$(5.3) \quad \mathcal{D}_1^{(\mathfrak{g})} P^{(\mathfrak{g})}(x; s) = t^{\rho_1} \hat{e}_1(s) P^{(\mathfrak{g})}(x; s).$$

These universal solutions have the expansion

$$(5.4) \quad P^{(\mathfrak{g})}(x; s) = q^{\mu \cdot \lambda} \sum_{\beta \in Q_+(R)} c_\beta^{(\mathfrak{g})}(s) x^{-\beta}, \quad c_0^{(\mathfrak{g})}(s) = 1,$$

with the coefficients $c_\beta^{(\mathfrak{g})}(s)$ uniquely determined by (5.3), up to the normalization $c_0^{(\mathfrak{g})}(s)$.

Remark 5.1. *The apparent discrepancy between Q_+ in (5.1) and $Q_+(R)$ in (5.4) is just an artifact of the specializations. For instance using the $C_N^{(1)}$ specialization $(a, b, c, d) = t^{1/2}(1, -1, q^{1/2}, -q^{1/2})$ leads to an operator $\mathcal{D}_1^{(C_N)}(x; q, t)$ which has an expansion as a series of the variables $\{x_{i+1}/x_i\}_{1 \leq i \leq N-1}$ and x_N^{-2} , as opposed to the generic case $\mathcal{D}_1^{(a,b,c,d)}(x; q, t)$ which has an expansion as a series of the variables $\{x_{i+1}/x_i\}_{1 \leq i \leq N-1}$ and x_N^{-1} . This simply means that the series solution of the Macdonald eigenvalue equation at the specific $C_N^{(1)}$ specialization is an even function of x_N^{-1} , i.e. the coefficients of odd powers of x_N^{-1} vanish.*

Remark 5.2. *Specializing $s = q^{\lambda t^\rho}$ to λ an integer partition, and using the uniqueness of the universal solution, we recover the Koornwinder polynomial*

$$(5.5) \quad P^{(a,b,c,d)}(x; s = q^{\lambda t^\rho}) = t^{-\rho^* \cdot \lambda} P_\lambda^{(a,b,c,d)}(x).$$

The specialization therefore truncates the series (5.1) to finitely many terms. Specializing the Koornwinder parameters to the values in Table 1,

$$(5.6) \quad P^{(\mathfrak{g})}(x; s = q^{\lambda t^\rho}) = t^{-\rho^* \cdot \lambda} P_\lambda^{(\mathfrak{g})}(x),$$

for λ an integer partition, $\rho = \rho^{(\mathfrak{g})}$ and $\rho^ = \rho^{(\mathfrak{g}^*)}$. Moreover, by uniqueness, (5.6) holds also for λ any \mathfrak{g} -partition, for example the non-integer partitions in the cases $\mathfrak{g} = D_N^{(1)}, B_N^{(1)}, D_{N+1}^{(2)}$. This presentation is therefore more economical, as a single universal function $P^{(a,b,c,d)}(x; s)$ contains the information on all the \mathfrak{g} -Macdonald polynomials as well.*

Another important specialization of the universal Koornwinder function $P^{(a,b,c,d)}(x; s)$ is the Koornwinder Baker-Akhiezer quasi-polynomials introduced by Chalykh [Cha02]. The latter correspond to specializing the parameters a, b, c, d, t to arbitrary negative integer powers of q . The effect is again a truncation of the series to finitely many terms.

5.1.2. Universal Pieri solutions. We also define the universal solution $Q^{(a,b,c,d)}(s; x)$ of the first Pieri equation (3.17):

$$(5.7) \quad Q^{(a,b,c,d)}(s; x) = q^{\lambda \cdot \mu} \sum_{\beta \in Q_+} \bar{c}_\beta^{(a,b,c,d)}(x) s^{-\beta}, \quad \bar{c}_0^{(a,b,c,d)}(x) = 1$$

subject to:

$$\hat{e}_1(x) Q^{(a,b,c,d)}(s; x) = \hat{\mathcal{H}}_1^{(a,b,c,d)}(s; q, t) Q^{(a,b,c,d)}(s; x),$$

where

$$\begin{aligned} \hat{\mathcal{H}}_1^{(a,b,c,d)}(s; q, t) &:= t^{-\rho^* \cdot \lambda} \mathcal{H}_1^{(a,b,c,d)}(s; q, t) t^{\rho^* \cdot \lambda} \\ (5.8) \qquad \qquad \qquad &= \frac{1}{at^{N-1}} \Delta^{(a^*, b^*, c^*, d^*)}(s)^{-1} \mathcal{D}_1^{(a^*, b^*, c^*, d^*)}(s; q, t) \Delta^{(a^*, b^*, c^*, d^*)}(s). \end{aligned}$$

The series expansion (5.7) exists because $\hat{\mathcal{H}}_1^{(a,b,c,d)}(s; q, t)$ has a series expansion in the variables $s^{-\alpha_i}$. As above, the normalization $\bar{c}_0^{(a,b,c,d)}(x) = 1$ uniquely fixes the solution.

Specializing the parameters (a, b, c, d) as in Table 1, we have the universal solutions to the first \mathfrak{g} -Pieri equations (3.30):

$$(5.9) \qquad \qquad \hat{e}_1(x) Q^{(\mathfrak{g})}(s; x) = \hat{\mathcal{H}}_1^{(\mathfrak{g})}(s; q, t) Q^{(\mathfrak{g})}(s; x),$$

where we have used the fact that $\hat{e}_1^{(R)}(x) = \hat{e}_1(x)$ for all R , and $\hat{\mathcal{H}}_1^{(\mathfrak{g})}(s = \Lambda t^\rho; q, t) := t^{-\rho^* \cdot \lambda} \mathcal{H}_1^{(\mathfrak{g})}(\Lambda; q, t) t^{\rho^* \cdot \lambda}$, with $\mathcal{H}_1^{(\mathfrak{g})}(\Lambda; q, t)$ as in (3.31). The solution $Q^{(\mathfrak{g})}(s; x)$ is a series of the form

$$Q^{(\mathfrak{g})}(s; x) = q^{\mu \cdot \lambda} \sum_{\beta \in Q_+^*} \bar{c}_\beta^{(\mathfrak{g})}(x) s^{-\beta}, \quad \bar{c}_0^{(\mathfrak{g})}(x) = 1,$$

where the sum extends over the positive root cone Q_+^* of R^* . Due to triangularity, the coefficients of the series are uniquely determined by (5.9).

5.2. Duality.

5.2.1. *Duality of universal functions.* The duality of Macdonald polynomials can be extended to the universal functions as follows.

Theorem 5.3. *The functions $Q^{(a,b,c,d)}(s; x)$ and $P^{(a,b,c,d)}(x; s)$ are related via*

$$(5.10) \qquad \qquad Q^{(a,b,c,d)}(s; x) = \frac{P^{(a,b,c,d)}(x; s)}{\Delta^{(a,b,c,d)}(x)},$$

with $\Delta^{(a,b,c,d)}$ as in (3.14).

Proof. The proof is similar to that of Theorem 2.15. The universal solution $P^{(a,b,c,d)}(x; s)$ also obeys the Pieri equation (5.8), as a consequence of the existence of a solution to the bispectral problem [Che95b, vD96, Sah99, Cha02], i.e. of both Koornwinder eigenvalue and Pieri equations, and of the uniqueness of the solution up to the overall normalization determined by the leading coefficient. Expanding the coefficients $c_\beta^{(a,b,c,d)}(s) = \sum_{\delta \in Q_+} c_{\beta, \delta}^{(a,b,c,d)} s^{-\delta}$ allows to write an expansion

$$P^{(a,b,c,d)}(x; s) = q^{\lambda \cdot \mu} \sum_{\delta \in Q_+} \hat{c}_\delta^{(a,b,c,d)}(x) s^{-\delta}, \quad \hat{c}_\delta^{(a,b,c,d)}(x) = \sum_{\beta \in Q_+} c_{\beta, \delta}^{(a,b,c,d)} x^{-\beta}.$$

The Pieri equation uniquely fixes the coefficients in this expansion, up to the overall factor $\hat{c}_0^{(a,b,c,d)}(x)$, hence $P^{(a,b,c,d)}(x; s) = \hat{c}_0^{(a,b,c,d)}(x) Q^{(a,b,c,d)}(s; x)$ as a series in $s^{-\alpha_i}$.

To compute $\hat{c}_0^{(a,b,c,d)}(x)$, we note that it can be extracted as the successive limits $s_1 \rightarrow \infty$, $s_2 \rightarrow \infty$, ..., $s_N \rightarrow \infty$ of $P^{(a,b,c,d)}(x; s)$. Let us examine the m eigenvalue equation (3.6), for $P^{(a,b,c,d)}(x; s) = q^{\lambda \cdot \mu} (\hat{c}_0^{(a,b,c,d)}(x) + O(\Lambda^{-\alpha_i}))$. Writing

$$\mathcal{D}_m^{(a,b,c,d)} = d_m^{(a,b,c,d)}(x) + \sum_{\substack{I \subset [1,N] \\ |I| \leq m}} \sum_{\substack{\epsilon_k = \pm 1 \\ k \in I}} d_{I,\epsilon,m}(x) \prod_{k \in I} \Gamma_k^{\epsilon_k},$$

we have

$$\left(d_m^{(a,b,c,d)}(x) - \theta_m \hat{e}_m(x) + \sum_{\substack{I \subset [1,N] \\ |I| \leq m}} \sum_{\substack{\epsilon_k = \pm 1 \\ k \in I}} d_{I,\epsilon,m}(x) \prod_{k \in I} \Lambda_k^{\epsilon_k} \Gamma_k^{\epsilon_k} \right) (\hat{c}_0^{(a,b,c,d)}(x) + O(\Lambda^{-\alpha_i})) = 0,$$

where $\theta_m = \sigma^m t^{m(N - \frac{m+1}{2})}$. Dividing by $\Lambda_1 \Lambda_2 \cdots \Lambda_m$, we find that all the terms have a factor of the form $\Lambda^{-\beta}$ for some $\beta \in Q_+$, and only those with $\beta = 0$ survive in the limit $|\Lambda_1| \gg |\Lambda_2| \gg \cdots \gg |\Lambda_m| \gg 1$. This leaves us with

$$(5.11) \quad \left(\Gamma_1 \Gamma_2 \cdots \Gamma_m - \frac{\theta_m^2}{d_{[1,m],(+)^m,m}(x)} \right) \hat{c}_0^{(a,b,c,d)}(x) = 0.$$

Let us compute $d_{[1,m],(+)^m,m}(x)$. From Definition A.8, and the explicit expression of the van Diejen operators (A.8-A.9), we easily see that the coefficient of $\Gamma_1 \Gamma_2 \cdots \Gamma_m$ in $\mathcal{D}_m^{(a,b,c,d)}$ comes from the highest order van Diejen operator $\mathcal{V}_m^{(a,b,c,d)}$, and more precisely from the term $s = 1$, $J_1 = J = \{1, 2, \dots, m\}$ in (A.8):

$$\begin{aligned} d_{[1,m],(+)^m,m}(x) &= \prod_{i=1}^m \frac{(1 - ax_i)(1 - bx_i)(1 - cx_i)(1 - dx_i)}{(1 - x_i^2)(1 - qx_i^2)} \prod_{1 \leq i < j \leq m} \frac{1 - tx_i x_j}{1 - x_i x_j} \frac{1 - qtx_i x_j}{1 - qx_i x_j} \\ &\quad \times \prod_{1 \leq i < m < j \leq N} \frac{1 - tx_i x_j}{1 - x_i x_j} \frac{tx_i - x_j}{x_i - x_j}. \end{aligned}$$

Using

$$\begin{aligned} \frac{\theta_m^2}{d_{[1,m],(+)^m,m}(x)} &= \prod_{i=1}^m \frac{(1 - \frac{1}{x_i^2})(1 - \frac{1}{qx_i^2})}{(1 - \frac{1}{ax_i})(1 - \frac{1}{bx_i})(1 - \frac{1}{cx_i})(1 - \frac{1}{dx_i})} \prod_{1 \leq i < j \leq m} \frac{1 - \frac{1}{x_i x_j}}{1 - \frac{1}{tx_i x_j}} \frac{1 - \frac{1}{qx_i x_j}}{1 - \frac{1}{qtx_i x_j}} \\ &\quad \times \prod_{1 \leq i < m < j \leq N} \frac{1 - \frac{1}{x_i x_j}}{1 - \frac{1}{tx_i x_j}} \frac{1 - \frac{x_j}{x_i}}{1 - \frac{x_j}{tx_i}} \end{aligned}$$

we conclude that $\hat{c}_0^{(a,b,c,d)}(x)$ and $\Delta^{(a,b,c,d)}(x)$ of (3.14) both obey (5.11) for $m = 1, 2, \dots, N$, so their ratio must be a constant as it is invariant under the action of each Γ_i . This constant

is 1, by noting that $q^{-\lambda\mu}P^{(a,b,c,d)}(x; s) \rightarrow 1$ and $\Delta^{(a,b,c,d)} \rightarrow 1$ in the limit when all $x^{-\alpha_i} \rightarrow 0$. The Theorem follows. \square

Using the specializations of the parameters (a, b, c, d) as in Table 1,

Corollary 5.4. *The functions $Q^{(\mathfrak{g})}(s; x)$ and $P^{(\mathfrak{g})}(x; s)$ are related via*

$$(5.12) \quad Q^{(\mathfrak{g})}(s; x) = \frac{P^{(\mathfrak{g})}(x; s)}{\Delta^{(\mathfrak{g})}(x)},$$

with $\Delta^{(\mathfrak{g})}$ as in (3.27).

It is now a simple exercise to relate the universal (a, b, c, d) Koornwinder-Pieri solution to the universal solution of the (a^*, b^*, c^*, d^*) Koornwinder-Macdonald eigenvalue equation. As a result we have the following duality relation between universal Pieri solutions.

Theorem 5.5. *We have the following duality formulas:*

$$(5.13) \quad Q^{(a^*, b^*, c^*, d^*)}(x; s) = Q^{(a, b, c, d)}(s; x),$$

and their \mathfrak{g} specializations:

$$(5.14) \quad Q^{(\mathfrak{g}^*)}(x; s) = Q^{(\mathfrak{g})}(s; x).$$

Proof. For conciseness, we omit the superscripts (a, b, c, d) and use the superscript $*$ to stand for (a^*, b^*, c^*, d^*) . Starting from the equation $\mathcal{H}_1(s)Q(s; x) = \hat{e}_1(x)Q(s; x)$, using (5.8), we have

$$\mathcal{D}_1^*(s)\Delta^*(s)Q(s; x) = \sigma t^{N-1}\hat{e}_1(s)\Delta^*(s)Q(s; x).$$

Interchanging the variables $x \leftrightarrow s$, we find that $\Delta^*(x)Q(x; s)$ is a solution to the (a^*, b^*, c^*, d^*) eigenvalue equation. Moreover, using the normalization of $Q(x; s)$ with s and x interchanged, we have for small $\{x^{-\alpha_i}\}$: $\Delta^*(x)Q(x; s) = q^{\lambda\mu}(1 + O(\{x^{-\alpha_i}\}))$. We conclude that $\Delta^*(x)Q(x; s) = P^*(x; s)$ by uniqueness of the solution. The Theorem follows from $P^*(x; s) = \Delta^*(x)Q^*(s; x)$ by Theorem 5.3 applied to (a^*, b^*, c^*, d^*) . \square

This can alternatively be rephrased as duality between Koornwinder-Macdonald eigenvalue universal solutions:

$$(5.15) \quad \Delta^{(a^*, b^*, c^*, d^*)}(s)P^{(a, b, c, d)}(x; s) = \Delta^{(a, b, c, d)}(x)P^{(a^*, b^*, c^*, d^*)}(s; x),$$

and their specializations:

$$(5.16) \quad \Delta^{(\mathfrak{g}^*)}(s)P^{(\mathfrak{g})}(x; s) = \Delta^{(\mathfrak{g})}(x)P^{(\mathfrak{g}^*)}(s; x).$$

Some of the above relations appear in different guises in the literature: explicitly in the type A case [NS12] (see also Section 2.13), implicitly for the other types [Che95b, Che97] where universal functions are obtained as x, s -symmetric reproducing kernels. We now detail the explicit link between the universal function duality relation (5.15) and the Koornwinder polynomial duality (3.12).

Theorem 5.6. *The universal function duality relation (5.15) reduces to the Koornwinder polynomial duality relation (3.12) upon specializing the variables $x = q^\mu t^{\rho^*}$ and $s = q^\lambda t^\rho$ for λ, μ integer partitions.*

Proof. Starting from the universal function $P^{(a,b,c,d)}(x; s)$, we use the specialization $s = q^\lambda t^\rho$, leading to the Koornwinder polynomial $P^{(a,b,c,d)}(x; q^\lambda t^\rho) = t^{-\rho^* \cdot \lambda} P_\lambda^{(a,b,c,d)}(x)$. Similarly, using the specialization $x = q^\mu t^{\rho^*}$ on the function $P^{(a^*,b^*,c^*,d^*)}(s; x)$ leads to $P^{(a^*,b^*,c^*,d^*)}(s, q^\mu t^{\rho^*}) = t^{-\rho \cdot \mu} P_\mu^{(a^*,b^*,c^*,d^*)}(s)$. The double specialization $s = q^\lambda t^\rho, x = q^\mu t^{\rho^*}$ of (5.15) results in

$$(5.17) \quad t^{-\rho^* \cdot \lambda} \Delta^{(a^*,b^*,c^*,d^*)}(q^\lambda t^\rho) P_\lambda^{(a,b,c,d)}(q^\mu t^{\rho^*}) = t^{-\rho \cdot \mu} \Delta^{(a,b,c,d)}(q^\mu t^{\rho^*}) P_\mu^{(a^*,b^*,c^*,d^*)}(q^\lambda t^\rho).$$

The relation (3.12) follows from the following identity:

$$(5.18) \quad \frac{\Delta^{(a^*,b^*,c^*,d^*)}(q^\lambda t^\rho)}{\Delta^{(a,b,c,d)}(q^\mu t^{\rho^*})} = t^{\rho^* \cdot \lambda - \rho \cdot \mu} \frac{P_\mu^{(a^*,b^*,c^*,d^*)}(t^\rho)}{P_\lambda^{(a,b,c,d)}(t^{\rho^*})},$$

itself a consequence of (3.15), and of the identity $\Delta^{(a^*,b^*,c^*,d^*)}(t^\rho) = \Delta^{(a,b,c,d)}(t^{\rho^*})$. The Theorem follows. \square

Similarly, under the suitable specialization, the \mathfrak{g} -Macdonald duality relation (5.16) reduces to the Macdonald polynomial duality (3.26) for $x = q^\mu t^{\rho^*}$ and $s = q^\lambda t^\rho$, where λ is any \mathfrak{g} -partition, and μ any \mathfrak{g}^* -partition.

5.2.2. Duality of universal solutions in the q -Whittaker limit. Universal solutions of the \mathfrak{g} -Macdonald eigenvector and \mathfrak{g} -Pieri equations simplify drastically in the $t \rightarrow \infty$ limit. They read:

$$(5.19) \quad \Pi^{(\mathfrak{g})}(x; \Lambda) := \lim_{t \rightarrow \infty} t^{\rho^* \cdot \lambda} P^{(\mathfrak{g})}(x; \Lambda t^\rho) = x^\lambda \sum_{\beta \in Q_+(R)} c_\beta^{(\mathfrak{g})}(\Lambda) x^{-\beta},$$

and

$$(5.20) \quad \mathbf{K}^{(\mathfrak{g})}(\Lambda; x) := \lim_{t \rightarrow \infty} t^{\rho^* \cdot \lambda} Q^{(\mathfrak{g})}(\Lambda t^\rho; x) = x^\lambda \sum_{\beta \in Q_+(R^*)} \bar{c}_\beta^{(\mathfrak{g})}(x) \Lambda^{-\beta},$$

with $c_0^{(\mathfrak{g})}(\Lambda) = \bar{c}_0^{(\mathfrak{g})}(x) = 1$. Note that the limit $t \rightarrow \infty$ has broken the previous symmetry $x \leftrightarrow s$, as the s variable itself contained a t -dependent factor. However, Corollary 5.4 of the previous section turns into the following.

Theorem 5.7. *The universal solutions $\Pi^{(\mathfrak{g})}(x; \Lambda)$ and $\mathbf{K}^{(\mathfrak{g})}(\Lambda; x)$ are related via:*

$$(5.21) \quad \Pi^{(\mathfrak{g})}(x; \Lambda) = \bar{\Delta}^{(\mathfrak{g})}(x) \mathbf{K}^{(\mathfrak{g})}(\Lambda; x), \quad \bar{\Delta}^{(\mathfrak{g})}(x) := \lim_{t \rightarrow \infty} \Delta^{(\mathfrak{g})}(x) = \prod_{\alpha \in \widehat{\mathfrak{R}}_{++}(\mathfrak{g})} (1 - x^{-\alpha}).$$

The eigenvalue and Pieri equations are:

$$(5.22) \quad D_a^{(\mathfrak{g})}(x) \Pi^{(\mathfrak{g})}(x; \Lambda) = \Lambda^{\omega_a^*} \Pi^{(\mathfrak{g})}(x; \Lambda),$$

$$(5.23) \quad H_m^{(\mathfrak{g})}(\Lambda) \mathbf{K}^{(\mathfrak{g})}(\Lambda; x) = \hat{e}_m^{(R)}(x) \mathbf{K}^{(\mathfrak{g})}(\Lambda; x).$$

Both equations for $a, m = 1$ turn into simple triangular recursion relations for the coefficients $c_\beta^{(\mathfrak{g})}(\Lambda)$ and $\bar{c}_\beta^{(\mathfrak{g})}(x)$ respectively.

5.3. Fourier transform and proof of the Macdonald- Q-system conjecture. Like in the case of type A, we may reformulate the Fourier transform (1.3) in the q -Whittaker limit in terms of universal solutions via:

$$f(x) \Pi^{(\mathfrak{g})}(x; \Lambda) = \bar{f}(\Lambda) \Pi^{(\mathfrak{g})}(x; \Lambda) \Leftrightarrow f(x) \mathbf{K}^{(\mathfrak{g})}(\Lambda; x) = \bar{f}(\Lambda) \mathbf{K}^{(\mathfrak{g})}(\Lambda; x).$$

The main result of Section 4 is the relation (4.23) which expresses the fact that $\bar{D}_{a;n}^{(\mathfrak{g})}(\Lambda)$ is the Fourier transform of $D_{a;n}^{(\mathfrak{g})}(x)$. In terms of the universal Pieri solution, we expect:

$$D_{a;n}^{(\mathfrak{g})}(x) \mathbf{K}^{(\mathfrak{g})}(\Lambda; x) = \bar{D}_{a;n}^{(\mathfrak{g})}(\Lambda) \mathbf{K}^{(\mathfrak{g})}(\Lambda; x).$$

As a consequence any relation satisfied by the \bar{D} 's is satisfied by the D 's in the opposite direction, thus proving the Macdonald Q-system conjecture. The proof of these identities is identical to that in Section 4, and relies on the Fourier duality between $g^{(\mathfrak{g})}(\Lambda)$ and $\gamma^{(\mathfrak{g})}(x)$, whose adjoint action respectively generates the discrete time translation in the Λ and x pictures:

$$\gamma^{(\mathfrak{g})}(x) \mathbf{K}^{(\mathfrak{g})}(\Lambda; x) = g^{(\mathfrak{g})}(\Lambda) \mathbf{K}^{(\mathfrak{g})}(\Lambda; x).$$

The proofs rely on a uniqueness argument which can be rephrased as follows. The universal Pieri solution $\mathbf{K}^{(\mathfrak{g})}(\Lambda; x)$ (5.20) is fixed by the Pieri equation (5.23) for $m = 1$, up to an overall multiplicative function independent of Λ , and fixed by the leading term normalization $\bar{c}_0^{(\mathfrak{g})}(x) = 1$. Any other series solution of this Pieri equation is therefore proportional to $\mathbf{K}^{(\mathfrak{g})}(\Lambda; x)$, by a factor independent of Λ .

As we saw in Section 4.3.3, the case of odd times $a = N$ for $\mathfrak{g} = B_N^{(1)}$ required the use of Rains operators, and the mapping to a companion theory $B_N^{(1)'}$ with its own q -Whittaker polynomials and quantum Q-system. Let us rephrase the action of Rains operators at finite t (3.8) in the $B_N^{(1)}$ specialization $(a, b, c, d) = (t, -1, q^{\frac{1}{2}}, -q^{\frac{1}{2}})$ of Table 1 in terms of universal Macdonald functions. Let us denote for short $P(x; s) \equiv P^{(B_N^{(1)})}(x; s)$ and $P'(x; s) \equiv P^{(B_N^{(1)'})}(x; s)$ the respective universal Macdonald solutions. As both $B_N^{(1)}, B_N^{(1)'}$ share the same $\xi_{\mathfrak{g}} = \xi_{\mathfrak{g}'} = \frac{1}{2}$, i.e. $\sigma = \sigma' = t^{\frac{1}{2}}$ and $R(\mathfrak{g}) = R(\mathfrak{g}')$ while $t^{\rho_i^*} = t^{N-i+1}$ and $t^{(\rho')_i^*} = q^{\frac{1}{2}} t^{N-i+1}$, we may rewrite (3.8) as:

$$\mathcal{R}_N^{(1,-1)}(x) P(x; s) = F(s) P'(x; s), \quad \mathcal{R}_N^{(t,-1)}(x) P'(x; s) = F'(s) P(x; s),$$

with

$$F(s) = \prod_{i=1}^N (1 + t^{-\frac{1}{2}} s_i), \quad F'(s) = t^{N(N+1)/2} \prod_{i=1}^N (1 + t^{-\frac{1}{2}} s_i^{-1}),$$

where we have used the same notations as in the proof of Lemma A.10, with $u = ab/\sigma = -t^{\frac{1}{2}}$, and taken into account the product formula $q^{|\lambda|} = t^{-N^2/2} \prod_{i=1}^N s_i$. Taking the limit $t \rightarrow \infty$ results in:

$$R_N^{(0)} \Pi(x; \Lambda) = (\Lambda_1 \Lambda_2 \cdots \Lambda_N)^{\frac{1}{2}} \left(1 + \frac{1}{\Lambda_N} \right) \Pi'(x; \Lambda), \quad R_N^{(1)} \Pi'(x; \Lambda) = (\Lambda_1 \Lambda_2 \cdots \Lambda_N)^{\frac{1}{2}} \Pi(x; \Lambda).$$

The steps of the proof can then be repeated identically, in particular establishing that $\gamma(x) \Pi'(x; \Lambda) = g'(\Lambda) \Pi'(x; \Lambda)$, and then $D'_{a;n} \Pi'(x; \Lambda) = \bar{D}'_{a;n} \Pi'(x; \Lambda)$ for all $a \in [1, N]$ and $n \in \mathbb{Z}$, as well as finally $D_{a;n} \Pi(x; \Lambda) = \bar{D}_{a;n} \Pi(x; \Lambda)$, from which the main Theorems follow.

6. DISCUSSION

6.1. Companion quantum Q systems. Part of the proof of our main theorem, concerning the short label in type $B_N^{(1)}$ (Section 4.3.3), revealed that acting with the Rains operators (3.8) and their τ_+ -translates on the q -Whittaker functions gives rise to a companion system $B_N^{(1)'} 4.20$ to the $B_N^{(1)}$ quantum Q-system. This new system has the same classical ($q \rightarrow 1$ limit) as the $B_N^{(1)}$ quantum Q-system but corresponds to a different specialization of the Koornwinder parameters (a, b, c, d) , and has different Pieri operators and time translation operator (4.30).

One can obtain two further companion systems using the Rains operators, illustrated in Figure 6.1, from the theories at specialization parameters corresponding to $\mathfrak{g} = A_{2N-1}^{(2)}, A_{2N}^{(2)}$. These also have the same classical limit as the respective quantum Q-systems, and may be considered as alternative quantizations. In type $A_{2N}^{(2)}$, which is not a cluster algebra mutation in the first place, the companion system is, in some sense, simpler and more natural from the quantization point of view, and appears to be the one related to the graded tensor product character formulas of KR-modules [Lin].

Using the definition of the Rains operators (3.7), together with the properties (3.8) of Section 3.1.3, from with the specializations of Table 1, one obtains from the \mathfrak{g} -Macdonald polynomials $P_\lambda^{(\mathfrak{g})}(x)$ companion polynomials $P_\lambda^{(\mathfrak{g}')} (x)$, using three different specializations of the Rains operators, $\mathcal{R}_N^{(0)} = \mathcal{R}_N^{(1,-1)}$, $\mathcal{R}_N^{(1)} = \mathcal{R}_N^{(t,-1)}$ and $\mathcal{R}_N^{(2)} = \mathcal{R}_N^{(t^{\frac{1}{2}}, -t^{\frac{1}{2}})}$. In the cases $\mathfrak{g} = D_N^{(1)}, D_{N+1}^{(2)}, C_N^{(1)}$, the Rains operators leave the \mathfrak{g} -Macdonald polynomials invariant up to some scalar, so that $\mathfrak{g}' = \mathfrak{g}$. In those cases, the Rains operator is itself a Koornwinder-Macdonald operator, as shown in Lemma A.13. Only the specializations $\mathfrak{g} = B_N^{(1)}, A_{2N-1}^{(2)}$ and $A_{2N}^{(2)}$ are mapped to new companion theories $\mathfrak{g}' \neq \mathfrak{g}$.

$$\begin{array}{ccc}
 B_N^{(1)} & (t, -1, q^{\frac{1}{2}}, -q^{\frac{1}{2}}) & A_{2N}^{(2)} & (t, -1, t^{\frac{1}{2}}q^{\frac{1}{2}}, -t^{\frac{1}{2}}q^{\frac{1}{2}}) \\
 \mathfrak{R}_N^{(1)} \left(\begin{array}{c} \nearrow \\ \searrow \end{array} \right) \mathfrak{R}_N^{(0)} & & \mathfrak{R}_N^{(1)} \left(\begin{array}{c} \nearrow \\ \searrow \end{array} \right) \mathfrak{R}_N^{(2)} & \\
 B_N^{(1)'} & (1, -1, tq^{\frac{1}{2}}, -q^{\frac{1}{2}}) & A_{2N}^{(2)'} & (q^{\frac{1}{2}}t, -q^{\frac{1}{2}}, t^{\frac{1}{2}}, -t^{\frac{1}{2}}) \\
 \\
 A_{2N-1}^{(2)} & (t^{\frac{1}{2}}, -t^{\frac{1}{2}}, q^{\frac{1}{2}}, -q^{\frac{1}{2}}) & \mathfrak{R}_N^{(0)} \curvearrowright D_N^{(1)} & (1, -1, q^{\frac{1}{2}}, -q^{\frac{1}{2}}) \\
 \mathfrak{R}_N^{(2)} \left(\begin{array}{c} \nearrow \\ \searrow \end{array} \right) \mathfrak{R}_N^{(0)} & & \mathfrak{R}_N^{(1)} \curvearrowright D_{N+1}^{(2)} & (t, -1, tq^{\frac{1}{2}}, -q^{\frac{1}{2}}) \\
 A_{2N-1}^{(2)'} & (1, -1, t^{\frac{1}{2}}q^{\frac{1}{2}}, -t^{\frac{1}{2}}q^{\frac{1}{2}}) & \\
 \\
 & & \mathfrak{R}_N^{(2)} \curvearrowright C_N^{(1)} & (t^{\frac{1}{2}}, -t^{\frac{1}{2}}, t^{\frac{1}{2}}q^{\frac{1}{2}}, -t^{\frac{1}{2}}q^{\frac{1}{2}})
 \end{array}$$

FIGURE 1. The nine families of Koornwinder-Macdonald operators/polynomials. We have indicated the Rains operators that intertwine the various theories, and the specializations of the Koornwinder parameters (a, b, c, d) .

In the q -Whittaker limit, the Rains operators tend to $R_N^{(0)}, R_N^{(1)}$ of (3.49-3.50) and to

$$R_N^{(2)} = \lim_{t \rightarrow \infty} t^{-N(N+1)/2} \mathfrak{R}_N^{(t^{\frac{1}{2}}, -t^{\frac{1}{2}})} = \sum_{\epsilon_1, \dots, \epsilon_N = \pm 1} \prod_{i=1}^N \frac{x_i^{2\epsilon_i}}{x_i^{2\epsilon_i} - 1} \prod_{1 \leq i < j \leq N} \frac{x_i^{\epsilon_i} x_j^{\epsilon_j}}{x_i^{\epsilon_i} x_j^{\epsilon_j} - 1} \prod_{i=1}^N \Gamma_i^{\epsilon_i/2}.$$

These map the q -Whittaker polynomials $\Pi_\lambda = \Pi_\lambda^{(g)}$ to a multiple of the monic companions $\Pi'_\lambda = \Pi_\lambda^{(g')} = \lim_{t \rightarrow \infty} P_\lambda^{(g')}$. More precisely (c.f. (4.28) for $B_N^{(1)}$), (3.8) implies the following mapping of polynomials:

$$\begin{array}{l}
 A_{2N-1}^{(2)} : R_N^{(0)} \Pi_\lambda = (\Lambda_1 \Lambda_2 \cdots \Lambda_N)^{\frac{1}{2}} \left(1 + \frac{1}{\Lambda_N} \right) \Pi'_\lambda, \quad R_N^{(2)} \Pi'_\lambda = (\Lambda_1 \Lambda_2 \cdots \Lambda_N)^{\frac{1}{2}} \Pi_\lambda, \\
 A_{2N}^{(2)} : R_N^{(2)} \Pi_\lambda = (\Lambda_1 \Lambda_2 \cdots \Lambda_N)^{\frac{1}{2}} \Pi'_\lambda, \quad R_N^{(1)} \Pi'_\lambda = (\Lambda_1 \Lambda_2 \cdots \Lambda_N)^{\frac{1}{2}} \Pi_\lambda.
 \end{array}$$

For the new companion cases $A_{2N-1}^{(2)'}, A_{2N}^{(2)'}$, we define companion Koornwinder-Macdonald operators by specializing Definition A.8 for $m = 1, 2, \dots, N$ at the appropriate values of the parameters as in Figure 6.1. Their q -Whittaker limits are denoted by $D'_m = D_m^{(g')}$, and their τ_+ -translates are defined for all $a \in [1, N]$ and $n \in \mathbb{Z}$ as follows:

$$\begin{array}{l}
 A_{2N-1}^{(2)'} : D'_{a;2n+i} = q^{-na} \gamma^{-n} D'_{a;i} \gamma^n, \quad (i = -1, 0) \\
 A_{2N}^{(2)'} : D'_{a;n} = q^{-na/2} \gamma^{-n} D'_a \gamma^n,
 \end{array}$$

where $D'_{a;-1}$ is defined as in (3.42) with the suitable function $\phi_{i,\epsilon}^{(\mathfrak{g}')} (x)$. Note that all labels of $A_{2N-1}^{(2)'}$ are short and all labels of $A_{2N}^{(2)'}$ are long, which is the reverse of the case for the un-primed theories. Using the methods of this paper, it can be shown that these new operators obey new quantum Q-systems, with the same q -commutation relations and recursion relations as their non-primed companions, except for the label N , where

$$\begin{aligned} A_{2N-1}^{(2)'} : \quad & q^N Q_{N,2n+2} Q_{N,2n} = Q_{N,2n+1}^2 - q^{-2n} Q_{N-1,2n+1}^2, \\ & q^N Q_{N,2n+1} Q_{N,2n-1} = (Q_{N,2n} - q^{-n} Q_{N-1,2n})(Q_{N,2n} + q^{1-n} Q_{N-1,2n}), \\ A_{2N}^{(2)'} : \quad & q^N Q_{N,n+1} Q_{N,n-1} = Q_{N,n}^2 - q^{-\frac{n}{2}} Q_{N,n} Q_{N-1,n}. \end{aligned}$$

for all $n \in \mathbb{Z}$. To derive these results, we first define the candidate Fourier transforms $\bar{D}'_{a;n}$ via the opposite quantum Q-systems and same initial data as the non-primed companions (that is, the same values of ω_a, ω_a^*), and compute the time translation operators:

$$\begin{aligned} A_{2N-1}^{(2)'} : \quad & g' = g_T g_\Lambda \frac{1}{(\Lambda_N^{-2}; q^2)_\infty} g_T g_\Lambda \frac{1}{(\Lambda_N^{-1}; q)_\infty (-q\Lambda_N^{-1}; q)_\infty}, \\ A_{2N}^{(2)'} : \quad & g' = g_T g_\Lambda \frac{1}{(\Lambda_N^{-1}; q)_\infty}, \end{aligned}$$

which commute, respectively, with the two Pieri operators or Toda Hamiltonians

$$\begin{aligned} A_{2N-1}^{(2)'} : \quad & H'_1(\Lambda) = T_1 + \sum_{i=2}^N \left(1 - \frac{\Lambda_i}{\Lambda_{i-1}}\right) T_i + \sum_{i=1}^{N-1} \left(1 - \frac{\Lambda_{i+1}}{\Lambda_i}\right) T_i^{-1} + \left(1 - \frac{1}{\Lambda_N}\right) \left(1 + \frac{q}{\Lambda_N}\right) T_N^{-1}, \\ A_{2N}^{(2)'} : \quad & H'_1(\Lambda) = T_1 + \sum_{i=2}^N \left(1 - \frac{\Lambda_i}{\Lambda_{i-1}}\right) T_i + \sum_{i=1}^{N-1} \left(1 - \frac{\Lambda_{i+1}}{\Lambda_i}\right) T_i^{-1} + \left(1 - \frac{1}{\Lambda_N}\right) T_N^{-1} - \frac{q^{-\frac{1}{2}}}{\Lambda_N} \end{aligned}$$

obtained from the q -Whittaker limit of the specialized operators of Theorem (B.1). After proving that $g'\Pi_\lambda = \gamma^{t_1} \Pi'_\lambda$ ($t_1 = 2$ for $A_{2N-1}^{(2)'}$, $t_1 = 1$ for $A_{2N}^{(2)'}$), one concludes that $D_{a;n} \Pi'_\lambda = \bar{D}'_{a;n} \Pi'_\lambda$ and the quantum Q-system relations for $D_{a;n}$ follow.

Finally, matching the specialization of the formula (3.14) for $\Delta^{(\mathfrak{g}')}$ with the form of the product over affine roots (3.27), we make the following identification of affine roots corresponding to \mathfrak{g}' :

$$\begin{aligned} B_N^{(1)'} : \quad & \widehat{R}_{++} = \{(n - \frac{1}{2})\delta + e_i : 1 \leq i \leq N, n \geq 1\} \sqcup \{n\delta + (e_i \pm e_j) : 1 \leq i < j \leq N, n \geq 1\} \\ A_{2N-1}^{(2)'} : \quad & \widehat{R}_{++} = \{(2n-1)\delta + 2e_i : 1 \leq i \leq N, n \geq 1\} \sqcup \{n\delta + (e_i \pm e_j) : 1 \leq i < j \leq N, n \geq 1\} \\ A_{2N}^{(2)'} : \quad & \widehat{R}_{++} = \{(n - \frac{1}{2})\delta + e_i, 2n\delta + 2e_i : 1 \leq i \leq N, n \geq 1\} \sqcup \{n\delta + (e_i \pm e_j) : 1 \leq i < j \leq N, n \geq 1\}. \end{aligned}$$

6.2. Universal solutions and Path models. In Section 5, we introduced universal solutions $P(x; s)$ and $Q(s; x)$ for the various eigenvalue equations and Pieri rules, by solving an infinite triangular linear system of equations for the expansion coefficients in each case. Remarkably, the solutions can all be put in the form of path models, such as that for the q -Whittaker vectors in the $A_{N-1}^{(1)}$ case [DFKT17], see also [GT21]. As shown in Example 2.16, this results in an interpretation of the solution of the triangular system as a partition function for weighted paths on the relevant positive root cone Q_+ or Q_+^* .

Let us start with the case of $Q(s; x) = q^{\lambda \cdot \mu} \sum_{\beta \in Q_+^*} \bar{c}_\beta(x) s^{-\beta}$, $\bar{c}_0(x) = 1$, the series solution of the Pieri equation $\mathcal{H}_1(s) Q(s; x) = \hat{e}_1(x) Q(s; x)$ (either the Koornwinder or \mathfrak{g} -Macdonald Pieri rule). The important fact is that $\mathcal{H}_1(s)$ is a q -difference operator in the variable s of the form $\mathcal{H}_1(s) = \sum_{i=1}^N \sum_{\epsilon=0, \pm 1} h_{i, \epsilon}(s) T_i^\epsilon$, where the coefficients $h_{i, \epsilon}$ are rational functions of the variables $\{s^{-\alpha_i^*}\}$, with α_i^* the simple roots of R^* . Factoring out the common denominator in the coefficients, $h_{i, \epsilon}(s) = p_{i, \epsilon}(s)/q(s)$ with $p_{i, \epsilon}(s)$ and $q(s)$ some *polynomials* in $\{s^{-\alpha_i^*}\}$, the Pieri equation is equivalent to $(\hat{e}_1(x) q(s) - \sum_{i=1}^N \sum_{\epsilon=0, \pm 1} p_{i, \epsilon}(s) x_i^\epsilon T_i^\epsilon) q^{-\lambda \cdot \mu} Q(s; x) = 0$. The difference operator can be written as $\sum_{i=1}^N \sum_{\epsilon=0, \pm 1} \sum_{\alpha \in U^*} h_{i, \epsilon, \alpha}(x) s^{-\alpha} T_i^\epsilon$ for some x -dependent coefficients $h_{i, \epsilon, \alpha}(x)$, where α is summed over a *finite* subset U^* of Q_+^* . Noting that $T_i^\epsilon s^{-\beta} = q^{-\epsilon e_i \cdot \beta} s^{-\beta}$ and that $s^{-\alpha} s^{-\beta} = s^{-(\alpha+\beta)}$, and collecting the coefficient of $s^{-\beta}$ in the Pieri equation, we obtain $\sum_{i=1}^N \sum_{\epsilon=0, \pm 1} \sum_{\alpha \in U^*} h_{i, \epsilon, \alpha}(x) q^{-\epsilon e_i \cdot (\beta - \alpha)} \bar{c}_{\beta - \alpha}(x) = 0$, or equivalently

$$\left(\sum_{i, \epsilon} h_{i, \epsilon, 0}(x) q^{-\epsilon e_i \cdot \beta} \right) \bar{c}_\beta(x) = - \sum_{i, \epsilon} \sum_{\alpha \in U^* \setminus \{0\}} h_{i, \epsilon, \alpha}(x) q^{-\epsilon e_i \cdot (\beta - \alpha)} \bar{c}_{\beta - \alpha}(x).$$

The factor on the left hand side is non-vanishing for generic x and is therefore invertible. The path model interpretation goes as follows. Given the initial data $\bar{c}_0(x) = 1$, the coefficient $\bar{c}_\beta(x)$ is the partition function of paths from 0 to β in Q_+^* , consisting of steps in the finite subset $U^* \setminus \{0\}$. Each path has a weight equal to the product of vertex and edge weights along the path, defined respectively as

$$w_\varphi = \left(\sum_{i, \epsilon} h_{i, \epsilon, 0}(x) q^{-\epsilon e_i \cdot \varphi} \right)^{-1}, \quad w_0 = 1, \quad w_{\varphi - \alpha, \varphi} = - \sum_{i, \epsilon} h_{i, \epsilon, \alpha}(x) q^{-\epsilon e_i \cdot (\varphi - \alpha)}$$

for any $\varphi \in Q_+^*$, $\alpha \in U^* \setminus \{0\}$. The weight of a path p is a product over its vertex and edge sets $v(p), e(p)$:

$$w(p) = \prod_{\varphi \in v(p)} w_\varphi \times \prod_{(\varphi - \alpha, \varphi) \in e(p)} w_{\varphi - \alpha, \varphi},$$

and the coefficient $\bar{c}_\beta(x)$ is the partition function of all such paths

$$\bar{c}_\beta(x) = \sum_{\substack{\text{paths } p \text{ in } Q_+^* \\ \text{from } 0 \rightarrow \beta}} w(p).$$

A similar argument leads to a formulation of the Macdonald eigenvalue equation universal solution $P(x; s)$ in terms of a path model on Q_+ .

The path model for the Pieri solution simplifies drastically in the q -Whittaker limit as the Hamiltonians are directly polynomials of the $\Lambda^{-\alpha_i^*}$, giving rise to a small set of steps $U^* \setminus \{0\}$. For instance, by inspection of (3.38-3.39), we find that for all \mathfrak{g} , $w_\varphi^{-1} = \sum_{i=1}^N \sum_{\epsilon=\pm 1} x_i^\epsilon (1 - q^{-\epsilon\beta \cdot e_i})$.

Example 6.1. *In the case $\mathfrak{g} = A_4^{(2)}$, the q -Whittaker limit of the Pieri equation is*

$$\begin{aligned} 0 &= \left(T_1 + \left(1 - \frac{\Lambda_2}{\Lambda_1} \right) (T_2 + T_1^{-1}) + \left(1 - \frac{1}{\Lambda_2} \right) T_2^{-1} - \frac{1}{\Lambda_2} - (x_1 + x_1^{-1} + x_2 + x_2^{-1}) \right) \\ &\times x_1^{\lambda_1} x_2^{\lambda_2} \sum_{n,m \geq 0} \tilde{c}_{n,m}(x) \left(\frac{\Lambda_2}{\Lambda_1} \right)^n \left(\frac{1}{\Lambda_2} \right)^m \end{aligned}$$

expressed in terms of the variables $\Lambda^{-\alpha_1} = \frac{\Lambda_2}{\Lambda_1}$, $\Lambda^{-\alpha_2} = \frac{1}{\Lambda_2}$, α_i the simple roots of B_2 , and with the normalization $\tilde{c}_{0,0}(x) = 1$. This is equivalent to the recursion relation

$$\tilde{c}_{n,m}(x) = \frac{q^{n-1}(q^{-m}x_2 + x_1^{-1})\tilde{c}_{n-1,m}(x) + (q^{m-n-1}x_2^{-1} + 1)\tilde{c}_{n,m-1}(x)}{x_1(q^{-n} - 1) + x_1^{-1}(q^n - 1) + x_2(q^{n-m} - 1) + x_2^{-1}(q^{m-n} - 1)},$$

which has the form $\tilde{c}_{n,m}(x) = a_{n,m}\tilde{c}_{n-1,m}(x) + b_{n,m}\tilde{c}_{n,m-1}(x)$, the solution is

$$\tilde{c}_{n,m}(x) = \sum_{\text{paths } p: (0,0) \rightarrow (n,m)} \prod_{\text{steps } s \in p} w_s,$$

where the sum is over all the paths on \mathbb{Z}_+^2 with steps $(1,0)$ and $(0,1)$, from the origin to (n,m) , of the product of step weights w_s where $w_s = a_{i,j}$ for a horizontal step $(i-1, j) \rightarrow (i, j)$ and $w_s = b_{i,j}$ for a vertical step $(i, j-1) \rightarrow (i, j)$.

6.3. Universal solutions and q -Whittaker functions. We have used the terminology q -Whittaker functions for the $t \rightarrow \infty$ limits of Koornwinder-Macdonald polynomials. Strictly speaking, $\Pi_\lambda^{(\mathfrak{g})}(x)$ is a class-1 q -Whittaker function for the quantum universal enveloping algebra of a simple Lie algebra only in the cases where the Pieri operators are known to be q -deformed quantum Toda Hamiltonians. These are related to the quantum Q -system of types $\mathfrak{g} = A_{N-1}^{(1)}, D_N^{(1)}, D_{N+1}^{(2)}, A_{2N-1}^{(2)}$, whose conserved quantities are the relativistic Toda Hamiltonians, of types A_{N-1}, D_N, B_N, C_N , respectively (see Remark 3.10). By analogy, we call all the eigenfunctions $\Pi_\lambda^{(\mathfrak{g})}(x)$ q -Whittaker functions for all \mathfrak{g} in this paper. These share a number of properties.

In the q -Whittaker limit, the universal series solutions of the \mathfrak{g} -Macdonald eigenvalue equations and the Pieri equations (5.19-5.20) of Section 5, when specialized to λ a \mathfrak{g} -partition, reduce respectively to the analogues of class-1 q -Whittaker functions $\Pi_\lambda^{(\mathfrak{g})}(x) = \Pi^{(\mathfrak{g})}(x; q^\lambda)$, and the analogues of fundamental q -Whittaker functions $\Psi_\lambda^{(\mathfrak{g})}(x) = K^{(\mathfrak{g})}(x; q^\lambda)$. The former is a Weyl-symmetric polynomial in x , whereas the latter is a non-symmetric series solution of the relativistic Toda equation with prescribed leading term. It is associated with the highest weight Verma module V_μ , where μ is obtained from the variable $x = q^{\rho+\mu}$. In the theory of Whittaker functions, the fundamental Whittaker function is convergent only in a particular Weyl chamber. The class-1 function is a linear combination of fundamental Whittaker functions, regular in all the Weyl chambers. It is obtained via a symmetrization over the Weyl group action on the fundamental solutions with suitable coefficients.

A subtlety arises in the study of convergence of the above series. In the definition of $\bar{\Delta}^{(\mathfrak{g})}(x)$ as an infinite product, it is assumed that $|q| < 1$ for convergence. On the other hand, the series $K^{(\mathfrak{g})}(q^\lambda; x)$ or $\Psi_\lambda^{(\mathfrak{g})}(x)$ is well-behaved for $|q| > 1$. Obviously $\Pi_\lambda^{(\mathfrak{g})}(x)$, being a polynomial, makes sense for both $|q| > 1$ and $|q| < 1$. Note that the infinite product

$$\tilde{\Delta}^{(\mathfrak{g})} = \frac{1}{\prod_{n=0}^{\infty} \prod_{\alpha \in R_+} (1 - q^{-n} x^{-\alpha})}$$

is another solution of (2.39), convergent for $|q| > 1$ instead of $|q| < 1$. (To see this, note that conjugating $\Gamma_i^{\pm 1}$ with $\bar{\Delta}^{(\mathfrak{g})}(x)$ or $\tilde{\Delta}^{(\mathfrak{g})}(x)$ yields the same result.) The remarkable fact is that as a convergent series for $|q| > 1$, $\tilde{\Delta}^{(\mathfrak{g})}(x) \Psi_\lambda^{(\mathfrak{g})}(x)$ is not equal to the polynomial class-1 q -Whittaker function $\Pi_\lambda^{(\mathfrak{g})}(x)$. We conjecture⁷ that for general \mathfrak{g} it requires a symmetrization over the Weyl group, valid for $|q| > 1$:

$$(6.1) \quad \Pi_\lambda^{(\mathfrak{g})}(x) = \sum_{w \in W} \tilde{\Delta}^{(\mathfrak{g})}(wx) \Psi_\lambda^{(\mathfrak{g})}(wx).$$

We end up with two characterizations of $K(\Lambda; x)$: (1) as a series with $|q| < 1$ in the variables $\Lambda^{-\alpha_i}$, equal to $\Pi^{(\mathfrak{g})}(x; \Lambda) / \bar{\Delta}^{(\mathfrak{g})}(x)$ (by Theorem 5.7) and (2) as a series with $|q| > 1$ equal to $\Psi_\lambda^{(\mathfrak{g})}(x)$ when λ is a \mathfrak{g} -partition. The polynomials $\Pi_\lambda^{(\mathfrak{g})}(x)$ are expressed in terms of both, but with very different formulas for $|q| < 1$ and $|q| > 1$.

⁷Such a symmetrization formula exists relating classical fundamental and class-1 Whittaker functions.

Example 6.2. *Let us illustrate the above in the simplest case of $A_1^{(1)}$. Denoting by $u = x_2/x_1$, $v = \Lambda_2/\Lambda_1$,*

$$p(u; v) := x_1^{-\lambda_1} x_2^{-\lambda_2} \Pi(x; \Lambda) = \sum_{n=0}^{\infty} u^n \prod_{i=0}^{n-1} \frac{1 - vq^i}{1 - q^{-i-1}},$$

$$k(v; u) := x_1^{-\lambda_1} x_2^{-\lambda_2} K(\Lambda; x) = \sum_{n=0}^{\infty} v^n \frac{q^{n^2} u^n}{\prod_{i=1}^n (1 - q^i)(1 - uq^i)}.$$

With $\bar{\Delta}(u) = (qu; q)_{\infty}$ and $\tilde{\Delta}(u) = 1/(u; q^{-1})_{\infty}$,

$$p(u; v) = \bar{\Delta}(u) k(v; u) \quad (|q| < 1),$$

$$\Pi_{\lambda_1, \lambda_2}(x) = x_1^{\lambda_1} x_2^{\lambda_2} \tilde{\Delta}(u) k(q^{\lambda_2 - \lambda_1}; u) + x_2^{\lambda_1} x_1^{\lambda_2} \tilde{\Delta}(u^{-1}) k(q^{\lambda_2 - \lambda_1}; u^{-1}) \quad (|q| > 1),$$

$$= x_1^{\lambda_1} x_2^{\lambda_2} p(u; q^{\lambda_2 - \lambda_1}) \quad (|q| < 1),$$

where the second and third lines hold for any integers $\lambda_1 \geq \lambda_2 \geq 0$.

6.4. Summary/Perspectives. We have proved the Macdonald operator-quantum Q-system conjectures, which state that suitably defined, τ_+ -translated \mathfrak{g} -Macdonald difference operators, in the q -Whittaker limit, obey the \mathfrak{g} -type quantum Q-system relations, and may as such be considered as cluster variables in a suitable quantum cluster algebra in all cases except $A_{2N}^{(2)}$. The proofs cover the cases $\mathfrak{g} = A_{N-1}^{(1)}, D_N^{(1)}, B_N^{(1)}, C_N^{(1)}, A_{2N-1}^{(2)}, A_{2N}^{(2)}, D_{N+1}^{(2)}$, and rely strongly on the duality between the Koornwinder/Macdonald eigenvalue equations and the associated Pieri rules.

We have proved that the conserved quantities of the \mathfrak{g} -quantum Q-systems are the \mathfrak{g} -Pieri operators in the q -Whittaker limit, which in a number of cases can themselves be identified with known q -difference Toda Hamiltonians. Our construction provides explicit expressions for all (higher) Hamiltonians as well. It would be interesting to explore the combinatorial content of these, possibly in the language of cluster integrable systems [GK13].

We have constructed time-translation operators $g(\Lambda)$ for all \mathfrak{g} that commute with the corresponding q -difference Toda Hamiltonians. More generally, in the spirit of [SS18], it would be desirable to construct commuting Baxter Q -operators $Q(u; \Lambda)$ that coincide with $g(\Lambda)^{-1}$ at $u = 1$, but are in general quantum-dilogarithmic generating functions for the Hamiltonians. In all cases but $A_{2N}^{(2)}$, such a construction should exist in terms of mutations of the corresponding quantum cluster algebra.

As another by-product of our proofs, we have unearthed a remarkable structure involving three more companion theories $B_N^{(1)'}$, $A_{2N-1}^{(2)'}$ and $A_{2N}^{(2)'}$ and their associated quantum Q-systems. It would be interesting to further investigate their quantum Laurent property, as well as their combinatorial content, in particular the meaning of the associated fermionic sums which in the known cases provide q -multiplicities of decompositions of KR modules onto irreducibles (see [DFK14, DFK18]).

The quantum Q-system relates to the q -Whittaker limit of Macdonald-Koornwinder theory, but it is natural to consider the extension of some of our results to finite t . The time-shifted Macdonald operators $\mathcal{D}_{a;n}^{(\mathfrak{g})}(x; q, t)$ by iterated conjugation by $\gamma(x)$ were considered in [DFK19] in type $A_{N-1}^{(1)}$, and are generators of the spherical DAHA or, in the limit $N \rightarrow \infty$, the quantum toroidal algebra of \mathfrak{gl}_1 or the elliptic Hall algebra. In the Koornwinder we again have elements of the corresponding spherical DAHA but the limit $N \rightarrow \infty$ is still to be understood.

Moreover, in type $A_{N-1}^{(1)}$ a t -deformed analogue of the time translation operator $g(\Lambda)$ was defined in [LNS20] (see Proposition 4.2), and conjectured to act diagonally on the so-called Non-stationary Ruijsenaars function, introduced in [Sh19] as the (conjectural) universal series solution to the elliptic Ruijsenaars operator eigenvalue equation, also related to the geometry of affine Laumon spaces. We expect this operator to play the same role for the finite t macdonald case as $g(\Lambda)$ for the q -Whittaker limit $t \rightarrow \infty$, i.e. to be the (Macdonald) Fourier transform of the Gaussian for finite t . We believe there should exist a cluster algebra formulation of this operator. It would also be extremely interesting to investigate this operator for general Koornwinder theory.

APPENDIX A. DERIVATION OF THE \mathfrak{g} -MACDONALD OPERATORS

We combine several constructions [Mac01, vD95, Rai05] of commuting difference operators corresponding to the affine algebras in Table 1. The goal is to construct an appropriate set of N commuting operators for each \mathfrak{g} , with eigenvalues proportional to the symmetric functions in Table 2, which form a basis for the space spanned by the irreducible fundamental characters of the Lie algebras R . This choice of \mathfrak{g} -Macdonald difference operators is designed such that their q -Whittaker limits satisfy the type \mathfrak{g} quantum Q-systems.

A.1. Macdonald's operators. For each affine algebra \mathfrak{g} in Table 1, except for the case of $A_{2n}^{(2)}$, and for each minuscule co-weight of S , Macdonald defines a difference operator with eigenvalue which is a fundamental character of R^* [Mac01].

Let $\{e_i\}_{1 \leq i \leq N}$ be the standard basis of \mathbb{R}^N with the standard inner product (\cdot, \cdot) . For the set of variables $x = (x_1, \dots, x_N)$, we denote $x^v = x_1^{v_1} \cdots x_N^{v_N}$ for any $v = \sum_i v_i e_i$.

There is a surjective map $*$: $R \rightarrow S$, $\alpha \mapsto \alpha_* = \alpha/u_\alpha \in S$, for some real u_α . In the case where $R \neq S$, $u_\alpha = \frac{(\alpha, \alpha)}{2}$, so that $u_\alpha = 1$ in all cases but for the short roots of type B_N or the long roots of type C_N , in which case it is equal to $\frac{1}{2}$ or 2 , respectively.

Let $\pi = \sum \pi_i e_i$, and define⁸

$$(A.1) \quad \Phi_\pi := \prod_{\substack{\alpha \in R_+ \\ (\pi, \alpha_*)=1}} \frac{1 - tx^\alpha}{1 - x^\alpha}, \quad \Gamma_\pi = \prod_i \Gamma_i^{\pi_i}.$$

⁸We choose Macdonald's parameters $t_\alpha = t$ for all α , independently of the length α . This allows us to obtain the dual q -Whittaker limit by simply taking $t \rightarrow \infty$.

| Algebra | Positive roots R_+ | fundamental weights ω_a |
|---------|--|--|
| B_N | $e_i, i \in [1, N];$ $e_i \pm e_j, 1 \leq i < j \leq N$ | $\omega_i = \sum_{k=1}^i e_k, i \in [1, N-1];$ $\omega_N = \frac{1}{2} \sum_{k=1}^N e_k$ |
| C_N | $e_i \pm e_j, 1 \leq i < j \leq N; 2e_i, i \in [1, N]$ | $\omega_i = \sum_{k=1}^i e_k, i \in [1, N]$ |
| D_N | $e_i \pm e_j, 1 \leq i < j \leq N$ | $\omega_i = \sum_{k=1}^i e_k, i \in [1, N-2];$ $\omega_N = \frac{1}{2} \sum_{k=1}^N e_k, \omega_{N-1} = \omega_N - e_N$ |

TABLE 4. Positive roots and fundamental weights of the finite dimensional algebras of types BCD.

For each minuscule weight π of $S^\vee = \{2\frac{\alpha}{(\alpha, \alpha)}, \alpha \in S\}$, i.e. a weight such that $(\pi, \alpha_*) \in \{0, \pm 1\}$ for all $\alpha \in R$, there is a Macdonald difference operator \mathcal{E}_π which acts on functions $f(x)$ by the symmetrization

$$(A.2) \quad \mathcal{E}_\pi f = \sum_{w \in W} w(\Phi_\pi \Gamma_\pi f).$$

We list below the explicit formulas for each case treated in [Mac01]. The construction refers to the positive roots and fundamental weights for the simple Lie groups of types BCD in Table 4.

A.1.1. *Macdonald operators for $D_N^{(1)}$.* Here, $R = S$ is the root system of type D_N . There are three minuscule weights: $\omega_1 = e_1, \omega_{N-1}$ and ω_N . Equation (A.1) becomes

$$\begin{aligned} \Phi_{\omega_1} &= \prod_{j=2}^N \frac{1 - tx_1 x_j}{1 - x_1 x_j} \frac{tx_1 - x_j}{x_1 - x_j}, & \Phi_{\omega_{N-1}} &= \prod_{1 \leq i < j \leq N-1} \frac{1 - tx_i x_j}{1 - x_i x_j} \prod_{i=1}^{N-1} \frac{tx_i - x_N}{x_i - x_N}, \\ \Phi_{\omega_N} &= \prod_{1 \leq i < j \leq N} \frac{1 - tx_i x_j}{1 - x_i x_j}. \end{aligned}$$

The Weyl group of D_N acts on the set (x_1, x_2, \dots, x_N) by permutations of the indices and inversions of an even number of variables. The three Macdonald operators are

$$(A.3) \quad \mathcal{E}_{\omega_1}^{(D_N^{(1)})} = \sum_{\epsilon=\pm 1} \sum_{i=1}^N \prod_{j \neq i} \frac{1 - tx_i^\epsilon x_j}{1 - x_i^\epsilon x_j} \frac{tx_i^\epsilon - x_j}{x_i^\epsilon - x_j} \Gamma_i^\epsilon,$$

$$(A.4) \quad \mathcal{E}_{\omega_{N-1}}^{(D_N^{(1)})} = \sum_{\substack{\epsilon_1, \dots, \epsilon_N = \pm 1 \\ \epsilon_1 \epsilon_2 \dots \epsilon_N = -1}} \prod_{1 \leq i < j \leq N} \frac{1 - tx_i^{\epsilon_i} x_j^{\epsilon_j}}{1 - x_i^{\epsilon_i} x_j^{\epsilon_j}} \prod_{i=1}^N \Gamma_i^{\frac{\epsilon_i}{2}},$$

$$(A.5) \quad \mathcal{E}_{\omega_N}^{(D_N^{(1)})} = \sum_{\substack{\epsilon_1, \dots, \epsilon_N = \pm 1 \\ \epsilon_1 \epsilon_2 \dots \epsilon_N = 1}} \prod_{1 \leq i < j \leq N} \frac{1 - tx_i^{\epsilon_i} x_j^{\epsilon_j}}{1 - x_i^{\epsilon_i} x_j^{\epsilon_j}} \prod_{i=1}^N \Gamma_i^{\frac{\epsilon_i}{2}}.$$

These will be identified below as $\mathcal{D}_a^{(D_N^{(1)})}(x; q, t) = \mathcal{E}_{\omega_a}^{(D_N^{(1)})}$ with $a = 1, N-1, N$.

A.1.2. *Macdonald operator for $B_N^{(1)}$.* Here, $R = S$ is the root system of B_N . There is a unique minuscule weight of $S^\vee = C_N$, $\pi = e_1 = \omega_1$, with

$$\Phi_{\omega_1} = \frac{1 - tx_1}{1 - x_1} \prod_{j=2}^N \frac{1 - tx_1 x_j}{1 - x_1 x_j} \frac{tx_1 - x_j}{x_1 - x_j}.$$

The Weyl group $W \simeq S_N \times \mathbb{Z}_2$ is generated by all permutations and inversions of the variables in the set $x = (x_1, x_2, \dots, x_N)$, so the corresponding difference operator is

$$\mathcal{E}_{\omega_1}^{(B_N^{(1)})} = \sum_{\epsilon=\pm 1} \sum_{i=1}^N \frac{1 - tx_i^\epsilon}{1 - x_i^\epsilon} \prod_{j \neq i} \frac{1 - tx_i^\epsilon x_j}{1 - x_i^\epsilon x_j} \frac{tx_i^\epsilon - x_j}{x_i^\epsilon - x_j} \Gamma_i^\epsilon.$$

This will be identified as $\mathcal{D}_1^{(B_N^{(1)})}(x; q, t) = \mathcal{E}_{\omega_1}^{(B_N^{(1)})}$.

A.1.3. *Macdonald operator for $\mathfrak{g} = C_N^{(1)}$.* Here, $R = S$ is the root system of type C_N . There is a unique minuscule weight of $S^\vee = B_N$, $\pi = \frac{1}{2} \sum_{i=1}^N e_i = \omega_N$, with

$$\Phi_{\omega_N} = \prod_{i=1}^N \frac{1 - tx_i^2}{1 - x_i^2} \prod_{1 \leq i < j \leq N} \frac{1 - tx_i x_j}{1 - x_i x_j}.$$

The Weyl group is the same as for type B_N , resulting in the difference operator

$$(A.6) \quad \mathcal{E}_{\omega_N}^{(C_N^{(1)})} = \sum_{\epsilon_1, \dots, \epsilon_N = \pm 1} \prod_{i=1}^N \frac{1 - tx_i^{2\epsilon_i}}{1 - x_i^{2\epsilon_i}} \prod_{1 \leq i < j \leq N} \frac{1 - tx_i^{\epsilon_i} x_j^{\epsilon_j}}{1 - x_i^{\epsilon_i} x_j^{\epsilon_j}} \prod_{i=1}^N \Gamma_i^{\frac{\epsilon_i}{2}}.$$

This will be identified as $\mathcal{D}_N^{(C_N^{(1)})}(x; q, t) = \mathcal{E}_{\omega_N}^{(C_N^{(1)})}$.

A.1.4. *Macdonald operator for $\mathfrak{g} = A_{2N-1}^{(2)}$.* Here, $(R, S) = (C_N, B_N)$. The map $*$: $R \rightarrow S$ is given by $(e_i \pm e_j)_* = e_i \pm e_j$, and $(2e_i)_* = e_i$. There is a unique minuscule weight of $S^\vee = C_N$, $\pi = e_1 = \omega_1$, and

$$\Phi_{\omega_1} = \frac{1 - tx_1^2}{1 - x_1^2} \prod_{j=2}^N \frac{1 - tx_1 x_j}{1 - x_1 x_j} \frac{tx_1 - x_j}{x_1 - x_j}.$$

Summing over the Weyl group of type C_N ,

$$\mathcal{E}_{\omega_1}^{(A_{2N-1}^{(2)})} = \sum_{\epsilon=\pm 1} \sum_{i=1}^N \frac{1 - tx_i^{2\epsilon}}{1 - x_i^{2\epsilon}} \prod_{j \neq i} \frac{1 - tx_i^\epsilon x_j}{1 - x_i^\epsilon x_j} \frac{tx_i^\epsilon - x_j}{x_i^\epsilon - x_j} \Gamma_i^\epsilon.$$

This will be identified as $\mathcal{D}_1^{(A_{2N-1}^{(2)})}(x; q, t) = \mathcal{E}_{\omega_1}^{(A_{2N-1}^{(2)})}$.

Remark A.1. *The algebra $A_{2N-1}^{(2)}$ is obtained from $A_{2N-1}^{(1)}$ by a folding procedure using the natural \mathbb{Z}_2 automorphism. Remarkably, this extends to the difference operators as follows. Consider the specialization τ of $x = (x_1, x_2, \dots, x_{2N})$ obtained by setting $x_{2N+1-i} = x_i^{-1}$, $i = 1, 2, \dots, N$, and accordingly $\Gamma_{2N+1-i} = \Gamma_i^{-1}$. We have*

$$\tau(\mathcal{D}_1^{(A_{2N-1}^{(1)})}(x; q, t)) = \mathcal{D}_1^{(A_{2N-1}^{(2)})}(x; q, t).$$

However, the $A_{2N-1}^{(1)}$ -Macdonald polynomials specialized via τ have a non-trivial decomposition onto the basis of $A_{2N-1}^{(2)}$ -Macdonald polynomials.

A.1.5. *Macdonald operator for $\mathfrak{g} = D_{N+1}^{(2)}$.* Here, $(R, S) = (B_N, C_N)$. The map $*$: $R \rightarrow S$ is given by $(e_i \pm e_j)_* = e_i \pm e_j$, and $(e_i)_* = 2e_i$. There is a unique minuscule weight $\pi = \frac{1}{2} \sum_{i=1}^N e_i = \omega_N$ of type $S^\vee = B_N$, so that

$$\Phi_{\omega_N} = \prod_{i=1}^N \frac{1 - tx_i}{1 - x_i} \prod_{1 \leq i < j \leq N} \frac{1 - tx_i x_j}{1 - x_i x_j}.$$

Summing over the Weyl group of type C_N gives

$$(A.7) \quad \mathcal{E}_{\omega_N}^{(D_{N+1}^{(2)})} = \sum_{\epsilon_1, \dots, \epsilon_N = \pm 1} \prod_{i=1}^N \frac{1 - tx_i^{\epsilon_i}}{1 - x_i^{\epsilon_i}} \prod_{1 \leq i < j \leq N} \frac{1 - tx_i^{\epsilon_i} x_j^{\epsilon_j}}{1 - x_i^{\epsilon_i} x_j^{\epsilon_j}} \prod_{i=1}^N \Gamma_i^{\epsilon_i/2}.$$

This will be identified as $\mathcal{D}_N^{(D_{N+1}^{(2)})}(x; q, t) = \mathcal{E}_{\omega_N}^{(D_{N+1}^{(2)})}$.

A.2. Higher order Koornwinder-Macdonald difference operators. For generic parameters (a, b, c, d, q, t) , Koornwinder defined the first order q -difference operator whose eigenfunctions are the Koornwinder polynomials, invariant under the Weyl group of type C . Consequently, van Diejen [vD95] defined a commuting family of higher order difference operators with the same eigenfunctions. We recall this construction in A.2.1. Using the spectrum of these operators, we construct in A.2.2 linear combinations of these, such that their eigenvalues are proportional to elementary symmetric functions $\hat{e}_a(s)$. We add to this certain higher order operators due to Rains [Rai05]. Upon specialization of the parameters (a, b, c, d) , we combine this in Sect. A.2.4, with Macdonald's construction of Section A.1.

A.2.1. *van Diejen's higher order Koornwinder difference operators.*

Definition A.2. *The van Diejen operator of order $m \in [1, N]$ is*

$$(A.8) \quad \mathcal{V}_m^{(a,b,c,d)} := \sum_{\substack{J \subset [1, N], |J|=m \\ \epsilon_j = \pm 1 \forall j \in J}} \sum_{s=1}^m (-1)^{s-1} \sum_{\emptyset \subsetneq J_1 \subsetneq \dots \subsetneq J_s = J} \prod_{r=1}^s V_{\{x\}, \{\epsilon\}; J_r \setminus J_{r-1}; K_r}^{(a,b,c,d)} \left(\prod_{j \in J_1} \Gamma_j^{\epsilon_j} - 1 \right),$$

where $J_0 = \emptyset$, $K_r = [1, N] \setminus J_r$ and

$$(A.9) \quad V_{\{x\}, \{\epsilon\}; J; K}^{(a,b,c,d)} := \prod_{i \in J} \frac{(1 - ax_i^{\epsilon_i})(1 - bx_i^{\epsilon_i})(1 - cx_i^{\epsilon_i})(1 - dx_i^{\epsilon_i})}{(1 - x_i^{2\epsilon_i})(1 - qx_i^{2\epsilon_i})} \\ \times \prod_{\substack{i < j \\ i, j \in J}} \frac{1 - tx_i^{\epsilon_i} x_j^{\epsilon_j}}{1 - x_i^{\epsilon_i} x_j^{\epsilon_j}} \frac{1 - qtx_i^{\epsilon_i} x_j^{\epsilon_j}}{1 - qx_i^{\epsilon_i} x_j^{\epsilon_j}} \prod_{\substack{i \in J \\ j \in K}} \frac{1 - tx_i^{\epsilon_i} x_j}{x_i^{\epsilon_i} - x_j},$$

where $J, K \subset [1, N]$ are such that $J \cap K = \emptyset$.

When $m = 1$, the van Diejen operator is the Koornwinder operators of Equation (3.2).

Example A.3. *Define*

$$(A.10) \quad A^{(a,b,c,d)}(x) = \frac{(1 - ax)(1 - bx)(1 - cx)(1 - dx)}{(1 - x^2)(1 - qx^2)}.$$

Then

$$\mathcal{V}_2^{(a,b,c,d)} = \sum_{\substack{1 \leq i_1 < i_2 \leq N \\ \epsilon_1, \epsilon_2 = \pm 1}} \prod_{\ell=1}^2 \left(A^{(a,b,c,d)}(x_{i_\ell}^{\epsilon_\ell}) \prod_{k \neq i_1, i_2} \frac{1 - tx_{i_\ell}^{\epsilon_\ell} x_k}{1 - x_{i_\ell}^{\epsilon_\ell} x_k} \frac{tx_{i_\ell}^{\epsilon_\ell} - x_k}{x_{i_\ell}^{\epsilon_\ell} - x_k} \right) \\ \times \left\{ \frac{1 - tx_{i_1}^{\epsilon_1} x_{i_2}^{\epsilon_2}}{1 - x_{i_1}^{\epsilon_1} x_{i_2}^{\epsilon_2}} \frac{1 - qtx_{i_1}^{\epsilon_1} x_{i_2}^{\epsilon_2}}{1 - qx_{i_1}^{\epsilon_1} x_{i_2}^{\epsilon_2}} (\Gamma_{i_1}^{\epsilon_1} \Gamma_{i_2}^{\epsilon_2} - 1) - \frac{1 - tx_{i_1}^{\epsilon_1} x_{i_2}}{1 - x_{i_1}^{\epsilon_1} x_{i_2}} \frac{tx_{i_1}^{\epsilon_1} - x_{i_2}}{x_{i_1}^{\epsilon_1} - x_{i_2}} (\Gamma_{i_1}^{\epsilon_1} - 1) \right. \\ \left. - \frac{1 - tx_{i_2}^{\epsilon_2} x_{i_1}}{1 - x_{i_2}^{\epsilon_2} x_{i_1}} \frac{tx_{i_2}^{\epsilon_2} - x_{i_1}}{x_{i_2}^{\epsilon_2} - x_{i_1}} (\Gamma_{i_2}^{\epsilon_2} - 1) \right\},$$

where the sum over s in (A.8) decomposes into three terms, with $s = 1, J_1 = J = \{i_1, i_2\}$, $s = 2, J_1 = \{i_1\}, J_2 = J = \{i_1, i_2\}$ and $s = 2, J_1 = \{i_2\}, J_2 = J = \{i_1, i_2\}$.

The operators (A.8) form a commuting family of difference operators with common eigenfunctions being the Koornwinder polynomials. We give a description of their eigenvalues. Let $\sigma = \sqrt{\frac{abcd}{q}}$. Recall the elementary and complete symmetric functions $e_k(x) = s_{1^k}(x)$ and $h_k(x) = s_m(x)$.

Definition A.4. For arbitrary $\lambda_1, \dots, \lambda_N$, and $k, m \in [1, N]$, we define the collections of variables

$$\begin{aligned} u^{(k)} &:= \{\sigma t^{k-i}\}_{1 \leq i \leq k}, \\ v &:= \{s_i + s_i^{-1}\}_{1 \leq i \leq N}, \\ w_{(m)} &:= \{\sigma t^{N-i} + \sigma^{-1} t^{-(N-i)}\}_{m \leq i \leq N}, \end{aligned}$$

with $s_i = \sigma t^{N-i} q^{\lambda_i}$ as usual, and the functions

$$(A.11) \quad d_m^{(k)} := \sigma^m t^{m(k-\frac{m+1}{2})} \hat{e}_m(u^{(k)}),$$

$$(A.12) \quad d_{\lambda; m} := \sigma^m t^{m(N-\frac{m+1}{2})} \hat{e}_m(s),$$

$$(A.13) \quad f_{\lambda; m} := \sigma^m t^{m(N-\frac{m+1}{2})} \sum_{j=0}^m (-1)^j e_{m-j}(v) h_j(w_{(m)}).$$

The spectral theorem for van Diejen operators is

Theorem A.5. [vD95] The (monic) symmetric Koornwinder polynomials $P_\lambda^{(a,b,c,d)}(x)$ satisfy

$$(A.14) \quad \mathcal{V}_m^{(a,b,c,d)} P_\lambda^{(a,b,c,d)}(x) = f_{\lambda; m} P_\lambda^{(a,b,c,d)}(x).$$

A.2.2. Koornwinder-Macdonald operators. Using the spectral theorem A.14, we can construct linear combinations of van Diejen's operators with eigenvalues equal to $d_{\lambda; m}$. To do this we prove two combinatorial lemmas about symmetric functions. Given a set of variables $x = \{x_1, \dots, x_N\}$, we define associated set $\tilde{x} = \{x_i + x_i^{-1}, i \in [1, N]\}$, and if $\beta \leq N$, define $x^{[\beta]} = \{x_1, \dots, x_\beta\}$ and $\tilde{x}^{[\beta]} = \{x_i + x_i^{-1}, i \in [1, \beta]\}$. In particular, $x^{[0]} = \emptyset$.

Lemma A.6. For all $r \geq 0$ and $n \geq 1$,

$$(A.15) \quad \theta_{r,n}(u) := \sum_{\ell=0}^r (-1)^\ell e_{r-\ell}(u^{[n-\ell]}) h_\ell(u^{[n-\ell+1]}) = \delta_{r,0}.$$

Proof. When $r = 0$ the sum is trivially equal to 1. We consider $r > 0$. We define two types of integer configurations. A fermionic configuration on $\{1, \dots, p\}$ with j particles, denoted by F , is a set of j distinct integers in the set $[1, p]$. A bosonic configuration on




FIGURE 2. An illustration of the involution Φ . Case (a) has $t_F = 6 < t_B = 7$, hence we move the topmost rightmost bosonic particle to a fermionic particle at position $t'_F = t_B$. Case (b) has $t_F = 6 \geq t_B = 5$, hence we move the fermionic particle to a bosonic one at position $t_{B'} = t_F$.

$\{1, \dots, p'\}$ with j' particles, denoted by B , is a sequence of j' integers in $[1, p']$ which are not necessarily distinct. We assign a weight $w_F = \prod_{i \in F} u_i$ to a fermionic configuration, and a weight $w_B = \prod_{i \in B} (-u_i)$ to a bosonic configuration. Moreover, to the pair (F, B) , we assign a weight $w_{F,B} = w_F w_B$. The partition function of j fermions on $[1, p]$ is $e_j(u^{[p]})$, and the partition function of j' bosons on $[1, p']$ is $h_{j'}(-u^{[p']}) = (-1)^{j'} h_{j'}(u^{[p']})$.

We define the following set of pairs of fermionic and bosonic configurations:

$$\mathcal{S}_{n,r} = \{(F, B) \mid t_F := \max(F) \leq n - |B|, t_B := \max(B) \leq n - |B| + 1, |F| + |B| = r\}$$

where $|X|$ is the cardinality of the set X , and if $|X| = 0$ we define $\max(X) = 0$. The identity (A.15) is an identity for the partition function:

$$Z_{n,r} := \sum_{(F,B) \in \mathcal{S}_{n,r}} w_{F,B} = \delta_{r,0}.$$

To prove this, for any $r \geq 1$ we construct a fixed point-free involution Φ on the set $\mathcal{S}_{n,r}$, such that $\Phi(w_F w_B) = -w_F w_B$. If such an involution exists, the partition function for any $r > 0$ vanishes:

$$\sum_{(F,B) \in \mathcal{S}_{n,r}} w_{(F,B)} = \frac{1}{2} \sum_{(F,B) \in \mathcal{S}_{n,r}} (w_{(F,B)} + w_{\Phi(F,B)}) = 0.$$

The involution Φ is illustrated in Figure 2 in the language of particles, namely by considering F, B as the sets of integer coordinates of particles along the integer line: e.g. in the case of Fig. 2(a) we have $F = \{2, 4, 5, 6\}$ and $B = \{1, 1, 1, 3, 4, 4, 7, 7, 7\}$. Let $(F, B) \in \mathcal{S}_{n,r}$. The map Φ acts by moving one particle between F and B , thus preserving $|F| + |B| = r$ and reversing the sign of the weight. It is defined as $\Phi(F, B) = (F', B')$, where

- (a) If $t_B > t_F$: $F' = F \cup t_B$, $B' = B \setminus t_B$. Since $|B'| = |B| - 1$, $t_{F'} = t_B \leq n - |B| + 1 = n - |B'|$ and $t_{B'} \leq t_B \leq n - |B| + 1 = n - |B'| < n - |B'| + 1$. Therefore, $(F', B') \in \mathcal{S}_{n,r}$, while $w_{F',B'} = -w_{F,B}$.
- (b) If $t_B \leq t_F$: $F' = F \setminus t_F$ and $B' = B \cup t_F$. Then $t_{F'} \leq t_F - 1 \leq n - |B| - 1 = n - |B'|$ and $t_{B'} = t_F \leq n - |B| = n - |B'| + 1$, so that $(F', B') \in \mathcal{S}_{n,r}$, and $w_{F',B'} = -w_{F,B}$.

The map Φ is clearly an involution. When $r > 0$, Φ has no fixed points, since one can always move a particle. The Lemma follows. \square

Lemma A.7. *There is an identity on symmetric functions:*

$$(A.16) \quad \hat{e}_m(x) = \sum_{j=0}^m \hat{e}_{m-j}(y^{[N-j]}) \sum_{k=0}^j (-1)^k e_{j-k}(\tilde{x}) h_k(\tilde{y}^{[N-j+1]}).$$

Proof. We rewrite (A.16) using the generating function $\hat{E}(z; x)$ of (3.1) as

$$(A.17) \quad \hat{E}(z; x)|_{z^m} = \sum_{j=0}^m \hat{E}(z; y^{[N-j]})|_{z^{m-j}} F(z; \tilde{x}, \tilde{y}^{[N-j+1]})|_{z^j},$$

where $f(z)|_{z^m}$ is the coefficient of z^m in the series expansion of $f(z)$ around 0, and

$$\hat{E}(z; y^{[\beta]}) = \prod_{i=1}^{\beta} (1 + \tilde{y}_i z + z^2), \quad F(z; \tilde{x}, \tilde{y}^{[\beta]}) = \frac{\prod_{i=1}^N (1 + z\tilde{x}_i)}{\prod_{i=1}^{\beta} (1 + z\tilde{y}_i)}.$$

One can further decompose

$$\prod_{i=1}^{N-j} (1 + z\tilde{y}_i + z^2) = (1 + z^2)^{N-j} \hat{E}\left(\frac{z}{1+z^2}; \tilde{y}\right) = \sum_{k=0}^{N-j} z^k (1 + z^2)^{N-j-k} e_k(\tilde{y}^{[N-j]}),$$

and

$$\frac{\prod_{i=1}^N (1 + z\tilde{x}_i)}{\prod_{i=1}^{N-j+1} (1 + z\tilde{y}_i)} = \prod_{i=1}^N (1 + z\tilde{x}_i) \sum_{\ell \geq 0} (-1)^\ell z^\ell h_\ell(\tilde{y}^{[N-j+1]}).$$

The right hand side of (A.17) is therefore

$$\begin{aligned}
 & \sum_{j=0}^m \sum_{k,\ell} (1+z^2)^{N-j-k} \Big|_{z^{m-j-k}} \prod_{i=1}^N (1+z\tilde{x}_i) \Big|_{z^{j-\ell}} (-1)^\ell e_k(\tilde{y}^{[N-j]}) h_\ell(\tilde{y}^{[N-j+1]}) \\
 = & \sum_{r=0}^m \sum_{\substack{k,\ell \geq 0 \\ k+\ell=r}} \sum_{j=\ell}^{m-k} (1+z^2)^{N-j-k} \Big|_{z^{m-j-k}} \prod_{i=1}^N (1+z\tilde{x}_i) \Big|_{z^{j-\ell}} (-1)^\ell e_k(\tilde{y}^{[N-j]}) h_\ell(\tilde{y}^{[N-j+1]}), \\
 = & \sum_{r=0}^m \sum_{j=0}^{m-r} (1+z^2)^{N-j-r} \Big|_{z^{m-j-r}} \prod_{i=1}^N (1+z\tilde{x}_i) \Big|_{z^j} \sum_{\ell=0}^r (-1)^\ell e_{r-\ell}(\tilde{y}^{[N-j-\ell]}) h_\ell(\tilde{y}^{[N-j-\ell+1]}),
 \end{aligned} \tag{A.18}$$

where we changed variables $k \mapsto r - \ell$ and $j \mapsto j - \ell$. Finally, using Lemma A.6 for the collection $u = \tilde{y}$ and $n = N - j$, the expression above drastically simplifies into

$$\sum_{j=0}^m (1+z^2)^{N-j} \Big|_{z^{m-j}} \prod_{i=1}^N (1+z\tilde{x}_i) \Big|_{z^j} = \prod_{i=1}^N (1+z\tilde{x}_i + z^2) \Big|_{z^m} = \hat{e}_m(x),$$

which implies (A.17). □

Definition A.8. We define the Koornwinder-Macdonald difference operators as

$$\mathcal{D}_m^{(a,b,c,d)}(x) := \sum_{j=0}^m d_j^{(N-m+j)} \mathcal{V}_{m-j}^{(a,b,c,d)}, \quad (m = 1, 2, \dots, N),$$

in terms of the van Diejen operators of Sect.A.2.1, with $d_j^{(k)}$ as in (A.11).

Theorem A.9. The Koornwinder-Macdonald polynomials are common eigenfunctions of \mathcal{D}_m with eigenvalue $d_{\lambda;m}$ defined in (A.12):

$$(A.19) \quad \mathcal{D}_m^{(a,b,c,d)}(x; q, t) P_\lambda^{(a,b,c,d)}(x) = d_{\lambda;m} P_\lambda^{(a,b,c,d)}(x).$$

Proof. Equation (A.19) follows from Theorem A.5 and the relation

$$(A.20) \quad d_{\lambda;m} = \sum_{j=0}^m d_j^{(N-m+j)} f_{\lambda;m-j}.$$

Equation (A.20) is obtained by specializing Lemma A.7 to the variables $x_i = \sigma t^{N-i} q^{\lambda_i}$ and $y_i = \sigma t^{N-i}$, and noting that the prefactors in (A.12-A.13) amount to an overall factor of $\sigma^m t^{m(N-\frac{m+1}{2})}$ on both sides of the equation. □

A.2.3. *Rains operators.* The Rains operator $\widehat{\mathcal{D}}_N^{(a,b,c,d)}(x; q, t)$ defined in (3.7,3.9) has eigenvalues (3.10). This operator is not linearly independent of the K-M operators, as is seen from the following Lemma.

Lemma A.10. *For generic (a, b, c, d) the following relation holds between the Koornwinder-Macdonald operators and the Rains operator:*

$$(A.21) \quad \widehat{\mathcal{D}}_N^{(a,b,c,d)}(x; q, t) = \mathcal{D}_N^{(a,b,c,d)}(x; q, t) + \sum_{m=1}^N (-1)^m \left(a^m b^m + \frac{c^m d^m}{q^m} \right) t^{m(m-1)/2} \mathcal{D}_{N-m}^{(a,b,c,d)}(x; q, t).$$

Proof. This is a consequence of the relation between the eigenvalues:

$$(A.22) \quad \hat{d}_{\lambda;N}^{(a,b,c,d)} = d_{\lambda;N} + \sum_{m=1}^N (-1)^m \left(a^m b^m + \frac{c^m d^m}{q^m} \right) t^{m(m-1)/2} d_{\lambda;N-m}.$$

Using the notation $\sigma = \sqrt{abcd/q}$, $u = ab/\sigma$, $u^{-1} = cd/q\sigma$ and $s_i = \sigma q^{\lambda_i} t^{N-i}$, as well as the expressions (A.12) for $d_{\lambda,m}$ and (3.10) for $\hat{d}_{\lambda;N}$, Equation (A.22) is

$$\begin{aligned} q^{-|\lambda|} \prod_{i=1}^N (1 - us_i)(1 - u^{-1}s_i) &= \sigma^N t^{N(N-1)/2} \left\{ \hat{e}_N(s) + \sum_{m=1}^N (-1)^m (u^m + u^{-m}) \hat{e}_{N-m}(s) \right\} \\ &= \frac{(-1)^N \sigma^N t^{N(N-1)/2}}{u^N} \sum_{m=0}^{2N} (-1)^m u^m \hat{e}_m(s) = \frac{(-1)^N \sigma^N t^{N(N-1)/2}}{u^N} \prod_{i=1}^N (1 - us_i)(1 - us_i^{-1}), \end{aligned}$$

where we have used (3.1) in the last line. The equality follows from $\prod_{i=1}^N s_i = q^{|\lambda|} \sigma^N t^{N(N-1)/2}$. \square

A.2.4. *Koornwinder-Macdonald operators and Macdonald's operators of Section A.1.* The first Koornwinder-Macdonald operator in Equation of Definition A.8 is

$$(A.23) \quad \begin{aligned} \mathcal{D}_1^{(a,b,c,d)}(x; q, t) &= \mathcal{V}_1^{(a,b,c,d)}(x; q, t) + \frac{1-t^N}{1-t} \left(1 + \frac{abcd}{q} t^{N-1} \right), \\ &= \mathcal{K}_1^{(a,b,c,d)}(x; q, t) + \frac{1-t^N}{1-t} \left(1 + \frac{abcd}{q} t^{N-1} \right). \end{aligned}$$

To relate these with some of the operators of Section A.1, we have the following Lemma.

Lemma A.11. *The first order Koornwinder-Macdonald operator can be written as*

$$\mathcal{D}_1^{(a,b,c,d)}(x; q, t) = \varphi^{(a,b,c,d)}(x) + \sum_{i=1}^N \sum_{\epsilon=\pm 1} \Phi_{i,\epsilon}^{(a,b,c,d)}(x) \Gamma_i^\epsilon,$$

where

$$(A.24) \quad \varphi^{(a,b,c,d)}(x) = \sum_{\epsilon=\pm 1} \frac{(1 - \frac{\epsilon}{q^{1/2}}a)(1 - \frac{\epsilon}{q^{1/2}}b)(1 - \frac{\epsilon}{q^{1/2}}c)(1 - \frac{\epsilon}{q^{1/2}}d)}{2(1-t)(1-q^{-1}t)} \times \left\{ \prod_{i=1}^N \frac{1 - \frac{\epsilon t}{q^{1/2}}x_i}{1 - \frac{\epsilon}{q^{1/2}}x_i} \frac{1 - \frac{\epsilon t}{q^{1/2}}x_i^{-1}}{1 - \frac{\epsilon}{q^{1/2}}x_i^{-1}} - t^N \right\}.$$

Proof. We have:

$$(A.25) \quad \varphi^{(a,b,c,d)}(x) = \frac{1-t^N}{1-t} \left(1 + \frac{abcd}{q} t^{N-1} \right) - \sum_{i=1}^N \sum_{\epsilon=\pm 1} \Phi_{i,\epsilon}^{(a,b,c,d)}(x).$$

Using the simple fraction decomposition of the function

$$(A.26) \quad \theta(z) := \frac{(1 - \frac{z}{a})(1 - \frac{z}{b})(1 - \frac{z}{c})(1 - \frac{z}{d})}{(1 - \frac{z^2}{q})(1 - \frac{z^2}{t})} \prod_{i=1}^N \frac{1 - \frac{z}{t}x_i}{1 - zx_i} \frac{1 - \frac{z}{t}x_i^{-1}}{1 - zx_i^{-1}}.$$

We find

$$\theta(z) = \frac{q}{abcd t^{2N-1}} + \sum_{\epsilon=\pm 1} \left\{ \sum_{i=1}^N \frac{A_{i,\epsilon}}{1 - zx_i^\epsilon} + \frac{B_\epsilon}{1 - \epsilon \frac{z}{q^{1/2}}} + \frac{C_\epsilon}{1 - \epsilon \frac{z}{t^{1/2}}} \right\},$$

where

$$\begin{aligned} A_{i,\epsilon} &= \frac{1-t}{t^{2N}} \frac{(1 - \frac{x_i^{-\epsilon}}{a})(1 - \frac{x_i^{-\epsilon}}{b})(1 - \frac{x_i^{-\epsilon}}{c})(1 - \frac{x_i^{-\epsilon}}{d})}{(1 - \frac{x_i^{-2\epsilon}}{q})(1 - \frac{x_i^{-2\epsilon}}{t})} \prod_{j \neq i} \frac{tx_i^\epsilon - x_j}{x_i^\epsilon - x_j} \frac{tx_i^\epsilon x_j - 1}{x_i^\epsilon x_j - 1}, \\ &= \frac{q(1-t)}{abcd t^{2N-1}} \Phi_{i,\epsilon}^{(a,b,c,d)}(x), \end{aligned}$$

and

$$\begin{aligned} B_\epsilon &= \frac{(1 - \epsilon \frac{a}{q^{1/2}})(1 - \epsilon \frac{b}{q^{1/2}})(1 - \epsilon \frac{c}{q^{1/2}})(1 - \epsilon \frac{d}{q^{1/2}})}{2abcd t^{2N-1} (1 - \frac{t}{q})} \prod_{i=1}^N \frac{1 - \frac{\epsilon t}{q^{1/2}}x_i}{1 - \frac{\epsilon}{q^{1/2}}x_i} \frac{1 - \frac{\epsilon t}{q^{1/2}}x_i^{-1}}{1 - \frac{\epsilon}{q^{1/2}}x_i^{-1}}, \\ C_\epsilon &= q \frac{(1 - \epsilon \frac{a}{t^{1/2}})(1 - \epsilon \frac{b}{t^{1/2}})(1 - \epsilon \frac{c}{t^{1/2}})(1 - \epsilon \frac{d}{t^{1/2}})}{2abcd t^{N-1} (1 - \frac{q}{t})}. \end{aligned}$$

By definition (A.26), $\theta(0) = 1$, therefore

$$(A.27) \quad \sum_{\epsilon=\pm 1} \left\{ \sum_{i=1}^N \Phi_{i,\epsilon}^{(a,b,c,d)}(x) + \frac{abcd t^{2N-1}}{q(1-t)} (B_\epsilon + C_\epsilon) \right\} = \frac{abcd t^{2N-1} - 1}{1-t}.$$

Using

$$\sum_{\epsilon=\pm 1} \left\{ \frac{t}{2} \left(1 - \epsilon \frac{a}{t^{1/2}}\right) \left(1 - \epsilon \frac{b}{t^{1/2}}\right) \left(1 - \epsilon \frac{c}{t^{1/2}}\right) \left(1 - \epsilon \frac{d}{t^{1/2}}\right) - \frac{q}{2} \left(1 - \epsilon \frac{a}{q^{1/2}}\right) \left(1 - \epsilon \frac{b}{q^{1/2}}\right) \left(1 - \epsilon \frac{c}{q^{1/2}}\right) \left(1 - \epsilon \frac{d}{q^{1/2}}\right) \right\} = \left(1 - \frac{q}{t}\right) \left(t - \frac{abcd}{q}\right),$$

we have:

$$\begin{aligned} \sum_{\epsilon=\pm 1} \frac{abcd t^{2N-1}}{q(1-t)} C_\epsilon &= \frac{t^N}{2(1-t)(1-\frac{q}{t})} \sum_{\epsilon=\pm 1} \left(1 - \epsilon \frac{a}{t^{1/2}}\right) \left(1 - \epsilon \frac{b}{t^{1/2}}\right) \left(1 - \epsilon \frac{c}{t^{1/2}}\right) \left(1 - \epsilon \frac{d}{t^{1/2}}\right), \\ &= \frac{t^N - \frac{abcd}{q} t^{N-1}}{1-t} - \frac{t^N}{2(1-t)(1-\frac{q}{t})} \sum_{\epsilon=\pm 1} \left(1 - \epsilon \frac{a}{q^{1/2}}\right) \left(1 - \epsilon \frac{b}{q^{1/2}}\right) \left(1 - \epsilon \frac{c}{q^{1/2}}\right) \left(1 - \epsilon \frac{d}{q^{1/2}}\right). \end{aligned}$$

The Lemma follows by substituting this into (A.27), and using the result to reexpress $\varphi^{(a,b,c,d)}(x)$ as given by (A.25). \square

When $c = q^{1/2}$ and $d = -q^{1/2}$, $\varphi^{(a,b,q^{1/2},-q^{1/2})}(x) = 0$. Using Table 1, we obtain the following.

Corollary A.12. *In the cases $\mathfrak{g} = D_N^{(1)}, B_N^{(1)}, A_{2N-1}^{(2)}$ the first Macdonald operator is equal to the first specialized Koornwinder-Macdonald operator:*

$$(A.28) \quad \mathcal{D}_1^{(\mathfrak{g})}(x; q, t) = \sum_{i=1}^N \sum_{\epsilon=\pm 1} \Phi_{i,\epsilon}^{(\mathfrak{g})}(x) \Gamma_i^\epsilon = \mathcal{E}_1^{(\mathfrak{g})}(x; q, t).$$

By direct inspection, the specialized Rains operators of (3.7) and (3.9) can also be expressed in terms Macdonald's operators.

Lemma A.13. *In the cases $\mathfrak{g} = D_N^{(1)}, C_N^{(1)}, D_{N+1}^{(2)}$ we have the identifications*

$$(A.29) \quad \begin{aligned} D_N^{(1)} : \quad & \mathcal{E}_{\omega_N}^{(D_N^{(1)})}(x; q, t) + \mathcal{E}_{\omega_{N-1}}^{(D_N^{(1)})}(x; q, t) = \mathcal{R}_N^{(1,-1)}(x; q, t), \\ & \widehat{\mathcal{D}}_N^{(1,-1,q^{1/2},-q^{1/2})} = \left(\mathcal{R}_N^{(1,-1)}\right)^2; \\ C_N^{(1)} : \quad & \mathcal{E}_{\omega_N}^{(C_N^{(1)})}(x; q, t) = \mathcal{R}_N^{(t^{1/2},-t^{1/2})}(x; q, t), \\ & \widehat{\mathcal{D}}_N^{(t^{1/2},-t^{1/2},t^{1/2}q^{1/2},-t^{1/2}q^{1/2})} = \left(\mathcal{R}_N^{(t^{1/2},-t^{1/2})}\right)^2; \\ D_{N+1}^{(2)} : \quad & \mathcal{E}_{\omega_{N+1}}^{(D_{N+1}^{(2)})}(x; q, t) = \mathcal{R}_N^{(t,-1)}(x; q, t), \\ & \widehat{\mathcal{D}}_N^{(t,-1,tq^{1/2},-q^{1/2})} = \left(\mathcal{R}_N^{(t,-1)}\right)^2. \end{aligned}$$

A.3. \mathfrak{g} -Macdonald operators and their eigenvalues. For each \mathfrak{g} , we have chosen a list of N linearly independent commuting difference operators (see Def. 3.5) using the operators in the preceding sections of this Appendix. These have the property that in the q -Whittaker limit, they and their time-translates provide solutions of the various quantum Q-systems. We first explore redundancies between the various definitions to justify Def. 3.5. Then we describe the eigenvalues of the \mathfrak{g} -Macdonald operators, and conclude with a remark on the choice leading to Def. 3.5.

A.3.1. Redundancies. Specializing the parameters (a, b, c, d) as in Table 1 in the difference operators of Theorem A.9 gives a list of N commuting difference operators for each \mathfrak{g} . As noted in Corollary A.12, some of these are Macdonald's operators listed in Section A.1. In addition, there are non-linear relations⁹:

$$\begin{aligned} \mathcal{E}_{\omega_N}^{(D_N^{(1)})} \mathcal{E}_{\omega_{N-1}}^{(D_N^{(1)})} &= \sum_{\alpha=0}^{\lfloor \frac{N-1}{2} \rfloor} t^{\alpha(2\alpha+1)} \mathcal{D}_{N-2\alpha-1}^{(a,b,c,d)}, \\ (\mathcal{E}_{\omega_N}^{(D_N^{(1)})})^2 + (\mathcal{E}_{\omega_{N-1}}^{(D_N^{(1)})})^2 &= \mathcal{D}_N^{(a,b,c,d)} + 2 \sum_{\alpha=0}^{\lfloor \frac{N}{2} \rfloor} t^{\alpha(2\alpha-1)} \mathcal{D}_{N-2\alpha}^{(a,b,c,d)}, \end{aligned}$$

for the $D_N^{(1)}$ specialization. Similarly, for the $C_N^{(1)}$ and $D_{N+1}^{(2)}$ specializations,

$$(\mathcal{E}_{\omega_N}^{(\mathfrak{g})})^2 = \mathcal{D}_N^{(a,b,c,d)} + 2 \sum_{\alpha=1}^N t^{\frac{\alpha(\alpha+1)}{2}} \mathcal{D}_{N-\alpha}^{(a,b,c,d)}.$$

In those cases, the Macdonald operators carry more information than the specialization of the Koornwinder-Macdonald operators $\mathcal{D}_m^{(a,b,c,d)}$. This justifies the choice in Def. 3.5.

A.3.2. The eigenvalues of \mathfrak{g} -Macdonald operators.

Definition A.14. If $m \leq N_{\mathfrak{g}}$, let $d_{\lambda,m}^{(\mathfrak{g})}$ to be the specialization of Equation A.12 to the parameters corresponding to \mathfrak{g} in Table 1:

$$(A.30) \quad d_{\lambda,m}^{(\mathfrak{g})} := d_{\lambda,m} = t^{m(N+\xi_{\mathfrak{g}}-\frac{m+1}{2})} \hat{e}_m(s), \quad m \in [1, N_{\mathfrak{g}}],$$

where $s = t^{\rho^{(\mathfrak{g})}} q^{\lambda}$, i.e. $s_i = q^{\lambda_i} t^{N-i+\xi_{\mathfrak{g}}}$. Otherwise, define

$$(A.31) \quad \begin{aligned} \mathfrak{g} = D_N^{(1)} : \quad & d_{\lambda;\beta}^{(D_N^{(1)})} = t^{\frac{N(N-1)}{4}} \hat{e}_{\beta}^{(D_N)}(s), \quad \beta = N-1, N, \\ \mathfrak{g} = C_N^{(1)}, D_{N+1}^{(2)} : \quad & d_{\lambda;N}^{(\mathfrak{g})} = t^{\frac{N(N+1)}{4}} \hat{e}_N^{(B_N)}(s), \end{aligned}$$

⁹These relations are proved using the same argument as in the proof of Theorem A.9, using identities between eigenvalues. Some are a direct consequence of the identifications with Rains operators from Theorem A.13.

where

$$(A.32) \quad \begin{aligned} \hat{e}_\beta^{(D_N)}(x) &:= \sum_{\substack{\epsilon_1, \dots, \epsilon_N = \pm 1 \\ \prod \epsilon_i = 2(\beta - N) + 1}} \prod_{i=1}^N x_i^{\epsilon_i/2} \quad (\beta = N - 1, N), \\ \hat{e}_N^{(B_N)}(x) &:= \prod_{i=1}^N \frac{1 + x_i}{\sqrt{x_i}} = \hat{e}_{N-1}^{(D_N)}(x) + \hat{e}_N^{(D_N)}(x). \end{aligned}$$

The functions $\hat{e}_m^{(R^*)}(s)$ have the property that their dominant monomial (in powers of t) is $s^{\omega_m^*}$, where ω_m^* are the fundamental weights of R^* .

Theorem A.15. *The \mathfrak{g} -specialized Koornwinder polynomials are common eigenvectors of the \mathfrak{g} -Macdonald operators, satisfying the eigenvalue equation*

$$(A.33) \quad \mathcal{D}_m^{(\mathfrak{g})}(x) P_\lambda^{(\mathfrak{g})} = d_{\lambda; m}^{(\mathfrak{g})} P_\lambda^{(\mathfrak{g})}, \quad m \in [1, N],$$

with $d_{\lambda; m}^{(\mathfrak{g})} = \theta_m^{(\mathfrak{g})} \hat{e}_m^{(R^*)}(s)$ ($\hat{e}_m^{(R^*)} = \hat{e}_m$ for $m \leq N_{\mathfrak{g}}$) and

$$(A.34) \quad \theta_m^{(\mathfrak{g})} = t^{m(N + \xi_{\mathfrak{g}} - \frac{m+1}{2})}, \quad m \leq N_{\mathfrak{g}},$$

$$(A.35) \quad \begin{aligned} \theta_\beta^{(D_N^{(1)})} &= t^{\frac{N(N-1)}{4}}, \quad \beta = N - 1, N, \\ \theta_N^{(\mathfrak{g})} &= t^{\frac{N(N+1)}{4}}, \quad \mathfrak{g} = C_N^{(1)}, D_{N+1}^{(2)}. \end{aligned}$$

A.3.3. *Remark about our choice of Macdonald operators.* We made the choice of \mathfrak{g} -Macdonald operators for simplicity. A more natural choice is to choose eigenvalues proportional to the fundamental characters of R^* . This leads to a more complicated choice of the difference operators, but in the q -Whittaker limit, they have the same limit as our \mathfrak{g} -Macdonald operators.

For Macdonald's operators in Section A.1, the eigenvalue equation is

$$\mathcal{D}_m^{(\mathfrak{g})}(x) P_\lambda^{(\mathfrak{g})}(x) = \theta_m^{(\mathfrak{g})} s_{\omega_m}^{(R^*)}(s) P_\lambda^{(\mathfrak{g})}(x),$$

where the Schur functions $s_{\omega_m}^{(R^*)}$ are the fundamental characters of R^* with highest weight ω_m^* . We may choose the set of difference operators $\tilde{\mathcal{D}}_m^{(\mathfrak{g})}(x)$, with eigenvalues proportional to the fundamental characters of R^* for all m as above. These are related to $\mathcal{D}_m^{(\mathfrak{g})}$ via a triangular change of basis, as can be seen from the relation between the fundamental characters and $e_m^{(R)}$:

$$\begin{aligned} D_N : \quad s_{\omega_m}^{(D_N)} &= \hat{e}_m^{(D_N)}, \quad m \in [1, N], \\ B_N : \quad s_{\omega_m}^{(C_N)} &= \hat{e}_m - \hat{e}_{m-2}, \quad m \in [1, N], \\ C_N : \quad s_{\omega_m}^{(B_N)} &= \hat{e}_m + \hat{e}_{m-1} \quad m \in [1, N - 1], \\ s_{\omega_N}^{(B_N)} &= \hat{e}_N^{(B_N)}. \end{aligned}$$

Therefore $\tilde{\mathcal{D}}^{(\mathfrak{g})}$ are

$$\begin{aligned}
 \mathfrak{g} = D_N^{(1)} : \quad \tilde{\mathcal{D}}_m^{(\mathfrak{g})} &:= \mathcal{D}_m^{(\mathfrak{g})} \quad (m = 1, 2, \dots, N), \\
 \mathfrak{g} = B_N^{(1)}, A_{2N-1}^{(2)}, A_{2N}^{(2)} : \quad \tilde{\mathcal{D}}_m^{(\mathfrak{g})} &:= \mathcal{D}_m^{(\mathfrak{g})} - (1 - \delta_{m,1}) \frac{\theta_m^{(\mathfrak{g})}}{\theta_{m-2}^{(\mathfrak{g})}} \mathcal{D}_{m-2}^{(\mathfrak{g})} \quad (m = 1, 2, \dots, N), \\
 \mathfrak{g} = C_N^{(1)}, D_{N+1}^{(2)} : \quad \tilde{\mathcal{D}}_m^{(\mathfrak{g})} &:= \mathcal{D}_m^{(\mathfrak{g})} + \frac{\theta_m^{(\mathfrak{g})}}{\theta_{m-1}^{(\mathfrak{g})}} \mathcal{D}_{m-1}^{(\mathfrak{g})} \quad (m = 1, 2, \dots, N-1), \\
 \text{(A.36)} \quad \tilde{\mathcal{D}}_N^{(\mathfrak{g})} &:= \mathcal{D}_N^{(\mathfrak{g})},
 \end{aligned}$$

with the convention that $\mathcal{D}_0^{(\mathfrak{g})} = 1$. This choice guarantees the following:

Theorem A.16. *For all $m = 1, 2, \dots, N$ and all \mathfrak{g} ,*

$$\text{(A.37)} \quad \tilde{\mathcal{D}}_m^{(\mathfrak{g})} P_\lambda(x) = \theta_m^{(\mathfrak{g})} s_{\omega_m^{(R^*)}}(s) P_\lambda(x), \quad s = q^\lambda t^{\rho^{(\mathfrak{g})}}.$$

A.4. q -Whittaker limit of the \mathfrak{g} -Macdonald operators. The q -Whittaker limit corresponds to sending $t \rightarrow \infty$. The quantity $A^{(a,b,c,d)}(x)$ of (A.10), which appears in the van Diejen operators (A.8), tends to $\sigma = t^{\xi_{\mathfrak{g}}}$ under the specialization of Table 1. All terms in (A.8) have the same leading behavior as that with $s = 1$ and $J = J_1 = \{1, 2, \dots, m\}$, namely a sum of three contributions from (A.9):

$$m\xi_{\mathfrak{g}} + m(m-1) + 2m(N-m) = m(2N - m - 1 + \xi_{\mathfrak{g}}).$$

Therefore, $\mathcal{V}_m^{(\mathfrak{g})} \sim (\theta_m^{(\mathfrak{g})})^2 V_m^{(\mathfrak{g})}$, where the difference operator $V_m^{(\mathfrak{g})}$ is independent of t . Similarly $\mathcal{D}_m^{(\mathfrak{g})} \sim (\theta_m^{(\mathfrak{g})})^2 D_m^{(\mathfrak{g})}$ where $D_m^{(\mathfrak{g})}$ independent of t , for $m = 1, 2, \dots, N_{\mathfrak{g}}$. By inspection, we find the same leading behavior for $N_{\mathfrak{g}} < m \leq N$, using (A.35). This leads to the definitions:

$$\text{(A.38)} \quad D_m^{(\mathfrak{g})}(x) := \lim_{t \rightarrow \infty} (\theta_m^{(\mathfrak{g})})^{-2} \mathcal{D}_m^{(\mathfrak{g})}(x), \quad V_m^{(\mathfrak{g})} := \lim_{t \rightarrow \infty} (\theta_m^{(\mathfrak{g})})^{-2} \mathcal{V}_m^{(\mathfrak{g})}$$

with $\theta_m^{(\mathfrak{g})}$ as in (A.34-A.35).

Remark A.17. *The limit $t \rightarrow \infty$ of $\mathcal{D}_m^{(\mathfrak{g})}$ is the same as that of $\tilde{\mathcal{D}}_m^{(\mathfrak{g})}$ of Sect. A.3.3: Using $\mathcal{D}_m^{(\mathfrak{g})} \simeq (\theta_m^{(\mathfrak{g})})^2 D_m^{(\mathfrak{g})}$ at leading order in t , the statement follows immediately from (A.36) by noting that for $m \leq N_{\mathfrak{g}}$:*

$$\begin{aligned}
 \lim_{t \rightarrow \infty} \frac{\theta_{m-1}^{(\mathfrak{g})}}{\theta_m^{(\mathfrak{g})}} &= 0 \quad (m \geq 1; \mathfrak{g} = C_N^{(1)}, D_{N+1}^{(2)}), \\
 \lim_{t \rightarrow \infty} \frac{\theta_{m-2}^{(\mathfrak{g})}}{\theta_m^{(\mathfrak{g})}} &= 0 \quad (m \geq 2; \mathfrak{g} = B_N^{(1)}, A_{2N-1}^{(2)}, A_{2N}^{(2)}).
 \end{aligned}$$

For any \mathfrak{g} , define $\Pi_\lambda^{(\mathfrak{g})} = \lim_{t \rightarrow \infty} t^{-\rho^* \cdot \lambda} P_\lambda^{(\mathfrak{g})}$. We refer to these as q -Whittaker functions, although they coincide with the usual definition only for the case of $\mathfrak{g} = D_N^{(1)}$ and the twisted algebras in Table 1. The spectral theorem for $D_m^{(\mathfrak{g})}$ is as follows.

Theorem A.18. *The functions $\Pi_\lambda^{(\mathfrak{g})}$ are common eigenfunctions of $D_m^{(\mathfrak{g})}$, with*

$$D_m^{(\mathfrak{g})}(x) \Pi_\lambda^{(\mathfrak{g})}(x) = \Lambda^{\omega_m^*} \Pi_\lambda^{(\mathfrak{g})}(x).$$

Proof. The eigenvalues are the limits

$$\delta_{\lambda; m}^{(\mathfrak{g})} = \lim_{t \rightarrow \infty} (\theta_m^{(\mathfrak{g})})^{-1} \hat{e}_m^{(R^*)}(s), \quad s_i = q^{\lambda_i} t^{N + \xi_{\mathfrak{g}} - i}.$$

The limit extracts the leading monomial of the s variables in the symmetric function, which is the term involving s_1, s_2, \dots, s_m with positive powers. For $m = 1, 2, \dots, N_{\mathfrak{g}}$ this gives, using (A.30) (A.34) and (A.12): $\delta_{\lambda; m}^{(\mathfrak{g})} = \lim_{t \rightarrow \infty} t^{-m(N + \xi_{\mathfrak{g}} - \frac{m+1}{2})} \hat{e}_m(s) = q^{\lambda_1 + \lambda_2 + \dots + \lambda_m}$. By inspection, using (A.31), (A.35) and (A.32),

$$\begin{aligned} D_N^{(1)} : \quad \delta_{\lambda; N}^{(D_N^{(1)})} &= q^{\frac{1}{2}(\lambda_1 + \lambda_2 + \dots + \lambda_{N-1} + \lambda_N)}, \quad \delta_{\lambda; N-1}^{(D_N^{(1)})} = q^{\frac{1}{2}(\lambda_1 + \lambda_2 + \dots + \lambda_{N-1} - \lambda_N)}, \\ C_N^{(1)} : \quad \delta_{\lambda; N}^{(C_N^{(1)})} &= q^{\frac{1}{2}(\lambda_1 + \lambda_2 + \dots + \lambda_{N-1} + \lambda_N)}, \\ D_{N+1}^{(2)} : \quad \delta_{\lambda; N}^{(D_{N+1}^{(2)})} &= q^{\frac{1}{2}(\lambda_1 + \lambda_2 + \dots + \lambda_{N-1} + \lambda_N)}. \end{aligned}$$

The result can be uniformly written as $\Lambda^{\omega_m^*} = q^{\omega_m^* \cdot \lambda}$, ω_m^* the fundamental weights of R^* . \square

Remark A.19. *Using the convention $V_0^{(\mathfrak{g})} = 1$, Def. A.8 implies*

$$(A.39) \quad D_m^{(\mathfrak{g})} = \sum_{j=0}^m V_j^{(\mathfrak{g})}, \quad m \leq N_{\mathfrak{g}}.$$

APPENDIX B. PIERI RULES

In this section, we list some of the Pieri operators implementing the first Pieri rule for Koornwinder and \mathfrak{g} -Macdonald polynomials. Their q -Whittaker limit gives q -difference Toda Hamiltonians for certain root systems.

B.1. First Pieri rule for Koornwinder polynomials. The first Pieri operator for generic (a, b, c, d) is the q -difference operator $\mathcal{H}_1^{(a, b, c, d)}(s)$ of Theorem (3.3).

Theorem B.1. *The first Pieri operator for Koornwinder polynomials is*

$$\begin{aligned} \mathcal{H}_1^{(a,b,c,d)}(s; q, t) &= a^{-1}t^{1-N}\varphi^{(a^*,b^*,c^*,d^*)}(s) + \sum_{i=1}^N \left[\left(\prod_{j=1}^{i-1} \frac{s_i - t^{-1}s_j}{s_i - s_j} \frac{qs_i - ts_j}{qs_i - s_j} \right) T_i \right. \\ &\quad + \frac{\prod_{u \in \{a^*,b^*,c^*,d^*\}} (1 - u^{-1}s_i)(q - us_i)}{(1 - s_i^2)(q - s_i^2)^2(q^2 - s_i^2)} \\ &\quad \left. \times \left(\prod_{j=i+1}^N \frac{t^{-1}s_i - s_j}{s_i - s_j} \frac{ts_i - qs_j}{s_i - qs_j} \prod_{j \neq i} \frac{q - ts_i s_j}{q - s_i s_j} \frac{1 - t^{-1}s_i s_j}{1 - s_i s_j} \right) T_i^{-1} \right], \end{aligned}$$

where

$$\begin{aligned} \varphi^{(a^*,b^*,c^*,d^*)}(x) &= \sum_{\epsilon=\pm 1} \frac{(1 - \frac{\epsilon}{q^{1/2}}a^*)(1 - \frac{\epsilon}{q^{1/2}}b^*)(1 - \frac{\epsilon}{q^{1/2}}c^*)(1 - \frac{\epsilon}{q^{1/2}}d^*)}{2(1-t)(1-q^{-1}t)} \\ &\quad \times \left\{ \prod_{i=1}^N \frac{1 - \frac{\epsilon t}{q^{1/2}}x_i}{1 - \frac{\epsilon}{q^{1/2}}x_i} \frac{1 - \frac{\epsilon t}{q^{1/2}}x_i^{-1}}{1 - \frac{\epsilon}{q^{1/2}}x_i^{-1}} - t^N \right\}, \end{aligned}$$

and (a^*, b^*, c^*, d^*) are the dual Koornwinder parameters (3.11).

Proof. We use the formula (3.18) with $(a, b, c, d) \rightarrow (a^*, b^*, c^*, d^*)$. For simplicity, we work with the values (a, b, c, d) and interchange them with their duals at the end. We need the following.

Lemma B.2.

$$\begin{aligned} &\Delta^{(a,b,c,d)}(x)^{-1} \Gamma_i \Delta^{(a,b,c,d)}(x) \\ &= \frac{(1 - \frac{1}{x_i^2})(1 - \frac{1}{qx_i^2})}{(1 - \frac{1}{ax_i})(1 - \frac{1}{bx_i})(1 - \frac{1}{cx_i})(1 - \frac{1}{dx_i})} \prod_{j=1}^{i-1} \frac{1 - \frac{qx_i}{tx_j}}{1 - \frac{qx_i}{x_j}} \prod_{j=i+1}^N \frac{1 - \frac{x_j}{tx_i}}{1 - \frac{x_j}{x_i}} \prod_{j \neq i} \frac{1 - \frac{1}{x_i x_j}}{1 - \frac{1}{tx_i x_j}} \Gamma_i, \\ &\Delta^{(a,b,c,d)}(x)^{-1} \Gamma_i^{-1} \Delta^{(a,b,c,d)}(x) \\ &= \frac{(1 - \frac{q}{ax_i})(1 - \frac{q}{bx_i})(1 - \frac{q}{cx_i})(1 - \frac{q}{dx_i})}{(1 - \frac{q}{x_i^2})(1 - \frac{q^2}{x_i^2})} \prod_{j=1}^{i-1} \frac{1 - \frac{x_i}{x_j}}{1 - \frac{x_i}{tx_j}} \prod_{j=i+1}^N \frac{1 - \frac{qx_j}{tx_i}}{1 - \frac{qx_j}{x_i}} \prod_{j \neq i} \frac{1 - \frac{q}{tx_i x_j}}{1 - \frac{q}{x_i x_j}} \Gamma_i^{-1}. \end{aligned}$$

Using Lemma A.11, the conjugation of the $\mathcal{D}_1^{(a,b,c,d)}$ operator boils down to that of the terms $\Phi_{i,\epsilon}^{(a,b,c,d)}\Gamma_i^\epsilon$. We have:

$$\begin{aligned} \Delta^{(a,b,c,d)}(x)^{-1}\Phi_{i,+}^{(a,b,c,d)}\Gamma_i\Delta^{(a,b,c,d)}(x) &= \frac{abcd}{q}t^{2N-2}\left(\prod_{j=1}^{i-1}\frac{1-\frac{x_j}{tx_i}}{1-\frac{x_j}{x_i}}\frac{1-\frac{qx_i}{tx_j}}{1-\frac{qx_i}{x_j}}\right)\Gamma_i, \\ \Delta^{(a,b,c,d)}(x)^{-1}\Phi_{i,-}^{(a,b,c,d)}\Gamma_i^{-1}\Delta^{(a,b,c,d)}(x) &= t^{2N-2}\frac{(1-\frac{a}{x_i})(1-\frac{q}{ax_i})(1-\frac{b}{x_i})(1-\frac{q}{bx_i})(1-\frac{c}{x_i})(1-\frac{q}{cx_i})(1-\frac{d}{x_i})(1-\frac{q}{dx_i})}{(1-\frac{1}{x_i^2})(1-\frac{q}{x_i^2})^2(1-\frac{q^2}{x_i^2})} \\ &\quad \times \left(\prod_{j=i+1}^N\frac{1-\frac{x_i}{tx_j}}{1-\frac{x_i}{x_j}}\frac{1-\frac{qx_j}{tx_i}}{1-\frac{qx_j}{x_i}}\prod_{j\neq i}\frac{1-\frac{q}{tx_ix_j}}{1-\frac{q}{x_ix_j}}\frac{1-\frac{x_ix_j}{t}}{1-x_ix_j}\right)\Gamma_i^{-1}. \end{aligned}$$

We finally use the formula (3.18) for $m = 1$ and substitute the above results with the change $(a, b, c, d) \rightarrow (a^*, b^*, c^*, d^*)$, and the Theorem follows. \square

B.2. First Pieri rules for \mathfrak{g} . We may now specialize the result of Theorem B.1 to \mathfrak{g} , according to Table 1. We first treat the cases where $\mathfrak{g} = D_N^{(1)}, C_N^{(1)}, A_{2N-1}^{(2)}$ for which the constant term $\varphi^{(\mathfrak{g}^*)}(x)$ vanishes¹⁰:

$$\begin{aligned} \mathcal{H}_1^{(D_N^{(1)})} &= \sum_{i=1}^N \left\{ \prod_{j<i} \frac{t^{i-j-1}\Lambda_j - \Lambda_i}{t^{i-j}\Lambda_j - \Lambda_i} \frac{t^{i-j+1}\Lambda_j - q\Lambda_i}{t^{i-j}\Lambda_j - q\Lambda_i} T_i \right. \\ &\quad \left. + \prod_{j>i} \frac{t^{j-i+1}\Lambda_i - q\Lambda_j}{t^{j-i}\Lambda_i - q\Lambda_j} \frac{t^{j-i-1}\Lambda_i - \Lambda_j}{t^{j-i}\Lambda_i - \Lambda_j} \prod_{j\neq i} \frac{1 - t^{2N-i-j-1}\Lambda_i\Lambda_j}{1 - t^{2N-i-j}\Lambda_i\Lambda_j} \frac{t^{2N-i-j+1}\Lambda_i\Lambda_j - q}{t^{2N-i-j}\Lambda_i\Lambda_j - q} T_i^{-1} \right\}, \\ \mathcal{H}_1^{(C_N^{(1)})} &= \sum_{i=1}^N \left\{ \prod_{j<i} \frac{t^{j-i+1}\Lambda_i - \Lambda_j}{t^{j-i}\Lambda_i - \Lambda_j} \frac{\Lambda_j - qt^{j-i-1}\Lambda_i}{\Lambda_j - qt^{j-i}\Lambda_i} T_i \right. \\ &\quad \left. + \frac{1 - t^{N-i}\Lambda_i}{1 - t^{N+1-i}\Lambda_i} \frac{t^{N+2-i}\Lambda_i - q}{t^{N+1-i}\Lambda_i - q} \prod_{j>i} \frac{t^{j-i+1}\Lambda_i - q\Lambda_j}{t^{j-i}\Lambda_i - q\Lambda_j} \frac{\Lambda_j - t^{j-i-1}\Lambda_i}{\Lambda_j - t^{j-i}\Lambda_i} \right. \\ &\quad \left. \times \prod_{j\neq i} \frac{1 - t^{2N+1-i-j}\Lambda_i\Lambda_j}{1 - t^{2N+2-i-j}\Lambda_i\Lambda_j} \frac{t^{2N+3-i-j}\Lambda_i\Lambda_j - q}{t^{2N+2-i-j}\Lambda_i\Lambda_j - q} T_i^{-1} \right\}. \end{aligned}$$

¹⁰Here we show the explicit dependence on Λ_i and t (rather than $s_i = \Lambda_i t^{\xi_{\mathfrak{g}} + N - i}$), as $\xi_{\mathfrak{g}}$ varies with \mathfrak{g} . This makes the limit $t \rightarrow \infty$ easier to follow.

$$\begin{aligned} \mathcal{H}_1^{(A_{2N-1}^{(2)})} &= \sum_{i=1}^N \left\{ \prod_{j<i} \frac{t^{j-i+1}\Lambda_i - \Lambda_j}{t^{j-i}\Lambda_i - \Lambda_j} \frac{\Lambda_j - qt^{j-i-1}\Lambda_i}{\Lambda_j - qt^{j-i}\Lambda_i} T_i \right. \\ &\quad + \frac{t^{2N-2i}\Lambda_i^2 - 1}{t^{2N+1-2i}\Lambda_i^2 - 1} \frac{t^{2N+2-2i}\Lambda_i^2 - q^2}{t^{2N+1-2i}\Lambda_i^2 - q^2} \prod_{j>i} \frac{t^{j-i+1}\Lambda_i - q\Lambda_j}{t^{j-i}\Lambda_i - q\Lambda_j} \frac{t^{j-i-1}\Lambda_i - \Lambda_j}{t^{j-i}\Lambda_i - \Lambda_j} \\ &\quad \left. \times \prod_{j\neq i} \frac{t^{2N-i-j}\Lambda_i\Lambda_j - 1}{t^{2N+1-i-j}\Lambda_i\Lambda_j - 1} \frac{t^{2N+2-i-j}\Lambda_i\Lambda_j - q}{t^{2N+1-i-j}\Lambda_i\Lambda_j - q} T_i^{-1} \right\}. \end{aligned}$$

We now treat the cases $\mathfrak{g} = B_N^{(1)}, D_{N+1}^{(2)}, A_{2N}^{(2)}$ for which $\varphi^{(\mathfrak{g}^*)}$ has a non-trivial contribution. We find:

$$\begin{aligned} \mathcal{H}_1^{(B_N^{(1)})} &= G^{(B_N^{(1)})}(\Lambda) + \sum_{i=1}^N \left\{ \prod_{j<i} \frac{t^{j-i+1}\Lambda_i - \Lambda_j}{t^{j-i}\Lambda_i - \Lambda_j} \frac{\Lambda_j - qt^{j-i-1}\Lambda_i}{\Lambda_j - qt^{j-i}\Lambda_i} T_i \right. \\ &\quad + \frac{1 - t^{2N-2i}\Lambda_i^2}{1 - t^{2N+1-2i}\Lambda_i^2} \frac{q - t^{2N-2i}\Lambda_i^2}{q - t^{2N+1-2i}\Lambda_i^2} \frac{t^{2N+2-2i}\Lambda_i^2 - q^2}{t^{2N+1-2i}\Lambda_i^2 - q^2} \frac{t^{2N+2-2i}\Lambda_i^2 - q}{t^{2N+1-2i}\Lambda_i^2 - q} \\ &\quad \left. \times \prod_{j>i} \frac{t^{j-i+1}\Lambda_i - q\Lambda_j}{t^{j-i}\Lambda_i - q\Lambda_j} \frac{\Lambda_j - t^{j-i-1}\Lambda_i}{\Lambda_j - t^{j-i}\Lambda_i} \prod_{j\neq i} \frac{1 - t^{2N-i-j}\Lambda_i\Lambda_j}{1 - t^{2N+1-i-j}\Lambda_i\Lambda_j} \frac{t^{2N+2-i-j}\Lambda_i\Lambda_j - q}{t^{2N+1-i-j}\Lambda_i\Lambda_j - q} T_i^{-1} \right\}, \end{aligned}$$

where

$$G^{(B_N^{(1)})}(\Lambda) = -1 + \frac{1}{2} \sum_{\epsilon=\pm 1} \prod_{i=1}^N \frac{t^{N+\frac{3}{2}-i}\Lambda_i - \epsilon q^{1/2}}{t^{N+\frac{1}{2}-i}\Lambda_i - \epsilon q^{1/2}} \frac{t^{N-\frac{1}{2}-i}\Lambda_i - \epsilon q^{-1/2}}{t^{N+\frac{1}{2}-i}\Lambda_i - \epsilon q^{-1/2}}.$$

$$\begin{aligned} \mathcal{H}_1^{(D_{N+1}^{(2)})} &= G^{(D_{N+1}^{(2)})}(\Lambda) + \sum_{i=1}^N \left\{ \prod_{j<i} \frac{t^{j-i+1}\Lambda_i - \Lambda_j}{t^{j-i}\Lambda_i - \Lambda_j} \frac{qt^{j-i-1}\Lambda_i - \Lambda_j}{qt^{j-i}\Lambda_i - \Lambda_j} T_i \right. \\ &\quad + \frac{1 - t^{N-i}\Lambda_i}{1 - t^{N+1-i}\Lambda_i} \frac{q^{1/2} - t^{N-i}\Lambda_i}{q^{1/2} - t^{N+1-i}\Lambda_i} \frac{q - t^{N+2-i}\Lambda_i}{q - t^{N+1-i}\Lambda_i} \frac{q^{1/2} - t^{N+2-i}\Lambda_i}{q^{1/2} - t^{N+1-i}\Lambda_i} \\ &\quad \left. \times \prod_{j\neq i} \frac{1 - t^{2N+1-i-j}\Lambda_i\Lambda_j}{1 - t^{2N+2-i-j}\Lambda_i\Lambda_j} \frac{q - t^{2N+3-i-j}\Lambda_i\Lambda_j}{q - t^{2N+2-i-j}\Lambda_i\Lambda_j} \prod_{j>i} \frac{t^{j-i-1}\Lambda_i - \Lambda_j}{t^{j-i}\Lambda_i - \Lambda_j} \frac{t^{j-i+1}\Lambda_i - q\Lambda_j}{t^{j-i}\Lambda_i - q\Lambda_j} T_i^{-1} \right\}, \end{aligned}$$

where

$$G^{(D_{N+1}^{(2)})}(\Lambda) = \frac{(1 - tq^{-1/2})(1 + q^{-1/2})}{1 - tq^{-1}} \left\{ \prod_{i=1}^N \frac{q^{1/2} - \Lambda_i t^{N+2-i}}{q^{1/2} - \Lambda_i t^{N+1-i}} \frac{1 - q^{1/2}\Lambda_i t^{N-i}}{1 - q^{1/2}\Lambda_i t^{N+1-i}} - 1 \right\},$$

and finally

$$\begin{aligned} \mathcal{H}_1^{(A_{2N}^{(2)})} &= G^{(A_{2N}^{(2)})}(\Lambda) + \sum_{i=1}^N \left\{ \prod_{j<i} \frac{t^{j-i+1}\Lambda_i - \Lambda_j}{t^{j-i}\Lambda_i - \Lambda_j} \frac{qt^{j-i-1}\Lambda_i - \Lambda_j}{qt^{j-i}\Lambda_i - \Lambda_j} T_i \right. \\ &+ \frac{1 - t^{N-i}\Lambda_i}{1 - t^{N+1-i}\Lambda_i} \frac{q - t^{2N-2i+1}\Lambda_i^2}{q - t^{2N+2-2i}\Lambda_i^2} \frac{q^2 - t^{2N+3-2i}\Lambda_i^2}{q^2 - t^{2N+2-2i}\Lambda_i^2} \frac{q - t^{N+2-i}\Lambda_i}{q - t^{N+1-i}\Lambda_i} \\ &\left. \times \prod_{j>i} \frac{t^{j-i-1}\Lambda_i - \Lambda_j}{t^{j-i}\Lambda_i - \Lambda_j} \frac{t^{j-i+1}\Lambda_i - q\Lambda_j}{t^{j-i}\Lambda_i - q\Lambda_j} \prod_{j\neq i} \frac{1 - t^{2N+1-i-j}\Lambda_i\Lambda_j}{1 - t^{2N+2-i-j}\Lambda_i\Lambda_j} \frac{q^2 - t^{2N+3-i-j}\Lambda_i\Lambda_j}{q^2 - t^{2N+2-i-j}\Lambda_i\Lambda_j} T_i^{-1} \right\}, \end{aligned}$$

where

$$G^{(A_{2N}^{(2)})}(\Lambda) = \sum_{\epsilon=\pm 1} \frac{(1 - \epsilon tq^{-1/2})(1 + \epsilon q^{-1/2})}{2(1 - tq^{-1})} \left\{ \prod_{i=1}^N \frac{q^{1/2} - \epsilon\Lambda_i t^{N+2-i}}{q^{1/2} - \epsilon\Lambda_i t^{N+1-i}} \frac{\epsilon - q^{1/2}\Lambda_i t^{N-i}}{\epsilon - q^{1/2}\Lambda_i t^{N+1-i}} - 1 \right\}.$$

Remark B.3. When we consider the q -Whittaker limit $t \rightarrow \infty$, the constant pieces $G^{(\mathfrak{g})}(\Lambda)$ in the latter three cases $\mathfrak{g} = B_N^{(1)}, D_{N+1}^{(2)}, A_{2N}^{(2)}$ provide an extra constant term in the limiting Hamiltonians $H_1^{(\mathfrak{g})}$. These read respectively:

$$\lim_{t \rightarrow \infty} G^{(B_N^{(1)})}(\Lambda) = \frac{q^{-1}}{\Lambda_{N-1}\Lambda_N} - \frac{1 + q^{-1}}{\Lambda_N^2}, \quad \lim_{t \rightarrow \infty} G^{(D_{N+1}^{(2)})}(\Lambda) = -\frac{1 + q^{-1/2}}{\Lambda_N}, \quad \lim_{t \rightarrow \infty} G^{(A_{2N}^{(2)})}(\Lambda) = -\frac{1}{\Lambda_N}.$$

APPENDIX C. PROOF OF THEOREM 4.10

Here, we provide the proof of Theorem 4.10, that the time translation operators $g^{(\mathfrak{g})}$ of Equation (4.12) commute with the limiting first Pieri operator/Toda Hamiltonian $H_1^{(\mathfrak{g})}$ of Equation (3.38):

$$(C.1) \quad g^{(\mathfrak{g})} H_1^{(\mathfrak{g})} = H_1^{(\mathfrak{g})} g^{(\mathfrak{g})}.$$

We divide each Hamiltonian into bulk and boundary pieces, study how g commutes with each, and then sum up the contributions to prove the commutation (C.1).

Lemma C.1. For all \mathfrak{g} ,

$$(C.2) \quad g_T (1 - \Lambda^{-\alpha_a}) = (1 - q\Lambda^{-\alpha_a}) T^{-\alpha_a} g_T \quad (a = 0, 1, \dots, N-1),$$

$$(C.3) \quad g_\Lambda (1 - \Lambda^{-\alpha_a}) T_a = (1 - q^{-1}\Lambda^{-\alpha_{a+1}}) T_{a+1} g_\Lambda \quad (a = 1, 2, \dots, N-2),$$

$$(C.4) \quad g_\Lambda (1 - \Lambda^{-\alpha_a}) T_a^{-1} = (1 - q^{-1}\Lambda^{-\alpha_{a-1}}) T_a^{-1} g_\Lambda \quad (a = 1, 2, \dots, N-1),$$

with the convention $\alpha_0 = 0$.

C.1. Bulk and boundary terms. Each of the first Pieri operators (3.38) is the sum of a bulk and boundary piece, where the bulk piece has the same form for all \mathfrak{g} :

$$H_1^{[m]} := \sum_{a=1}^m \left(1 - \frac{\Lambda_a}{\Lambda_{a-1}}\right) T_a + \left(1 - \frac{\Lambda_{a+1}}{\Lambda_a}\right) \frac{1}{T_a},$$

with the convention that $\frac{1}{\Lambda_0} = 0$, for suitable value of $m = m_{\mathfrak{g}}$: $m_{\mathfrak{g}} = N - 2$ for $\mathfrak{g} = D_N^{(1)}, B_N^{(1)}, A_{2N}^{(2)}$, and $m_{\mathfrak{g}} = N - 1$ for all other cases.

The operators $g^{(\mathfrak{g})}$ of (4.12) also have some common structures. For $\mathfrak{g} = B_N^{(1)}, A_{2N-1}^{(2)}, D_{N+1}^{(2)}$, $g = g_T^{\frac{1}{2}} \alpha g_T^{\frac{1}{2}} g_{\Lambda} \beta$ where α, β are functions of Λ_N only. For $\mathfrak{g} = D_N^{(1)}$ $g = g_T g_{\Lambda} \beta$ with β a function of $\Lambda_{N-1} \Lambda_N$ only. The cases $\mathfrak{g} = C_N^{(1)}, A_{2N}^{(2)}$ are different, and have a g operator of the form $g = g_T g_{\Lambda} \alpha g_T g_{\Lambda} \beta$, where α, β are functions of Λ_N only.

Lemma C.2. *For all \mathfrak{g} and $m \leq m_{\mathfrak{g}}$, and $t_1 = 2$ for $\mathfrak{g} = C_N^{(1)}, A_{2N}^{(2)}$, $t_1 = 1$ otherwise,*

$$g^{(\mathfrak{g})} H_1^{[m]} (g^{(\mathfrak{g})})^{-1} = (g_T g_{\Lambda})^{t_1} H_1^{[m]} (g_T g_{\Lambda})^{-t_1}.$$

Proof. The only subtleties arise in the cases $\mathfrak{g} = B_N^{(1)}, A_{2N}^{(2)}$. When $\mathfrak{g} = B_N^{(1)}$, with $g = g_T^{\frac{1}{2}} \alpha g_T^{\frac{1}{2}} g_{\Lambda} \beta$, the action of $g_T^{\frac{1}{2}} g_{\Lambda} \beta$ on $H_1^{[m]}$ creates a term proportional to T_{m+1} , as a consequence of

$$(C.5) \quad g_T g_{\Lambda} \left(1 - \frac{\Lambda_a}{\Lambda_{a-1}}\right) T_a = \left(T_a - \frac{\Lambda_{a+1}}{\Lambda_a} T_{a+1}\right) g_T g_{\Lambda}$$

for $a = m$. This term commutes with α only if $m + 1 \leq N - 1$, hence we set $m_{\mathfrak{g}} = N - 2$. Similarly, when $\mathfrak{g} = A_{2N}^{(2)}$, with $g = g_T g_{\Lambda} \alpha g_T g_{\Lambda} \beta$, the action of $g_T g_{\Lambda} \beta$ on $H_1^{[m]}$ creates a term proportional to T_{m+1} which commutes with α only if $m + 1 \leq N - 1$, hence we set $m_{\mathfrak{g}} = N - 2$ as well. \square

Lemma C.3. *We have the commutation relations for $m = 1, 2, \dots, N - 1$:*

$$g_T g_{\Lambda} H_1^{[m]} = \left\{ H_1^{[m]} + \frac{\Lambda_{m+1}}{\Lambda_m} \left(\frac{1}{T_m} - T_{m+1} \right) \right\} g_T g_{\Lambda}.$$

Proof. We simply sum over the relevant values of a the commutation relations (C.5), as well as the following:

$$(C.6) \quad g_T g_{\Lambda} \left(1 - \frac{\Lambda_{a+1}}{\Lambda_a}\right) \frac{1}{T_a} = \left(\frac{1}{T_a} - \frac{\Lambda_a}{\Lambda_{a-1}} \frac{1}{T_{a-1}} \right) g_T g_{\Lambda},$$

obtained by combining (C.2-C.4). \square

The cases $\mathfrak{g} = C_N^{(1)}$ and $A_{2N}^{(2)}$ are special, as we will have to use Lemma C.3 *twice*. This is deferred to Section C.6 below.

For all the other cases, we apply Lemmas C.2 and C.3 to commute g through the relevant bulk contributions to $H_1^{(\mathfrak{g})}$ corresponding to $a = 1, 2, \dots, m_{\mathfrak{g}}$. The remaining terms of $H_1^{(\mathfrak{g})}$ are the “boundary” terms, corresponding to $m_{\mathfrak{g}} < a \leq N$ and possibly to constant terms independent of $T_a^{\pm 1}$. We address them individually in the following Sections C.2 through C.5. In all cases, the Lemma 4.10 follows from summing all contributions from the bulk $a = 1, 2, \dots, m_{\mathfrak{g}}$ and the boundary $a = m_{\mathfrak{g}} + 1, \dots, N$ plus constant terms.

C.2. Case $\mathfrak{g} = D_N^{(1)}$. Recall that $g = g_T g_{\Lambda} \prod_{n=0}^{\infty} \left(1 - \frac{q^n}{\Lambda_{N-1} \Lambda_N}\right)^{-1}$, and that $m_{\mathfrak{g}} = N - 2$. For $a = N - 1$, we have:

$$\begin{aligned} g \left(1 - \frac{\Lambda_{N-1}}{\Lambda_{N-2}}\right) T_{N-1} &= \left(T_{N-1} - \frac{\Lambda_N}{\Lambda_{N-1}} T_N - \frac{1}{\Lambda_{N-1} \Lambda_N} T_N^{-1} + \frac{1}{\Lambda_{N-1}^2} T_{N-1}^{-1}\right) g, \\ g \left(1 - \frac{\Lambda_N}{\Lambda_{N-1}}\right) \left(1 - \frac{1}{\Lambda_{N-1} \Lambda_N}\right) T_{N-1}^{-1} &= \left(T_{N-1}^{-1} - \frac{\Lambda_{N-1}}{\Lambda_{N-2}} T_{N-2}^{-1}\right) g, \end{aligned}$$

and for $a = N$:

$$\begin{aligned} g \left(1 - \frac{\Lambda_N}{\Lambda_{N-1}}\right) T_N &= \left(T_N - \frac{1}{\Lambda_{N-1} \Lambda_N} T_{N-1}^{-1}\right) g, \\ g \left(1 - \frac{1}{\Lambda_{N-1} \Lambda_N}\right) T_N^{-1} &= \left(T_N^{-1} - \frac{\Lambda_N}{\Lambda_{N-1}} T_{N-1}^{-1}\right) g. \end{aligned}$$

Eq. (C.1) follows by summing all bulk and boundary contributions.

C.3. Case $\mathfrak{g} = B_N^{(1)}$. Recall that $g = g_1 g_2$, $g_1 = g_T^{\frac{1}{2}} \alpha$, $g_2 = g_1 g_{\Lambda}$ with $\alpha = \prod_{n=0}^{\infty} \left(1 - \frac{q^n}{\Lambda_N^2}\right)^{-1}$, and that $m_{\mathfrak{g}} = N - 2$. The terms $a = N - 1, N$ and the constant term read

$$\begin{aligned} g \left(1 - \frac{\Lambda_N}{\Lambda_{N-1}}\right) T_{N-1}^{-1} &= T_{N-1}^{-1} g_T^{\frac{1}{2}} \alpha g_T^{\frac{1}{2}} \left(1 - \frac{\Lambda_{N-1}}{\Lambda_{N-2}}\right) \alpha g_{\Lambda} = \left(T_{N-1}^{-1} - \frac{\Lambda_{N-1}}{\Lambda_{N-2}} T_{N-2}^{-1}\right) g, \\ g \left(1 - \frac{\Lambda_{N-1}}{\Lambda_{N-2}}\right) T_{N-1} &= T_{N-1} g_1 g_T^{\frac{1}{2}} \left(1 - \frac{\Lambda_N}{\Lambda_{N-1}}\right) \alpha g_{\Lambda} = g_1 \left(T_{N-1} - q^{-1} \frac{\Lambda_N}{\Lambda_{N-1}} T_N^{\frac{1}{2}} T_{N-1}^{\frac{1}{2}}\right) g_2, \\ g \left(1 - \frac{\Lambda_N}{\Lambda_{N-1}}\right) T_N &= g_1 \left(1 - \frac{1}{\Lambda_N^2} T_N^{-1}\right) \left(1 - q^{-2} \frac{1}{\Lambda_N^2} T_N^{-1}\right) T_N g_2, \\ g \left(1 - \frac{1}{\Lambda_N^2}\right) \left(1 - q^2 \frac{1}{\Lambda_N^2}\right) T_N^{-1} &= g_1 \left(T_N^{-1} - q^{-1} \frac{\Lambda_N}{\Lambda_{N-1}} T_{N-1}^{-\frac{1}{2}} T_N^{-\frac{1}{2}}\right) g, \\ g \left(\frac{q^{-2}}{\Lambda_{N-1} \Lambda_N} - \frac{1 + q^{-2}}{\Lambda_N^2}\right) &= g_1 \left(\frac{q^{-1}}{\Lambda_{N-1} \Lambda_N} T_{N-1}^{-\frac{1}{2}} T_N^{-\frac{1}{2}} - q^2 \frac{1 + q^{-2}}{\Lambda_N^2} T_N^{-1}\right) g_2. \end{aligned}$$

Summing these five terms gives

$$\begin{aligned}
 & g \left\{ \left(1 - \frac{\Lambda_N}{\Lambda_{N-1}}\right) T_{N-1}^{-1} + \left(1 - \frac{\Lambda_{N-1}}{\Lambda_{N-2}}\right) T_{N-1} + \left(1 - \frac{\Lambda_N}{\Lambda_{N-1}}\right) T_N \right. \\
 & \quad \left. + \left(1 - \frac{1}{\Lambda_N^2}\right) \left(1 - q^2 \frac{1}{\Lambda_N^2}\right) T_N^{-1} + \frac{q^{-2}}{\Lambda_{N-1}\Lambda_N} - \frac{1+q^{-2}}{\Lambda_N^2} \right\} \\
 & = \left\{ T_{N-1} + \left(1 - \frac{\Lambda_N}{\Lambda_{N-1}}\right) T_N - \frac{\Lambda_{N-1}}{\Lambda_{N-2}} T_{N-2}^{-1} + \left(1 - \frac{\Lambda_N}{\Lambda_{N-1}}\right) T_{N-1}^{-1} \right. \\
 & \quad \left. + \left(1 - \frac{1}{\Lambda_N^2}\right) \left(1 - q^2 \frac{1}{\Lambda_N^2}\right) T_N^{-1} + \frac{q^{-2}}{\Lambda_{N-1}\Lambda_N} - \frac{1+q^{-2}}{\Lambda_N^2} \right\} g.
 \end{aligned}$$

Eq. (C.1) follows by summing all bulk and boundary contributions.

C.4. **Case** $\mathfrak{g} = A_{2N-1}^{(2)}$. Recall that $g = g_T g_\Lambda \alpha$, with $\alpha = \prod_{n=0}^{\infty} \left(1 - \frac{q^{2n}}{\Lambda_N^2}\right)^{-1}$, and $m_{\mathfrak{g}} = N - 1$. For the two boundary terms for $a = N$ we have:

$$\begin{aligned}
 g \left(1 - \frac{\Lambda_N}{\Lambda_{N-1}}\right) T_N &= g_T \left(1 - q^{-2} \frac{1}{\Lambda_N^2}\right) T_N g_\Lambda \alpha = \left(T_N - \frac{1}{\Lambda_N^2} T_N^{-1}\right) g, \\
 g \left(1 - \frac{1}{\Lambda_N^2}\right) T_N^{-1} g^{-1} &= g_T \left(1 - q^{-1} \frac{\Lambda_N}{\Lambda_{N-1}}\right) T_N^{-1} g_\Lambda \alpha = \left(T_N^{-1} - \frac{\Lambda_N}{\Lambda_{N-1}} T_{N-1}^{-1}\right) g.
 \end{aligned}$$

Eq. (C.1) follows by summing all bulk and boundary contributions.

C.5. **Case** $\mathfrak{g} = D_{N+1}^{(2)}$. Recall that $g = g_T g_\Lambda \alpha$, with $\alpha = \prod_{n=0}^{\infty} \left(1 - \frac{q^{\frac{n}{2}}}{\Lambda_N}\right)^{-1}$, and $m_{\mathfrak{g}} = N - 1$. For the two boundary terms for $a = N$ and the constant term we have:

$$\begin{aligned}
 g \left(1 - \frac{\Lambda_N}{\Lambda_{N-1}}\right) T_N &= \left(T_N - \frac{1+q^{-\frac{1}{2}}}{\Lambda_N} + \frac{q}{\Lambda_N^2} T_N^{-1}\right) g, \\
 g \left(1 - \frac{1}{\Lambda_N}\right) \left(1 - \frac{q^{\frac{1}{2}}}{\Lambda_N}\right) T_N^{-1} &= g_T \left(1 - q^{-1} \frac{\Lambda_N}{\Lambda_{N-1}}\right) T_N^{-1} g_\Lambda \alpha = \left(T_N^{-1} - \frac{\Lambda_N}{\Lambda_{N-1}} T_{N-1}^{-1}\right) g, \\
 g \left(-\frac{1+q^{-\frac{1}{2}}}{\Lambda_N}\right) &= -\frac{1+q^{\frac{1}{2}}}{\Lambda_N} T_N^{-1} g.
 \end{aligned}$$

Eq. (C.1) follows by summing all bulk and boundary contributions.

C.6. **Cases** $\mathfrak{g} = C_N^{(1)}, A_{2N}^{(2)}$. Recall the general structure of the operator $g = g_1 g_2$ with $g_1 = g_T g_\Lambda \alpha$ and $g_2 = g_T g_\Lambda \beta$, where α, β are functions of Λ_N only. We note that g_2 is the

same for both cases, with $\beta = \prod_{n=0}^{\infty} \left(1 - \frac{q^n}{\Lambda_N}\right)^{-1}$. We note also that $H_1^{[N-1]}$ is the same for both cases, and Lemma C.3 gives the commutation of g_2 through $H_1^{[N-1]}$:

$$(C.7) \quad g_2 H_1^{[N-1]} = \left\{ H_1^{[N-1]} + \frac{\Lambda_N}{\Lambda_{N-1}} \left(\frac{1}{T_{N-1}} - T_N \right) \right\} g_2.$$

We now need to add the boundary and constant terms. For $\mathfrak{g} = C_N^{(1)}$ we have for $a = N$ (and no constant term):

$$g_2 \left(1 - \frac{\Lambda_N}{\Lambda_{N-1}}\right) T_N = \left(T_N - \frac{q^{-\frac{1}{2}}}{\Lambda_N}\right) g_2, \quad g_2 \left(1 - \frac{1}{\Lambda_N}\right) T_N^{-1} = \left(T_N^{-1} - \frac{\Lambda_N}{\Lambda_{N-1}} T_{N-1}^{-1}\right) g_2.$$

For $\mathfrak{g} = A_{2N}^{(2)}$ we have for $a = N$ and the constant term:

$$\begin{aligned} g_2 \left(1 - \frac{\Lambda_N}{\Lambda_{N-1}}\right) T_N &= g_T \left(1 - \frac{q^{-1}}{\Lambda_N}\right) T_N g_\Lambda \beta = \left(T_N - \frac{q^{-\frac{1}{2}}}{\Lambda_N}\right) g_2, \\ g_2 \left(1 - \frac{1}{\Lambda_N}\right) T_N^{-1} &= \left(T_N^{-1} - \frac{\Lambda_N}{\Lambda_{N-1}} T_{N-1}^{-1}\right) g_2, \\ g_2 \left(1 - \frac{1}{\Lambda_N}\right) &= \left(1 - \frac{q^{\frac{1}{2}}}{\Lambda_N} T_N^{-1}\right) g_2. \end{aligned}$$

Adding these to the bulk contributions (C.7), we deduce the following commutations:

$$(C.8) \quad \mathfrak{g} = C_N^{(1)} : \quad g_2 H_1 = \left\{ H_1 + \frac{1}{\Lambda_N} (T_N^{-1} - q^{-\frac{1}{2}}) \right\} g_2,$$

$$(C.9) \quad \mathfrak{g} = A_{2N}^{(2)} : \quad g_2 H_1 = \left\{ H_1 + \frac{1 - q^{\frac{1}{2}}}{\Lambda_N} (T_N^{-1} - q^{-\frac{1}{2}}) \right\} g_2.$$

We must now commute g_1 through this. For both cases, we split again H_1 into a bulk piece $H_1^{[N-1]}$ and boundary pieces and constant terms. For the bulk contribution, we use again Lemma C.3 to write in both cases

$$(C.10) \quad g_1 H_1^{[N-1]} = \left\{ H_1^{[N-1]} + \frac{\Lambda_N}{\Lambda_{N-1}} \left(\frac{1}{T_{N-1}} - T_N \right) \right\} g_1.$$

Then for $\mathfrak{g} = C_N^{(1)}$ and $a = N$:

$$\begin{aligned} g_1 \left(1 - \frac{\Lambda_N}{\Lambda_{N-1}}\right) T_N &= T_N g_1, \\ g_1 \left(1 - \frac{1}{\Lambda_N}\right) T_N^{-1} &= \left(1 - \frac{q^{\frac{1}{2}}}{\Lambda_N} T_N^{-1}\right) \left(T_N^{-1} - \frac{\Lambda_N}{\Lambda_{N-1}} T_{N-1}^{-1}\right) g_1, \end{aligned}$$

and for $\mathfrak{g} = A_{2N}^{(2)}$, $a = N$ and the constant term:

$$g_1 \left(1 - \frac{\Lambda_N}{\Lambda_{N-1}} \right) T_N = g_T \left(1 - \frac{q^{-\frac{1}{2}}}{\Lambda_N} \right) T_N g_\Lambda \alpha = \left(T_N - \frac{1}{\Lambda_N} \right) g_1,$$

$$g_1 \left(1 - \frac{q^{\frac{1}{2}}}{\Lambda_N} \right) T_N^{-1} = \left(T_N^{-1} - \frac{\Lambda_N}{\Lambda_{N-1}} T_{N-1}^{-1} \right) g_1,$$

$$g_1 \left(1 - \frac{q^{-\frac{1}{2}}}{\Lambda_N} \right) = \left(1 - \frac{1}{\Lambda_N} T_N^{-1} \right) g_1.$$

Summing these with the bulk contributions (C.8) and (C.9) respectively, we arrive at

$$(C.11) \quad \mathfrak{g} = C_N^{(1)} : \quad g_1 \left\{ H_1 + \frac{1}{\Lambda_N} (T_N^{-1} - q^{-\frac{1}{2}}) \right\} = H_1 g_1,$$

$$(C.12) \quad \mathfrak{g} = A_{2N}^{(2)} : \quad g_1 \left\{ H_1 + \frac{1 - q^{\frac{1}{2}}}{\Lambda_N} (T_N^{-1} - q^{-\frac{1}{2}}) \right\} = H_1 g_1.$$

Finally, Lemma 4.10 follows for $\mathfrak{g} = C_N^{(1)}, A_{2N}^{(2)}$ by combining the commutation relations (C.8-C.9) and (C.11-C.12). In all cases, Eq. (C.1) follows by summing all bulk and boundary contributions.

REFERENCES

- [BFN19] Alexander Braverman, Michael Finkelberg, and Hiraku Nakajima. Coulomb branches of $3d$ $N = 4$ quiver gauge theories and slices in the affine Grassmannian. *Adv. Theor. Math. Phys.*, 23(1):75–166, 2019. With two appendices by Braverman, Finkelberg, Joel Kamnitzer, Ryosuke Kodera, Nakajima, Ben Webster and Alex Weekes.
- [BZ05] Arkady Berenstein and Andrei Zelevinsky. Quantum cluster algebras. *Adv. Math.*, 195(2):405–455, 2005.
- [Cha02] Oleg A. Chalykh. Macdonald polynomials and algebraic integrability. *Adv. Math.*, 166(2):193–259, 2002.
- [Che95a] Ivan Cherednik. Double affine Hecke algebras and Macdonald’s conjectures. *Ann. of Math. (2)*, 141(1):191–216, 1995.
- [Che95b] Ivan Cherednik. Macdonald’s evaluation conjectures and difference Fourier transform. *Invent. Math.*, 122(1):119–145, 1995.
- [Che97] Ivan Cherednik. Difference Macdonald-Mehta conjecture. *Internat. Math. Res. Notices*, (10):449–467, 1997.
- [Che05] Ivan Cherednik. *Double affine Hecke algebras*, volume 319 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2005.
- [Che09] Ivan Cherednik. Whittaker limits of difference spherical functions. *Int. Math. Res. Not. IMRN*, (20):3793–3842, 2009.
- [CM06] Vyjayanthi Chari and Adriano Moura. The restricted Kirillov-Reshetikhin modules for the current and twisted current algebras. *Comm. Math. Phys.*, 266(2):431–454, 2006.

- [CNV] S. Cecotti, A. Neitzke, and C. Vafa. R-twisting and 4d/2d correspondences. Preprint arXiv:1006.3435.
- [DFK09] Philippe Di Francesco and Rinat Kedem. Q -systems as cluster algebras. II. Cartan matrix of finite type and the polynomial property. *Lett. Math. Phys.*, 89(3):183–216, 2009.
- [DFK10] Philippe Di Francesco and Rinat Kedem. Q -systems, heaps, paths and cluster positivity. *Comm. Math. Phys.*, 293(3):727–802, 2010.
- [DFK11] Philippe Di Francesco and Rinat Kedem. Non-commutative integrability, paths and quasi-determinants. *Adv. Math.*, 228(1):97–152, 2011.
- [DFK14] Philippe Di Francesco and Rinat Kedem. Quantum cluster algebras and fusion products. *Int. Math. Res. Not. IMRN*, (10):2593–2642, 2014.
- [DFK17a] Philippe Di Francesco and Rinat Kedem. Quantum Q systems: from cluster algebras to quantum current algebras. *Lett. Math. Phys.*, 107(2):301–341, 2017.
- [DFK17b] Philippe Di Francesco and Rinat Kedem. Quantum Q systems: from cluster algebras to quantum current algebras. *Lett. Math. Phys.*, 107(2):301–341, 2017.
- [DFK18] P. Di Francesco and R. Kedem. Difference equations for graded characters from quantum cluster algebra. *Transform. Groups*, 23(2):391–424, 2018.
- [DFK19] Philippe Di Francesco and Rinat Kedem. (q, t) -Deformed Q -Systems, DAHA and Quantum Toroidal Algebras via Generalized Macdonald Operators. *Comm. Math. Phys.*, 369(3):867–928, 2019.
- [DFK21] Philippe Di Francesco and Rinat Kedem. Macdonald Operators and Quantum Q -Systems for classical types. 2021. Preprint arXiv:1908.00806 [math-ph].
- [DFKT17] Philippe Di Francesco, Rinat Kedem, and Bolor Turmunkh. A path model for Whittaker vectors. *J. Phys. A*, 50(25):255201, 35, 2017.
- [Eti99] Pavel Etingof. Whittaker functions on quantum groups and q -deformed Toda operators. In *Differential topology, infinite-dimensional Lie algebras, and applications*, volume 194 of *Amer. Math. Soc. Transl. Ser. 2*, pages 9–25. Amer. Math. Soc., Providence, RI, 1999.
- [FJMM09] Boris Feigin, Michio Jimbo, Tetsuji Miwa and Evgeny Mukhin. Fermionic formulas for eigenfunctions of the difference Toda Hamiltonian. *Lett. Math. Phys.*, 88(1-3), 39, 2009.
- [FL99] Boris Feigin and Sergey Loktev. On generalized Kostka polynomials and the quantum Verlinde rule. In *Differential topology, infinite-dimensional Lie algebras, and applications*, volume 194 of *Amer. Math. Soc. Transl. Ser. 2*, pages 61–79. Amer. Math. Soc., Providence, RI, 1999.
- [FT19] Michael Finkelberg and Alexander Tsymbaliuk. Multiplicative slices, relativistic Toda and shifted quantum affine algebras. In *Representations and nilpotent orbits of Lie algebraic systems*, volume 330 of *Progr. Math.*, pages 133–304. Birkhäuser/Springer, Cham, 2019.
- [GK13] Alexander B. Goncharov and Richard Kenyon. Dimers and cluster integrable systems. *Ann. Sci. Éc. Norm. Supér. (4)*, 46(5):747–813, 2013.
- [GSV11] Michael Gekhtman, Michael Shapiro, and Alek Vainshtein. Generalized Bäcklund-Darboux transformations for Coxeter-Toda flows from a cluster algebra perspective. *Acta Math.*, 206(2):245–310, 2011.
- [GT21] Roman Gonin and Alexander Tsymbaliuk. On Sevostyanov’s construction of quantum difference Toda lattices. *Int. Math. Res. Not. IMRN*, (12):8885–8945, 2021.
- [HKKR00] Tim Hoffmann, Johannes Kellendonk, Nadja Kutz, and Nicolai Reshetikhin. Factorization dynamics and Coxeter-Toda lattices. *Comm. Math. Phys.*, 212(2):297–321, 2000.
- [HKO⁺99] G. Hatayama, A. Kuniba, M. Okado, T. Takagi, and Y. Yamada. Remarks on fermionic formula. In *Recent developments in quantum affine algebras and related topics (Raleigh, NC, 1998)*, volume 248 of *Contemp. Math.*, pages 243–291. Amer. Math. Soc., Providence, RI, 1999.

- [HKO⁺02] Goro Hatayama, Atsuo Kuniba, Masato Okado, Taichiro Takagi, and Zengo Tsuboi. Paths, crystals and fermionic formulae. In *MathPhys odyssey, 2001*, volume 23 of *Prog. Math. Phys.*, pages 205–272. Birkhäuser Boston, Boston, MA, 2002.
- [Kac90] Victor G. Kac. *Infinite-dimensional Lie algebras*. Cambridge University Press, Cambridge, third edition, 1990.
- [Ked08] Rinat Kedem. Q -systems as cluster algebras. *J. Phys. A*, 41(19):194011, 14, 2008.
- [KN99] A. N. Kirillov and M. Noumi. q -difference raising operators for Macdonald polynomials and the integrality of transition coefficients. In *Algebraic methods and q -special functions (Montréal, QC, 1996)*, volume 22 of *CRM Proc. Lecture Notes*, pages 227–243. Amer. Math. Soc., Providence, RI, 1999.
- [Koo92] Tom H. Koornwinder. Askey-Wilson polynomials for root systems of type BC . In *Hypergeometric functions on domains of positivity, Jack polynomials, and applications (Tampa, FL, 1991)*, volume 138 of *Contemp. Math.*, pages 189–204. Amer. Math. Soc., Providence, RI, 1992.
- [KR87] A. N. Kirillov and N. Yu. Reshetikhin. Representations of Yangians and multiplicities of the inclusion of the irreducible components of the tensor product of representations of simple Lie algebras. *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)*, 160(Anal. Teor. Chisel i Teor. Funktsii. 8):211–221, 301, 1987.
- [LNS20] Edwin Langmann, Masatoshi Noumi and Junichi Shiraishi. Basic properties of non-stationary Ruijsenaars functions. *SIGMA* 16, 105, 2020.
- [Lin] Mingyan Simon Lin. Quantum twisted q -systems and graded fermionic sums. Preprint 2021.
- [Lin21] Mingyan Simon Lin. Quantum Q -systems and fermionic sums—the non-simply laced case. *Int. Math. Res. Not. IMRN*, (2):805–854, 2021.
- [Mac01] I. G. Macdonald. Orthogonal polynomials associated with root systems. *Sém. Lothar. Combin.*, 45:Art. B45a, 40, 2000/01.
- [Mac95] I. G. Macdonald. *Symmetric functions and Hall polynomials*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, second edition, 1995. With contributions by A. Zelevinsky, Oxford Science Publications.
- [Nou95] Masatoshi Noumi. Macdonald-Koornwinder polynomials and affine Hecke rings. Number 919, pages 44–55. 1995. Various aspects of hypergeometric functions (Japanese) (Kyoto, 1994).
- [NS12] Masatoshi Noumi and Jun'ichi Shiraishi. A direct approach to the bispectral problem for the ruijsenaars-macdonald q -difference operators, 2012. arXiv:1206.5364 [math.QA].
- [Rai05] Eric M. Rains. BC_n -symmetric polynomials. *Transform. Groups*, 10(1):63–132, 2005.
- [Res03] N. Reshetikhin. Integrability of characteristic Hamiltonian systems on simple Lie groups with standard Poisson Lie structure. *Comm. Math. Phys.*, 242(1-2):1–29, 2003.
- [Sah99] Siddhartha Sahi. Nonsymmetric Koornwinder polynomials and duality. *Ann. of Math. (2)*, 150(1):267–282, 1999.
- [Sh19] Junichi Shiraishi. Affine screening operators, affine Laumon spaces and conjectures concerning non-stationary Ruijsenaars functions. *Journal of Integrable Systems*, 3, 1–30, 2019.
- [SS18] Gus Schrader and Alexander Shapiro. On b -whittaker functions. 2018. Preprint arXiv:1806.00747 [math-ph].
- [SS19] Gus Schrader and Alexander Shapiro. K -theoretic coulomb branches of quiver auge theories. 2019. Preprint arXiv:1910.03186.
- [S14] Jasper Stokman. Connection coefficients for basic Harish-Chandra series. *Adv. Math.*, 250:351–386, 2014.
- [vD95] J. F. van Diejen. Commuting difference operators with polynomial eigenfunctions. *Compositio Math.*, 95(2):183–233, 1995.

- [vD96] J. F. van Diejen. Self-dual Koornwinder-Macdonald polynomials. *Invent. Math.*, 126(2):319–339, 1996.
- [vDE15] J. F. van Diejen and E. Emsiz. Integrable boundary interactions for Ruijsenaars’ difference Toda chain. *Comm. Math. Phys.*, 337(1):171–189, 2015.
- [Vic18] Panupong Vichitkunakorn. Conserved quantities of Q -systems from dimer integrable systems. *Electron. J. Combin.*, 25(1):Paper No. 1.36, 43, 2018.
- [Wil15] Harold Williams. Q -systems, factorization dynamics, and the twist automorphism. *Int. Math. Res. Not. IMRN*, (22):12042–12069, 2015.
- [YY21] Kohei Yamaguchi and Shintarou Yanagida. Specializing koornwinder polynomials to macdonald polynomials of type b, c, d and bc . 2021. Preprint arXiv:2105.00936 [math.QA].

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS MC-382, URBANA, IL 61821, U.S.A.
E-MAIL: PHILIPPE@ILLINOIS.EDU AND INSTITUT DE PHYSIQUE THÉORIQUE, UNIVERSITÉ PARIS SACLAY,
CEA, CNRS, F-91191 GIF-SUR-YVETTE, FRANCE
E-MAIL: PHILIPPE.DI-FRANCESCO@CEA.FR

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS MC-382, URBANA, IL 61821, U.S.A.
E-MAIL: RINAT@ILLINOIS.EDU