

Behavioral assessment of visual acuity in a new multifactorial model of schizophrenia

Caroline Lahogue, Olivia Karche, Théo Guezennec, Michel Boulouard,

Valentine Bouet, Solenn Percelay

▶ To cite this version:

Caroline Lahogue, Olivia Karche, Théo Guezennec, Michel Boulouard, Valentine Bouet, et al.. Behavioral assessment of visual acuity in a new multifactorial model of schizophrenia. Motion Sickness Workshop, Jun 2022, Caen, France. . hal-04216508v2

HAL Id: hal-04216508 https://hal.science/hal-04216508v2

Submitted on 12 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Behavioral assessment of visual acuity

in a new multifactorial model of schizophrenia



optomotor

Caroline Lahogue¹, @, Olivia Karche², Théo Guezennec², Michel Boulouard¹, Valentine Bouet¹, Solenn Percelay¹

@:caroline.lahogue@unicaen.fr

1 : COMETE, UMR 1075 INSERM/Université de Caen Normandie 2 : Université de Caen Normandie

Introduction

Most behavioral tests used to assess cognitive functions (touchscreen, object recognition...) in animal models, require normal vision functioning. However, vision which could be affected by genetic alteration or brain insult, is rarely assessed, possibly leading to misinterpretation.

We are currently studying a new animal model of schizophrenia constructed on the combination of two factors (2-hit) : a genetic factor (deletion of serine racemase altered glutamate transmission) and an environmental factor (maternal separation for 24h at post-natal day 9, early stress). Serine racemase synthesizes the NMDA receptor co-agonist, D-serine and is present not only in the brain but also in the retina (Stevens et al., 2003). Its absence in SRKO mice could disturb vision development. Moreover, the second factor, maternal separation, would have long-term effects on visual functions and the retina (Grigoruta et al., 2020).

Methods



Based on virtual optomotor system of Prusky et al. (2000), our aim was herein to do a longitudinal checking of optomotor response development between 4 and 10 weeks of age in 2-hit mice.

response **Optomotor response:** head movement in the direction of the movement of vertical black and white bars, without body moving.

Visual acuity threshold: the highest frequency (cycle/deg) where the optomotor movement is visible



Longitudinal experiment to follow the evolution of visual acuity.



Number of mice that validated the criterion. Control male and female mice predominantly validated the criterion at 4 weeks, as do the SRKO male and female and MS female groups. However, in the 2-hit male and female groups, only one mouse passed the criterion.

Maximal visual acuity is normally reached during mouse's 6 weeks (Prusky et al. 2000)





Longitudinal tracking of visual acuity in male mice.

In both control, SRKO and MS male mice an increase of visual acuity is observed until the maximal acuity at 6 weeks. After 7 weeks a decrease of visual capacity is found. For the 2-hit group, no improvement of visual acuity is found.

Median +/- interquartile range



No difference between groups

Comparaison of the visual acuity at 6 weeks. The maximal visual acuity for male control and MS mice in both clockwise and counter-clockwise directions is around 0.35 cycle/deg corresponding at the normal vision in adult mice. SRKO female and male mice are a maximal visual acuity of 0.3 cycle/deg. In contrast control and MS female mice have a maximal visual acuity of 0.2 cycle/deg. The 2-hit male and female mice, in both directions, seem to have a deleterious visual acuity (about 0.1 cycle/deg for male mice in clockwise direction and 0.2 cycle/deg in counter-clockwise and lower than 0.1 cycle/deg for female 2-hit mice in both directions).

Conclusion



acuity in female mice.

The optomotor test was used to monitor the visual acuity of juvenile 2-hit mice. The results suggest that the combination of genetic and environmental factors has an impact on visual acuity. It will therefore be important to adapt the behavioral tests to be used to assess cognitive processes and avoid misinterpretation of measurements.

Prusky GT et al. "Behavioral assessment of visual acuity in mice and rats." Vision research vol. 40,16 (2000): 2201-9. doi:10.1016/s0042-6989(00)00081-x Gustafson EC et al. "Endogenous D-serine contributes to NMDA-receptor-mediated light-evoked responses in the vertebrate retina." Journal of neurophysiology vol. 98,1 (2007): 122-30. doi:10.1152/jn.00057.2006 Grigoruta M et al. "Maternal separation induces retinal and peripheral blood mononuclear cell alterations across the lifespan of female rats." Brain research vol. 1749 (2020): 147117. doi:10.1016/j.brainres.2020.147117