



HAL
open science

Simple Lie rings of Morley rank 4

Adrien Deloro, Jules Tindzogho Ntsiri

► **To cite this version:**

Adrien Deloro, Jules Tindzogho Ntsiri. Simple Lie rings of Morley rank 4: The Spanish Inquisition. 2023. hal-04216593

HAL Id: hal-04216593

<https://hal.science/hal-04216593>

Preprint submitted on 25 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Simple Lie rings of Morley rank 4 (The Spanish Inquisition)

Adrien Deloro and Jules Tindzogho Ntsiri

25th September 2023

– *Ici, Chimère ! Arrête-toi !*
– *Non ! Jamais !*

Abstract. We prove the Lie ring equivalent of the Cherlin-Zilber conjecture \bullet in characteristic 0, for any rank and \bullet in characteristic $\neq 2, 3$, for rank ≤ 4 . Both are open in the group case.

§ 1. Introduction — § 2. Generalities — § 3. Proofs — § 4. Questions

1 Introduction

For first glance, our main result is here; an explanation will follow. Just consider Morley rank as a form of abstract dimension not related to any linear structure.

Theorem. *Let \mathfrak{g} be a simple Lie ring of finite Morley rank.*

- *If the characteristic is 0, then \mathfrak{g} is a Lie algebra over a definable field (and one of Lie-Chevalley type).*
- *If the characteristic is $\neq 2, 3$, then \mathfrak{g} does not have Morley rank 4.*

The first ad should be considered folklore; as a matter of fact, it was announced without a proof in [Zil82, Theorem 2]. Later [Nes89] did not explicitly prove nor state it, but he certainly knew it. This is classical model theoretic-algebra, and not the core of the present work. Like the unpublished [Ros91] we shall focus on positive characteristic, where the topic is more algebraic and can be appreciated without knowing mathematical logic. Experts in modular Lie algebras should understand our results, our methods, and most of our final questions.

The second ad is a partial analogue in model theory of [Str07, Theorem 2.2]. We emphasize that one starts with a Lie ring devoid of any linear structure or gradation, which highly complicates matters: paradoxically, Lie rings bear extremely little geometric information. Therefore our proof does not proceed through Cartan subalgebras, but through Borel subalgebras, with obvious influences from group-theoretic ‘local analysis’. Finally we mention that the equivalent for simple *groups* of finite Morley rank is notoriously open and challenging.

Returning to simple Lie rings of finite Morley rank, one conjectures that classifying them would be to the Block-Premet-Strade-Wilson theorem (see § 1.1), what the theory of groups of finite Morley rank is to the classification of the finite simple groups: an abstract, simpler sketch (see § 1.2). This will deserve future work.

1.1 Abstract Lie rings

Lie-Chevalley functors, Lie-Cartan functors. Chevalley’s celebrated basis theorem [Hum75, § 25 and notably § 25.4] brought into Lie algebras the idea of functoriality, with all its power and might. Let Φ be a simple root system, and fix a Chevalley basis of Φ ; this gives rise to the Lie ring $\mathfrak{G}_\Phi(\mathbb{Z})$. (\mathfrak{G} is upper-case fraktur g.) Now for R an associative ring with identity, let $\mathfrak{G}_\Phi(R) = \mathfrak{G}_\Phi(\mathbb{Z}) \otimes_{\mathbb{Z}} R$; this is functorial in R . The construction \mathfrak{G}_Φ could be called a (simple) Lie-Chevalley functor; our terminology may evolve in the future. It generates phrases such as ‘Lie rings of Lie-Chevalley type’, for images of Lie-Chevalley functors, in Chevalley’s notation A_n, \dots, G_2 . (Experts call ‘Lie-Chevalley type’ the *classical* algebras. This clashes with group-theoretic terminology, for which *classical* only means A_n, B_n, C_n, D_n , while E_6, E_7, E_8, F_4, G_2 are referred to as *exceptional*.)

The Lie-Chevalley functors, familiar in algebraic geometry, contrast sharply with other constructions yielding simple Lie algebras, usually called ‘of Cartan type’: Witt $W(m; \underline{1})$, special $S(m; \underline{1})^{(1)}$, Hamiltonian $H(m; \underline{1})^{(2)}$, contact $K(m; \underline{1})^{(1)}$. We prefer ‘of Lie-Cartan type’ to prevent confusion with Cartan’s maximal toral subalgebras.

While infinite-dimensional in general, simple Lie algebras of Lie-Cartan type become finite-dimensional in positive characteristic. Thus finite-dimensional, simple Lie algebras over algebraically closed fields of positive characteristics do not all fall into Chevalley’s families.

Finite-dimensional, simple Lie algebras over algebraically closed fields. A key fact is the Block-(Premet-Strade)-Wilson [BW88; PS06] positive answer to the Kostrikin-Shavarevitch conjecture [KS69], with the following consequence.

Fact. *Let \mathfrak{g} be a finite-dimensional, simple Lie algebra over an algebraically closed field of characteristic ≥ 5 . Then :*

- *either \mathfrak{g} is of Lie-Chevalley type,*
- *or \mathfrak{g} is of Lie-Cartan type,*
- *or the characteristic is 5 and \mathfrak{g} is ‘of Melikian type’ [Mel80].*

One recommends [PS06] or [Str04, Introduction]. No understanding of the proof nor of the result itself is needed here, as the only simple Lie algebra encountered in the present work is \mathfrak{sl}_2 .

Abstract Lie rings. A *Lie ring* is an abelian group equipped with a bi-additive map satisfying Jacobi’s identity and called the *bracket*. Subrings and ideals are defined in these terms; so are nilpotence, solubility, and simplicity. A *Lie morphism* is a morphism of Lie rings, viz. an additive morphism preserving the bracket. There is a legitimate tradition of calling Lie rings ‘Lie algebras over \mathbb{Z} ’.

The *characteristic* of a Lie ring is the exponent of the underlying additive group; ‘characteristic 0’ means torsion-free divisible. Every simple Lie ring has characteristic 0 or a prime. Every such is therefore a simple Lie algebra over either \mathbb{Q} or \mathbb{F}_p ; but the dimension is in general infinite. We reserve the phrase *Lie algebra* for the presence of an interesting field acting compatibly. (For us, interesting means ‘infinite and definable’, in the sense below.)

Simple Lie rings are unclassifiable, unless mathematical logic itself brings some form of ‘tameness’.

1.2 Lie rings in model theory

The present work aims at showing that some simple Lie rings provided by model theory are *finite-dimensional* Lie algebras. We restrict our study to finite Morley rank.

Morley rank (for non-logicians). When doing model-theoretic algebra, Morley rank is best construed as a dimension on definable sets; see for instance [BN94, chapters 2, 3, 4]. Alternative discussions are [Poi01, Introduction] or [ABC08, § A.I.2].

Let $(\mathfrak{g}; +, [\cdot])$ be a Lie ring. A subset $X \subseteq \mathfrak{g}^n$ is *definable* if there is a first-order formula, viz. one using the algebraic operations, elements of \mathfrak{g} as parameters, equations, negations, conjunctions, quantification on elements, and giving X as a solution set. (This is the natural generalisation of *constructible* when losing the Chevalley-Tarski theorem that the constructible class is closed under projections.) For good measure and up to abusing terminology, allow quotients of definable sets by definable equivalence relations.

Now \mathfrak{g} has *finite Morley rank* if there is a dimension function assigning to each non-empty definable set an integer, and satisfying the following:

- for definable X , $\dim X \geq n + 1$ iff there are infinitely many, pairwise disjoint $Y_i \subseteq X$ with $\dim Y_i \geq n$;
- for definable $f: X \rightarrow Y$ and integer k , the set $Y_k = \{y \in Y : \dim f^{-1}(\{y\}) = k\}$ is definable;
- as above, if $Y = Y_k$, then $\dim X = \dim Y + k$;
- as above, there is an integer m such that $Y_0 = \{y \in Y : \text{card } f^{-1}(\{y\}) \leq m\}$.

The definition (actually the Borovik-Poizat axiomatisation of groups of finite Morley rank) was given for self-containedness. In practice, there is a dimension provided by mathematical logic, which obeys algebraically predictable rules. For example, ‘finite non-empty’ is equivalent to ‘0-dimensional’. If \mathfrak{g} is a \mathbb{K} -Lie algebra and \mathbb{K} is definable and infinite, then $\dim \mathfrak{g} = \dim \mathbb{K} \cdot \text{ldim}_{\mathbb{K}}(\mathfrak{g})$, where $\text{ldim}_{\mathbb{K}}$ denotes the \mathbb{K} -linear dimension. If \mathbb{K} is finite but \mathfrak{g} is not, the formula makes no sense as $0 \cdot \infty$ is not defined. If \mathbb{K} is not definable, the formula makes no sense at all since only definable sets bear a dimension.

Lie rings of finite Morley rank. So far they received insufficient attention. Nesin’s seminal [Nes89], from which the first author learnt a lot, and Rosengarten’s thesis [Ros91], are not widely cited. However Baudisch’s groups (non-algebraic, nilpotent groups of Morley rank 2) are built from Lie rings [Bau96], so interactions between Lie rings and model theory are not to be despised. As a matter of fact, an ambitious model-theoretic study of nilpotent Lie algebras and the *Lazard(-Malcev) correspondence* with nilpotent groups is in preparation [DE1+23] under neo-stability assumptions. But we focus on model-theoretic algebra of finite Morley rank and the following.

Conjecture (‘log CZ’; implicit in [Ros91]). *Let \mathfrak{g} be an infinite simple Lie ring of finite Morley rank. Suppose the characteristic is sufficiently large. Then \mathfrak{g} is a simple Lie algebra over an algebraically closed field.*

Remarks.

- Applying the Block-Premet-Strade-Wilson theorem, one would even know the isomorphism type of \mathfrak{g} .
- It remains to explain ‘sufficiently large’. The lazy version is: characteristic 0, which we establish in Theorem 0. The modest version is: $> f(d)$ where d is the Morley rank, and f would be determined while working on the conjecture. The optimistic version is: > 3 . There is an intermediate version in characteristic $> d$, where one would also prove that \mathfrak{g} is of Lie-Chevalley type.

Relations to groups and to Lie algebras.

1. Relation to the Cherlin-Zilber conjecture. The conjecture is a clear analogue of the Cherlin-Zilber conjecture on infinite simple groups of finite Morley rank [ABC08]. We do *not* believe there is a general method for obtaining a Lie ring and an adjoint action from an abstract group of finite Morley rank. Conversely, we do *not* believe there is a general method for retrieving a definable group acting on an abstract Lie ring of finite Morley rank. In short we do *not* believe in a ‘Lie-Chevalley correspondence’ at this level of generality, so the two lines of thought should remain independent. We do not know whether adding a Zariski geometry [HZ96] would favorably change the landscape but this is worth asking. See final questions in § 4.
2. Relation to the Block-Premet-Strade-Wilson theorem. Work on the Cherlin-Zilber conjecture is supposed to provide a simplified skeleton of the classification of the finite simple groups. Though the latter is settled but not the former, short versions are always interesting to have. Likewise, we ask experts whether work on ‘log CZ’ could give a blueprint of the Block-Premet-Strade-Wilson classification [Str04; Str09; Str13].

Rosengarten’s work. The ‘log CZ’ conjecture implicitly motivated Rosengarten’s early and unpublished (and in our opinion, underrated) study [Ros91] of Lie rings of Morley rank ≤ 3 . Directed by Cherlin, Rosengarten proved the following in characteristic $\neq 2, 3$:

- a connected Lie ring of Morley rank 1 is abelian [Ros91, Corollary 4.1.1];
- a connected Lie ring of Morley rank 2 is soluble; if non-nilpotent, it covers $\mathfrak{ga}_1(\mathbb{K})$ [Ros91, Theorem 4.2.1];
- a connected, non-soluble Lie ring of Morley rank 3 covers $\mathfrak{sl}_2(\mathbb{K})$ [Ros91, Theorem 4.4.1].

These statements are discussed further in §§ 3.1–3.3.

Our results. We prove the following.

Theorem. *Let \mathfrak{g} be a simple Lie ring of finite Morley rank.*

- *If the characteristic is 0, then \mathfrak{g} is a Lie algebra over a definable field (and one of Lie-Chevalley type).*

- If the characteristic is $\neq 2, 3$, then \mathfrak{g} does not have Morley rank 4.

Group theorists will deem these results merely inspirational insofar as they should have no impact on the Cherlin-Zilber conjecture, other than a renewed challenge. However seen from the legitimate Lie-theoretic perspective we do believe the connection with the Block-Premet-Strade-Wilson theorem deserves serious investigations in model-theoretic algebra.

2 Notation, facts, lemmas

This section contains general notation and terminology (§ 2.1), some facts from the theory of groups of finite Morley rank (§ 2.2), and then a couple of lemmas on Lie rings of finite Morley rank (§2.3).

2.1 Notation; Borel and Cartan subrings

- We occasionally combine bracket and product notations, as in Jacobi's identity:

$$[a, bc] = [ab, c] + [b, ac].$$

- To distinguish between mere additive subgroups, subrings, and ideals, we reserve:

$$\leq \text{ for subgroups, } \sqsubseteq \text{ for subrings, } \trianglelefteq \text{ for ideals.}$$

Throughout, \oplus is used at the group-theoretic level.

- If \mathfrak{g} is a Lie ring and $x \in \mathfrak{g}$, we denote by ad_x the derivation $y \mapsto [x, y]$.
- We use capital B for 'bracket', not for 'Borel'. If \mathfrak{g} is a Lie ring and $x \in \mathfrak{g}$, we let $B_x = [x, \mathfrak{g}] = \text{im ad}_x \leq \mathfrak{g}$, a mere *subgroup*. If \mathfrak{g} is connected (see § 2.2), so is B_x . (Notation generalised below.)
- For Borel subrings we favour $\mathfrak{b} \sqsubseteq \mathfrak{g}$. By definition, a Borel subring is one definable, connected, soluble, and maximal as such.
- We use capital C for 'centraliser', not for 'Cartan'. If \mathfrak{g} is a Lie ring with connected components (see § 2.2) and $x \in \mathfrak{g}$, we let $C_x = C_{\mathfrak{g}}^{\circ}(x) = \{y \in \mathfrak{g} : [x, y] = 0\}^{\circ} \sqsubseteq \mathfrak{g}$, a *subring*. (Notation generalised below.)
- For Cartan subrings we favour $\mathfrak{c} \sqsubseteq \mathfrak{g}$. By definition, a Cartan subring is one definable, connected, nilpotent, and quasi-self-normalising, viz. of finite index in its normaliser.
- For n an integer and $x \in \mathfrak{g}$ we let:

$$B_x^n = \text{im ad}_x^n \leq \mathfrak{g} \quad \text{and} \quad C_x^n = \ker^{\circ} \text{ad}_x^n \leq \mathfrak{g}.$$

Be careful that for $n > 1$, C_x^n need not be a subring. If \mathfrak{g} is connected, these are series of definable, connected *subgroups* (however see Lemma A); the latter is increasing while the former is decreasing. When one series stabilises, so does the other.

- If $x \in \mathfrak{g}$ and k is an integer (modulo the characteristic of \mathfrak{g}), we let $E_k(x) = \{y \in \mathfrak{g} : [x, y] = ky\}^{\circ} = \ker^{\circ}(\text{ad}_x - k \text{Id})$. It is a subgroup, and $[E_k(x), E_{\ell}(x)] \leq E_{k+\ell}(x)$. Often x will remain implicit. For fixed x , the various E_k are in direct sum, being well-understood that indices are integers modulo the characteristic.

2.2 Facts from the theory of groups of finite Morley rank

We need only basic results, used with no reference.

Fact. *Work inside a group of finite Morley rank.*

- *Every descending chain of definable subgroups is stationary [BN94, § 5.1].*
- *Every definable subgroup A has a connected component A° , viz. a smallest definable subgroup of finite index; as a subgroup it is characteristic in A [BN94, § 5.2]. In case \mathfrak{g} is a Lie ring, then \mathfrak{g}° is an ideal of \mathfrak{g} [Nes89, Lemma 1]. In particular, if \mathfrak{g} is simple and infinite then it is connected.*
- *The image of a definable, connected group under a definable group morphism remains definable and connected.*
- *The sum of two definable, connected subgroups remains definable and connected; the sum of infinitely many such reduces to a finite sum.*
- *(Strong) Cherlin-Macintyre-Shelah property: infinite definable skew-fields are algebraically closed fields [BN94, Theorem 8.10].*

Remark (more on fields). Augmented fields abound in model theory, even in universes of finite Morley rank. The existence of fields of finite Morley rank with non-minimal multiplicative group (at least in characteristic 0, [Bau+09]) complicates considerably the study of groups of finite Morley rank. They play no role here. However, fields of finite Morley rank with non-minimal additive groups (which exist only in positive characteristic, [BMZ07]) make matters non-trivial. See Steps 3.1.2, 4.4.3.

We turn to Lie rings of finite Morley rank. The following are straightforward adaptations of their group counterparts. More details can be found in [Ros91, § 3.3].

Fact. *Let \mathfrak{g} be a Lie ring of finite Morley rank.*

- *For $x \in \mathfrak{g}$, one has $\dim \mathfrak{g} = \dim B_x + \dim C_x$.
In particular, if \mathfrak{g} is connected, soluble and infinite, then all centralisers are infinite; otherwise $\mathfrak{g} = B_x \leq \mathfrak{g}'$, a contradiction. Without solubility there are some challenging open problems; see the questions in § 4.*
- *If \mathfrak{g} is connected, then the ideal $\mathfrak{g}' \trianglelefteq \mathfrak{g}$ generated by commutators is definable and connected.
(This is actually trivial, as abelianity completely bypasses the Chevalley-Zilber ‘indecomposable generation’ lemma of [BN94, § 5.4]. We take it as an indication that all the strength of finite Morley rank may not be required. See the final questions in § 4.)*
- *If \mathfrak{g} is connected and the centre $Z(\mathfrak{g})$ is finite, then $\mathfrak{g}/Z(\mathfrak{g})$ is centreless [BN94, Lemma 6.1].*
- *If \mathfrak{g} is non-abelian and definably simple (viz. has no definable ideals other than $\{0\}$ and \mathfrak{g}), then \mathfrak{g} is simple.
(To our surprise, this is absent from Rosengarten’s work and only implicit in Nesin’s. However the argument is present in [Nes89, top of p. 130].)*

- If \mathfrak{g} is nilpotent and infinite, then $Z^\circ(\mathfrak{g})$ is infinite; every definable, infinite subring meets $Z^\circ(\mathfrak{g})$, and the normaliser condition holds [BN94, Lemmas 6.2 and 6.3].
- Borel subrings are quasi-self-normalising. (Relies on Fact 1 below.)

We move to modules, linearisation principles, and their consequences. For V an abelian group, we let $\text{DefEnd}(V)$ be the Lie ring of *definable* endomorphisms. Be careful that $\text{DefEnd}(V)$ itself need not be definable. The bracket is the usual one: $[[f_1, f_2]] = f_1 \circ f_2 - f_2 \circ f_1$.

Definition. A definable Lie ring action is a triple (\mathfrak{g}, V, \cdot) where:

- \mathfrak{g} is a definable Lie ring,
- V is a definable, connected abelian group,
- $\cdot: \mathfrak{g} \times V \rightarrow V$ is a definable map inducing a Lie morphism $\rho: \mathfrak{g} \rightarrow \text{DefEnd}(V)$.

When the context is clear we simply say *definable module*.

A definable module is *faithful* if $\ker \rho = \{0\}$, and *\mathfrak{g} -irreducible* if V has no definable, connected, infinite, proper $0 < W < V$ which is \mathfrak{g} -invariant. (Also called ‘*g-minimal*’.)

The following fact extends Zilber’s original ‘definabilisation’ of the Schur covariance field.

Fact (consequence of [Del23]). *Let (\mathfrak{g}, V) be a definable, faithful, \mathfrak{g} -irreducible module in a finite-dimensional universe. Suppose that \mathfrak{g} and $C_{\text{DefEnd}(V)}(\mathfrak{g})$ are infinite. Then there is a definable field \mathbb{K} such that V is a finite-dimensional \mathbb{K} -vector space and \mathfrak{g} acts \mathbb{K} -linearly.*

2.3 General lemmas

We first discuss iterated centralisers.

Lemma A. *Let \mathfrak{g} be a Lie ring of finite Morley rank and $h \in \mathfrak{g}$. Let n be an integer such that C_h^n is abelian. Then C_h^{n+1} is a subring.*

Proof. Let $\delta = \text{ad}_h$, so that $C_n = \ker^\circ \delta^n$. Let $x, y \in C_{n+1}$. Then:

$$\delta^{n+1}([x, y]) = \sum_{k=0}^{n+1} \binom{n}{k} [\delta^k x, \delta^{n+1-k} y] = \sum_{k=1}^n \binom{n}{k} [\delta^k x, \delta^{n+1-k} y].$$

Now for $1 \leq k \leq n$, one has $\delta^k x \in C_{n+1-k} \leq C_n$ and $\delta^{n+1-k} y \in C_k \leq C_n$. By abelianity, all brackets vanish, so $[x, y] \in \ker \delta^{n+1}$. Since $[C_h^{n+1}, C_h^{n+1}]$ is connected, it is contained in $\ker^\circ \delta^{n+1} = C_h^{n+1}$. \square

There is a form of converse.

Lemma B ([Ros91, Lemma 3.2.6]). *Let \mathfrak{g} be a Lie ring with no additive 2-torsion and $a \in \mathfrak{g}$ be such that $C_a^2 = \mathfrak{g}$. Then B_a is an abelian subring of C_a , and actually an ideal of C_a .*

Proof. By assumption, $B_a \leq C_a$. Now let $b_1, b_2 \in B_a$, say $b_i = [a, x_i]$. Then:

$$\begin{aligned}
[b_1, b_2] &= [b_1, [a, x_2]] = [b_1 a, x_2] + [a, b_1 x_2] \\
&= [a, b_1 x_2] = [a, [a, x_1] x_2] \\
&= [a, [a x_2, x_1] + [a, x_1 x_2]] \\
&= [a, [b_2, x_1]] + [a, \underbrace{[a, x_1 x_2]}_{\in B_a \leq C_a}] \\
&= [a b_2, x_1] + [b_2, a x_1] \\
&= [b_2, b_1].
\end{aligned}$$

Since there is no 2-torsion, we find $[b_1, b_2] = 0$, as wished.

Last, if $b = [a, x] \in B_a$ and $c \in C_a$, then $[b, c] = [[a, x], c] = [a c, x] + [a, x c] = [a, x c] \in B_a$, meaning $B_a \leq C_a$. \square

The following plays a key role when removing nilpotent Borel subrings in Propositions 3.1 and 4.4.

Lemma C ([Ros91, Claim 4.3.2]). *Let \mathfrak{g} be a Lie ring with no additive 2-torsion and $\mathfrak{a} \subseteq \mathfrak{g}$ be an abelian subring. For $n \geq 1$ and $x \in \mathfrak{g}$, let $A_n = [\mathfrak{a}, \dots, [a, x] \dots] \leq \mathfrak{g}$, where \mathfrak{a} appears n times. Then $[A_n, A_n] \leq [\mathfrak{a}, \mathfrak{g}]$.*

Proof. Compute modulo $[\mathfrak{a}, \mathfrak{g}]$. For any $a \in \mathfrak{a}$ and $x, y \in \mathfrak{g}$ one has:

$$0 \equiv [a, [x, y]] = [a x, y] + [x, a y],$$

whence $[a x, y] \equiv -[x, a y]$. Let f be a product of n \mathfrak{a} -derivations, say $f = \text{ad}_{a_1} \circ \dots \circ \text{ad}_{a_n}$ with the a_i all in \mathfrak{a} . By abelianity, the order does not matter. Then $[f(x), y] \equiv (-1)^n [x, f(y)]$. If f' is another product, of say n' \mathfrak{a} -derivations, then keeping abelianity in mind:

$$\begin{aligned}
[f(x), f'(x)] &\equiv (-1)^n [x, f f'(x)] \equiv (-1)^n [x, f' f(x)] \\
&\equiv (-1)^{n+n'} [f'(x), f(x)] \equiv (-1)^{n+n'+1} [f(x), f'(x)].
\end{aligned}$$

Hence if $n + n'$ is even then $2[f(x), f'(x)] \equiv 0$ and $[f(x), f'(x)] \equiv 0$. This is the case when $n' = n$, implying $[A_n, A_n] \leq [\mathfrak{a}, \mathfrak{g}]$. \square

The following extremely useful principle is simply additivity of dim.

Lemma D ('lifting the eigenspace'). *Let \mathfrak{g} be a Lie ring of finite Morley rank and X be a subquotient module, viz. $X = V/V'$ where $V' \leq V \leq \mathfrak{g}$ are \mathfrak{g} -submodules. If for some $h \in \mathfrak{g}$ and some integer k modulo the characteristic, $E_k^X(h) = \{x \in X : [h, x] = x\}^\circ$ is non-trivial, then $E_k^{\mathfrak{g}}(h) \leq \mathfrak{g}$ is non-trivial.*

Proof. Let $\varphi(x) = [h, x] - kx$, which stabilises both V and V' . Let $\pi: V \rightarrow X$ be the quotient map and $\bar{\varphi}: X \rightarrow X$ be induced by φ . Notice $E_k^X = \ker^\circ \bar{\varphi}$. Let $W = (\pi^{-1}(\ker^\circ \bar{\varphi}))^\circ > V'$. Then $\varphi(W) \leq V'$, so $\ker^\circ \varphi > 0$. \square

We now turn to the action of a definable subring on the quotient group.

Lemma E. *Let \mathfrak{g} be a simple Lie ring of finite Morley rank and $\mathfrak{h} \sqsubset \mathfrak{g}$ be a definable, connected, proper subring. Then $I = C_{\mathfrak{h}}^\circ(\mathfrak{g}/\mathfrak{h})$ is nilpotent.*

Proof. Of course $I \trianglelefteq \mathfrak{h}$. Let $I^{[n]}$ be the descending nilpotence series, so that $I^{[n]} \trianglelefteq \mathfrak{h}$. We claim that for $n \geq 0$, one has $[I^{[n+1]}, \mathfrak{g}] \trianglelefteq I^{[n]}$. Indeed when $n = 0$, one has:

$$[I^{[1]}, \mathfrak{g}] = [II, \mathfrak{g}] \trianglelefteq [I, I\mathfrak{g}] \trianglelefteq [I, \mathfrak{h}] \trianglelefteq I,$$

and then by induction:

$$[I^{[n+2]}, \mathfrak{g}] = [II^{[n+1]}, \mathfrak{g}] \trianglelefteq [I, I^{[n+1]}\mathfrak{g}] + [I^{[n+1]}, I\mathfrak{g}] \trianglelefteq [I, I^{[n]}] + [I^{[n+1]}, \mathfrak{h}] \trianglelefteq I^{[n+1]}.$$

By the descending chain condition, there is n such that $I^{[n+1]} = I^{[n]}$. Then $[I^{[n]}, \mathfrak{g}] \trianglelefteq I^{[n]}$ so $I^{[n]} \trianglelefteq \mathfrak{g}$. Since $I^{[n]} < \mathfrak{g}$, we get $I^{[n]} = 0$ by simplicity. \square

To our own surprise we could easily obtain a self-normalisation criterion.

Lemma F. *Let \mathfrak{g} be a definable, connected Lie ring and $\mathfrak{h} \sqsubset \mathfrak{g}$ be a definable, connected subring of codimension 1. Then either $\mathfrak{h} \triangleleft \mathfrak{g}$ is an ideal, or $N_{\mathfrak{g}}(\mathfrak{h}) = \mathfrak{h}$.*

Proof. Let $I = C_{\mathfrak{g}}(\mathfrak{g}/\mathfrak{h})$. If $I = \mathfrak{h}$ then \mathfrak{h} is an ideal of \mathfrak{g} ; so suppose not. Then \mathfrak{h}/I acts non-trivially on $\mathfrak{g}/\mathfrak{h}$. Linearising in dimension 1 (see § 3.1), the action is scalar; hence free.

Let $h \in \mathfrak{h} \setminus I$. Let $n \in N_{\mathfrak{g}}(\mathfrak{h})$. Then $[h, n] = -[n, h] \in \mathfrak{h}$, so the non-trivial, scalar action of h on $\mathfrak{g}/\mathfrak{h}$ sends the image of n to 0. By freeness, $n \in \mathfrak{h}$ already. \square

Last is an important observation on definable derivations.

Lemma G. *An infinite field of finite Morley rank has no non-trivial definable derivation.*

Proof. Recall that the Cherlin-Macintyre-Shelah property holds: \mathbb{K} , or any infinite definable subfield, is algebraically closed.

Let $\delta: \mathbb{K} \rightarrow \mathbb{K}$ be a definable derivation, and \mathbb{K}_0 be the field of constants. Suppose the characteristic is not 2. Let $x \in \mathbb{K}_0$, and let $y \in \mathbb{K}$ such that $y^2 = x$. Then $0 = \delta(x) = 2y\delta(y)$, so $y \in \mathbb{K}_0$. So \mathbb{K}_0 is closed under taking square roots: hence infinite. In characteristic 2, it is however closed under taking cubic roots: hence infinite as well. So in either case, \mathbb{K}_0 is infinite and definable: this proves $\mathbb{K}_0 = \mathbb{K}$, viz. $\delta = 0$. \square

The same holds in o -minimal universes, but seems unclear in the general finite-dimensional case. It however appears like a desirable property for a more systematic investigation of Lie rings in model theory.

3 The proofs

§ 3.0 handles characteristic 0; Theorem 0 will not surprise people familiar with Zilber's early work. Turning to positive characteristic, §§ 3.1, 3.2, 3.3, 3.4 deal with increasing dimensions, and theorems are numbered accordingly. Since we do not reprove Rosengarten's rank 1 analysis, we call it *Fact 1*; Corollary 1 should be folklore. Theorem 2 is already in [Ros91], but the too partial (though important) Corollary 2 is new. Theorem 3 too is already in [Ros91]; its simple Corollary 3 is not. Theorem 4 is definitely new.

3.0 Characteristic 0

The following entirely settles matters in characteristic 0. It is essentially Zilber’s definability of the Schur covariance field and was known to Zilber [Zil82, Theorem 2]. We learnt about this paper from Belegradek, whom we quote (personal correspondence).

This is one-page paper containing only formulations of results and some hints of proofs. It was submitted in 1979 but published only in 1982. The rule of that journal was that they could publish a short paper containing no proofs in case if the author also submitted a manuscript with detailed proofs; the referee wrote a report based on the short paper and manuscript together. I don’t know whether that manuscript is still available. But I think it makes sense for people working on groups of finite Morley rank to rediscover the proofs of the paper, and to advertise the paper (which seems not well-known even though its author is the famous Zilber; maybe this is because that second rate Russian journal was not translated into English, and the paper was not reviewed in MR).

Although we found no further trace of the topic in the literature, the result must also have been known to Nesin [Nes89] among others. Why Rosengarten does not discuss it, is a mystery to us.

Theorem 0. *Work in a theory of finite Morley rank.*

- (i) *Let (\mathfrak{g}, V, \cdot) be a definable Lie ring action with \mathfrak{g} infinite. Suppose it is irreducible and faithful. Suppose further that V does not have bounded exponent. Then the configuration is linear: there is a definable field \mathbb{K} such that V is a \mathbb{K} -vector space and \mathfrak{g} is a \mathbb{K} -Lie subalgebra of $\mathfrak{gl}_{\mathbb{K}}(V)$.*
- (ii) *If in addition \mathfrak{g} is a Lie algebra over some algebraically closed field \mathbb{L} , then $\mathbb{L} \simeq \mathbb{K}$ definably.*
- (iii) *If \mathfrak{g} is a simple Lie ring of finite Morley rank and characteristic 0, then there is a definable field \mathbb{K} such that \mathfrak{g} is the ring of \mathbb{K} -points of one of the Lie-Chevalley functors A_n, \dots, G_2 .*

Proof.

- (i) The action of \mathfrak{g} commutes to $\mathbb{Z}\text{Id}_V$, so we may linearise (see § 2.2). There is a definable field \mathbb{K} such that V a finite-dimensional vector space over \mathbb{K} and $\mathfrak{g} \hookrightarrow \mathfrak{gl}_{\mathbb{K}}(V)$, a mere embedding of Lie rings. Of course $\text{char } \mathbb{K} = 0$. It suffices to show that \mathfrak{g} is a vector subspace of $\mathfrak{gl}_{\mathbb{K}}(V)$. Indeed $N = \{\lambda \in \mathbb{K} : (\forall x \in \mathfrak{g})(\lambda \cdot x \in \mathfrak{g})\}$ is an infinite, definable subring of \mathbb{K} , hence equal to \mathbb{K} by the Cherlin-Macintyre-Shelah property.
- (ii) Let \models denote interpretability. The above and definability of linear structures over a pure field imply $(\mathbb{K}; +, \cdot) \models (\mathfrak{g}, V; +, [\cdot, \cdot], \cdot)$. Now using the adjoint action of a Cartan subalgebra, one has $(\mathfrak{g}; +, [\cdot, \cdot]) \models (\mathbb{L}; +, \cdot)$. It follows $(\mathbb{K}; +, \cdot) \models (\mathbb{L}; +, \cdot)$ and one may conclude by Poizat’s monosomy theorem [Poi01, Theorem 4.15] that $\mathbb{K} \simeq \mathbb{L}$, even \mathbb{K} -definably.
- (iii) Let \mathfrak{g} act on $V = (\mathfrak{g}; +)$ by the adjoint representation: the module is faithful and irreducible. By (i), \mathfrak{g} is a Lie algebra over some algebraically closed field \mathbb{K} . It remains simple as such. Since \mathbb{K} has characteristic 0, the isomorphism type of \mathfrak{g} is given by the classification into Chevalley families. \square

Nothing similar is possible in positive characteristic—and one should also expect objects of Lie-Cartan type.

3.1 Dimension 1

We state Rosengarten’s ‘minimal’ result (which we do not reprove), and then a number of consequences in Corollary 1.

Fact 1 (Rosengarten; [Ros91, Theorem 4.1.1]). *Let \mathfrak{g} be an infinite Lie ring of finite Morley rank and characteristic $\neq 2, 3$. Then \mathfrak{g} contains an infinite, definable, connected, abelian subring.*

This is the analogue of Reineke’s theorem on ‘definably minimal’ groups [Rei75]. The existing proof is non-trivial and relies on results in the finite case. (Moreover the problem is open in characteristic 3). See the section on open questions, § 4.

We prove a couple of consequences, some already mentioned in § 2. In general they will be used without mention.

Corollary 1. *Let $\mathfrak{g} \neq 0$ be a connected Lie ring of finite Morley rank and characteristic $\neq 2, 3$.*

- (i) *Borel subrings are infinite and quasi-self-normalising.*
- (ii) *Let V be a definable, faithful module. Suppose that V is definably minimal as a group (viz. $\{0\}$ -irreducible). Then $\dim \mathfrak{g} \leq \dim V$.*
- (iii) *Let V be a faithful 1-dimensional \mathfrak{g} -module. Then there is a field \mathbb{K} with $V \simeq \mathbb{K}_+$ and $\mathfrak{g} \simeq \mathbb{K} \text{Id}_V$. In particular, there is $h \in \mathfrak{g}$ acting like Id_V .*

We call the last ad ‘linearising in dimension 1’.

Proof.

- (i) By Fact 1, there exist infinite definable abelian subrings, so Borel subrings are non-trivial. Let $\mathfrak{b} \subseteq \mathfrak{g}$ be one such. If $N_{\mathfrak{g}}(\mathfrak{b})/\mathfrak{b}$ is infinite, then it contains an infinite definable abelian subring $\bar{\mathfrak{a}}$. Lifting and taking the connected component, the soluble subring $\pi^{-1}(\bar{\mathfrak{a}})^{\circ} > \mathfrak{b}$ contradicts the definition of \mathfrak{b} .
- (ii) Let V be a faithful \mathfrak{g} -module which is $\{0\}$ -irreducible. Let $v \in V \setminus \{0\}$ and $\mathfrak{h} = C_{\mathfrak{g}}^{\circ}(v)$. Suppose that \mathfrak{h} is infinite. By Fact 1, it contains an infinite, definable, connected, abelian $\mathfrak{a} \subseteq \mathfrak{h}$. Now V remains faithful as an \mathfrak{a} -module, and is \mathfrak{a} -irreducible. Linearising the abelian action (see end of § 2.2), V is a vector space over a definable field \mathbb{K} , and the action of \mathfrak{a} is by scalars, hence free. But \mathfrak{a} acts trivially on $v \neq 0$: a contradiction. So \mathfrak{h} is finite. In particular, $\dim(\mathfrak{g} \cdot v) = \dim \mathfrak{g} \leq \dim V$.
- (iii) Let V be a 1-dimensional, faithful \mathfrak{g} -module. By (ii), $\dim \mathfrak{g} = 1$. By Fact 1, \mathfrak{g} is abelian and we may linearise. Clearly $V \simeq \mathbb{K}_+$ and $\mathfrak{g} \simeq \mathbb{K} \text{Id}_V$. \square

3.2 Dimension 2

We reprove Rosengarten’s dimension 2 analysis (Theorem 2), and derive an important, though too partial, form of Lie’s theorem *in dimension 2* (Corollary 2). The following is already in [Ros91]; our proof is as short but more conceptual.

Theorem 2 (Rosengarten; [Ros91, Theorem 4.2.1]). *Let \mathfrak{g} be a connected Lie ring of Morley rank 2 and characteristic $\neq 2, 3$. Then:*

(i) \mathfrak{g} is soluble;

(ii) if \mathfrak{g} is non-nilpotent, then:

- there is a Cartan subring $\mathfrak{c} \subseteq \mathfrak{g}$ with $\mathfrak{g} = \mathfrak{c} \oplus \mathfrak{g}'$;
- there is $h \in \mathfrak{c}$ with $(\text{ad}_h)|_{\mathfrak{g}'} = \text{Id}_{\mathfrak{g}'}$ (in particular, $Z(\mathfrak{g}) \leq \mathfrak{c}$);
- $Z(\mathfrak{g})$ is finite and there is a definable field \mathbb{K} with $\mathfrak{g}/Z(\mathfrak{g}) \simeq \mathfrak{g}\mathfrak{a}_1(\mathbb{K})$.

Proof.

- (i) Let $\mathfrak{h} \subseteq \mathfrak{g}$ be an infinite, definable, connected subring of minimal dim. By Fact 1, \mathfrak{h} is abelian, so we may assume it is proper; $\dim \mathfrak{h} = 1$. Consider the action of \mathfrak{h} on $V = \mathfrak{g}/\mathfrak{h}$. If it is trivial, then $\mathfrak{h} \triangleleft \mathfrak{g}$ is an ideal, so \mathfrak{g} is soluble and we are done. So we suppose that said action is non-trivial. Linearising in dimension 1, there is $h \in \mathfrak{h}$ acting on V as Id_V . Lifting the eigenspace with Lemma D, $E_1(h) \neq 0$. Now \mathfrak{h} is abelian so $\mathfrak{h} \leq E_0(h)$. Since $\dim \mathfrak{g} = 2$, we get $\mathfrak{g} = E_0 \oplus E_1$; finally $[E_1, E_1] \leq E_2 = 0$ so E_1 is an ideal of \mathfrak{g} and we are done.
- (ii) Suppose \mathfrak{g} is non-nilpotent. Then \mathfrak{g}' has dimension 1, so it is abelian by Fact 1. By non-nilpotence, the action of \mathfrak{g} on \mathfrak{g}' is non-trivial. Linearising in dimension 1, there is $h \in \mathfrak{g}$ acting on \mathfrak{g}' like $\text{Id}_{\mathfrak{g}'}$; hence $E_1 \neq 0$. Since \mathfrak{g} is soluble, one has $E_0 \neq 0$. So we find $\mathfrak{g} = E_0 \oplus E_1$ and $E_1 = \mathfrak{g}'$. Always by non-nilpotence, E_1 is the only 1-dimensional, connected ideal of \mathfrak{g} . It follows that $\mathfrak{c} = E_0$ is abelian, and has finite index in its normaliser: it is a Cartan subring. Clearly $Z(\mathfrak{g}) \leq E_0$. Last, $Z(\mathfrak{g})$ is finite, so $\mathfrak{g}/Z(\mathfrak{g})$ is centreless. We now suppose \mathfrak{g} centreless. The action of E_0 on E_1 is then faithful, so linearisation directly gives the required isomorphism. \square

Remark. A finite centre is unavoidable, as the following example shows. Let:

$$\mathfrak{g} = \left\{ \begin{pmatrix} x & 0 & 0 \\ 0 & \alpha(x) & y \\ 0 & 0 & 0 \end{pmatrix} : (x, y) \in \mathbb{K}^2 \right\},$$

for any additive $\alpha: \mathbb{K}_+ \rightarrow \mathbb{K}_+$, such as for instance $x \mapsto x^p$. Then $Z(\mathfrak{g})$ is isomorphic to $\ker \alpha$.

The following is very unsatisfactory and one should generalise it to $\dim V$ less than the characteristic, with no mention of $\dim \mathfrak{b}$. It plays an important role in the proof of Theorem 4.

Corollary 2. *Work in finite Morley rank. Let \mathfrak{b} be a non-nilpotent, connected Lie ring of characteristic $\neq 2, 3$. Let V be an irreducible \mathfrak{b} -module. Suppose $\dim \mathfrak{b} = \dim V = 2$. Then \mathfrak{b}' centralises V .*

Proof. By Theorem 2, $\mathfrak{a} = \mathfrak{b}' > 0$; moreover there is a Cartan subring $\mathfrak{t} < \mathfrak{b}$ such that $\mathfrak{b} = \mathfrak{t} \oplus \mathfrak{a}$ and $\mathfrak{a} = [\mathfrak{t}, \mathfrak{a}]$.

Let $\Delta = \text{DefEnd}(V)$, which is an associative ring with induced Lie bracket $\llbracket f, g \rrbracket = fg - gf$. Let $\rho: \mathfrak{b} \rightarrow \Delta$, a Lie morphism. We avoid it from notation mostly, with exceptions for enhanced clarity. To prove that $\mathfrak{a} \leq \ker \rho$, we suppose otherwise. Notice that $C_V^\circ(\mathfrak{a})$ is \mathfrak{b} -invariant, so we may suppose $C_V^\circ(\mathfrak{a}) = 0$.

Step 2.1. V is not \mathfrak{a} -irreducible.

Proof. Suppose it is. Linearising the abelian action, $\mathbb{K} = C_{\text{DefEnd}(V)}(\rho(\mathfrak{a}))$ is a definable field containing $\rho(\mathfrak{a})$. We claim that \mathfrak{t} acts by definable derivations of \mathbb{K} . Indeed, let $\lambda \in \mathbb{K}$ and $a \in \mathfrak{a}$. Also let $h \in \mathfrak{t}$ and $a' = [h, a]$. Then:

$$\begin{aligned} \llbracket [\rho(h), \lambda], \rho(a) \rrbracket &= (\rho(h)\lambda - \lambda\rho(h))\rho(a) - \rho(a)(\rho(h)\lambda - \lambda\rho(h)) \\ &= \rho(h)\rho(a)\lambda - \lambda\rho(h)\rho(a) - \rho(a)\rho(h)\lambda + \lambda\rho(a)\rho(h) \\ &= \llbracket \rho(h), \rho(a) \rrbracket \lambda - \lambda \llbracket \rho(h), \rho(a) \rrbracket \\ &= \rho([h, a])\lambda - \lambda\rho([h, a]) \\ &= \llbracket \rho(a'), \lambda \rrbracket \\ &= 0, \end{aligned}$$

proving that $\rho(h)$ does act on \mathbb{K} . The action is clearly that of a derivation. By Lemma G, $\rho(\mathfrak{t})$ must act trivially; in particular $\rho(\mathfrak{t})$ centralises $\rho(\mathfrak{a}) \leq \mathbb{K}$. So $\mathfrak{a} = [\mathfrak{t}, \mathfrak{a}] \leq \ker \rho$, as wanted. \diamond

Step 2.2. V is \mathfrak{a} -irreducible.

Proof. Otherwise let $W < V$ be a 1-dimensional \mathfrak{a} -submodule. If \mathfrak{a} acts trivially on W , then $C_V^\circ(\mathfrak{a}) \neq 0$, a contradiction. So \mathfrak{a} acts non-trivially on W .

Linearising, $\mathbb{K} = C_{\text{DefEnd}(W)}(\mathfrak{a})$ is a definable field containing $\rho(\mathfrak{a})$. We fix $h \in \mathfrak{t}$ ad-acting on \mathfrak{a} like $\text{Id}_{\mathfrak{a}}$, viz. $[h, a] = a$ for $a \in \mathfrak{a}$.

We claim $hW \neq 0$ and $(W \cap hW)^\circ = 0$, so that $V = W + hW$. Suppose $w_1 = hw_2$ for some $w_1, w_2 \in W$. Then applying an arbitrary $a \in \mathfrak{a}$:

$$aw_1 = ahw_2 = haw_2 - aw_2,$$

so $haw_2 = a(w_1 + w_2) \in W$. But if $w_2 \notin C_V(\mathfrak{a})$, then $aw_2 = W$, which proves $hW = W$. So h acts on W , and as before on \mathbb{K} as well. Now it induces a derivation of $\mathbb{K} \geq \mathfrak{a}$, although it acts non-trivially on \mathfrak{a} ; against Lemma G. So $w_2 \in C_V(\mathfrak{a})$, which is finite. So $w_1 = hw_2$ has finitely many solutions, which proves all claims.

We derive a contradiction. Let $w \in W$. Then there are $w_1, w_2 \in W$ such that $h^2w = w_1 + hw_2$. Let $a \in \mathfrak{a}$ act on W like Id_W (this is the element going to $1_{\mathbb{K}}$). Remember $[h, a] = a$; in particular $ah^2 = h^2a - 2ha + a$. Applying to w one finds:

$$\begin{aligned} ah^2w &= aw_1 + ahw_2 = w_1 + haw_2 - aw_2 = w_1 - w_2 + hw_2 \\ &= (h^2a - 2ha + a)w = h^2w - 2hw + w = w_1 + w + h(w_2 - 2w). \end{aligned}$$

This proves $hw \in W \cap hW$ which is finite, so by connectedness $hW = 0$, a contradiction. \diamond

This completes the proof of Corollary 2. \square

3.3 Dimension 3

Theorem 3 (Rosengarten; [Ros91, Theorem 4.4.1]). *Let \mathfrak{g} be a simple Lie ring of Morley rank 3 and characteristic $\neq 2, 3$. Then $\mathfrak{g} \simeq \mathfrak{sl}_2(\mathbb{K})$.*

Our three reasons to give a proof are: 1. Rosengarten's own argument was never published; 2. ours is sufficiently different from Rosengarten's, and more conceptual; 3. parts of it make sense in the group setting; 4. some key ideas reappear in the dimension 4 argument of § 3.4.

The proof first removes nilpotent Borel subrings (Proposition 3.1), then explicitly identifies \mathfrak{g} (Proposition 3.2).

Proposition 3.1. *No Borel subring is nilpotent.*

Proof. Let $\mathfrak{b} \sqsubseteq \mathfrak{g}$ be a nilpotent Borel subring. Let $Z = Z^\circ(\mathfrak{b}) > 0$. First we prove that \mathfrak{b} acts trivially on subquotients of rank 1 (Step 3.1.1), and has rank 1 itself. Then we find a field \mathbb{L} with rank 2, such that \mathfrak{b} embeds properly into \mathbb{L}_+ (Step 3.1.2). Then we derive a contradiction (Step 3.1.3).

Step 3.1.1. Let X be a 1-dimensional subquotient \mathfrak{b} -module. Then \mathfrak{b} centralises X .

Proof. Suppose not. Linearising in dimension 1, there is $h \in \mathfrak{b}$ acting on X like Id_X . Lifting the eigenspace, $E_1 = E_1(h) \leq \mathfrak{g}$ is non-trivial. It is proper since $E_0 \geq Z > 0$.

For each integer $k \neq 0$ one has $E_k \cap \mathfrak{b} = 0$. Indeed, if $a \in E_k \cap \mathfrak{b}$, then $a = \frac{1}{k}[h, a] \in \mathfrak{b}'$, and for every integer n one has $a \in \mathfrak{b}^{[n]}$, the descending nilpotent series. Hence $a = 0$.

We claim $\dim \mathfrak{b} = 1$. Otherwise $\dim \mathfrak{b} = 2$ and $\mathfrak{g} = \mathfrak{b} \oplus E_1$. But then $E_2 = 0$, so E_1 is an abelian subring. Now $Z \leq E_0$ normalises E_1 . If the action is trivial, then $E_1 \leq C_{\mathfrak{g}}^\circ(Z) = \mathfrak{b}$, a contradiction. So the action of Z on E_1 is non-trivial. Hence we may assume $h \in Z$; in that case, $E_0 = \mathfrak{b}$ normalises E_1 , which is an ideal of \mathfrak{g} . This is a contradiction; hence $\dim \mathfrak{b} = 1$.

It follows that \mathfrak{b} is abelian and $E_0 = \mathfrak{b}$. If $E_2 \neq 0$ then $\mathfrak{g} = E_0 \oplus E_1 \oplus E_2$ and $E_2 \triangleleft \mathfrak{g}$; a contradiction. So $E_2 = 0$ and E_1 is an abelian subring. But then $E_0 \oplus E_1$ is a subring properly containing $E_0 = \mathfrak{b}$. By definition of a Borel subring, $\mathfrak{g} = E_0 \oplus E_1$, which contains E_1 as an ideal: a contradiction. \diamond

It follows $\dim \mathfrak{b} = 1$. Otherwise $\dim \mathfrak{b} = 2$ and $\dim \mathfrak{g}/\mathfrak{b} = 1$. By Step 3.1.1, \mathfrak{b} centralises $\mathfrak{g}/\mathfrak{b}$, meaning $\mathfrak{b} \triangleleft \mathfrak{g}$, a contradiction. Hence $\dim \mathfrak{b} = 1$ and in particular \mathfrak{b} is abelian. It follows that for $a \in \mathfrak{b} \setminus \{0\}$ one has $C_a = \mathfrak{b}$.

Let $B_{\mathfrak{b}} = [\mathfrak{b}, \mathfrak{g}]$.

Step 3.1.2. $\mathfrak{g} = \mathfrak{b} + B_{\mathfrak{b}}$. There is a definable field \mathbb{L} of dimension 2 such that $B_{\mathfrak{b}} \simeq \mathbb{L}_+$ and $\mathfrak{b} \hookrightarrow \mathbb{L} \text{Id}_{B_{\mathfrak{b}}}$.

Proof. First, $\dim B_{\mathfrak{b}} = 2$. Indeed, let $a, a' \in \mathfrak{b}$ be non-zero. Clearly $\dim B_a = 2$; by abelianity, B_a is \mathfrak{b} -invariant. By Step 3.1.1, \mathfrak{b} centralises \mathfrak{g}/B_a . In particular $B_{a'} \leq B_a$ and equality follows. So $B_{\mathfrak{b}} = B_a$ for any $a \in \mathfrak{b} \setminus \{0\}$, which proves $\dim B_{\mathfrak{b}} = 2$.

Suppose $\mathfrak{b} \leq B_{\mathfrak{b}}$. Then for $a \in \mathfrak{b} \setminus \{0\}$ one has $C_a = \mathfrak{b} \leq B_{\mathfrak{b}} = B_a$. In particular $B_a^2 < B_a$, and $C_a^2 > C_a$. However, $C_a = \mathfrak{b}$ is abelian, so by Lemma A, C_a^2 is a subring. But $\dim C_a^2 \leq 2 \dim C_a = 2 \dim \mathfrak{b} < \dim \mathfrak{g}$, so C_a^2 is a definable, connected, proper subring containing $C_a = \mathfrak{b}$. Hence $C_a^2 = \mathfrak{b} = C_a$, a contradiction proving $\mathfrak{b} \not\leq B_{\mathfrak{b}}$.

$B_{\mathfrak{b}}$ is \mathfrak{b} -irreducible. Otherwise there is a 1-dimensional \mathfrak{b} -submodule $V_1 < B_{\mathfrak{b}}$. By Step 3.1.1, \mathfrak{b} centralises V_1 , so $V_1 \leq C_{\mathfrak{g}}^\circ(\mathfrak{b}) = \mathfrak{b}$. Always by Step 3.1.1, \mathfrak{b} centralises $B_{\mathfrak{b}}/\mathfrak{b}$, so $B_{\mathfrak{b}} \leq N_{\mathfrak{g}}^\circ(\mathfrak{b}) = \mathfrak{b}$, a contradiction.

The action of \mathfrak{b} on $B_{\mathfrak{b}}$ is faithful. Otherwise there is $a \in \mathfrak{b} \setminus \{0\}$ with $B_{\mathfrak{b}} \leq C_a = \mathfrak{b}$, a contradiction again.

Linearising the abelian action, there is a definable field \mathbb{L} with $B_{\mathfrak{b}} \simeq \mathbb{L}_+$ and $\mathfrak{b} \hookrightarrow \mathbb{L} \text{Id}_{B_{\mathfrak{b}}}$. \diamond

Step 3.1.3. Contradiction.

Proof. Fix any $a_0 \in \mathfrak{b} \setminus \{0\}$ and $x_0 \in B_{\mathfrak{b}} \setminus \{0\}$.

Because the action of \mathfrak{b} on $B_{\mathfrak{b}}$ is by scalars, $\dim[a_0, [\mathfrak{b}, x_0]] = \dim[\mathfrak{b}, x_0] = 1$ (the subgroups need not be equal). We claim that $\dim[\mathfrak{b}, [\mathfrak{b}, x_0]] = 1$. Let $A_2 = [\mathfrak{b}, [\mathfrak{b}, x_0]] \leq B_{\mathfrak{b}}$. By Lemma C, $[A_2, A_2] \leq B_{\mathfrak{b}}$. If $\dim A_2 = 2$ then dimensions match and $A_2 = B_{\mathfrak{b}}$. Now $[B_{\mathfrak{b}}, B_{\mathfrak{b}}] \leq B_{\mathfrak{b}}$, so $B_{\mathfrak{b}}$ is a subring. It also is \mathfrak{b} -invariant, but $\mathfrak{b} \not\leq B_{\mathfrak{b}}$ by Step 3.1.2. So $B_{\mathfrak{b}} \triangleleft \mathfrak{g}$, a contradiction. Hence $\dim A_2 = 1$.

Thus $\dim[a_0, [\mathfrak{b}, x_0]] = \dim[\mathfrak{b}, [\mathfrak{b}, x_0]]$; the groups are equal. So for $a_1, a_2 \in \mathfrak{b}$, there is $a_3 \in \mathfrak{b}$ with $[a_0, [a_3, x_0]] = [a_1, [a_2, x_0]]$. Returning to the field \mathbb{L} , this means $a_0 a_3 = a_1 a_2$ as elements of \mathbb{L} . So the image of \mathfrak{b} in \mathbb{L} is closed under the map $(a_1, a_2) \mapsto \frac{a_1 a_2}{a_0}$.

Let $R = \{\lambda \in \mathbb{L} : \lambda \mathfrak{b} \leq \mathfrak{b}\}$, a definable subring. We just proved $\frac{1}{a_0} \mathfrak{b} \leq R$, so R is infinite. By the Cherlin-Macintyre-Shelah property, every infinite definable domain is an algebraically closed field. So $R = \mathbb{L}$ and $\mathfrak{b} \neq 0$ is an ideal of \mathbb{L} . Hence $\mathfrak{b} = \mathbb{L}$ but dimensions do not match. \diamond

This eliminates nilpotent Borel subrings in a 3-dimensional, simple Lie ring: Proposition 3.1 is proved. (The argument will be extended to the 4-dimensional case in Proposition 4.4.) \square

By Theorem 2, the structure of Borel subrings of \mathfrak{g} is well-understood.

Proposition 3.2. $\mathfrak{g} \simeq \mathfrak{sl}_2(\mathbb{K})$.

Proof. The proof starts by some local analysis, viz. the study of intersections of Borel subrings (Step 3.2.1). Then we find the desired weight space decomposition (Step 3.2.2), which allows final coordinatisation (Step 3.2.3).

Step 3.2.1. If $\mathfrak{b}_1 \neq \mathfrak{b}_2$ are distinct Borel subrings, then $\mathfrak{c} = (\mathfrak{b}_1 \cap \mathfrak{b}_2)^\circ$ is a Cartan subring of both.

Proof. Clearly $\dim \mathfrak{c} = 1$. By Proposition 3.1, \mathfrak{b}_1 and \mathfrak{b}_2 are non-nilpotent; it is enough to prove that \mathfrak{c} is normal in neither. Suppose it is, say in \mathfrak{b}_1 ; then $\mathfrak{c} = \mathfrak{b}'_1 \leq \mathfrak{b}_2$. Considering $N_{\mathfrak{g}}^\circ(\mathfrak{c}) < \mathfrak{g}$, the ring \mathfrak{c} is not normal in \mathfrak{b}_2 , so it is a Cartan subring of \mathfrak{b}_2 .

Let \mathfrak{t} be a Cartan subring of \mathfrak{b}_1 . Let $t \in \mathfrak{t}$ act on $\mathfrak{b}'_1 = \mathfrak{c}$ like the identity. Now let $h \in \mathfrak{c}$ act on \mathfrak{b}'_2 like the identity. Notice that $[t, h] = \mathfrak{b}'_1 = \mathfrak{c}$, and $[h, \mathfrak{b}'_2] = \mathfrak{b}'_2$, so $B_h \geq \mathfrak{c} + \mathfrak{b}'_2 = \mathfrak{b}_2$. Equality follows since $C_h \geq \mathfrak{b}'_1 > 0$.

Now for $f \in \mathfrak{b}'_2$ one has:

$$[t, f] = [t, [h, f]] = [th, f] + [h, tf] = [h, f] + [h, tf].$$

The right-hand is in $B_h = \mathfrak{b}_2$. Therefore $[t, f] \in \mathfrak{b}_2$, and the right-hand is in \mathfrak{b}'_2 . Therefore $[t, f] \in \mathfrak{b}'_2$, implying $[h, tf] = [t, f]$. So there remains $[h, f] = 0$, a contradiction. \diamond

Step 3.2.2. Weight decomposition: there is $h_1 \in \mathfrak{g}$ such that $\mathfrak{g} = E_{-2} \oplus E_0 \oplus E_2$, and $[E_{-2}, E_2] = E_0$. Moreover, for $e \in E_2 \setminus \{0\}$ one has $C_{\mathfrak{g}}(e) = E_2$.

Proof. Let $\mathfrak{b} \sqsubseteq \mathfrak{g}$ be a Borel subring. By Proposition 3.1, it is non-nilpotent. In particular, by Theorem 2, there is a definable field such that $\mathfrak{b}/Z(\mathfrak{b}) \simeq \mathfrak{ga}_1(\mathbb{K})$. Let a temporary $h_1 \in \mathfrak{b}$ act on \mathfrak{b}' like $\text{Id}_{\mathfrak{b}'}$. (It will be rescaled at the end of the proof of the Step.) Of course $E_0 = E_0(h_1) = C_{h_1} > 0$ and $E_1 \geq \mathfrak{b}' > 0$. Let $e \in \mathfrak{b}' \setminus \{0\}$.

We claim that $C_e = \mathfrak{b}'$ and $B_e = \mathfrak{b}$. Clearly $C_e \geq \mathfrak{b}'$. If equality does not hold, then $\dim C_e = 2$. Now C_e is a Borel subring distinct from \mathfrak{b} since $e \notin Z(\mathfrak{b})$. But

$C_e \geq \mathfrak{b}'$ which is not Cartan in \mathfrak{b} , against Step 3.2.1. So $C_e = \mathfrak{b}'$ and $\dim B_e = 2$. Now consider the action of \mathfrak{b} on $\mathfrak{g}/\mathfrak{b}$. Linearising in dimension 1, \mathfrak{b}' acts trivially so $B_e \leq \mathfrak{b}$ and equality holds.

We now claim that $E_{-1} = E_{-1}(h_1) \neq 0$. For suppose $E_{-1} = 0$. The additive morphism $\varphi(x) = [h_1, x] + x$ then has a finite kernel, and is surjective. Therefore:

$$\text{im ad}_{h_1} \text{ad}_e = \text{im ad}_e(1 + \text{ad}_{h_1}) = \text{im ad}_e = B_e = \mathfrak{b}.$$

But $\text{im ad}_{h_1} \text{ad}_e = [h_1, B_e] = [h_1, \mathfrak{b}] < \mathfrak{b}$, a contradiction proving $E_{-1} \neq 0$.

It follows $\mathfrak{g} = E_{-1} \oplus E_0 \oplus E_1$, and $\mathfrak{c} = E_0$ normalises each term. Also, $\mathfrak{b} = E_0 \oplus E_1$ and $\mathfrak{b}' = E_1$. If $[E_{-1}, E_1] = 0$, then $E_{-1} \oplus E_1$ is a subring, hence a Borel subring. However it intersects \mathfrak{b} in $E_1 = \mathfrak{b}'$, against Step 3.2.1. So $[E_{-1}, E_1] = E_0$.

There remains to prove that for any $e \in E_1 \setminus \{0\}$, one has $C_{\mathfrak{g}}(e) = E_1$ (notice the absence of a connected component). Since $\mathfrak{g} = E_{-1} \oplus E_0 \oplus E_1$, it is enough to prove $C_{E_0}(e) = C_{E_{-1}}(e) = 0$.

- Let $h \in E_0$ centralise e ; then it centralises E_0 and $[E_0, e] = E_1$. But also, for $y \in E_{-1}$, one has $[e, y] \in E_0 \leq C_h$, so:

$$[e, [h, y]] = [eh, y] + [h, ey] = 0.$$

Therefore $[h, E_{-1}] \leq C_e = E_1$. However $[h, E_{-1}] \leq E_{-1}$, proving $h \in Z(\mathfrak{g}) = 0$.

- Now let $f \in E_{-1}$ centralise e . Then it normalises $C_e = E_1$ and $N_{\mathfrak{g}}^{\circ}(E_1) = E_0 + E_1$. In particular, $[f, h_1] \in E_0 + E_1$, but $[f, h_1] = -[h_1, f] = f \in E_{-1}$ so $f = 0$.

This shows $C_{\mathfrak{g}}(e) = E_1$.

We finally rescale and replace h_1 by $\frac{1}{2}h_1$. ◇

Step 3.2.3. Coordinatisation.

Proof. Let $\mathfrak{c} = E_0$, $\mathfrak{u}_+ = E_2$, and $\mathfrak{u}_- = E_{-2}$. By Step 3.2.2, there are $e_1 \in \mathfrak{u}_+$ and $f_1 \in \mathfrak{u}_-$ with $[e_1, f_1] = h_1$.

We simultaneously coordinatise \mathfrak{c} and \mathfrak{u}_+ as follows. Linearising in dimension 1, there is a definable field \mathbb{K} such that $\mathfrak{c} \simeq \mathbb{K} \text{Id}$ acts by *doubled* scalars on $\mathfrak{u}_+ \simeq \mathbb{K}_+$, viz. $[h_{\lambda}, e_{\mu}] = 2e_{\lambda\mu}$. Notice that h_1 was defined consistently. Since the multiplicative unit is not fixed, we choose it to be e_1 and consistently write $\mathfrak{u}_+ = \{e_{\lambda} : \lambda \in \mathbb{K}\}$. Now we let $f_{\lambda} = -\frac{1}{2}[h_{\lambda}, f_1]$, so $\mathfrak{u}_- = \{f_{\lambda} : \lambda \in \mathbb{K}\}$. Notice that f_1 was defined consistently.

We prove $\mathfrak{g} \simeq \mathfrak{sl}_2(\mathbb{K})$. It suffices to check a couple of identities. First, by definition, $[h_{\lambda}, e_{\mu}] = \frac{1}{2}[h_{\lambda}, [h_{\mu}, e_1]] = [h_{\mu}, e_{\lambda}]$; and $[h_{\lambda}, f_{\mu}] = [h_{\mu}, f_{\lambda}]$ likewise.

Then $[e_{\mu}, f_1] = h_{\mu}$, because:

$$[[e_{\mu}, f_1], e_1] = [e_{\mu}e_1, f_1] + [e_{\mu}, f_1e_1] = [h_1, e_{\mu}] = 2e_{\mu} = [h_{\mu}, e_1],$$

meaning that $[e_{\mu}, f_1] - h_{\mu} \in C_{\mathfrak{c}}(e_1) = 0$ by Step 3.2.2.

It follows $[e_{\mu}, f_{\nu}] = h_{\mu\nu}$. Indeed,

$$-2[e_{\mu}, f_{\nu}] = [e_{\mu}, [h_{\nu}, f_1]] = [e_{\mu}h_{\nu}, f_1] + [h_{\nu}, e_{\mu}f_1] = -2[e_{\mu\nu}, f_1] = -2h_{\mu\nu}.$$

Last, $[h_{\mu}, f_{\nu}] = -2f_{\mu\nu}$ since:

$$[[h_{\mu}, f_{\nu}], e_1] = [h_{\mu}e_1, f_{\nu}] + [h_{\mu}, f_{\nu}e_1] = 2[e_{\mu}, f_{\nu}] = 2h_{\mu\nu} = -2[f_{\mu\nu}, e_1],$$

implying $[h_{\mu}, f_{\nu}] + 2f_{\mu\nu} \in C_{\mathfrak{u}_-}(e_1) = 0$ by Step 3.2.2 again.

The isomorphism $\mathfrak{g} \simeq \mathfrak{sl}_2(\mathbb{K})$ is now obvious. ◇

This completes the recognition and proves Theorem 3. \square

The extension problem is easy, although not considered by Rosengarten.

Corollary 3. *Let \mathfrak{g} be a connected, non-soluble Lie ring of Morley rank 3 and characteristic $\neq 2, 3$. Then $\mathfrak{g} \simeq \mathfrak{sl}_2(\mathbb{K})$ for some definable field \mathbb{K} .*

Proof. It is enough to show simplicity of \mathfrak{g} , which we shall derive from the isomorphism type of $\mathfrak{g}/Z(\mathfrak{g})$. Let $Z = Z(\mathfrak{g})$, a definable ideal of \mathfrak{g} , and $\bar{\mathfrak{g}} = \mathfrak{g}/Z$.

We first prove that $\bar{\mathfrak{g}}$ is simple. It is enough to prove that it is definable simple (see § 2.2). Let $I \triangleleft \mathfrak{g}$ be any proper definable ideal; then $I^\circ \triangleleft \mathfrak{g}$ as well. If $I^\circ > 0$ then by Theorem 2 both I° and \mathfrak{g}/I° are soluble, against the assumption. Therefore I is finite. By connectedness, $[I, \mathfrak{g}] = 0$ so I is central. This shows that the largest proper definable ideal of \mathfrak{g} is Z . In particular, $\bar{\mathfrak{g}}$ is definably simple; being non-abelian, it is simple.

We now conclude $Z = 0$. By Theorem 3, $\bar{\mathfrak{g}} \simeq \mathfrak{sl}_2(\mathbb{K})$ for some definable field \mathbb{K} . So there is $\eta \in \bar{\mathfrak{g}}$ such that $\bar{\mathfrak{g}} = E_{-1}(\eta) \oplus E_0(\eta) \oplus E_1(\eta)$. Lift η to $h \in \mathfrak{g}$; lifting eigenspaces, one has $\mathfrak{g} = E_{-1}(h) \oplus E_0(h) \oplus E_1(h)$, and each has dimension 1. It now follows that $Z \leq E_0(h)$. Moreover, $[E_{-1}(h), E_1(h)] = E_0(h)$. Let $z \in Z$. Then there are $a_{-1} \in E_{-1}$ and $a_1 \in E_1$ with $z = [a_{-1}, a_1]$. Let $\alpha_i = (a_i \bmod Z) \in E_i(\eta)$. Then $[\alpha_{-1}, \alpha_1] = 0$ in $\mathfrak{sl}_2(\mathbb{K})$, so at least one α_i is zero. Thus $a_i \in E_i \cap Z = 0$ and $z = 0$. This proves $Z = 0$ and $\mathfrak{g} \simeq \bar{\mathfrak{g}}$. \square

3.4 Dimension 4

Theorem 4. *There is no simple Lie ring of Morley rank 4 and characteristic $\neq 2, 3$.*

The main idea is elementary: use Borel subrings as flintstones. We need at least two, and at least one of them should be large enough. It is unreasonable to have more upper-triangular matrices than lower-triangular ones, and this is the contradictory picture. For this we need two Borel subrings of distinct dimensions intersecting over a 1-dimensional, ‘diagonal’ subring, creating friction between unbalanced dimensions.

So we shall produce a 3-dimensional Borel subring \mathfrak{b}_1 , and another Borel subring \mathfrak{b}_2 . With rudimentary proxies for ‘a toral subring’ and ‘the nilpotent radical’, \mathfrak{b}_1 and \mathfrak{b}_2 will intersect in a 1-dimensional, toral subring \mathfrak{c} , and have disjoint nilpotent radicals $I_1 \neq I_2$; also we shall see $\dim I_1 = 2$. Then $[I_1, I_2]$ will be too large to fit into \mathfrak{c} . This creates centralisation phenomena between nilpotent radicals in a tight configuration, where nilpotence is not for sharing: a contradiction.

Of course the above strategy has to be implemented without weight theory, since there is no linear structure to begin with. So we are left with basic means inspired by the group-theoretic analysis of $\mathrm{PGL}_2(\mathbb{K})$ in terms of Borel subgroups and unipotent radicals [CJ04; Del07; DJ16]. In short the proof does *not* proceed by picking a Cartan subring and studying the root decomposition. To do this one would need a number of tools yet dubious; see final questions in § 4.

The proof starts. Let \mathfrak{g} be a simple Lie ring of Morley rank 4. There are three main stages. We first prove minimal simplicity of \mathfrak{g} in Proposition 4.1. We derive that \mathfrak{g} cannot be a sum of integral weight spaces in Proposition 4.2; Proposition 4.3 is an important consequence. Changing topic, we remove nilpotent Borel subrings in Proposition 4.4; Proposition 4.5 then allows us to find a 3-dimensional Borel subring,

whose fine structure is described in Proposition 4.6. Proposition 4.7 gives the final contradiction.

Proposition 4.1. \mathfrak{g} is minimal simple, viz. every definable, connected, proper subring is soluble.

Proof. Let $\mathfrak{h} \sqsubset \mathfrak{g}$ be a definable, connected, proper subring. By Theorem 2, we may assume $\dim \mathfrak{h} = 3$. Let $I = C_{\mathfrak{h}}(\mathfrak{g}/\mathfrak{h})$, an ideal of \mathfrak{h} . Notice $I < \mathfrak{h}$ since otherwise $\mathfrak{h} \triangleleft \mathfrak{g}$, a contradiction. In particular I° is soluble by Theorem 2. Linearising in dimension 1 forces abelianity of \mathfrak{h}/I . Thus \mathfrak{h} is soluble. \square

Much later, Proposition 4.6 will return to the analysis of 3-dimensional subrings.

Lack of weights. We shall now forbid \mathfrak{g} to be a sum of integral weight spaces. While mostly trivial, the proof needs attention as we also allow for characteristic 5. There, dimension 4 comes extremely close to ‘Witt-type’ root distribution and requires some care. Moreover, the final case of two Borel subrings intersecting over a 2-dimensional Cartan subring (Step 4.2.3) is a key configuration to eliminate.

Proposition 4.2. Let $h \neq 0$. Then \mathfrak{g} is not a sum of eigenspaces, viz. $\sum_{i \in \mathbb{F}_p} E_i(h) < \mathfrak{g}$.

Proof. Unless otherwise indicated, all eigenspaces in this proof are with respect to h . Let p be the characteristic; by assumption $p \neq 2, 3$, but 5 is allowed.

Suppose $\sum_{i \in \mathbb{F}_p} E_i = \mathfrak{g}$. Bear in mind that the sum is direct.

Step 4.2.1. One has $h \in E_0$. Allowing trivial terms, we may suppose $\mathfrak{g} = E_{-1} \oplus E_0 \oplus E_1 \oplus E_k$ for some $k \notin \{-1, 0, 1\}$.

Proof. A priori, and up to allowing trivial terms, $\mathfrak{g} = E_i \oplus E_j \oplus E_k \oplus E_\ell$ for i, j, k, ℓ distinct integers modulo p .

We claim $h \in E_0 \neq 0$ and $\mathfrak{g} = E_0 \oplus E_i \oplus E_j \oplus E_k$. Indeed, write $h = e_i + e_j + e_k + e_\ell$ in obvious notation; we may assume $e_\ell \neq 0$. Applying ad_h , we find $ie_i + je_j + ke_k + le_\ell = 0$. In particular, $le_\ell = 0$ and therefore $\ell = 0$. Since the integers were distinct, $e_i = e_j = e_k = 0$ and $h = e_\ell \in E_\ell = E_0$.

We may assume E_i, E_j, E_k all non-trivial and proper. Properness is since $h \in E_0$. Now if $E_i = 0$, then let $h' = \frac{1}{j}h$. Notice how $E_\ell(h') = E_{j\ell}(h)$, so $\mathfrak{g} = E_0(h') \oplus E_1(h') \oplus E_{j^{-1}k}(h')$. Up to considering h' instead of h (which we call *up to dividing*) we have the desired form.

We may assume $i + j, i + k, j + k$ all non-zero. Indeed if $j = -i$, then up to dividing we have the desired form.

We may assume $i + j \neq k$ and $i + k \neq j$. Indeed if $i + j - k = i + k - j = 0$, then $2i = 0$ and $i = 0$; a contradiction. So at most one the three equalities of type $i + j = k$ holds.

We may assume $2i \in \{j, k\}$. Indeed since $i + j \neq 0$ and $i + j \neq k$, we have $i + j \notin \{0, i, j, k\}$, so $[E_i, E_j] \leq E_{i+j} = 0$. By our assumptions, $[E_i, E_k] = 0$ likewise. So $0 < E_i < \mathfrak{g}$ is normalised by $E_0 + E_j + E_k$. Since E_i is no ideal of \mathfrak{g} , we therefore have $[E_i, E_i] \neq 0$ and $2i \in \{0, i, j, k\}$. But $2i \notin \{0, i\}$ since $i \neq 0$.

We have $2j \in \{i, k\}$ and $2k \in \{i, j\}$. For suppose $j + k = i$. With $2i = j$ this gives $i + k = 0$, a contradiction; the case $2i = k$ is similarly excluded. So $j + k \neq i$, and the previous paragraph also gives $2j \in \{i, k\}$ and $2k \in \{i, j\}$.

Conclusion. We may assume $2i = j$; the other case is similar. If $2j = i$ then $4 = 1$, against $p \neq 3$. So $2j = k$. Similarly, $2k = i$. Therefore $8 = 1$ and $p = 7$. Up to dividing for readability, we now have $\mathfrak{g} = E_0 \oplus E_1 \oplus E_2 \oplus E_4$. But now $E_3 = E_5 = E_6 = 0$ and $E_8 = E_1$, so $E_1 \oplus E_2 \oplus E_4 \triangleleft \mathfrak{g}$, a contradiction. \diamond

Step 4.2.2. We may assume $\mathfrak{g} = E_{-1} \oplus E_0 \oplus E_1$. Moreover E_{-1}, E_0, E_1 are non-trivial and proper.

Proof. By Step 4.2.1 we may assume $\mathfrak{g} = E_{-1} \oplus E_0 \oplus E_1 \oplus E_k$ for some $k \notin \{-1, 0, 1\}$.

We first show that we may assume $k = 2$. If $k = -2$, then up to considering $-h$, we have the desired form. Therefore suppose $k \neq \pm 2$, so that $k \notin \{0, \pm 1, \pm 2\}$. In particular $E_{-2} = E_2 = 0$, implying that E_{-1} and E_1 are abelian subrings. Let $\mathfrak{h} = E_{-1} + E_0 + E_1$, a definable, connected subring of \mathfrak{g} . If $\mathfrak{h} = \mathfrak{g}$, then up to taking $k = 2$ as a dummy we have the desired form. So we may assume $\mathfrak{h} < \mathfrak{g}$. By minimal simplicity (Proposition 4.1), \mathfrak{h} is soluble. By Step 4.2.1, $h \in E_0 \leq \mathfrak{h}$, so $E_1 = [h, E_1] \leq \mathfrak{h}'$, and $E_{-1} \leq \mathfrak{h}'$ likewise. By solubility, $\mathfrak{h}' < \mathfrak{h}$, implying $[E_{-1}, E_1] < E_0$. If $\dim E_0 = 2$ then since $\mathfrak{h} < \mathfrak{g}$ is proper, at least one of E_{-1}, E_1 is trivial; hence $[E_{-1}, E_1] = 0$. If $\dim E_0 = 1$ then $[E_{-1}, E_1] = 0$. So $[E_{-1}, E_1] = 0$ in either case. Moreover $k \pm 1 \notin \{-1, 0, 1, k\}$, so E_{-1} and E_1 are ideals of \mathfrak{g} . Since $E_0 \neq 0$ they are proper, hence trivial. There remains $\mathfrak{g} = E_0 \oplus E_k$ and E_k is a proper ideal of \mathfrak{g} , hence trivial. Finally $\mathfrak{g} = E_0$, so $h \in Z(\mathfrak{g}) = 0$, a contradiction. So we may assume $k = 2$. Hence $\mathfrak{g} = E_{-1} \oplus E_0 \oplus E_1 \oplus E_2$.

We prove that E_{-1}, E_0, E_1 are non-trivial and proper.

- We already know $0 < E_0 < \mathfrak{g}$.
- If $E_{-1} = 0$ then (even when $p = 5$) E_2 is an ideal of \mathfrak{g} ; hence trivial. Then E_1 is an ideal of \mathfrak{g} ; hence trivial as well, and this is a contradiction.
- If $E_1 = 0$ then $[E_{-1}, E_2] = 0$ so (even when $p = 5$) E_{-1} is an ideal of \mathfrak{g} ; hence trivial, a contradiction.

It remains to show that $E_2 = 0$. Suppose not. Then $\mathfrak{g} = E_{-1} \oplus E_0 \oplus E_1$, and each has dimension 1.

We claim $[E_{-1}, E_1] = E_0$, $[E_1, E_1] = E_2$, and $[E_{-1}, E_2] = E_1$.

- If $[E_{-1}, E_1] = 0$ then (even when $p = 5$) $E_{-1} + E_1 + E_2$ is an ideal of \mathfrak{g} , a contradiction.
- If $[E_1, E_1] = 0$, then $\mathfrak{h} = E_{-1} + E_0 + E_1$ is a proper subring, hence soluble by Proposition 4.1. But $h \in E_0$ so $\mathfrak{h}' \geq E_{-1} + E_1 + [E_{-1}, E_1] = \mathfrak{h}$, a contradiction.
- Suppose $[E_{-1}, E_2] = 0$. Write $h \in E_0 = [E_{-1}, E_1]$ as $h = [e_{-1}, e_1]$; let $e_2 \in E_2$. Then:

$$2e_2 = [h, e_2] = [[e_{-1}, e_1], e_2] = [e_{-1}e_2, e_1] + [e_{-1}, e_1e_2] = 0,$$

a contradiction. (If the characteristic is not 5 there is a shorter argument: E_2 is an ideal of \mathfrak{g} , a contradiction.)

We claim that for any $x_0 \in E_0$ there is $y_2 \in E_2 \setminus \{0\}$ such that $[x_0, y_2] = 0$. Let $e_{-1} \in E_{-1}$ be such that $[e_{-1}, E_1] = E_0$ and $[e_{-1}, E_2] = E_1$; there is one such since otherwise $C_{E_{-1}}(E_1) \cup C_{E_{-1}}(E_2) = E_{-1}$, a contradiction. Let $x_0 \in E_0$. Then there is

$y_1 \in E_1$ with $[y_1, e_{-1}] = x_0$. Now there is $y_2 \in E_2$ with $[e_{-1}, y_2] = y_1$. On the other hand, $[E_1, E_2] = 0$. Altogether, one has:

$$\begin{aligned} [x_0, y_2] &= [[y_1, e_{-1}], y_2] \\ &= [y_1 y_2, e_{-1}] + [y_1, e_{-1} y_2] \\ &= [0, e_{-1}] + [y_1, y_1] \\ &= 0. \end{aligned}$$

Conclusion. Since $h \in E_0$, the action of E_0 on E_2 is non-trivial. Linearising in dimension 1, there is a finite ideal $F \triangleleft E_0$ such that E_0/F acts by scalars, hence freely, on E_2 . This contradicts the last claim. \diamond

Step 4.2.3. Contradiction.

Proof. By Step 4.2.2, exactly one of E_{-1}, E_0, E_1 has dimension 2; moreover $E_{\pm 1}$ are abelian. For the end of the argument, bear in mind that E_{-1} does not normalise E_1 nor vice-versa, since the normalised one would be an ideal of \mathfrak{g} . There are essentially two cases.

- Suppose $\dim E_1 = 2$. Let $a \in E_{-1}$. Since ad_a takes E_1 to E_0 , the subgroup $X_a = C_{E_1}^\circ(a)$ is infinite; since E_1 is abelian, X_a is actually a subring. Notice that both a and h normalise X_a . Moreover $C_{\mathfrak{g}}^\circ(X_a) \geq E_1$. If $C_{\mathfrak{g}}^\circ(X_a) > E_1$, then it is a 3-dimensional Borel subring. In that case, by Lemma F, $h \in C_{\mathfrak{g}}^\circ(X_a)$, a contradiction as the action of h on X_a is non-trivial. Hence $C_{\mathfrak{g}}^\circ(X_a) = E_1$ is normalised by a . This shows that E_{-1} normalises E_1 , a contradiction.

This rules out $\dim E_{-1} = 2$ as well, by considering $-h$.

- Suppose $\dim E_0 = 2$. (It is unclear whether E_0 is abelian.) Since E_0 contains h , it is trivial neither on E_1 nor on E_{-1} . Linearising in dimension 1, both $\mathfrak{t}_1 = C_{E_0}^\circ(E_1)$ and $\mathfrak{t}_{-1} = C_{E_0}^\circ(E_{-1})$ are subrings of dimension 1. In particular they are abelian.

We show $\mathfrak{t}_1 \neq \mathfrak{t}_{-1}$. Otherwise denote it by $\mathfrak{t}_{\pm 1}$ and let $\mathfrak{h} = C_{\mathfrak{g}}^\circ(\mathfrak{t}_{\pm 1})$, a proper subring of \mathfrak{g} . By assumption, $\mathfrak{h} \geq E_{-1} + E_1 + \mathfrak{t}_{\pm 1}$; so equality holds. In particular $[E_{-1}, E_1] \leq (\mathfrak{h} \cap E_0)^\circ = \mathfrak{t}_{\pm 1}$. Now h normalises E_{-1}, E_1 , and their bracket $\mathfrak{t}_{\pm 1}$; so h normalises \mathfrak{h} . By Lemma F, one gets $h \in \mathfrak{h} = E_{-1} \oplus E_0^{\mathfrak{h}} \oplus E_1$, and finally $h \in E_0^{\mathfrak{h}} = \mathfrak{t}_{\pm 1}$, a contradiction.

Let $\mathfrak{c} = [E_{-1}, E_1] \neq 0$. Since E_0 normalises \mathfrak{c} , this is a subring. We prove $\mathfrak{t}_1 \not\leq \mathfrak{c}$; suppose inclusion holds. Then $\mathfrak{t}_1 \neq \mathfrak{t}_{-1}$ acts non-trivially on E_{-1} , so linearising in dimension 1, there is $t_1 \in \mathfrak{t}_1$ acting on E_{-1} as 1; by definition it acts trivially on E_1 . Then by assumption, $t_1 \in \mathfrak{c} = [E_{-1}, E_1]$. At most two elements $e_{-1}, e'_{-1} \in E_{-1}$ suffice to write $\mathfrak{c} = \text{ad}_{e_{-1}}(E_1) + \text{ad}_{e'_{-1}}(E_1)$. So there are $e_1, e'_1 \in E_1$ with $t_1 = [e_{-1}, e_1] + [e'_{-1}, e'_1]$. But then:

$$\begin{aligned} 0 &= [t_1, t_1] = [t_1, [e_{-1}, e_1]] + [t_1, [e'_{-1}, e'_1]] \\ &= [t_1 e_{-1}, e_1] + [e_{-1}, t_1 e_1] + [t_1 e'_{-1}, e'_1] + [e'_{-1}, t_1 e'_1] \\ &= [e_{-1}, e_1] + [e'_{-1}, e'_1] = t_1, \end{aligned}$$

a contradiction. Notice $\mathfrak{t}_{-1} \not\leq \mathfrak{c}$ for the same reasons.

Conclusion. This implies $\dim \mathfrak{c} = 1$. Indeed, $\dim \mathfrak{c} = 2$ would contradict $\mathfrak{t}_1 \not\leq \mathfrak{c}$. Finally let $\mathfrak{h} = E_{-1} + \mathfrak{c} + E_1$, a subring. If \mathfrak{c} centralises E_1 , then $\mathfrak{c} = \mathfrak{t}_1$, a contradiction. For the same reason, \mathfrak{c} does not centralise E_{-1} . So $\mathfrak{h}' = \mathfrak{h}$, against Proposition 4.1. \diamond

This proves Proposition 4.2. \square

As a consequence we derive a severe control on sizes of weight spaces.

Proposition 4.3. *Suppose $h \in \mathfrak{g}$ is such that $E_0(h) \neq 0$ and $E_1(h) \neq 0$. Then $\dim E_0(h) = \dim E_1(h) = 1$ and $E_2(h) = 0$.*

Proof. All eigenspaces in this proof are with respect to h .

Let $\mathfrak{b} = E_0 + E_1 + E_2$, a priori a definable, connected subgroup. By Proposition 4.2, $E_3 = E_4 = 0$, so \mathfrak{b} is actually a proper subring. We must prove $\dim \mathfrak{b} = 2$ and suppose not; then \mathfrak{b} is a 3-dimensional Borel subring. By Lemma F, it is self-normalising in \mathfrak{g} . Therefore $h \in N_{\mathfrak{g}}(\mathfrak{b}) = \mathfrak{b}$.

Since $h \in \mathfrak{b}$, for $i \in \{1, 2\}$ one has $E_i = [h, E_i] \leq \mathfrak{b}'$. It follows $\mathfrak{b}' = E_1 + E_2 + E_0^{\mathfrak{b}'}$, where each of the last two terms is allowed to be trivial. Then $[h, \mathfrak{b}'] \leq E_1 + E_2$.

Consider the action of \mathfrak{b} on $\mathfrak{g}/\mathfrak{b}$. Linearising in dimension 1, \mathfrak{b}' centralises $\mathfrak{g}/\mathfrak{b}$. So for $i \in \{1, 2\}$ and $e \in E_i$ one has $B_e \leq \mathfrak{b}$. By Proposition 4.2 again, $E_{-i} = 0$, so $\text{ad}_h + i$ is surjective and:

$$[h, B_e] = \text{im ad}_h \text{ad}_e = \text{im ad}_e(\text{ad}_h + i) = \text{im ad}_e = B_e.$$

Since $B_e \leq \mathfrak{b}$ and $h \in \mathfrak{b}$ one has $B_e = [h, B_e] \leq \mathfrak{b}'$. Now $B_e = [h, B_e] \leq [h, \mathfrak{b}'] \leq E_1 + E_2$. Summing over $e \in E_i$ and $i \in \{1, 2\}$, for $a \in E_1 + E_2$ one has $B_a \leq E_1 + E_2$. Hence $E_1 + E_2 \triangleleft \mathfrak{g}$. But $E_0 \neq 0$ and $E_1 \neq 0$: a contradiction to simplicity. \square

A first analysis of Borel subrings. We now change thread and study abstract Borel subrings.

Proposition 4.4 (see Proposition 3.1). *No Borel subring is nilpotent.*

Remark. Some of the ideas in the following proof could be pushed further. One suspects that if \mathfrak{g} is minimal simple, then no Borel subring has dimension 1.

Proof. Towards a contradiction, let \mathfrak{b} be one such. Let $Z = Z^\circ(\mathfrak{b}) > 0$. It is a possibility that $Z = \mathfrak{b}$. For $z \in Z \setminus \{0\}$, one has $\mathfrak{b} \subseteq C_z$; by minimal simplicity, equality holds.

Step 4.4.1 (see Step 3.1.1). Let X be a 1-dimensional subquotient \mathfrak{b} -module. Then \mathfrak{b} centralises X . In particular, $\dim \mathfrak{b} \leq 2$.

Proof. The former claim implies the latter. Indeed, if $\dim \mathfrak{b} = 3$ then \mathfrak{b} acts trivially on $\mathfrak{g}/\mathfrak{b}$, so $[\mathfrak{b}, \mathfrak{g}] \leq \mathfrak{b}$ and \mathfrak{b} is an ideal of \mathfrak{g} , a contradiction. So we deal with the former claim.

We first prove that if Y is a 1-dimensional subquotient Z -module, then Z centralises Y . (Both the conclusion and assumption are weaker.) Suppose not. Linearising in dimension 1, there is $z \in Z$ acting on Y like Id_Y . In the present paragraph, eigenspaces are with respect to z . Lifting the eigenspace, $E_1 \leq \mathfrak{g}$ is non-trivial. Since $p \geq 5$, the multiplicative order of $2 \in \mathbb{F}_p^\times$ is at least 3; on the other hand $E_0 > 0$. So there is $\ell \in \mathbb{F}_p^\times$ such that $E_\ell \neq 0$ but $E_{2\ell} = 0$. Then $E_0 + E_\ell$ is a subring properly

containing $E_0 = C_z = \mathfrak{b}$. It follows $\mathfrak{g} = E_0 + E_\ell$ and $E_\ell \triangleleft \mathfrak{g}$, a contradiction. So the claim about 1-dimensional Z -subquotient modules holds. (We lost z .)

To prove our full claim, we may suppose that \mathfrak{b} is non-abelian, implying $\dim \mathfrak{b} \geq 2$, and that \mathfrak{b} does not act trivially on X . Linearising in dimension 1, there is $h \in \mathfrak{b}$ acting on X as the identity. *From now on, all eigenspaces are with respect to h .* Lifting the eigenspace, there is $h \in \mathfrak{b}$ with $E_1 \leq \mathfrak{g}$ non-trivial. We aim for a contradiction.

We claim that $\mathfrak{g} = \mathfrak{b} \oplus E_1$ where $\dim \mathfrak{b} = \dim E_1 = 2$, and $\mathfrak{g}/\mathfrak{b} \simeq E_1$ as Z -modules. (Not ‘as \mathfrak{b} -modules’, since \mathfrak{b} need not normalise E_1 .) First, $\mathfrak{b} \cap E_1 = 0$. This is because $\mathfrak{b} \cap E_1$ is contained in each term of the descending nilpotence series, which vanishes by nilpotence of \mathfrak{b} . Second, $\dim E_1 \geq 2$. Otherwise $\dim E_1 = 1$ and the action of Z on E_1 is trivial, so $E_1 \leq C_{\mathfrak{g}}^{\circ}(Z) = \mathfrak{b}$, a contradiction. In particular, $\dim \mathfrak{b} = \dim E_1 = 2$. Now $\mathfrak{g} = \mathfrak{b} \oplus E_1$ and $E_1 \simeq \mathfrak{g}/\mathfrak{b}$ as Z -modules.

We show that Z acts freely and irreducibly on $\mathfrak{g}/\mathfrak{b}$. If E_1 is not Z -irreducible, there is a 1-dimensional Z -module $Y < E_1$. Now Z acts trivially on Y so $Y \leq C_{\mathfrak{g}}^{\circ}(Z) = \mathfrak{b}$ and $Y \leq E_1 \cap \mathfrak{b}$, a contradiction again. Hence E_1 is Z -irreducible. If some $z \in Z \setminus \{0\}$ acts trivially then $E_1 \leq C_z = \mathfrak{b}$, a contradiction. Linearising the abelian action, there is a field \mathbb{K} with $E_1 \simeq \mathbb{K}_+$ and $Z \curvearrowright \mathbb{K} \text{Id}_{E_1}$. Therefore Z acts freely on $\mathfrak{g}/\mathfrak{b} \simeq E_1$.

We finish the proof by turning to the action of \mathfrak{b} on $\mathfrak{g}/\mathfrak{b}$. Let $I = C_{\mathfrak{b}}(\mathfrak{g}/\mathfrak{b})$. Since Z acts freely on $\mathfrak{g}/\mathfrak{b}$, one has $I \cap Z = 0$. In particular $\dim I < 2$. If $\dim I = 1$ then $\mathfrak{b} = I + Z$ is a sum of two abelian ideals, hence abelian itself: a contradiction. Therefore $\dim I = 0$. Notice that $\mathfrak{g}/\mathfrak{b}$ is \mathfrak{b} -irreducible as it was Z -irreducible. Now ZI/I is infinite, and can be used to linearise the action of \mathfrak{b}/I (see § 2.2). Hence \mathfrak{b}/I is a linear nilpotent Lie ring, and $\mathfrak{g}/\mathfrak{b}$ is \mathfrak{b} -irreducible. This forces the linear dimension to be 1, and therefore \mathfrak{b}/I is abelian. But then \mathfrak{b}' is finite and connected: \mathfrak{b} is abelian, a contradiction. \diamond

Let $W = \mathfrak{g}/\mathfrak{b}$, a \mathfrak{b} -module, and $I = C_{\mathfrak{b}}(W)$, an ideal of \mathfrak{b} . The proof of Proposition 3.1 cannot be followed literally but some ideas will look familiar.

Step 4.4.2. $\dim \mathfrak{b} = 1$; in particular, \mathfrak{b} is abelian.

Proof. Suppose $\dim \mathfrak{b} = 2$.

We claim that W is \mathfrak{b} -irreducible. Otherwise there is a 3-dimensional \mathfrak{b} -module V_3 with $\mathfrak{b} < V_3 < \mathfrak{g}$. By Step 4.4.1, \mathfrak{b} acts trivially on V_3/\mathfrak{b} , so $V_3 \leq N_{\mathfrak{g}}^{\circ}(\mathfrak{b})$: a contradiction.

Now we show $\dim I = 1$ and $I^{\circ} \leq Z$. The former implies the latter. Indeed if \mathfrak{b} is abelian, we are done; while if \mathfrak{b} is not, then its only 1-dimensional, connected ideal is Z , so $I^{\circ} = Z$. So we focus on proving $\dim I = 1$. If $\dim I = 2$ then \mathfrak{b} is an ideal of \mathfrak{g} : a contradiction. We suppose $\dim I = 0$ and give a contradiction. First, \mathfrak{b}/I can be linearised thanks to $ZI/I > 0$. Since \mathfrak{b} is nilpotent and W is \mathfrak{b} -irreducible, this forces the linear dimension to be 1 and \mathfrak{b} to be abelian. Moreover there is a field structure with $\mathfrak{b}/I \simeq \mathbb{K} \text{Id}_W$ and $W \simeq \mathbb{K}_+$. In particular there is $h \in \mathfrak{b}$ acting on $\mathfrak{g}/\mathfrak{b}$ as the identity. We lift the eigenspace; thus $E_1 > 0$, but also $E_0 = \mathfrak{b}$ by abelianity. We prove $\dim E_1 \geq 2$ as follows (covering properties are unclear to us; $\dim E_1^{\mathfrak{g}/\mathfrak{b}}(h) = 2$ does not seem to be a sufficient argument). Let $z \in Z \setminus \{0\}$; for $x \in \mathfrak{g}$ there is $b \in \mathfrak{b}$ such that $[h, x] = x + b$, so:

$$[h, [z, x]] = [hz, x] + [z, hx] = [z, x + b] = [z, x].$$

Therefore $\text{ad}_z : \mathfrak{g} \rightarrow E_1$. However $\ker^\circ \text{ad}_z = C_z = \mathfrak{b}$, so $\dim E_1 \geq \dim \mathfrak{g} - \dim \mathfrak{b} = 2$, as wished. Equality follows as $\mathfrak{b} \cap E_1 = 0$ by abelianity. So $\mathfrak{g} = \mathfrak{b} \oplus E_1$ and $\mathfrak{b} = E_0$ normalises E_1 , an ideal of \mathfrak{g} ; this is a contradiction. The claim is proved.

It follows $N_{\mathfrak{g}}(\mathfrak{b}) = \mathfrak{b}$. Indeed, \mathfrak{b}/I is abelian so we may linearise the action of \mathfrak{b}/I on $\mathfrak{g}/\mathfrak{b}$: it is by scalars, hence free. Now if $n \in N_{\mathfrak{g}}(\mathfrak{b})$ and $b \in \mathfrak{b} \setminus I$, then $[b, n] \in \mathfrak{b}$ implies $n \in \mathfrak{b}$. (This is the argument of Lemma F, which we cannot invoke as stated.)

We derive a contradiction. Let $z \in I^\circ \leq Z$ be non-trivial. Since $C_z = \mathfrak{b}$, one has $\dim B_z = 2$. By definition of I , one also has $B_z \leq \mathfrak{b}$, so equality follows. Hence there is $x \in \mathfrak{g}$ with $z = [x, z]$. For arbitrary $b \in \mathfrak{b}$ one has:

$$[[b, x], z] = [bz, x] + [b, xz] = [b, z] = 0,$$

so $[b, x] \leq C_z = \mathfrak{b}$. Hence $x \in N_{\mathfrak{g}}(\mathfrak{b}) = \mathfrak{b}$, and $z = [x, z] = 0$, a contradiction. \diamond

Step 4.4.3 (see Step 3.1.3). Contradiction.

Proof. Throughout, h will stand for an arbitrary non-zero element of \mathfrak{b} . Let $B_{\mathfrak{b}} = [\mathfrak{b}, \mathfrak{g}]$, a \mathfrak{b} -module containing each B_h for $h \in \mathfrak{b} \setminus \{0\}$.

We first show that $B_h = B_{\mathfrak{b}}$ and $\dim B_h = 3$. Indeed, by abelianity, B_h is \mathfrak{b} -invariant; now $C_h = \mathfrak{b}$, so $\dim(\mathfrak{g}/B_h) = 1$. By Step 4.4.1, \mathfrak{b} acts trivially on \mathfrak{g}/B_h . So $[\mathfrak{b}, \mathfrak{g}] \leq B_h$. Hence for any $h' \in \mathfrak{b}$ one has $B_{h'} \leq B_h$ and equality follows. Thus B_h does not depend on $h \in \mathfrak{b} \setminus \{0\}$; it equals $B_{\mathfrak{b}}$.

We claim that $C_h^2 = C_h = \mathfrak{b}$ and $B_h^2 = B_{\mathfrak{b}}$. Indeed $C_h = \mathfrak{b}$ is abelian, so by Lemma A, C_h^2 is a subring of \mathfrak{g} . But $\dim C_h^2 \leq 2 \dim C_h < \dim \mathfrak{g}$, so C_h^2 is a proper subring containing $C_h = \mathfrak{b}$. Hence $C_h^2 = \mathfrak{b} = C_h$ and therefore $B_h^2 = B_{\mathfrak{b}} = \mathfrak{b}$.

We deduce that $\mathfrak{g} = \mathfrak{b} + B_{\mathfrak{b}}$ and the action of \mathfrak{b} on $B_{\mathfrak{b}}$ is faithful. Otherwise $\mathfrak{b} \leq B_{\mathfrak{b}}$; hence $C_h \leq B_h$, which forces $B_h^2 < B_h$, a contradiction. Moreover, if $h \in \mathfrak{b}$ centralises $B_{\mathfrak{b}}$, it also centralises $\mathfrak{b} + B_{\mathfrak{b}} = \mathfrak{g}$ and is therefore zero.

We now contend that $B_{\mathfrak{b}}$ is \mathfrak{b} -irreducible. Suppose it is not. If there is a 1-dimensional \mathfrak{b} -submodule $V_1 < B_{\mathfrak{b}}$, then by Step 4.4.1 one has $V_1 \leq C_{\mathfrak{g}}^\circ(\mathfrak{b}) = \mathfrak{b}$, so $\mathfrak{b} = V_1 \leq B_{\mathfrak{b}}$, a contradiction. If there is a 2-dimensional \mathfrak{b} -submodule $V_2 < B_{\mathfrak{b}}$, then by Step 4.4.1 again, one has $[b, B_{\mathfrak{b}}] \leq V_2$, so $B_{\mathfrak{b}}^2 < B_{\mathfrak{b}}$, a contradiction.

We may thus linearise the action of \mathfrak{b} on $B_{\mathfrak{b}}$; there is a field \mathbb{K} such that $B_{\mathfrak{b}} \simeq \mathbb{K}_+$ and \mathfrak{b} acts by scalars on $B_{\mathfrak{b}}$. Let us fix some notation: there are $\varphi : B_{\mathfrak{b}} \simeq \mathbb{K}_+$ and $\chi : \mathfrak{b} \rightarrow \mathbb{K}_+$ such that, for $h \in \mathfrak{b}$ and $x \in B_{\mathfrak{b}}$:

$$\varphi([h, x]) = \chi(h) \cdot \varphi(x).$$

Fix arbitrary $x \in B_{\mathfrak{b}}$ and let $A_n = [\mathfrak{b}, \dots, [b, x] \dots]$. Also let $A'_n = \varphi(A_n)$. These definable, connected subgroups of \mathbb{K}_+ need not form an ascending series but their dimensions do. Let n be such that $\dim A'_n = \dim A'_{n+1}$. If $h \in \mathfrak{b}$ and $q \in A'_n$, say $q = \varphi(y)$ with $y \in A_n$, then:

$$\chi(h) \cdot q = \varphi([h, y]) \in A'_{n+1}.$$

But dimensions match, so $\chi(h)A'_n = A'_{n+1}$ as subgroups of \mathbb{K}_+ .

So fix $h_0 \in \mathfrak{b} \setminus \{0\}$. For arbitrary $h_1 \in \mathfrak{b}$, one has $\frac{1}{\chi(h_0)}\chi(h_1) \in N_{\mathbb{K}}(A'_n) = \{\lambda \in \mathbb{K} : \lambda A'_n \leq A'_n\}$. The latter is therefore an infinite definable subring, implying $N_{\mathbb{K}}(A'_n) = \mathbb{K}$. Hence A'_n is a non-trivial ideal of a field, and $A'_n = \mathbb{K}$. Going back through φ , we have $A_n = B_{\mathfrak{b}}$. But $[A_n, A_n] \leq B_{\mathfrak{b}}$ by Lemma C. So $B_{\mathfrak{b}}$ is a subring. As it is normalised by \mathfrak{b} and $\mathfrak{g} = \mathfrak{b} + B_{\mathfrak{b}}$, we have a final contradiction. \diamond

This completes the proof of Proposition 4.4. \square

Using non-nilpotence of Borel subrings (Proposition 4.4) and the impossibility of a global weight decomposition (Proposition 4.2), we derive the existence of a 3-dimensional Borel subring.

Proposition 4.5. *There is a 3-dimensional Borel subring.*

Proof. Suppose not. The proof has something common with the weight decomposition of Proposition 3.2, Step 3.2.2.

Let \mathfrak{b} be a Borel subring; by Proposition 4.4, $\dim \mathfrak{b} = 2$ and \mathfrak{b} is non-nilpotent. Using Theorem 2 we let $h \in \mathfrak{b}$ act on \mathfrak{b}' as $\text{Id}_{\mathfrak{b}'}$. Let $e \in \mathfrak{b}' \setminus \{0\}$.

We claim that $C_e = \mathfrak{b}'$. Otherwise $\mathfrak{b}' \subset C_e \subset \mathfrak{g}$, so C_e is a 2-dimensional Borel subring; it is non-nilpotent by Proposition 4.4. Now $C_e \neq \mathfrak{b} = N_{\mathfrak{g}}^{\circ}(\mathfrak{b}')$ so $\mathfrak{b}' \neq C'_e$. By Theorem 2, \mathfrak{b}' is a Cartan subring of C_e . Moreover h normalises C_e since for $c \in C_e$ one has:

$$[[h, c], e] = [he, c] + [h, ce] = [e, c] = 0.$$

Hence h also normalises C'_e . Since \mathfrak{b}' is a Cartan subring of the non-nilpotent, 2-dimensional ring C_e , Theorem 2 gives $\eta \in \mathfrak{b}'$ acting on C'_e as $\text{Id}_{C'_e}$. Then for any $a \in C'_e$ one has $[h, a] \in C'_e$ and therefore:

$$[h, a] = [h, [\eta, a]] = [h\eta, a] + [\eta, ha] = [\eta, a] + [h, a] = a + [h, a].$$

So $a = 0$, a contradiction.

We derive $B_e^2 = \mathfrak{b}$. Indeed $C_e = \mathfrak{b}'$ is abelian. By Lemma A, C_e^2 is a subring of \mathfrak{g} ; since $\dim C_e^2 \leq 2 \dim C_e = 2$, it is proper. Now $\mathfrak{b} \subseteq C_e^2$ so equality holds. Hence $\mathfrak{b}' = C_e \subset C_e^2 = \mathfrak{b}$. In particular, $\dim B_e^2 = 2$. We turn to the action of \mathfrak{b} on $\mathfrak{g}/\mathfrak{b}$. If $\mathfrak{g}/\mathfrak{b}$ is \mathfrak{b} -irreducible, then by Corollary 2, \mathfrak{b}' centralises $\mathfrak{g}/\mathfrak{b}$; in particular $B_e^2 \leq B_e \leq \mathfrak{b}$. If $\mathfrak{g}/\mathfrak{b}$ is \mathfrak{b} -reducible, we still get $B_e^2 \leq \mathfrak{b}$ by linearising in dimension 1 *twice*. In either case, $B_e^2 = \mathfrak{b}$.

We now consider eigenspaces with respect to h and show $\mathfrak{g} = E_{-2} + E_{-1} + E_0 + E_1$. Let us turn to the adjoint action of h . First, $E_0 \neq 0$ since $h \in \mathfrak{b}$ which is soluble. By construction, $\mathfrak{b}' \leq E_1$. Suppose $E_{-1} = 0$. Then by surjectivity of $\text{ad}_h + 1$ one gets:

$$B_e = \text{im ad}_e = \text{im ad}_e(\text{ad}_h + 1) = \text{im ad}_h \text{ad}_e = [h, B_e].$$

However $B_e \geq B_e^2 = \mathfrak{b}$ intersects C_h by solubility, a contradiction. We apply just the same argument to prove $E_{-2} \neq 0$; otherwise:

$$B_e^2 = \text{im ad}_e^2(\text{ad}_h + 2) = \text{im ad}_h \text{ad}_e^2 = [h, B_e^2],$$

but $B_e^2 = \mathfrak{b}$, a contradiction.

Thus $\mathfrak{g} = E_{-2} + E_{-1} + E_0 + E_1$, against Proposition 4.2. \square

The analysis of 3-dimensional Borel subrings. We now combine the two main threads (weights, and large Borel subrings) into the analysis of 3-dimensional Borel subrings.

Proposition 4.6. *Let $\mathfrak{b} \subset \mathfrak{g}$ be a 3-dimensional Borel subring. Then:*

- (i) $I = C_{\mathfrak{b}}^{\circ}(\mathfrak{g}/\mathfrak{b}) \triangleleft \mathfrak{b}$ is a 2-dimensional, nilpotent ideal of \mathfrak{b} containing \mathfrak{b}' ;

- (ii) if $J \leq I$ is a 1-dimensional subring, then $J \leq Z(I)$;
- (iii) if $h \in \mathfrak{b}$ is such that I is a sum of eigenspaces, then either $I \leq E_0$ or $I = E_k \oplus E_{-k}$ is a sum of 1-dimensional subrings, with $k \neq 0$;
- (iv) $Z^\circ(\mathfrak{b}) = 0$;
- (v) \mathfrak{b} is the only Borel subring containing I ;
- (vi) I is abelian; for $i \in I \setminus Z(\mathfrak{b})$ one has $C_i = I$.

Be however careful that until (vi) is proved, a subgroup of I need not be a subring.

Proof.

- (i) By Lemma E, I is nilpotent; it is proper in \mathfrak{b} since \mathfrak{b} is no ideal of \mathfrak{g} . Consider the action of \mathfrak{b} on $\mathfrak{g}/\mathfrak{b}$. Linearising in dimension 1, $\dim(\mathfrak{b}/I) \leq 1$ and equality holds.
- (ii) By nilpotence, $Z^\circ(I) \neq 0$. If $J \not\leq Z^\circ(I)$ then $I = J + Z^\circ(I)$ and I is abelian, a contradiction.
- (iii) *All eigenspaces in this proof are with respect to h .* Suppose $I \not\leq E_0$, so there is $k \neq 0$ with $E_k^I \neq 0$. Since $E_0^b \neq 0$ by solubility, Proposition 4.3 implies $\dim E_0 = \dim E_k = 1$. Therefore write $I = E_k \oplus E_\ell$; possibly $\ell = 0$, but even so $\dim E_\ell = 1$.

We may assume $E_{-k} \neq 0$. Indeed, suppose $E_{-k} = 0$. Fix $e \in E_k = E_k^b \leq I$. Notice $B_e \leq \mathfrak{b}$. Now $E_{-k} = 0$ implies $[h, B_e] = \text{im ad}_h \text{ad}_e = \text{im ad}_e(\text{ad}_h + k) = \text{im ad}_e = B_e$. Therefore $B_e \leq \mathfrak{b}' \leq I$. If $\ell = 0$ then $B_e \leq [h, I] = E_k$, and summing over $e \in E_k$ we obtain $E_k \triangleleft \mathfrak{g}$, a contradiction. Hence $\ell \neq 0$. If $E_{-\ell} = 0$, then for $e \in E_\ell$ we also have $B_e \leq I$; summing over $e \in E_k \cup E_\ell$ we obtain $I \triangleleft \mathfrak{g}$, a contradiction. Hence $\ell \neq 0$ and $E_{-\ell} \neq 0$. Up to exchanging, we may assume $E_{-k} \neq 0$.

Let $\mathfrak{h} = E_{-k} + E_0 + E_k$. By Proposition 4.3, each term has dimension 1. By Proposition 4.2, $E_{\pm 2k} = 0$, so \mathfrak{h} is a 3-dimensional subring, therefore self-normalising by Lemma F. In particular, $h \in \mathfrak{h}$; hence $E_{-k} + E_k \leq \mathfrak{h}'$. But \mathfrak{h} is soluble by Proposition 4.1, so $[E_{-k}, E_k] = 0$. However $I \leq C_{\mathfrak{g}}^\circ(E_k)$ by (ii). If $I < C_{\mathfrak{g}}^\circ(E_k)$, then the latter is a 3-dimensional subring, hence self-normalising; then $h \in C_{\mathfrak{g}}^\circ(E_k)$, a contradiction. So $I = C_{\mathfrak{g}}^\circ(E_k) \geq E_{-k}$, and the claim is proved.

- (iv) Let $Z = Z^\circ(\mathfrak{b})$ and suppose $Z \neq 0$. Since \mathfrak{b} is not nilpotent by Proposition 4.4, $\dim Z = 1$ and \mathfrak{b}/Z is non-nilpotent. In particular $Z \leq I$. By Theorem 2, there is $\eta \in \mathfrak{b}/Z$ acting on $(\mathfrak{b}/Z)'$ like the identity. We lift to $h \in \mathfrak{b}$ with the same action, and now lift the eigenspace. *All eigenspaces in this proof are with respect to h .* Thus $E_1^b \neq 0$. By Proposition 4.3, one has $\dim E_0 = \dim E_1 = 1$, so $E_0 = E_0^b = Z \leq I$; of course $E_1 = E_1^b \leq I$ as well, contradicting (iii).
- (v) Let $\mathfrak{b}_2 \neq \mathfrak{b}$ be another Borel subring containing I . Clearly $\dim \mathfrak{b}_2 = 3$, so our earlier analysis applies. Let $I_2 = C_{\mathfrak{b}_2}^\circ(\mathfrak{g}/\mathfrak{b}_2)$. Since $I_2 \triangleleft \mathfrak{b}_2$ one has $I_2 \neq I$. So $X = (I \cap I_2)^\circ$ has dimension exactly 1. Now X is a subring of both I and I_2 , so by (ii) one has $C_{\mathfrak{g}}^\circ(X) \geq I + I_2$ and therefore $C_{\mathfrak{g}}^\circ(X)$ is a 3-dimensional Borel subring with an infinite centre, against (iv).

(vi) Suppose not. Then $J = I' \triangleleft \mathfrak{b}$ is a 1-dimensional ideal of \mathfrak{b} .

There is $h \in \mathfrak{b}$ with $J = E_1(h)$. Indeed, by (iv), the action of \mathfrak{b} on J is non-trivial. Linearising in dimension 1, there is $h \in \mathfrak{b}$ acting on J like the identity. *All eigenspaces are now with respect to h .*

We may assume $E_{-1} \neq 0$. If not then for $j \in J = E_1$ one has $[h, B_j] = B_j$ and quickly $B_j \leq I$. Since $J \leq Z(I)$ by (ii), one also has $C_j \geq I$. Now by (v), \mathfrak{b} is the only Borel subring containing I and $j \notin Z(\mathfrak{b})$ because of the action of h , so $C_j = I$. It follows $B_j = I = C_j$. By Lemma B, $B_j = I$ is abelian and we are done.

Let $\mathfrak{h} = E_{-1} + E_0 + E_1$, which by Proposition 4.2 is a 3-dimensional Borel subring. By Lemma F, $h \in \mathfrak{h}$. In particular $E_{-1} + E_1 \leq \mathfrak{h}' < \mathfrak{h}$, implying $[E_{-1}, E_1] = 0$. However $I \leq C_{\mathfrak{g}}^{\circ}(E_1)$ by (ii), and equality must hold since otherwise $h \in C_{\mathfrak{g}}^{\circ}(E_1)$ by Lemma F. So $E_{-1} \leq C_{\mathfrak{g}}^{\circ}(E_1) = I$ and $I = E_{-1} + E_1$, hence abelian.

It remains to take $i \in I \setminus Z(\mathfrak{b})$ and prove $C_i = I$. One inclusion is clear. If $C_i > I$ then by (v), $C_i = \mathfrak{b}$ and $i \in Z(\mathfrak{b})$, a contradiction. \square

The final contradiction.

Proposition 4.7. *The configuration is inconsistent.*

Proof. Let $\mathfrak{b} \subset \mathfrak{g}$ be a 3-dimensional Borel subring, given by Proposition 4.5. Since \mathfrak{b} is no ideal of \mathfrak{g} , the action of \mathfrak{b} on $\mathfrak{g}/\mathfrak{b}$ is non-trivial. Linearising in dimension 1, there is $h \in \mathfrak{b}$ acting on $\mathfrak{g}/\mathfrak{b}$ like the identity. *All eigenspaces in this proof are with respect to h .* Lifting the eigenspace, $E_1 \neq 0$. By solubility, $E_0 \neq 0$. By Proposition 4.3, $\dim E_0 = \dim E_1 = 1$ and $E_2 = 0$, so E_1 is a 1-dimensional, abelian subring. As in Proposition 4.6, which will be used heavily, we let $I = C_{\mathfrak{g}}^{\circ}(\mathfrak{g}/\mathfrak{b})$.

Step 4.7.1. We may assume $h \in E_0$.

Proof. Consider the action of E_0 on $\mathfrak{g}/\mathfrak{b}$.

Suppose E_0 centralises $\mathfrak{g}/\mathfrak{b}$, viz. $E_0 \leq I$. By Proposition 4.3, one has $\dim E_0 = \dim E_1 = 1$. If $E_1 \leq \mathfrak{b}$, then $I = E_0 + E_1$ where neither is trivial, against Proposition 4.6 (iii). Hence $E_1 \not\leq \mathfrak{b}$ and $(E_1 \cap \mathfrak{b})^{\circ} = 0$. However $E_0 \leq I$, so $[E_0, E_1] \leq (\mathfrak{b} \cap E_1)^{\circ} = 0$ and E_0 centralises E_1 . Then $E_1 \leq C_{\mathfrak{g}}^{\circ}(E_0)$. The latter contains I ; by Proposition 4.6 (v), it equals either I or \mathfrak{b} . This is a contradiction in either case.

Therefore $E_0 \not\leq I$, and $\mathfrak{b} = E_0 + I$. Write $h = e_0 + i$ in obvious notation. Then e_0 also acts on $\mathfrak{g}/\mathfrak{b}$ like the identity; hence $E_1(e_0) \neq 0$. Moreover $E_0 = E_0(h) \leq E_0(e_0)$ by abelianity. By Proposition 4.3, one has $\dim E_0(e_0) = 1$, so $E_0(e_0) = E_0(h)$ contains e_0 . Up to considering e_0 we are done. \diamond

Step 4.7.2. $E_1 \not\leq \mathfrak{b}$.

Proof. Suppose $E_1 \leq \mathfrak{b}$. Let $f = \text{ad}_h - 1$, an additive endomorphism of \mathfrak{g} . Let $K_1 = \ker^{\circ} f = E_1$ and $K_2 = \ker^{\circ}(f^2) \geq K_1$. By definition of h , one has $\text{im } f \leq \mathfrak{b}$. Since $\dim \ker f = 1$, one has $\text{im } f = \mathfrak{b}$, viz. $f: \mathfrak{g} \rightarrow \mathfrak{b}$ is surjective. Since $K_1 = E_1 \leq \mathfrak{b}$, the restriction of $f: K_2 \rightarrow K_1$ remains surjective. Therefore $\dim K_2 = 2 \dim K_1 = 2$.

We claim $K_2 = I$. Let $k_2 \in K_2$, $k_1 = f(k_2) \in K_1$, and $e \in E_1$. Then:

$$[h, [e, k_2]] = [he, k_2] + [e, hk_2] = [e, k_2] + [e, k_2 + k_1] = 2[e, k_2] + [e, k_1].$$

However $e, k_1 \in E_1 = K_1$ while $E_2 = 0$, so $[e, k_1] = 0$. There remains $[e, k_2] \in E_2 = 0$, whence $K_2 \leq C_{\mathfrak{g}}^{\circ}(E_1)$. Now $E_1 \subset I$ so $I \leq C_{\mathfrak{g}}^{\circ}(E_1)$. If the latter has dimension 3, then it is a 3-dimensional Borel subring normalised by h . By Lemma F, $h \in C_{\mathfrak{g}}^{\circ}(E_1)$, a contradiction. Therefore $C_{\mathfrak{g}}^{\circ}(E_1) = I = K_2$. In particular, $B_{k_2} \leq \mathfrak{b}$.

In the notation above, also let $x \in \mathfrak{g}$. By definition of h , there is $b \in \mathfrak{b}$ with $[h, x] = x + b$. Therefore:

$$[h, [k_2, x]] = [hk_2, x] + [k_2, hx] = [k_2 + k_1, x] + [k_2, x + b] = 2[k_2, x] + [k_1, x] + [k_2, b].$$

The left-hand is in \mathfrak{b}' . So is $[k_2, b]$. If $E_{-1} = 0$ then $[h, B_{k_1}] = B_{k_1}$ so $B_{k_1} \leq \mathfrak{b}'$. In that case, $[k_2, x] \in \mathfrak{b}' \leq I$, and therefore $K_2 = I$ is an ideal of \mathfrak{g} , a contradiction. So $E_{-1} \neq 0$. Let $\mathfrak{h} = E_{-1} + E_0 + E_1$, which must be soluble. Hence $E_{-1} \leq C_{\mathfrak{g}}^{\circ}(E_1) = I$ and $\mathfrak{b} = E_0 + E_{-1} + E_1$. But then for $e_{-1} \in E_{-1} \leq I = K_2$, one has $f(e_{-1}) = -2e_{-1} \in E_1$, whence $e_{-1} = 0$, a contradiction.

This proves $E_1 \not\leq \mathfrak{b}$. ◇

Let $\mathfrak{a} = E_1$; by Step 4.7.2 one has $\mathfrak{g} = \mathfrak{a} + \mathfrak{b}$. Let $\mathfrak{b}_2 = \mathfrak{a} + E_0$, a soluble subring. By Step 4.7.1 we could assume $h \in E_0$; so one has $\mathfrak{b}' = [E_0, \mathfrak{a}] = \mathfrak{a}$. Let $\mathfrak{b}_2 < V \leq \mathfrak{g}$ be a \mathfrak{b}_2 -irreducible module; V need not be a subring. Linearising in dimension 1 or by Corollary 2, $[\mathfrak{b}'_2, V] \leq \mathfrak{b}_2$.

Let $Y = (I \cap V)^{\circ}$, an E_0 -module. Since I is abelian by Proposition 4.6 (vi), Y is actually a subring. A priori, $0 < Y \leq I$. Notice $[\mathfrak{a}, Y] \leq (\mathfrak{b} \cap \mathfrak{b}_2)^{\circ} = E_0$. We also claim $[E_0, Y] = Y$ since otherwise h centralises Y , so $Y \leq E_0$ and equality holds, forcing $h \in I$, a contradiction.

If $\dim Y = 2$ then $Y = I$. Then for $a \in \mathfrak{a}$, the map ad_a takes I to E_0 . But by Proposition 4.6 (iv) one has $\dim Z(\mathfrak{b}) = 0$, so there is $i \in I \setminus Z(\mathfrak{b})$ with $[a, i] = 0$. By Proposition 4.6 (vi), a normalises $C_i = I$ and therefore a normalises $N_{\mathfrak{g}}^{\circ}(I) = \mathfrak{b}$. Hence $a \in \mathfrak{b}$ by Lemma F. This proves $\mathfrak{a} \leq \mathfrak{b}$, against Step 4.7.2. Therefore $\dim Y = 1$.

Let $\mathfrak{h} = \mathfrak{a} + E_0 + Y$, a 3-dimensional subring. By Proposition 4.1, \mathfrak{h} is soluble. Since $[E_0, Y] = Y$ and $[E_0, \mathfrak{a}] = \mathfrak{a}$, solubility implies $[\mathfrak{a}, Y] = 0$. So $\mathfrak{a} \leq C_{\mathfrak{g}}^{\circ}(Y) = I \leq \mathfrak{b}$ by Proposition 4.6 (v), a final contradiction. □

It would be interesting to push this line further and try to classify minimal simple (or better: N_{\circ}° -)Lie rings of finite Morley rank. Care will be needed since there is no clear indication what unipotence theory will become in this context, and since the Witt algebra is N_{\circ}° .

4 Appendix: questions

We list a number of questions, some already in [Del16b].

Extending the result

Question 1. *What happens to our theorem in characteristic 3?*

This is likely to be a question for pure algebraists, and a challenging one. Fact 1 is open in characteristic 3.

Abstract Lie rings and their representations

Question 2. *State and prove an analogue of the ‘Borel-Tits’ theorem [BT73], viz. describe automorphisms of Lie rings of Lie-Chevalley type.*

One could start or content oneself over algebraically closed fields (see the elegant model-theoretic proof for groups in [Poi88]), and even ask the same in Lie-Cartan type.

Question 3. *Devise identification methods à la Curtis-Phan-Tits [Tim04; Gra09] for Lie rings of Lie-Chevalley type.*

Question 4. *Study Lie ring representations of Lie rings of Lie-Chevalley type $\mathfrak{G}_\Phi(\mathbb{K})$. (Started in [Del16a; Del17]. Also see Question 6.)*

One could also ask about Lie rings of Lie-Chevalley type (or even Lie-Cartan type) up to elementary equivalence, à la Malcev [Mal61; Bum06]. Such questions usually rely on retrieving the base field, which should be easier here than in groups. For more on elementary properties of Chevalley groups, see [BM02].

Lie modules of finite Morley rank

In the proof of Theorem 4 (more specifically, when analysing 3-dimensional Borel subrings), results like relevant analogues of Maschke’s Theorem [Tin17, Proposition 2.13] or Lie’s Theorem (in characteristic larger than Morley rank, generalising Corollary 2) would come handy.

Question 5. *Develop basic tools for Lie modules of finite Morley rank.*

Question 6. *Let \mathbb{K} be an algebraically closed field of characteristic $p > 0$ and \mathfrak{g} be a finite-dimensional, simple \mathbb{K} -lie algebra. Assume that \mathfrak{g} acts definably and irreducibly on an elementary abelian p -group V of finite Morley rank. Prove that:*

(i) *V has a structure of a finite dimensional \mathbb{K} -vector space compatible with the action of \mathfrak{g} [see [Bor23, Theorem 3]];*

(ii) *\mathfrak{g} is a \mathbb{K} -Lie subalgebra of $\mathfrak{gl}_{\mathbb{K}}(V)$. [More dubious.]*

The original result was for *connected algebraic groups*, which correspond more to Lie rings of Lie-Chevalley type. And for that reason, we are more confident in (i) than in (ii).

Question 7. *Work in a theory of finite Morley rank. Prove a ‘Steinberg tensor product theorem’ [Ste63] for definable $\mathfrak{G}_\Phi(\mathbb{K})$ -modules. [See [Bor23, Theorem 3].]*

A more abstract direction makes no assumptions on \mathfrak{g} .

Question 8. *Classify faithful, irreducible \mathfrak{g} -modules of Morley rank 2. [See [Del09].]*

Abstract Lie rings of finite Morley rank

There is of course the question of field interpretation in a non-abelian nilpotent Lie ring, where unreasonable hopes are shattered by [Bau96].

Returning to the Reineke phenomenon, some topics are quite unclear.

Question 9.

- Is there a more model-theoretic proof of Fact 1 (viz. one not using Block-Premet-Strade-Wilson)?
- Let $\mathfrak{g} > 0$ be a connected Lie ring of finite Morley rank and $x \in \mathfrak{g}$. Is C_x infinite? [See [BBC07, Proposition 1.1].]
- When is $x \in C_x$? [One should be careful with such questions, the Baudisch algebras could be counter-examples.]

At the soluble level, we have reasonable expectations.

Question 10. *Develop a theory of soluble Lie rings of finite Morley rank: Fitting theory and Cartan subring theory. [See [Fr 00].]*

Of course we do not aim for ‘conjugacy’ (if at all meaningful since there is no group around; fails anyway in Witt’s algebra)—but existence and self-normalisation. Notice that existence could be hoped without the solubility assumption [see [FJ05]], though we currently see no way to attack this.

Then of course, what matters is the simple case.

Let \mathbb{K} be an algebraically closed field of characteristic p . Then Witt’s algebra $W_{\mathbb{K}}(1; \underline{1})$ is a p -dimensional, simple Lie algebra over \mathbb{K} , hence definable in the pure field. Moreover, it has a subalgebra of codimension 1. Mind the assumption on the characteristic in the following.

Question 11. *Let \mathfrak{g} be a simple Lie ring of finite Morley rank d and characteristic $p > d$. Suppose \mathfrak{g} has a definable subring of corank 1. Then $\mathfrak{g} \simeq \mathfrak{sl}_2(\mathbb{K})$. [See [Hru89].]*

Last but not least in this direction, is of course the question of finding a strategy for the ‘log CZ’ conjecture. For CZ itself, the positive solution in so-called ‘even type’ [ABC08] was a form of the classification of the finite simple groups (CFSG) seen from the *reducing* lens of model-theoretic arguments for the tame infinite. At the high end of the classification, viz. in high Lie rank, are ‘generic identification results’.

Question 12. *Devise generic identification methods for simple Lie rings of finite Morley rank. [See Question 3, see [BB11].]*

As said in the introduction, we are curious whether model theory may provide a reduced sketch of the Block-Premet-Strade-Wilson theorem, like it did for the CFSG.

Model theory

We see two questions here; one aims at broadening the model-theoretic frame of study of Lie rings, and the other at tightening connections between groups and Lie rings, presumably under new assumptions.

First, it is unclear to us whether the full strength of finite Morley rank was used; dimensionality could suffice, but this may depend on the behaviour of differential fields, if any. (It is unclear to us whether non-trivial differential fields would give counter-examples to the dimensional version of our results. Reasoning by loose analogy, fields with non-minimal \mathbb{G}_m have not provided counter-examples to the Cherlin-Zilber conjecture, at least not yet.)

Question 13. *Determine in which model-theoretic setting the present paper took place.*

Second, by Lie-Chevalley correspondence, we mean the ability to attach to some groups a Lie ring. We do not believe in one for abstract groups of finite Morley rank, as categorical nature allows no infinitesimal methods. (This contrasts sharply with \mathcal{o} -minimal nature, where infinitesimals are provided by elementary extensions; in the tame *ordered* case, model-theoretic functoriality proves geometrically sufficient.) One could add a stronger model-theoretic assumption with geometric flavour.

Question 14. *Is there a Lie-Chevalley correspondence for groups definable in Zariski geometries [HZ96]?*

To our understanding, the Cherlin-Zilber conjecture is still open for groups definable in Zariski geometries. A positive answer to Question 14 would certainly give a natural proof.

Acknowledgements

We thank Gregory Cherlin for useful comments.

Very special thanks: the CIRM. The collaboration started virtually during a lockdown; but for serious research one must meet in person. This was made possible thanks to the CIRM-AIMS residence programme (no longer restricted to host AIMS countries), allowing the second author to visit France for one week and the first author to visit Gabon for two, all in the Summer of 2023. Both stays proved highly productive and pleasant. Our very special thanks to Olivia Barbarroux, Pascal Hubert and Carolin Pfaff.

References

- [ABC08] Tuna Altinel, Alexandre Borovik and Gregory Cherlin. *Simple groups of finite Morley rank*. Vol. 145. Mathematical Surveys and Monographs. Providence, RI: American Mathematical Society, 2008. xx+556.
- [Bau+09] Andreas Baudisch et al. ‘Die böse Farbe’. *J. Inst. Math. Jussieu* 8.3 (2009), pp. 415–443.
- [Bau96] Andreas Baudisch. ‘A new uncountably categorical group’. *Trans. Amer. Math. Soc.* 348.10 (1996), pp. 3889–3940.
- [BB11] Ayse Berkman and Alexandre Borovik. ‘A Generic Identification Theorem for Groups of Finite Morley Rank, Revisited’. arXiv 1111.6037. 2011.
- [BBC07] Alexandre Borovik, Jeffrey Burdges and Gregory Cherlin. ‘Involutions in groups of finite Morley rank of degenerate type’. *Selecta Math. (N.S.)* 13.1 (2007), pp. 1–22.
- [BM02] Elena Bunina and Aleksandr Mikhalev. ‘Elementary properties of linear and algebraic groups’. Vol. 110. 3. *Algebra*, 18. 2002, pp. 2595–2659.
- [BMZ07] Andreas Baudisch, Amador Martin-Pizarro and Martin Ziegler. ‘Red fields’. *J. Symbolic Logic* 72.1 (2007), pp. 207–225.

- [BN94] Alexandre Borovik and Ali Nésin. *Groups of finite Morley rank*. Vol. 26. Oxford Logic Guides. The Clarendon Press — Oxford University Press, New York, 1994. xviii+409.
- [Bor23] Alexandre Borovik. ‘Finite group actions on abelian groups of finite Morley rank’. 13 pages. Submitted. 2023.
- [BT73] Armand Borel and Jacques Tits. ‘Homomorphismes “abstraites” de groupes algébriques simples’. *Ann. of Math. (2)* 97 (1973), pp. 499–571.
- [Bun06] Elena Bunina. ‘Elementary equivalence of Chevalley groups over fields’. *Fundam. Prikl. Mat.* 12.8 (2006), pp. 29–77.
- [BW88] Richard Block and Robert Wilson. ‘Classification of the restricted simple Lie algebras’. *J. Algebra* 114.1 (1988), pp. 115–259.
- [CD12] Gregory Cherlin and Adrien Deloro. ‘Small representations of SL_2 in the finite Morley rank category’. *J. Symbolic Logic* 77.3 (2012), pp. 919–933.
- [CJ04] Gregory Cherlin and Éric Jaligot. ‘Tame minimal simple groups of finite Morley rank’. *J. Algebra* 276.1 (2004), pp. 13–79.
- [DEl+23] Christian D’Elbée et al. ‘Model-theoretic properties of nilpotent groups and Lie algebras’. In preparation. 2023.
- [Del07] Adrien Deloro. ‘Groupes simples connexes minimaux de type impair’. PhD thesis. Paris: Université Paris 7, 2007. 93 pp.
- [Del09] Adrien Deloro. ‘Actions of groups of finite Morley rank on small abelian groups’. *Bull. Symb. Log.* 15.1 (2009), pp. 70–90.
- [Del16a] Adrien Deloro. ‘Symmetric powers of $\text{Nat } \mathfrak{sl}_2(\mathbb{K})$ ’. *Comm. Algebra* 44.5 (2016), pp. 1981–2008.
- [Del16b] Adrien Deloro. ‘Un regard élémentaire sur les groupes algébriques’. Mémoire d’habilitation à diriger des recherches. Université Paris 6. Paris, 2016.
- [Del17] Adrien Deloro. ‘Locally quadratic modules and minuscule representations’. *Algebr. Represent. Theory* 20.4 (2017), pp. 977–1005.
- [Del23] Adrien Deloro. ‘Zilber’s skew-field theorem’. 13 pages. Submitted. 2023.
- [DJ16] Adrien Deloro and Éric Jaligot. ‘Involutive Automorphisms of N_{\circ}° -groups of finite Morley rank’. *Pacific J. Math.* 285.1 (2016), pp. 111–184.
- [FJ05] Olivier Frécon and Éric Jaligot. ‘The existence of Carter subgroups in groups of finite Morley rank’. *J. Group Theory* 8.5 (2005), pp. 623–633.
- [Fré00] Olivier Frécon. ‘Sous-groupes anormaux dans les groupes de rang de Morley fini résolubles’. *J. Algebra* 229.1 (2000), pp. 118–152.
- [Gra09] Ralf Gramlich. ‘Developments in finite Phan theory’. *Innov. Incidence Geom.* 9 (2009), pp. 123–175.
- [Hru89] Ehud Hrushovski. ‘Almost orthogonal regular types’. *Ann. Pure Appl. Logic* 45.2 (1989). Stability in model theory, II (Trento, 1987), pp. 139–155.
- [Hum75] James Humphreys. *Linear algebraic groups*. Vol. 21. Graduate Texts in Mathematics. Springer-Verlag, New York, 1975, pp. xiv+247.

- [HZ96] Ehud Hrushovski and Boris Zilber. ‘Zariski geometries’. *J. Amer. Math. Soc.* 9.1 (1996), pp. 1–56.
- [KS69] Aleksei Kostrikin and Igor Shafarevich. ‘Graded Lie algebras of finite characteristic’. *Izv. Akad. Nauk SSSR Ser. Mat.* 33 (1969), pp. 251–322.
- [Mal61] Anatoly Malcev. ‘The elementary properties of linear groups’. *Certain Problems in Mathematics and Mechanics (In Honor of M. A. Lavrentev) (Russian)*. Izdat. Sibirsk. Otdel. Akad. Nauk SSSR, Novosibirsk, 1961, pp. 110–132.
- [Mel80] Hayk Melikjan. ‘Simple Lie algebras of characteristic 5’. *Uspekhi Mat. Nauk* 35.1(211) (1980), pp. 203–204.
- [Nes89] Ali Nesin. ‘Nonassociative rings of finite Morley rank’. Vol. 11. 1989, pp. 117–137.
- [Poi01] Bruno Poizat. *Stable groups*. Vol. 87. Mathematical Surveys and Monographs. Translated from the 1987 French original by Moses Gabriel Klein. Providence, RI: American Mathematical Society, 2001. xiv+129.
- [Poi88] Bruno Poizat. ‘MM. Borel, Tits, Zilber et le général nonsense’. *J. Symbolic Logic* 53.1 (1988), pp. 124–131.
- [PS06] Alexander Premet and Helmut Strade. ‘Classification of finite dimensional simple Lie algebras in prime characteristics’. *Representations of algebraic groups, quantum groups, and Lie algebras*. Vol. 413. Contemp. Math. Amer. Math. Soc., Providence, RI, 2006, pp. 185–214.
- [Rei75] Joachim Reineke. ‘Minimale Gruppen’. *Z. Math. Logik Grundlagen Math.* 21.4 (1975), pp. 357–359.
- [Ros91] Richard Rosengarten. ‘Aleph-zero stable Lie algebras’. PhD thesis. New Brunswick: Rutgers, The State University of New Jersey, 1991. 55 pp.
- [Ste63] Robert Steinberg. ‘Representations of algebraic groups’. *Nagoya Math. J.* 22 (1963), pp. 33–56.
- [Str04] Helmut Strade. *Simple Lie algebras over fields of positive characteristic I*. Vol. 38. De Gruyter Expositions in Mathematics. Structure theory. Walter de Gruyter & Co., Berlin, 2004, pp. viii+540.
- [Str07] Helmut Strade. ‘Lie algebras of small dimension’. *Lie algebras, vertex operator algebras and their applications*. Vol. 442. Contemp. Math. Amer. Math. Soc., Providence, RI, 2007, pp. 233–265.
- [Str09] Helmut Strade. *Simple Lie algebras over fields of positive characteristic II*. Vol. 42. De Gruyter Expositions in Mathematics. Classifying the absolute toral rank two case. Walter de Gruyter & Co., Berlin, 2009, pp. vi+385.
- [Str13] Helmut Strade. *Simple Lie algebras over fields of positive characteristic III*. Vol. 57. De Gruyter Expositions in Mathematics. Completion of the classification. Walter de Gruyter GmbH & Co. KG, Berlin, 2013, pp. x+239.
- [Tim04] Franz Georg Timmesfeld. ‘The Curtis-Tits-presentation’. *Adv. Math.* 189.1 (2004), pp. 38–67.
- [Tin17] Jules Tindzogho Ntsiri. ‘The structure of an SL_2 -module of finite Morley rank’. *Math. Log. Q.* 63.5 (2017), pp. 364–375.

- [Zil82] Boris Iossifovitch Zilber. ‘Uncountably categorical nilpotent groups and Lie algebras’. *Izv. Vyssh. Uchebn. Zaved.* 26.5 (1982), p. 75.