



HAL
open science

The effects of a new immersive multidomain training on cognitive, dual-task and physical functions in older adults

Néva Béraud-Peigné, Pauline Maillot, Alexandra Perrot

► To cite this version:

Néva Béraud-Peigné, Pauline Maillot, Alexandra Perrot. The effects of a new immersive multidomain training on cognitive, dual-task and physical functions in older adults. *GeroScience*, 2023, 10.1007/s11357-023-00952-w . hal-04217390

HAL Id: hal-04217390

<https://hal.science/hal-04217390>

Submitted on 2 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The effects of a new immersive multidomain training on cognitive, dual-task and physical functions in older adults

Néva Béraud-Peigné ^{1,2,3}, Pauline Maillot ³, Alexandra Perrot ^{1,2}

¹ CIAMS, Université Paris Saclay, Orsay, France

² CIAMS, Université Paris Saclay, Orléans, France

³ I3SP, Université Paris-Cité, Paris, France

E-mail address of the corresponding author: nevabp1@live.fr

Orcid: [0000-0001-7142-540X](https://orcid.org/0000-0001-7142-540X)

Information of other authors: pauline.maillot@u-paris.fr (Orcid: [0000-0002-9101-8921](https://orcid.org/0000-0002-9101-8921));

alexandra.perrot@universite-paris-saclay.fr (Orcid: [0000-0003-1246-2221](https://orcid.org/0000-0003-1246-2221)).

1 Abstract

2 The aim of this study was to assess the potential of multidomain training using Immersive and
3 Interactive Wall Exergames (I2WE) to improve the cognitive and physical functions of older adults.
4 This new generation of exergames offers specific advantageous characteristics (e.g., immersion,
5 virtual cognitive solicitation, high physical stimulation, complex motor skills, multiple social
6 interactions) that could make I2WE an enjoyable multidomain training medium for older adults. A
7 pilot study was set up with 34 participants (M = 69.91 years old). It used the pre-tests – training (3
8 months, 2 one-hour sessions per week for the 2 groups) – post-tests method to compare: a) the
9 cognitive and the Dual-Task (DT) effects (primary outcomes), b) the physical effects and perceived
10 pleasure (secondary outcomes) between an experimental group following an I2WE program (n = 19)
11 to an active control group performing a Walking and Muscle-Strengthening (WMS) program (n = 15).
12 While visuospatial short-term memory was improved for both groups, only I2WE training enhanced
13 visuospatial working memory, inhibition, and DT. All physical functions, except upper body strength,
14 were enhanced for the 2 groups. The perceived pleasure was higher for the I2WE group than the WMS
15 group. The results of this first study have important clinical implications, showing that I2WE can
16 optimize strategies to improve older adults' physical and cognitive health. Virtual and immersive
17 cognitive stimulation combined with varied physical activity (i.e., aerobic, muscle-strengthening,
18 complex motor skills) appear to be major assets of these new exergames. Moreover, the high level of
19 perceived pleasure to I2WE makes it a promising tool for engaging older adults in sustained
20 multidomain practice.

21 **Keywords:** exergames, physical activity, healthy aging, cognition, cognitive-motor training, dual-task

22

1 Introduction

2 Cognitive-motor integrity and Dual-Task (DT) capacities are crucial for maintaining walking and
3 preventing falls [1], and thus, autonomy and quality of life of older adults. Despite this, as their
4 physical and cognitive functions decline [2] and even though novelty and change are beneficial to
5 maintain them, training with one modality (aerobic, muscle-strengthening) or usually combined
6 (aerobic and/or muscle-strengthening and/or balance and/or flexibility) are usually offered to them.
7 They improve cognitive functions [3,4], however, some researchers have pointed out [5] the necessity
8 to extend this field to other forms of physical activity (e.g., complex motor skills, open-ended
9 activities). An optimal combination that includes cognitive training based on enjoyable activities is
10 also essential to anchor participants in sustainable practice. Indeed, multidomain training that
11 combines different components (e.g., physical, motor, cognitive) can generate a synergistic effect,
12 increasing the efficiency of training in enhancing cognitive functions [6].

13 New technologies, like exergames, could offer efficient and enjoyable training support. However, their
14 effects on older adults' physical and cognitive functions are mixed and debated. Several studies have
15 shown that they can be effective [7], greater than physical programs alone [8–10], and even better than
16 programs combining physical and motor activities in improving mental flexibility [11], overall
17 cognitive level [12], and DT skills [13]. However, reviews and meta-analyses [8–10] highlight a great
18 deal of heterogeneity, which negatively impacts the conclusions. Moreover, some studies have failed
19 to demonstrate the superiority of exergames over conventional programs [14–16] or to be ineffective
20 [16,17]. In contrast, the development of a new generation of Immersive and Interactive Wall
21 Exergames (I2WE) could offer the support of an efficient combined training [18], maybe more than
22 training with one modality. However, due to their recent development, few studies have been
23 conducted (*Play Lü* [19], *ExerCube* [20], *Neo One* [18]). I2WE creates simultaneous multidomain
24 training, with complete physical stimulation (i.e., aerobic, muscle strengthening, and complex motor
25 skills) and virtual continuous cognitive solicitation. More specifically, the cognitive task is

1 incorporated into the physical task in accordance with the *Moving While Thinking* approach [21].
2 Thus, the 2 tasks are linked and interactive, which could benefit older adults' cognitive functions since
3 it can increase the synergistic effect of the combined training tenfold [22,23].

4 I2WE offers strong immersive capabilities that can increase session intensity [24], create lower
5 perceived exertion [19], generate better performance [25], and amplify the activity of brain regions
6 associated with executive functions [26]. Indeed, the cognitive stimulation of I2WE triggers greater
7 mental concentration than conventional physical training [20] and potentiates the effects on cognition.

8 These new exergames could generate higher physical intensity than their older counterparts (e.g., *Wii*)
9 or conventional physical activity. The large play area encourages energy expenditure, and participants
10 cannot use energy-saving strategies. They must unconsciously perform complex movements during
11 the sessions, using the whole body to achieve their goals (e.g., using the upper body to throw a ball,
12 moving quickly to reach targets, bending to retrieve a ball from the floor). A recent study on
13 adolescents using the *Play Lii* device [19] shows that it generates moderate to high intensity. *Neo One*
14 wall [18] also leads to moderate to high intensity on older adults, the levels of physical intensity
15 recommended by the ACSM [27] to generate benefits on cognitive functions [28,29].

16 Some I2WE can be played with up to 10 participants, naturally encouraging situations of cooperation
17 and/or opposition [19]. These activities hold promise for improving cognitive functions in older adults
18 [30]. Collective exergames also lead to higher intensity [19] and better results on improving functional
19 capacities in older adults [31] compared to exergames played alone. Thus, I2WE may be a promising
20 option with advantageous modalities to counter the limitations of conventional training and
21 simultaneously improve cognitive, dual-task, and physical functions in the elderly.

22 To our knowledge, the effects of I2WE compared to Walking and Muscle-Strengthening (WMS) on
23 cognitive, DT, and physical parameters is currently unknown, with no evidence describing which
24 exercise is superior to improve cognitive, DT, and physical functions. Therefore, this study aimed to

1 compare the effects of I2WE and WMS on a) cognitive functions and DT (primary outcomes) and b)
2 physical parameters and perceived pleasure (secondary outcomes). We hypothesized that both groups
3 would improve their cognitive, DT, and physical measures, whereas cognitive functions would
4 enhanced in the I2WE group to a higher degree, consistent with Kraft's theory [6] of the synergistic
5 effect of combined training.

6 **Methods**

7 **Population**

8 Potential participants were contacted via mail, flyers, and call for participation in the Montreuil city
9 (France) newspaper, and recruitment of participants was conducted between December 2021 and
10 February 2022. A power analysis, using G*Power version 3.1.9.7, based on the primary statistical test
11 planned for this study (2×2 within-between interaction ANOVA) revealed that a sample size of 34
12 participants was needed ($\alpha = .05$; power = .80; effect size: $\eta^2 = .06$). The dropout level was assumed to
13 be 15 % so 40 community-dwelling older adults were recruited from Seine-Saint-Denis, France,
14 volunteered to participate. All participants gave their written informed consent and were not
15 compensated for their participation. The study was approved by the Polethis research ethics committee
16 (CER-PS n°462). Participants met the following inclusion criteria: (a) be over 60 years old, (b) have a
17 MMSE [32] score above 27, (c) engage in less than 2.5 hours of physical activity per week assessed by
18 the QAPPA [33], (d) score 3 or higher on a subjective health scale (1= very poor to 5= very good), (e)
19 be retired, (f) have no major pathology, (g) have normal or corrected vision and hearing, (h) have no
20 difficulty bending or using their shoulders, (i) do not play video games or exergames, (j) present a
21 medical certificate not contraindicating physical activity). They did not know in advance which group
22 they would be included in.

23 **Study design**

1 This study was a single-center interventional study wherein participants were randomized into one of
2 two groups (I2WE or WMS). The randomization was carried out in single-blind conditions:
3 participants did not know their group in advance, and the researcher was not blinded. The allocation of
4 participants between groups was done as the participants were included in the study according to 2
5 criteria: a) the equality in terms of age between the 2 groups and b) the availability of participants
6 (randomization was not performed with a software). The pre- and post-tests were carried out just
7 before the start and just after the end of the program, so there was a 13-week gap. The researcher who
8 administered the tests had a bachelor's and a master's degree, specialized in physical activity for aging,
9 and was trained to administer the tests.

10 The cognitive, dual-task, and physical tests took around 1h30 per participant. Cognitive tests were
11 performed first, followed by physical tests. They were carried out in a quiet room. A break of about 3
12 minutes was observed between each test to manage the participant's mental fatigue and attention
13 effect. In addition, the participant was regularly asked if they needed an extra break and if they felt
14 ready to start another test. All participants took the tests in the same order, before and after the
15 programs, to control the mental fatigue effect.

16 **Outcomes**

17 **Primary outcome measures**

18 *The cognitive and DT battery*

19 The French version [34] of **the Zoo Map Test** is from the Behavioural Assessment of the
20 Dysexecutive Syndrome [35], a validated ecological test battery assessing planning ability [36]. To
21 our knowledge, no test-retest coefficients are available for the population studied in this research. The
22 participant indicated the route to take to visit a series of places on a map, avoiding making mistakes. In
23 the *formulation of the plan* part, the participant organized their own travel. In the *execution of the plan*

1 part, they followed instructions. For each condition, the *total time* and the *sequence score* out of 8
2 points were collected for a total score out of 16 points.

3 **The Spatial Span Test** is a reliable and valid subtest of the Wechsler Memory Scale [37], with an
4 acceptable internal consistency (Cronbach's $\alpha = .61$) [38] and test-retest reliability for 60 to 79 years
5 old ranging from .52 to .65 [39]. In the first part, which assesses visuospatial short-term memory, the
6 participant reproduces a sequence of cubes designated by the examiner in the same order. In the
7 second, which assesses visuospatial working memory, the participant reproduces a sequence of
8 movements in reverse order. The number of cubes increases progressively and determines the *spatial*
9 *span* and the *backward spatial span*. For each level, the participant always performed 2 trials and
10 scored 1 point for each successful trial. The test ended when 2 errors were made at the same level.

11 **The Stroop Test** [40] was used to assess inhibition. It has good reliability (.75 for Board 1, .82 for
12 Boards 2 and 3 [41]) and validity [42]. Each board contained 100 items divided into 10 columns. Only
13 the first 3 boards were kept: the *naming condition*, which contained colored rectangles, the *reading*
14 *condition*, with color words appearing in black ink and the *interference condition*, which contained color
15 words, appearing in a color different from their meaning. In all 3 conditions, the performance measure
16 was the number of correct responses in 45 seconds. The interference score measure was calculated as
17 follows: board 3 score/(mean of board 1 and 2 scores).

18 **The Mental Rotation Test** [43] assesses visuospatial abilities. This valid test displayed substantial
19 internal consistency (KR-20 = .88) and a high test-retest reliability ($r = .83$) [43]. Twenty figures were
20 presented as 2-dimensional visual images. A model was placed at the left end of the line, and 4 structures
21 were located to the right of the model. The participant was asked to indicate which ones were the same
22 figure as the model, taking rotation into account. The score was the number of correct items given in 6
23 minutes.

1 **The Trail Making Test** (TMT) [44] assesses cognitive flexibility, with high test-retest reliability for
2 Part A ($r = .74$), part B ($r = .85$), and B-A ($r = .74$) [45]. Part A involves linking a series of increasing
3 numbers from 1 to 25 by selecting the appropriate number. In part B, the participants connect 2 series
4 of symbols in alternation: 1 of numbers and 1 of letters. This had to be done as quickly as possible
5 without lifting the pen from the paper. If mistakes were made, the examiner informed the participant,
6 who had to correct them. Following Corrigan and Hinkeldey's method [46], flexibility was highlighted
7 specifically by analyzing the time difference in the result B-A, a purer measure of flexibility. Most
8 recently, Sánchez-Cubillo and collaborators confirmed that B-A time minimizes the visuo-perceptual
9 and working memory demand of TMT A and TMT B, providing a pure indicator of task-switching
10 [47].

11 **Dual-Task Test (DT)**. The *single recall task* and *TUG in the dual task (TUG-dt)* tests were used to
12 evaluate the DT. This test was designed for study purposes, so there were no data available to confirm
13 its validity and/or reliability. For the *single recall task*, the examiner said 7 numbers aloud, at a rate of
14 1 number per second, which the participant repeated immediately after. For the *TUG-dt*, the participant
15 had to simultaneously perform the single recall task and the TUG test (detailed below). Two scores
16 were calculated: the difference between digits given in DT and Single Task (ST) and the difference
17 between the time to complete the task in DT and ST.

18 **Secondary outcome measures**

19 ***The physical battery***

20 Physical tests were carried out in the following order: a) Chair stand, b) Chair-sit-and-reach, c)
21 Handgrip, d) TUG Test, e) Back scratch, f) Seated Medicine Ball Throw, g) 6-minutes Walking Test.
22 This order ensures that no two functions (e.g., strength, flexibility) and/or body parts (e.g., upper body,
23 lower body) were required in succession. Five tests [i.e., chair-stand, chair-sit-and-reach, Timed Up
24 and Go (*TUG*), Back scratch, 6-minutes Walking Test (*6WT*)], derived from the French version [48] of

1 the *Senior Fitness Test* [49] were used to assess respectively lower body strength, flexibility, dynamic
2 balance, and endurance.

3 ***Chair-stand Test.*** The participant had to stand and sit up as many times as possible in 30 seconds. The
4 arms were crossed on the chest, the feet always remained on the ground. The number of chair-stands
5 was recorded.

6 ***Chair-sit-and-reach Test.*** Sitting on the edge of a chair, one leg straight, the other bent, the participant
7 had to bring their hand as close as possible to the foot of the straight leg. The number of centimeters
8 between the hand and the foot was measured. The values were negative if the fingers of the hand
9 extended beyond the toes.

10 ***TUG Test.*** The participant had to get up from a chair, walk around a block located in front of them, 3
11 meters away, and sit down again as quickly as possible. The time to complete this course was
12 recorded.

13 ***Back scratch Test.*** The participant had to bring their two hands as close together as possible (one hand
14 reaching over the shoulder, palm against the back, the other reaching to the middle of the back, back of
15 the hand against the back). The number of centimeters between both hands was measured. The values
16 were negative if the fingers of the two hands crossed.

17 ***6WT.*** The participant was asked to walk as far as possible, in 30-meter round trips, for 6 minutes. The
18 distance was recorded, as well as the heart rate, to determine the intensity of the effort.

19 Two upper-body strength tests (i.e., Handgrip test [50], Seated Medicine Ball Throw [51]) were added
20 to complete the physical assessment.

21 ***Handgrip Test.*** The participant was standing and had to squeeze the dynamometer with maximum
22 isometric effort, maintained for about 5 seconds. The force created, measured in kilograms, was
23 recorded for both arms. The experimental protocol included a practice trial, followed by three attempts

1 for each side in alternation, with 30 seconds rest between them. The best of the three trials for each
2 hand was considered for analysis.

3 **Seated Medicine Ball Throw Test.** Sitting in a chair, back against the back of the chair, feet flat on the
4 floor, the subject had to throw a 3kg medicine ball as far as possible with both hands, starting from the
5 chest. The distance between the subject and the medicine ball was recorded.

6 ***The questionnaires***

7 The French version [52] of the **Physical Activity Enjoyment Scale** [53] with a scale from *absolutely*
8 *disagree* (1) to *completely agree* (7) was used to evaluate the perceived pleasure. The score was
9 between 7 and 70.

10 **The feeling questionnaire.** It was completed at the end of the program. It consisted of 4 questions in
11 the form of a scale from 1 to 5 concerning (a) the difficulty of the sessions, (b) their physical and (c)
12 cognitive developments, and (d) the desire to continue a program of this type.

13 **Intervention**

14 The 2 programs were implemented between February and May 2022 in a sports center in Montreuil
15 (France). The workload was the same between groups: programs were composed of bi-weekly one-
16 hour group sessions over 12 weeks (24 sessions) at a moderate intensity. The Heart Rate (HR) was
17 measured during all sessions using the Polar Verity Sense heart rate monitor placed on the biceps and
18 the perceived exertion using the modified Borg scale [54]. The Heart Rate Reserve (HRR), the
19 difference between the Maximum Theoretical Heart Rate ($MTHR = 192 - 0.007 \times \text{age}^2$, [55]) and the
20 RHR, is used to calculate exercise intensity. For all sessions, the mean intensity was calculated
21 $[(\text{Average Heart Rate} - \text{RHR})/\text{HRR}] \times 100$. A sport therapist supervised the sessions, explaining and
22 demonstrating the exercises, monitoring the participants' safety and correct posture, and encouraging
23 and providing feedback to the participants. There were no activities to increase compliance.

1 ***The Immersive and Interactive Wall Exergames (I2WE) program*** was performed using the *Neo One*
2 device described in Figure 1 and more detailed in this recent study [18]. This I2WE was chosen
3 because of the variety and quality of the games offered, which are accessible to older people [18]. The
4 sessions consisted of 10 minutes of warmup, 40 minutes of the main session using the *Neo One*
5 device, and 10 minutes of cooldown with full body stretching exercises in groups of 4 participants
6 (defined according to the availability of the participants, age, physical and cognitive capacities). The
7 intensity of body workout was moderate (46.97% HHR). Five to 7 five-minutes games were selected
8 from among 15 games [see Additional file 1 for a description of the games]. Various materials were
9 used (e.g., basketball and handball balls, balloons, chairs, hoops, rhythm scales). The games were
10 made more complex by selecting levels (easy, normal, hard) and by adding extra exercises to create
11 cooperative and/or oppositional situations to the games where they were not initially implemented and
12 to increase the physical and/or motor complexity [see Additional file 2].

13 In every other session, participants of the I2WE group played the *Break It Handball* game individually
14 to measure their capacity to progress in *Neo One* games and to evaluate if they were improving in the
15 training underlying their learning. For 2 minutes, the players had to throw the ball at different targets
16 on the wall. They received immediate feedback on their scores.

17 ***The Walking and Muscle-Strengthening (WMS) program*** was held in groups of 5 to 8 participants
18 (split according to the same criteria as the I2WE group) in the parks of the city of Montreuil, with flat
19 and gradient courses that were used alternately. Participants spent 45 minutes walking and 15 minutes
20 in muscle-strengthening exercises, during which they performed an average of 4881.59 steps
21 (SD = 218.73), measured by standard pedometers. The intensity was 43.90% of the HRR (moderate
22 intensity). The muscle-strengthening exercises were performed using either body weights or small
23 equipment (e.g., balls, elastic bands, rhythm scales) or street furniture (e.g., benches, trees, steps).
24 Approximately 3 to 5 exercises were performed, consisting of 2 to 3 sets of 10 to 20 repetitions.

25 **Data analysis**

1 The analyses were performed using JASP software. Before completing any inferential statistical tests,
2 all relevant assumptions were checked, and if they were met, statistical analysis was allowed to
3 proceed. Measures of variance used were Standard Deviation (SD). The equivalent of I2WE and WMS
4 groups before the program was checked using a *t*-test. For cognitive, DT, and physical tests, repeated
5 ANOVA measures were used to test the interaction between pre-test and post-test x training group
6 (I2WE, WMS) and were completed by 95% Confidence Intervals for mean difference (between pre-
7 and post-tests or between I2WE and WMS). All statistical analyses with a *p*-value < .05 were
8 considered significant. A Bonferroni Post Hoc test was realized if a significant interaction was found.
9 Only the *p*-values from the Post Hoc tests were Bonferroni-corrected. Effect Sizes were calculated to
10 study the power of the results: Eta-Squared for ANOVA and Cohen's *d* for *t* test. To test the reliability
11 of results, Reliable Change Indexes (RCIs) were calculated according to the method presented in
12 Duff's work [56]. The calculation was as follows:

$$13 \quad RCI = \frac{T_2 - T_1}{\sqrt{(2S_{21}(1 - r_{12}))}}$$

14 T_1 = mean score at pre-test, T_2 = mean score at post-test, S_1 = standard deviation at pre-test, r_{12} =
15 correlation between T_1 and T_2 . Calculation of the RCI results in a z-score, which needs to be
16 compared with a normal distribution table to be interpreted. Within the existing literature, a z-score of
17 +1.645 or -1.645 would be considered a "reliable change". Correlation analysis was also performed
18 using Spearman correlation, and the potential difference in perceived pleasure between the two groups
19 was tested with a *t*-test.

20 **Results**

21 **Participants**

22 The study workflow is shown in Figure 2. 34 participants (26 women) completed the study, and there
23 were no significant differences between the 2 groups in age, years of education, BMI, Resting Heart

1 Rate (RHR), subjective health, MMSE, QAPPA scores (Table 1) and between all pre-test (Additional
2 file 3 and 4).

3 **Effects of the programs**

4 **Primary outcome**

5 **Cognitive and DT assessments**

6 The results obtained in the pre- and post-tests for cognitive functions, DT, and the results on simple
7 time and time*group interaction effects from ANOVA are given in Table 2, the results on Post-Hoc
8 Tests are given in Additional file 5, and those of RCI in Additional file 6.

9 **Spatial Span test.** Both groups significantly improved their *spatial span*: there was a significant time
10 effect ($p = .02$), with no time*group interaction or group effect ($p = .44$). Post Hoc Comparison for
11 time effect was significant ($p = .02$, $d = .43$). For the *backward spatial span*, only participants in the
12 I2WE group improved their score. There was a significant time ($p < .001$) and a time*group
13 interaction ($p = .01$) effect, with no group effect. Post Hoc analysis revealed a significant difference
14 between pre- and post-tests for the I2WE group ($p < .001$, $d = 1.04$) and none for the other
15 comparisons.

16 **Stroop Test.** The I2WE group improved its *interference score* and *words in board 3* after the
17 program, while the WMS group did not, confirmed by the significant score*group interaction for the
18 interference score ($p = .01$), with no simple time or group effect. Post Hoc analysis indicated a
19 significant difference between the pre- and post-test for the I2WE group ($p = .05$, $d = .50$) and none for
20 the WMS group. For the *words in board 3*, there was also a score*group interaction ($p = .03$). Post
21 Hoc analysis showed a significant difference between pre- and post-tests for the I2WE group ($p = .05$,
22 $d = .33$).

1 **Dual-Task Test.** The I2WE group improved its time on DT compared to ST after the program, while
2 the WMS group did not. There was a significant time effect ($p = .01$) and a significant score*group
3 interaction ($p = .04$) with no group effect. Post Hoc analysis revealed a significant difference between
4 pre- and post-tests for the I2WE group ($p = .01$, $d = 1.02$) and none for the other comparisons.

5 **TMT.** The score*group interaction effect was nearly significant ($p = .06$), but Post Hoc analysis
6 revealed no significant differences.

7 For all cognitive tests, RCI values indicated that changes were unreliable (none RCIs were between -
8 ∞ and -1.645 or between ∞ and 1.645).

9 **Secondary Outcomes**

10 **Physical assessment**

11 The results of the pre- and post-tests for physical abilities and the results on simple time effects from
12 ANOVA are given in Table 3 and those of RCI in Additional file 7. Seven measures had a time effect
13 ($p < .01$ or $p < .001$) but no group effect or time*group interaction effect.

14 For all physical tests, RCI values indicated that changes were unreliable (none RCIs were between -
15 ∞ and -1.645 or between ∞ and 1.645).

16 **Evolution of Break It Handball performance**

17 The I2WE group improved their scores, with a significant time effect ($F(11,66) = 7.27$, $p < .001$,
18 $\eta^2 = 0.55$). Post Hoc analysis indicated a significant difference between score 1 and score 9 ($p = .04$),
19 scores 1 and 10 ($p = .01$), 1 and 11 ($p < .001$), and 1 and 12 ($p < .001$) as indicated in Additional file 8.

20 **The link between the evolution of DT performance and Break It Handball performance.**

1 The evolution score of the game performance (score 12–score 1) score was calculated to test the
2 correlation with the change in cognitive test scores (post-test–pre-test). The change in the percentage
3 of time spent in DT versus ST was positively correlated with the change in game performance ($r = .56$,
4 $p = .01$). There were no other correlations with changes in cognitive tests.

5 **Compliance**

6 *Compliance* was 91.01% (415 sessions out of 456) for the I2WE program versus 85.28% (307 sessions
7 out of 360) for the WMS program. *Perceived pleasure* was significantly higher ($t(32) = 2.34$, $p = .03$,
8 $d = 0.81$) for the I2WE group who scored 63.74 ± 6.04 to 59.57 ± 4.82 for the WMS group. All
9 participants wanted to continue their respective programs. For the I2WE group, 10.53% had a
10 moderate, 36.84% a strong, and 52.63% a very strong desire to continue. For the WMS group, 33.33%
11 had a moderate and 66.67% a strong desire. For physical evolution, 84.21% of the participants in the
12 I2WE group (21.05% very positive) and 93.33% in the WMS group (6.67% neutral) felt a positive
13 change. For cognitive evolution, 78.95% of the I2WE group (21.05% neutral) compared to 20% (80%
14 neutral) of the WMS group felt a positive change. As shown in Table 4, there were no significant
15 differences between effort intensities of the WMS and I2WE groups.

16 **Discussion**

17 This study aimed to compare the cognitive, dual-task, and physical effects of an Immersive and
18 Interactive Wall Exergames program (I2WE) to a Walking and Muscle-Strengthening (WMS) in
19 healthy but inactive older adults.

20 The effects of programs differ on cognitive and DT capacities, with an advantage for the I2WE group
21 with small to medium effects. Two of the 4 executive functions tested (i.e., visuospatial working
22 memory, inhibition) and DT capacity were enhanced only for the I2WE group. The increase of mental
23 flexibility specific to I2WE group participants was also nearly significant ($p = .06$), and visuospatial

1 short-term memory improved for participants in both groups. However, these improvements were not
2 confirmed by RCIs.

3 Since physical activity, moderate-intensity aerobic, and muscle strengthening were the only same
4 stimulation between the 2 groups, their similar improvement in visuospatial short-term memory could
5 be the consequence. Indeed, physical activity enhances cognitive functions [4] and memory [57]. The
6 benefits of aerobic programs on improving cardiorespiratory health are already known, allowing for
7 better cognitive efficiency [58–60] thanks to better oxygenation of the central nervous system and
8 increased neurogenesis through neurotrophic growth factors. Erickson and his collaborators [57]
9 showed a 2% increase in hippocampal volume and BDNF levels, associated with improvement in
10 spatial memory performance after an aerobic program. Thus, changes in visuospatial short-term
11 memory in both groups may have been driven by aerobic physical activity. Several researchers have
12 emphasized that the aerobic effect is more important for executive functions [4], whereas here, only
13 visuospatial short-term memory was improved for both groups. It may have been sensitive more
14 quickly, and the programs may not have been long/high frequency enough to improve the other
15 functions, whereas these 2 are fundamental to have an effect on executive efficiency [61]. In addition,
16 the strength training that all participants received could also mediate the positive changes in
17 visuospatial short-term memory [62,63]. Cassilhas and collaborators [64] found an increase in IGF-1
18 related to resistance training coinciding with improved cognition.

19 Except for one function, the benefits of aerobic physical activity combined with muscle-strengthening
20 did not explain the superior cognitive variation of the I2WE group in inhibition, visuospatial working
21 memory, and DT. Other multidimensional mechanisms (i.e., complex physical training, rich cognitive
22 training, combined training, cooperative and oppositional activity) inherent to the characteristics of
23 I2WE (see Figure 3) could have been responsible.

24 I2WE are complex physical training that solicited coordination, dynamic balance as well as flexibility
25 through large freedom of movement triggered different structural adaptations linked to distinct

1 cognitive functions [65]. Already in 2003, Colcombe and Kramer [4] stated that programs combining
2 endurance, muscle strengthening, flexibility, and balance were the most effective for improving
3 executive functions. The richness of physical stimulation and motor complexity could produce greater
4 cognitive benefits [4,5] and encourage transfer effects [21].

5 In addition to complex physical stimulation, I2WE provide rich cognitive solicitation. Fissler and
6 colleagues [66] stipulated that "*novelty interventions*", including video games that were novel and
7 optimally calibrated in difficulty, combined with high variability of the training task stimulated broad
8 processes. These benefits could be increased by the addition of physical stimulation [66], generating
9 effects on non-specific cognitive skills such as those found in this study. Indeed, the I2WE games
10 solicited diverse cognitive processes, with more than 15 games associated with different gameplay. For
11 example, players were required to retain information to be as efficiently as possible (short-term and
12 working memory), to combine a motor and a cognitive task (DT), to ignore certain details to focus on
13 critical information (inhibition), all while analyzing multiple visuospatial stimuli (visuospatial
14 functions), to switch quickly from one information to another (mental flexibility) as well as to plan
15 their action and strategies (planning). However, these last 3 functions were not improved while these
16 are usually sensitive to training using video games [3] and exergames [10]. Cognitive enrichment
17 could be dependent on the specificity of the video game [3], which implies that these 3 functions were
18 not engaged intensively and/or frequently enough by I2WE. In 2008, Basak and collaborators [67] did
19 not obtain the same increases in visuospatial skills using a strategy game as Green and Bavelier [68]
20 did with action video games. Thus, the characteristics of the games could be directly related to the
21 types of processes that were modified [69]. The transfer was not as general as claimed by Fissler and
22 collaborators' theory [66]. This could explain the discordance in the ability of exergames to improve
23 cognitive functions. Given low levels of evidence from meta-analysis, studies on exergames should
24 focus on an accurate description of the games used and on their specific cognitive stimulation. In
25 addition to cognitive stimulation, the strong immersive capabilities of I2WE provided an adequate
26 training environment. The combination of exergames and immersive virtual environments could

1 increase the efficiency of training on brain health [70]. Immersion could amplify the activity of brain
2 regions associated with executive functions [26] and create higher mental focus than a conventional
3 physical activity session [20]. Furthermore, I2WE is an alternative to head-mounted virtual reality,
4 which can negatively affect older adults, including nausea, dizziness, and malaise [71]. These could be
5 particularly dangerous for older persons, who have a greater propensity for falls and loss of balance.
6 No negative events, including those related to cybersickness, occurred during the program despite
7 repeated exposure to the immersive wall, and the User Experience is positive [18]. This mixed reality
8 offered by the I2WE represent a compromise to increase the efficiency of multidomain training
9 without generating negative events.

10 I2WE offered a physical and cognitive combination through the DT. Several studies have shown that
11 DT performance could be improved when stimulated and strengthened by the transfer of learning to
12 new tasks [72]. In the present study, DT skills were enhanced in the I2WE group, with a high size
13 effect, and were correlated to improved game performance. Thus, it is enhanced by the I2WE practice
14 and necessary to perform in the games. They generated positive effects even when the tasks used in
15 the test differed from the training situations, so-called “*learning to learn*” [73]. These DT abilities are
16 crucial to the quality of life of older adults, especially to maintain walking and preventing falls [1].
17 Combined training created by I2WE could have a synergistic effect and be more efficient in improving
18 cognitive functions in older adults [6]. Indeed, physical activity could “facilitate” neuroplasticity,
19 reflected by both groups' improvement in visuospatial short-term memory. The additional cognitive
20 stimulation of the I2WE group may have “guided” these effects [66], resulting in enhancement in
21 other cognitive processes, such as executive functions and DT. An additional advantage is that the
22 cognitive task was embedded simultaneously with the motor task and could be more promising to
23 increase the cognitive reserve [21,66]. This *Moving While Thinking* approach prevented older adults
24 from using adaptive strategies to prioritize one task over another (posture first phenomenon). They
25 often protect their motor function at the expense of the cognitive task when a situation involves a

1 threat to balance [74], which could affect the intensity of physical and/or motor stimulation. With
2 I2WE, participants were forced to engage in a motor task to perform the cognitive task and vice versa.

3 The original characteristics of this new type of exclusively collective exergames created a social bond
4 between players, thanks to cooperative and oppositional situations. Participants must communicate,
5 develop strategies, and plan together to perform in the games, creating important cognitive
6 engagement.

7 Thanks to its specific characteristics, this new generation of exergames generated cognitive benefits
8 superior to those of conventional physical activity programs. It can be applied for prevention, as was
9 done in this study, and could also be used in rehabilitation, re-education environments (e.g., hospitals),
10 and care facilities (e.g., residences) for older adults.

11 At the physical level, almost all capacities benefited from the programs, with no differences in
12 evolution between the 2 groups. Like conventional exergames [8,75] and physical activity, I2WE
13 improved multiple physical functions. Thus, they are an interesting successor to the insufficient first
14 generation of exergames, some of which did not solicit physical capacities sufficiently to lead to
15 improvements [76,77]. As Janhunen and collaborators [78] reported, exergames have the same
16 physical effect as physical activity. I2WE involve complex motor skills, which could have led to
17 greater improvements in physical function or higher intensity than a conventional program.

18 Nevertheless, the intensity was the desired one (i.e., moderate), and an intensity too high could have
19 negative effects, such as the impossibility of adequately performing the physical task and the cognitive
20 task.

21 Only upper body strength was unchanged for both groups, which may have been due to the lack of
22 solicitation: ball throws for the I2WE group and swinging movements of the arms during walking for
23 the WMS group. Still, as 82.21% of the participants felt, this new exergame improved a wide range of
24 physical functions, like conventional physical training. This is positive for maintaining the daily

1 functioning and quality of life of older adults since muscle strength and physical performance are
2 predictive of decline in daily living activities [79].

3 I2WE training is effective in improving the participants' performance in the games, as they showed a
4 significant increase in the *Break It Handball* game. These encouraging results are accompanied by a
5 very high level of compliance and the desire of all participants, very strong for 52.63%, to continue
6 their program. This may have been due to the perceived pleasure provided by I2WE. While a
7 preliminary study [18] had already shown that these games created a high level of perceived pleasure
8 (60.71/70), these results were confirmed here with an even higher score of perceived pleasure
9 evaluated by the same questionnaire and significantly higher than the WMS group. The collective [80]
10 and the playful aspect of video games were fun factors. In line with the findings of Graves and
11 collaborators [81] showing that the Wii generated more pleasure than conventional physical activity,
12 I2WE are enjoyable and promising alternatives for training.

13 The main limitation of this study concerned the small number of comparative groups, which made it
14 difficult to determine the specific benefits of I2WE multidimensional characteristics. Moreover, to our
15 knowledge, no studies have evaluated the effects of this type of program on older adults. The results
16 obtained were compared with those in the literature on physical activity, cognitive, exergames, and
17 combined training, but not specifically with I2WE. This limited the extent to which the results could
18 be compared to the literature. Furthermore, it would have been interesting to perform retention tests to
19 determine whether this new type of exergame could extend the benefits to a greater extent and/or for a
20 longer time than a conventional program. Sustained effects after the programs need further research, as
21 they play a key role in the context of clinical relevance. Finally, the lack of a passive control group did
22 not counteract the practice effects of the tests. Nevertheless, validated and reliable tests commonly
23 used in this population were selected, and there was a long delay between pre- and post-tests. In
24 addition, differences between pre and post-tests were evaluated with mean comparison tests and
25 interpreted by effect sizes, confidence intervals, and RCIs. However, the results of inferential tests and

1 RCIs were contradictory. On the one hand, inferential statistics and effect sizes can overestimate
2 results, notably due to measurement error and/or practice effects. On the other hand, RCIs can be
3 criticized, mainly because they do not consider the practice effects, test reliability, and variability of
4 results in post-tests. Therefore, the results of this study must be interpreted with caution. More
5 research is needed to determine what constitutes real change. Should change be assessed using
6 inferential statistics or RCI? In any case, improvements were noted in this study, particularly about the
7 dual task, which represents a real-life ability and could be a "true" change, as Duff and colleagues [56]
8 indicated.

9 This study is the first to show the effects of this new generation of exergames on the cognitive and
10 physical health of older adults, although the reliability of the results has yet to be confirmed by further
11 studies. I2WE produced benefits across a wide range of physical functions comparable to a
12 conventional physical activity program and superior in DT, visuospatial working memory, and
13 inhibition with small to medium effects. It suggests that I2WE has advantageous characteristics
14 representing a solution to introduce the practice of regular physical activity into the lives of initially
15 inactive older adults while stimulating their cognitive functions.

1 **Statements and Declarations**

2 **Sources of Funding**

3 No funding was received for this study.

4 **Competing interests**

5 The authors declare no competing interests.

6 **Acknowledgments**

7 The authors thank the city of Montreuil and the company NeoXperiences for providing the premises
8 and the *Neo One* wall.

References

- [1] Pichierri G, Murer K, De Bruin ED. A cognitive-motor intervention using a dance video game to enhance foot placement accuracy and gait under dual task conditions in older adults: a randomized controlled trial. *BMC Geriatr* 2012;12:1–14. <https://doi.org/10.1186/1471-2318-12-74>.
- [2] Daley MJ, Spinks WL. Exercise, Mobility and Aging. *Sports Med* 2000;29:1–12. <https://doi.org/10.2165/00007256-200029010-00001>.
- [3] Bediou B, Adams DM, Mayer RE, Tipton E, Green CS, Bavelier D. Meta-analysis of action video game impact on perceptual, attentional, and cognitive skills. *Psychol Bull* 2018;144:77–110. <https://doi.org/10.1037/bul0000130>.
- [4] Colcombe S, Kramer AF. Fitness Effects on the Cognitive Function of Older Adults: A Meta-Analytic Study. *Psychol Sci* 2003;14:125–30. <https://doi.org/10.1111/1467-9280.t01-1-01430>.
- [5] Torre MM, Temprado J-J. A Review of Combined Training Studies in Older Adults According to a New Categorization of Conventional Interventions. *Front Aging Neurosci* 2022;13:808539. <https://doi.org/10.3389/fnagi.2021.808539>.
- [6] Kraft E. Cognitive function, physical activity, and aging: Possible biological links and implications for multimodal interventions. *Aging Neuropsychol Cogn* 2012;19:248–63. <https://doi.org/10.1080/13825585.2011.645010>.
- [7] Maillot P, Perrot A, Hartley A. Effects of interactive physical-activity video-game training on physical and cognitive function in older adults. *Psychol Aging* 2012;27:589–600. <https://doi.org/10.1037/a0026268>.
- [8] Howes SC, Charles DK, Marley J, Pedlow K, McDonough SM. Gaming for Health: Systematic Review and Meta-analysis of the Physical and Cognitive Effects of Active Computer Gaming in Older Adults. *Phys Ther* 2017;97:1122–37. <https://doi.org/10.1093/ptj/pzx088>.
- [9] Soares VN, Yoshida HM, Magna TS, Sampaio RAC, Fernandes PT. Comparison of exergames versus conventional exercises on the cognitive skills of older adults: a systematic review with meta-analysis. *Arch Gerontol Geriatr* 2021;97:104485. <https://doi.org/10.1016/j.archger.2021.104485>.
- [10] Stanmore E, Stubbs B, Vancampfort D, de Bruin ED, Firth J. The effect of active video games on cognitive functioning in clinical and non-clinical populations: A meta-analysis of randomized controlled trials. *Neurosci Biobehav Rev* 2017;78:34–43. <https://doi.org/10.1016/j.neubiorev.2017.04.011>.
- [11] Kayama H, Nishiguchi S, Yamada M, Aoyama T, Okamoto K, Kuroda T. Effect of a Kinect-based exercise game on improving executive cognitive performance in community-dwelling elderly. *J Med Internet Res* 2014;16:e61. <https://doi.org/10.4108/icst.pervasivehealth.2013.252253>.
- [12] Htut TZC, Hiengkaew V, Jalayondeja C, Vongsirinavarat M. Effects of physical, virtual reality-based, and brain exercise on physical, cognition, and preference in older persons: a randomized controlled trial. *Eur Rev Aging Phys Act* 2018;15:10. <https://doi.org/10.1186/s11556-018-0199-5>.
- [13] Peng H-T, Tien C-W, Lin P-S, Peng H-Y, Song C-Y. Novel Mat Exergaming to Improve the Physical Performance, Cognitive Function, and Dual-Task Walking and Decrease the Fall Risk of Community-Dwelling Older Adults. *Front Psychol* 2020;11:1620. <https://doi.org/10.3389/fpsyg.2020.01620>.
- [14] Gavelin HM, Dong C, Minkov R, Bahar-Fuchs A, Ellis KA, Lautenschlager NT, et al. Combined physical and cognitive training for older adults with and without cognitive impairment: A systematic review and network meta-analysis of randomized controlled trials. *Ageing Res Rev* 2021;66:101232. <https://doi.org/10.1016/j.arr.2020.101232>.
- [15] Ogawa EF, You T, Leveille SG. Potential Benefits of Exergaming for Cognition and Dual-Task Function in Older Adults: A Systematic Review. *J Aging Phys Act* 2016;24:332–6. <https://doi.org/10.1123/japa.2014-0267>.
- [16] Sala G, Semir K, Gobet F. Still No Evidence That Exergames Improve Cognitive Ability: A Commentary on Stanmore et al. (2017). *Neurosci Biobehav Rev* 2021;123:352–3. <https://doi.org/10.1016/j.neubiorev.2019.11.015>.

- [17] Ordnung M, Hoff M, Kaminski E, Villringer A, Ragert P. No Overt Effects of a 6-Week Exergame Training on Sensorimotor and Cognitive Function in Older Adults. A Preliminary Investigation. *Front Hum Neurosci* 2017;11:160. <https://doi.org/10.3389/fnhum.2017.00160>.
- [18] Béraud-Peigné N, Maillot P, Perrot A. The User Experience of an Immersive and Interactive Wall Exergame in Older Adults. *Games Health J* 2022. <https://doi.org/10.1089/g4h.2022.0075>.
- [19] Roure C, Lentillon-Kaestner V, Fargier P. The effects of a Play Lü inter-team competition exergame on adolescents' moderate-to-vigorous physical activity and situational interest, XChange Virtually, England: 2021.
- [20] Martin-Niedecken AL, Rogers K, Turmo Vidal L, Mekler ED, Márquez Segura E. ExerCube vs. Personal Trainer: Evaluating a Holistic, Immersive, and Adaptive Fitness Game Setup. *Proc. 2019 CHI Conf. Hum. Factors Comput. Syst., Glasgow Scotland Uk: ACM; 2019, p. 1–15.* <https://doi.org/10.1145/3290605.3300318>.
- [21] Herold F, Hamacher D, Schega L, Müller NG. Thinking While Moving or Moving While Thinking – Concepts of Motor-Cognitive Training for Cognitive Performance Enhancement. *Front Aging Neurosci* 2018;10:228. <https://doi.org/10.3389/fnagi.2018.00228>.
- [22] Anderson-Hanley C, Barcelos NM, Zimmerman EA, Gillen RW, Dunnam M, Cohen BD, et al. The Aerobic and Cognitive Exercise Study (ACES) for Community-Dwelling Older Adults With or At-Risk for Mild Cognitive Impairment (MCI): Neuropsychological, Neurobiological and Neuroimaging Outcomes of a Randomized Clinical Trial. *Front Aging Neurosci* 2018;10:76. <https://doi.org/10.3389/fnagi.2018.00076>.
- [23] Pellegrini-Laplagne M, Dupuy O, Sosner P, Bosquet L. Effect of simultaneous exercise and cognitive training on executive functions, baroreflex sensitivity, and pre-frontal cortex oxygenation in healthy older adults: a pilot study. *GeroScience* 2022. <https://doi.org/10.1007/s11357-022-00595-3>.
- [24] Finkelstein S, Suma EA. Astrojumper: Motivating Exercise with an Immersive Virtual Reality Exergame. *Presence Teleoperators Virtual Environ* 2011;20:78–92. https://doi.org/10.1162/pres_a_00036.
- [25] Murray EG, Neumann DL, Moffitt RL, Thomas PR. The effects of the presence of others during a rowing exercise in a virtual reality environment. *Psychol Sport Exerc* 2016;22:328–36. <https://doi.org/10.1016/j.psychsport.2015.09.007>.
- [26] Kober SE, Kurzmann J, Neuper C. Cortical correlate of spatial presence in 2D and 3D interactive virtual reality: An EEG study. *Int J Psychophysiol* 2012;83:365–74. <https://doi.org/10.1016/j.ijpsycho.2011.12.003>.
- [27] Swain DP, ACSM, Brawner CA. ACSM's Resource Manual for Guidelines for Exercise Testing and Prescription. Lippincott Williams & Wilkins; 2012.
- [28] Audiffren M, André N, Albinet C. Effets positifs de l'exercice physique chronique sur les fonctions cognitives des seniors : bilan et perspectives. *Rev Neuropsychol* 2011;3:207–25. <https://doi.org/10.3917/rne.034.0207>.
- [29] Smith PJ, Blumenthal JA, Hoffman BM, Cooper H, Strauman TA, Welsh-Bohmer K, et al. Aerobic Exercise and Neurocognitive Performance: A Meta-Analytic Review of Randomized Controlled Trials. *Psychosom Med* 2010;72:239–52. <https://doi.org/10.1097/PSY.0b013e3181d14633>.
- [30] Béraud-Peigné N, Maillot P, Perrot A. Impact d'un programme d'activité physique de coopération et d'opposition sur les fonctions cognitives de personnes âgées. *NPG Neurol - Psychiatr - Gériatrie* 2022;22:149–57. <https://doi.org/10.1016/j.npg.2021.08.002>.
- [31] Da Silva Júnior JLA, Biduski D, Bellei EA, Becker OHC, Daroit L, Pasqualotti A, et al. A Bowling Exergame to Improve Functional Capacity in Older Adults: Co-Design, Development, and Testing to Compare the Progress of Playing Alone Versus Playing With Peers. *JMIR Serious Games* 2021;9:e23423. <https://doi.org/10.2196/23423>.
- [32] Folstein MF, Folstein SE, McHugh PR. "Mini-mental state." *J Psychiatr Res* 1975;12:189–98. [https://doi.org/10.1016/0022-3956\(75\)90026-6](https://doi.org/10.1016/0022-3956(75)90026-6).
- [33] de Souto Barreto P, Ferrandez A-M, Saliba-Serre. B. Questionnaire d'activité physique pour les personnes âgées (QAPPA) : validation d'un nouvel instrument de mesure en langue française. *Sci Sports* 2011;26:11–8. <https://doi.org/10.1016/j.scispo.2010.09.006>.

- [34] Allain P, Chaudet H, Nicoleau S, Etcharry-Bouyx F, Barré J, Dubas F, et al. Étude de la planification de l'action au moyen du test du plan du zoo dans la maladie d'Alzheimer. *Rev Neurol (Paris)* 2007;163:222–30. [https://doi.org/10.1016/S0035-3787\(07\)90393-8](https://doi.org/10.1016/S0035-3787(07)90393-8).
- [35] Wilson BA, Evans JJ, Emslie H, Alderman N, Burgess P. The Development of an Ecologically Valid Test for Assessing Patients with a Dysexecutive Syndrome. *Neuropsychol Rehabil* 1998;8:213–28. <https://doi.org/10.1080/713755570>.
- [36] Oosterman JM, Wijers M, Kessels RPC. Planning or Something Else? Examining Neuropsychological Predictors of Zoo Map Performance. *Appl Neuropsychol Adult* 2013;20:103–9. <https://doi.org/10.1080/09084282.2012.670150>.
- [37] Wechsler D. WMS-III: Wechsler memory scale administration and scoring manual. Harcourt Brace&Co. San Antonio: The Psychological Corporation; 1997.
- [38] Busch R, Farrell K, Lisdahl-Medina K, Krikorian R. Corsi Block-Tapping Task Performance as a Function of Path Configuration. *J Clin Exp Neuropsychol* 2005;27:127–34. <https://doi.org/10.1080/138033990513681>.
- [39] Lo AHY, Humphreys M, Byrne GJ, Pachana NA. Test-retest reliability and practice effects of the Wechsler Memory Scale-III: Test-retest reliability and practice effect in WMS-III. *J Neuropsychol* 2012;6:212–31. <https://doi.org/10.1111/j.1748-6653.2011.02023.x>.
- [40] Stroop JR. Studies of interference in serial verbal reactions. *J Exp Psychol* 1935;18:643–62. <https://doi.org/10.1037/h0054651>.
- [41] Lemay S, Bédard M-A, Rouleau I, Tremblay P-L. Practice Effect and Test-Retest Reliability of Attentional and Executive Tests in Middle-Aged to Elderly Subjects. *Clin Neuropsychol* 2004;18:284–302. <https://doi.org/10.1080/13854040490501718>.
- [42] MacLeod CM. Half a Century of Research on the Stroop Effect: An Integrative Review 1991.
- [43] Vandenberg SG, Kuse AR. Mental Rotations, a Group Test of Three-Dimensional Spatial Visualization. *Percept Mot Skills* 1978;47:599–604. <https://doi.org/10.2466/pms.1978.47.2.599>.
- [44] Reitan RM. Manual for administration and scoring of the Halstead–Reitan of neuropsychological test battery for adults and children. 1979.
- [45] Giovagnoli AR, Del Pesce M, Mascheroni S, Simoncelli M, Laiacona M, Capitani E. Trail making test: normative values from 287 normal adult controls. *Ital J Neurol Sci* 1996;17:305–9.
- [46] Corrigan JD, Hinkeldey NS. Relationships between Parts A and B of the Trail Making Test. *J Clin Psychol* 1987;43:402–9. [https://doi.org/10.1002/1097-4679\(198707\)43:4<402::AID-JCLP2270430411>3.0.CO;2-E](https://doi.org/10.1002/1097-4679(198707)43:4<402::AID-JCLP2270430411>3.0.CO;2-E).
- [47] Sánchez-Cubillo I, Periáñez JA, Adrover-Roig D, Rodríguez-Sánchez JM, Ríos-Lago M, Tirapu J, et al. Construct validity of the Trail Making Test: Role of task-switching, working memory, inhibition/interference control, and visuomotor abilities. *J Int Neuropsychol Soc* 2009;15:438–50. <https://doi.org/10.1017/S1355617709090626>.
- [48] Fournier J, Vuillemin A, Le Cren F. Mesure de la condition physique chez les personnes âgées. Évaluation de la condition physique des seniors : adaptation française de la batterie américaine « Senior Fitness Test ». *Sci Sports* 2012;27:254–9. <https://doi.org/10.1016/j.scispo.2012.07.005>.
- [49] Rikli RE, Jones CJ. Development and Validation of a Functional Fitness Test for Community-Residing Older Adults. *J Aging Phys Act* 1999;7:129–61. <https://doi.org/10.1123/japa.7.2.129>.
- [50] Bohannon RW, Schaubert KL. Test–Retest Reliability of Grip-strength Measures Obtained over a 12-week Interval from Community-dwelling Elders. *J Hand Ther* 2005;18:426–8. <https://doi.org/10.1197/j.jht.2005.07.003>.
- [51] Harris C, Wattles AP, DeBeliso M, Sevene-Adams PG, Berning JM, Adams KJ. The Seated Medicine Ball Throw as a Test of Upper Body Power in Older Adults. *J Strength Cond Res* 2011;25:2344–8. <https://doi.org/10.1519/JSC.0b013e3181ecd27b>.
- [52] Delignières D, Perez S. Le plaisir perçu dans la pratique des APS : élaboration d'un outil d'évaluation. *Rev STAPS* 1998;45:7–18.
- [53] Kendzierski D, DeCarlo KJ. Physical Activity Enjoyment Scale: Two Validation Studies. *J Sport Exerc Psychol* 1991;13:50–64. <https://doi.org/10.1123/jsep.13.1.50>.
- [54] Borg GAV. Psychophysical bases of perceived exertion: *Med Sci Sports Exerc* 1982;14:377–81. <https://doi.org/10.1249/00005768-198205000-00012>.

- [55] Gellish RL, Goslin BR, Olson RE, McDonald A, Russi GD. Longitudinal Modeling of the Relationship between Age and Maximal Heart Rate. *Med Sci Sports Exerc* 2007;39:822–9. <https://doi.org/10.1097/mss.0b013e31803349c6>.
- [56] Duff K. Evidence-Based Indicators of Neuropsychological Change in the Individual Patient: Relevant Concepts and Methods. *Arch Clin Neuropsychol* 2012;27:248–61. <https://doi.org/10.1093/arclin/acr120>.
- [57] Erickson KI, Voss MW, Prakash RS, Basak C, Szabo A, Chaddock L, et al. Exercise training increases size of hippocampus and improves memory. *Proc Natl Acad Sci* 2011;108:3017–22. <https://doi.org/10.1073/pnas.1015950108>.
- [58] Dustman RE, Ruhlman RO, Russell EM, Shearer DE, Bonekat WH, Shigeoka JW, et al. Aerobic Exercise Training and Improved Neuropsychological Function of Older Individuals. *Neurobiology Aging* 1984;5:35–42. [https://doi.org/10.1016/0197-4580\(84\)90083-6](https://doi.org/10.1016/0197-4580(84)90083-6).
- [59] Kramer AF, Hahn S, Cohen NJ, Banich MT, McAuley E, Harrison CR, et al. Ageing, fitness and neurocognitive function. *Nature* 1999;400:418–9. <https://doi.org/10.1038/22682>.
- [60] Rikli RE, Edwards DJ. Effects of a Three-Year Exercise Program on Motor Function and Cognitive Processing Speed in Older Women. *Res Q Exerc Sport* 1991;62:61–7. <https://doi.org/10.1080/02701367.1991.10607519>.
- [61] Hertzog C, Kramer AF, Wilson RS, Lindenberger U. Enrichment Effects on Adult Cognitive Development: Can the Functional Capacity of Older Adults Be Preserved and Enhanced? *Psychol Sci Public Interest* 2008;9:1–65. <https://doi.org/10.1111/j.1539-6053.2009.01034.x>.
- [62] Bherer L, Erickson KI, Liu-Ambrose T. A Review of the Effects of Physical Activity and Exercise on Cognitive and Brain Functions in Older Adults. *J Aging Res* 2013;2013:1–8. <https://doi.org/10.1155/2013/657508>.
- [63] Kelly ME, Loughrey D, Lawlor BA, Robertson IH, Walsh C, Brennan S. The impact of exercise on the cognitive functioning of healthy older adults: A systematic review and meta-analysis. *Ageing Res Rev* 2014;16:12–31. <https://doi.org/10.1016/j.arr.2014.05.002>.
- [64] Cassilhas RC, Viana VAR, Grassmann V, Santos RT, Santos RF, Tufik S, et al. The Impact of Resistance Exercise on the Cognitive Function of the Elderly. *Med Sci Sports Exerc* 2007;39:1401–7. <https://doi.org/10.1249/mss.0b013e318060111f>.
- [65] Voelcker-Rehage C, Niemann C. Structural and functional brain changes related to different types of physical activity across the life span. *Neurosci Biobehav Rev* 2013;37:2268–95. <https://doi.org/10.1016/j.neubiorev.2013.01.028>.
- [66] Fissler P, Küster O, Schlee W, Kolassa I-T. Novelty Interventions to Enhance Broad Cognitive Abilities and Prevent Dementia. *Prog Brain Res* 2013;207:403–34. <https://doi.org/10.1016/B978-0-444-63327-9.00017-5>.
- [67] Basak C, Boot WR, Voss MW, Kramer AF. Can training in a real-time strategy video game attenuate cognitive decline in older adults? *Psychol Aging* 2008;23:765–77. <https://doi.org/10.1037/a0013494>.
- [68] Green CS, Bavelier D. Action video game modifies visual selective attention. *Nature* 2003;423:534–7. <https://doi.org/10.1038/nature01647>.
- [69] Achtman RL, Green CS, Bavelier D. Video games as a tool to train visual skills. *Restor Neurol Neurosci* 2008;26:435–46.
- [70] Huang K-T. Exergaming Executive Functions: An Immersive Virtual Reality-Based Cognitive Training for Adults Aged 50 and Older. *Cyberpsychology Behav Soc Netw* 2020;23:143–9. <https://doi.org/10.1089/cyber.2019.0269>.
- [71] Campo-Prieto P, Cancela JM, Rodríguez-Fuentes G. Immersive virtual reality as physical therapy in older adults: present or future (systematic review). *Virtual Real* 2021;25:801–17. <https://doi.org/10.1007/s10055-020-00495-x>.
- [72] Bherer L, Gagnon C, Langeard A, Lussier M, Desjardins-Crépeau L, Berryman N, et al. Synergistic Effects of Cognitive Training and Physical Exercise on Dual-Task Performance in Older Adults. *J Gerontol Ser B* 2020;76:1533–41. <https://doi.org/10.1093/geronb/gbaa124>.
- [73] Zhang R-Y, Chopin A, Shibata K, Lu Z-L, Jaeggi SM, Buschkuhl M, et al. Action video game play facilitates “learning to learn.” *Commun Biol* 2021;4:1154. <https://doi.org/10.1038/s42003-021-02652-7>.

- [74] Schaefer S, Schumacher V. The Interplay between Cognitive and Motor Functioning in Healthy Older Adults: Findings from Dual-Task Studies and Suggestions for Intervention. *Gerontology* 2011;57:239–46. <https://doi.org/10.1159/000322197>.
- [75] Pacheco TBF, de Medeiros CSP, de Oliveira VHB, Vieira ER, de Cavalcanti FAC. Effectiveness of exergames for improving mobility and balance in older adults: a systematic review and meta-analysis. *Syst Rev* 2020;9:163. <https://doi.org/10.1186/s13643-020-01421-7>.
- [76] Anderson-Hanley C, Maloney M, Barcelos N, Striegnitz K, Kramer A. Neuropsychological Benefits of Neuro-Exergaming for Older Adults: A Pilot Study of an Interactive Physical and Cognitive Exercise System (iPACES). *J Aging Phys Act* 2017;25:73–83. <https://doi.org/10.1123/japa.2015-0261>.
- [77] Taylor LM, Kerse N, Frakking T, Maddison R. Active Video Games for Improving Physical Performance Measures in Older People: A Meta-analysis. *J Geriatr Phys Ther* 2018;41:108–23. <https://doi.org/10.1519/JPT.0000000000000078>.
- [78] Janhunen M, Karner V, Katajapuu N, Niiranen O, Jaakko I, Karvanen J, et al. Effectiveness of Exergame Intervention on Walking in Older Adults: A Systematic Review and Meta-Analysis of Randomized Controlled Trials 2021;101:pzab152. <https://doi.org/10.1093/ptj/pzab152>.
- [79] Wang DXM, Yao J, Zirek Y, Reijnierse EM, Maier AB. Muscle mass, strength, and physical performance predicting activities of daily living: a meta-analysis. *J Cachexia Sarcopenia Muscle* 2020;11:3–25. <https://doi.org/10.1002/jcsm.12502>.
- [80] Freed SA, Sprague BN, Stephan AT, Doyle CE, Tian J, Phillips CB, et al. Feasibility and Enjoyment of Exercise Video Games in Older Adults. *Front Public Health* 2021;9:751289. <https://doi.org/10.3389/fpubh.2021.751289>.
- [81] Graves LEF, Ridgers ND, Williams K, Stratton G, Atkinson G, Cable NT. The Physiological Cost and Enjoyment of Wii Fit in Adolescents, Young Adults, and Older Adults. *J Phys Act Health* 2010;7:393–401. <https://doi.org/10.1123/jpah.7.3.393>.

Figure Legends

Fig. 1. Legend text: *Neo One* device is equipped with a projector system and wall-mounted infrared sensors and is played without a head-mounted display or other equipment. Games are selected via a remote touch screen connected to a computer. Speakers are used to play the music.

Fig. 2. Legend text: I2WE: Immersive and interactive wall exergames; WMS: walking and strengthening.

Fig. 3. No legend text.

Tables

Table 1. Means of variables for I2WE and WMS groups

	I2WE (n = 19)		WMS (n = 15)		t	p
	M	SD	M	SD		
Age (years)	69.63	5.31	70.27	5.82	-0.33	.74
Years of education	12.89	3.91	13.87	3.72	-0.73	.47
BMI (m ² /kg)	25.09	3.67	26.40	4.12	-0.91	.37
Self-rated health (score)	3.68	0.58	3.40	0.63	1.36	.18
RHR (bpm)	59.31	3.92	59.33	6.04	-0.01	.99
HRR (bpm)	98.56	6.54	97.88	7.36	0.28	.78
MMSE (score)	28.95	1.08	28.80	1.01	0.41	.69
Physical activity (MET/week)	287.37	171.56	238.67	225.95	1.37	.18

Abbreviations: BMI = Body Mass Index; HRR = Heart Rate Reserve; I2WE = Immersive and Interactive Wall Exergames; M = Mean; METS = Metabolic Equivalent of Task; MMSE = Mini-Mental State Examination; RHR: Resting Heart Rate; SD = Standard Deviation; WMS = Walking and Muscle-Strengthening.

Table 2. Scores on cognitive and dual-task tests

	I2WE		WMS		Mean difference	95% CI for Mean Difference*		ANOVA			
	Pre-test	Post-test	Pre-test	Post-test		Lower	Upper	Main effect	Interaction effect		
	M (SD)	M (SD)	M (SD)	M (SD)				p	η ²	p	η ²
Zoo Map Test											
<i>Plan formulation</i>											
Total time (s)	214.10 (127.02)	205.13 (144.61)	202.34 (122.55)	196.01 (139.77)	-2.64	-86.83	81.56	.71	.00	.95	.00
Sequence score (n)	3.42 (3.27)	3.52 (8.19)	4.20 (3.49)	5.07 (3.28)	-0.76	-3.37	1.85	.45	.01	.56	.00
<i>Plan execution</i>											
Total time (s)	84.42 (51.07)	81.17 (28.30)	70.98 (38.86)	71.76 (41.90)	-4.02	-31.99	29.94	.86	.00	.77	.00
Sequence score (n)	7.10 (2.23)	6.89 (2.02)	7.67 (0.82)	7.73 (0.46)	-0.28	-1.04	0.49	.70	.00	.47	.00
Total score (n)	10.53 (4.41)	10.42 (3.79)	11.87 (3.58)	12.80 (3.34)	-1.04	-3.56	1.48	.51	.00	.41	.00
Spatial Span test											
Spatial span (n)	6.05 (1.18)	6.68 (1.12)	6.07 (0.88)	6.40 (1.06)	0.30	-0.48	1.08	.02	.05	.44	.00
Backward spatial span (n)	4.53 (1.58)	6.26 (1.66)	4.60 (2.03)	5.00 (1.36)	1.34	0.33	2.34	<.001	.09	.01	.04
Stroop test											
Board 1 (n)	68.79 (19.93)	68.58 (15.84)	63.60 (15.00)	65.80 (15.90)	-2.41	-11.22	6.40	.65	.00	.58	.00
Board 2 (n)	96.58 (26.04)	99.26 (19.99)	99.00 (18.78)	98.67 (18.09)	3.02	-5.41	11.45	.57	.00	.47	.00
Board 3 (n)	37.05 (16.09)	41.42 (14.18)	33.87 (10.53)	32.73 (12.37)	5.66	0.73	10.59	.17	.00	.03	.01
Interference score (n)	0.43 (0.13)	0.49 (0.11)	0.42 (0.09)	0.39 (0.11)	0.08	0.02	0.14	.28	.00	.01	.03
Mental Rotation test											
Score (n)	12.95 (4.43)	14.42 (5.39)	12.07 (5.19)	12.87 (4.94)	0.67	-2.44	3.79	.15	.01	.66	.00
Trail Making Test											
Time A (s)	36.12 (22.03)	35.93 (16.35)	37.94 (13.83)	40.72 (18.86)	-2.97	-12.54	6.60	.59	.00	.53	.00
Times B (s)	106.14 (87.75)	77.96 (38.83)	103.13 (72.68)	109.97 (70.09)	-35.02	-72.70	2.65	.26	.01	.07	.02
Time B-A (s)	70.02 (69.27)	42.03 (31.54)	65.19 (61.37)	69.25 (60.73)	-32.06	-65.93	1.82	.16	.01	.06	.02
Dual Task test											
TUG-dt (n) DT-ST	-0.79 (0.92)	-1.05 (1.22)	-0.93 (1.22)	-1.33 (1.17)	0.14	-0.96	1.23	0.23	.02	.80	.00
TUG-dt (time) DT - ST	2.80 (2.57)	1.17 (1.16)	1.71 (0.93)	1.44 (0.73)	-1.36	-2.71	-0.01	0.01	.08	0.04	.04

Bold values are significant. *95% CI for the mean difference in evolution (post-test – pre-test) between the two groups (I2WE, WMS).

Abbreviations: CI = Confidence Interval; Interaction effect = time*group effect; I2WE = Immersive and Interactive Wall Exergames; Main effect = time effect; M = Mean; n = numbers; SD = Standard Deviation; s = seconds; TUG-dt = Timed Up and Go Dual Task; WMS = Walking and Muscle Strengthening.

Table 3. Scores on physical tests

	I2WE					WMS					ANOVA	
	<i>Pre-test</i>	<i>Post-test</i>	<i>Mean difference</i>	<i>95% CI for Mean Difference</i>		<i>Pre-test</i>	<i>Post-test</i>	<i>Mean difference</i>	<i>95% CI for Mean Difference</i>		<i>Time effect</i>	
	M (SD)	M (SD)		Lower	Upper	M (SD)	M (SD)		Lower	Upper	<i>p</i>	η^2
Chair stand (number)	11.32 (2.43)	13.21 (3.15)	-1.89	-3.46	-0.33	10.20 (2.68)	11.13 (2.20)	-0.93	-2.70	0.83	<.01	.06
Handgrip left (kg)	24.21 (8.19)	25.63 (7.31)	-1.42	-3.27	0.44	22.95 (5.07)	23.25 (5.01)	-0.30	-2.39	1.79	.09	.00
Handgrip right (kg)	23.49 (7.85)	24.19 (7.63)	-0.70	-2.52	1.18	20.64 (3.77)	20.79 (4.11)	-0.15	-2.19	1.90	.39	.00
Chair-sit-and-reach left (cm)	3.10 (11.35)	-2.63 (8.19)	5.74	2.36	9.12	-1.67 (8.63)	-4.60 (10.51)	2.93	-0.87	11.38	<.001	.05
Chair-sit-and-reach right (cm)	3.95 (11.06)	-1.26 (9.24)	5.21	2.46	7.96	-1.36 (8.84)	-4.64 (10.35)	3.29	0.08	6.49	<.001	.04
Timed up and go (s)	5.84 (1.70)	5.44 (1.72)	0.41	-0.05	0.87	5.78 (0.97)	5.39 (0.85)	0.39	-0.13	0.91	<.01	.02
Back scratch right (cm)	7.68 (12.20)	4.00 (9.76)	3.68	-0.29	7.66	7.00 (13.57)	3.20 (11.57)	3.80	-0.67	8.27	<.001	.03
Back scratch left (cm)	13.13 (13.26)	8.68 (11.54)	4.58	1.70	7.46	10.13 (13.21)	8.27 (13.44)	1.87	-1.37	5.10	<.001	.02
Seated Medicine Ball Throw (cm)	261.84 (67.47)	272.32 (65.60)	-10.47	-26.07	5.12	239.27 (41.75)	239.00 (43.48)	0.27	-17.29	17.82	.23	.00
6WT Distance (m)	551.45 (111.17)	588.16 (128.80)	-36.71	-76.86	3.44	525.00 (91.72)	559.00 (88.01)	-34.00	-79.19	11.19	<.01	.03
6WT Intensity (% HRR)	0.52 (0.14)	0.53 (0.15)	-0.01	-0.09	0.06	0.48 (0.14)	0.51 (0.14)	-0.03	-0.12	0.15	.29	.01

Bold values are significant.

Abbreviations: CI = Confidence Interval; HRR = Heart Rate Reserve; I2WE = Immersive and Interactive Wall Exergames; M = Mean; SD = Standard Deviation; WMS = Walking and Muscle-Strengthening; 6WT = 6-minutes Walking Test.

Table 4. Objective and subjective intensities

	I2WE		WMS		t	p
	M	SD	M	SD		
Mean HR (bpm)	105.42	14.13	102.08	13.02	.71	.48
Peak HR (bpm)	127.21	16.94	125.33	16.20	.33	.75
Mean intensity (% HRR)	46.97	12.55	43.90	11.32	.74	.46
Peak intensity (% HRR)	69.01	15.16	67.83	14.60	.23	.82
Mean modified Borg (number)	4.75	0.89	4.59	0.69	.58	.57

Abbreviations: HR = Heart Rate; HRR = Heart Rate Reserve; I2WE = Immersive and Interactive Wall Exergames; M = Mean; SD = Standard Deviation; WMS = Walking and Muscle-Strengthening

Figure 1. Neo One device



Figure 2. Study workflow




Figure 3. *Multidimensional mechanisms from I2WE to explain cognitive improvement*

