

Morse frames Gilles Bertrand, Laurent Najman

▶ To cite this version:

Gilles Bertrand, Laurent Najman. Morse frames. International Conference on Discrete Geometry and Mathematical Morphology (DGMM), S. Brunetti; A. Frosini; S. Rinaldi, Apr 2024, Florence, Italy. hal-04217818v1

HAL Id: hal-04217818 https://hal.science/hal-04217818v1

Submitted on 11 Oct 2023 (v1), last revised 3 Feb 2024 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Morse frames

Gilles Bertrand and Laurent Najman

Univ Gustave Eiffel, CNRS, LIGM, F-77454 Marne-la-Vallée, France {gilles.bertrand,laurent.najman}@esiee.fr

Abstract. In the context of discrete Morse theory, we introduce Morse frames, which are maps that associate a set of critical simplexes to all simplexes. The main example of Morse frames are the Morse references. In particular, these Morse references allow computing Morse complexes, an important tool for homology. We highlight the link between Morse references and gradient flows. We also propose a novel presentation of the Annotation algorithm for persistent cohomology, as a variant of a Morse frame. Finally, we propose another construction, that takes advantage of the Morse reference for computing the Betti numbers in mod 2 arithmetic.

Keywords: Homology \cdot Cohomology \cdot Discrete Morse Theory.

1 Introduction

In this paper, we aim at developing new concepts and algorithm schemes for computing topological invariants for simplicial complexes, such as cycles, cocycles and Betti numbers (Sec. 2). In [2], one of the authors of the present paper, introduces a novel, sequential, presentation of discrete Morse theory [8], termed *Morse sequences.* By adding enough information to Morse sequences, we can compute cycles and cocycles that detect "holes". We achieve our goal by introducing Morse frames (Sec. 3), *i.e.*, maps that associate a set of critical simplexes to each simplex. The main one is called the Morse reference (Sec. 4), and is a by-product of Morse sequences. We discuss the link between reference maps and gradient flows. This leads us to the Morse complex. We then see (Sec. 5) that Morse frames allows for a novel presentation of Annotations [6] for computing persistent cohomology. Then, inspired by the annotation technique, we propose (Sec. 6) a novel, efficient construction for computing Betti numbers in mod 2 arithmetic. We then discuss (Sec. 7) how to implement the notions presented in the paper. Finally, we conclude the paper.

2 Simplicial complexes, homology, and cohomology

2.1 Simplicial complexes

Let K be a finite family composed of finite sets. The family K is a *(simplicial)* complex if we have $\sigma \in K$ whenever $\sigma \neq \emptyset$ and $\sigma \subseteq \tau$ for some $\tau \in K$.

G. Bertrand and L. Najman

An element of a simplicial complex K is a simplex of K or a face of K. A facet of K is a simplex of K that is maximal for inclusion. The dimension of $\sigma \in K$, written $\dim(\sigma)$, is the number of its elements minus one. If $\dim(\sigma) = p$, we say that σ is a *p*-simplex. The dimension of K, written $\dim(K)$, is the largest dimension of its simplices, the dimension of \emptyset , the void complex, being defined to be -1. We denote by $K^{(p)}$ the set composed of all *p*-simplexes of K.

We recall the definitions of the collapses/expansions operators [15].

Let K, L be simplicial complexes. Let $\sigma \in K^{(p)}$, $\tau \in K^{(p+1)}$. The couple (σ, τ) is a free pair for K or a free p-pair for K if τ is the only face of K that contains σ . Thus, τ is necessarily a facet of K. If (σ, τ) is a free pair for K, then $L = K \setminus (\sigma, \tau)$ is an elementary (p-)collapse of K, and K is an elementary (p-)expansion of L. We say that K collapses onto L, or that L expands onto K, if there exists a sequence $\langle K = K_0, \ldots, K_k = L \rangle$ such that K_i is an elementary collapse of $K_{i-1}, i \in [1, k]$.

2.2 Homology and cohomology

Let K be a simplicial complex. We write K[p] for the set composed of all subsets of $K^{(p)}$. Each element of K[p] is a *p*-chain of K. The symmetric difference of two elements of K[p] endows K[p] with the structure of a vector space over the field $\mathbb{Z}_2 = \{0, 1\}$. The set $K^{(p)}$ is a basis for this vector space. With this structure, a chain $c \in K[p]$ may be written as a sum $\sum_{\sigma \in c} \sigma$, with the convention that $\sum_{\sigma \in c} \sigma = 0$ whenever $c = \emptyset$. The sum of two chains is obtained using the modulo 2 arithmetic.

Let K be a simplicial complex. As we are dealing with finite simplicial complex, boundary and coboundary operators can be defined as operators on K[p]. If $\sigma \in K^{(p)}$ we set:

 $\delta(\sigma) = \{ \tau \in K^{(p+1)} \mid \sigma \subset \tau \} \text{ and } \partial(\sigma) = \{ \tau \in K^{(p-1)} \mid \tau \subset \sigma \}.$

The boundary operator $\partial_p : K[p] \to K[p-1]$ is such that, for each $c \in K[p]$, we have $\partial_p(c) = \sum_{\sigma \in c} \partial(\sigma)$. The coboundary operator $\delta^p : K[p] \to K[p+1]$ is such that, for each $c \in K[p]$, we have $\delta^p(c) = \sum_{\sigma \in c} \delta(\sigma)$. We have $\partial_p \circ \partial_{p+1} = 0$ and $\delta^p \circ \delta^{p-1} = 0$.

We define four subsets of K[p] which are vector spaces over \mathbb{Z}_2 :

- the set $Z_p(K)$ of *p*-cycles of K, $Z_p(K)$ is the kernel of ∂_p ;
- the set $B_p(K)$ of *p*-boundaries of K, $B_p(K)$ is the image of ∂_{p+1} ;
- the set $Z^p(K)$ of *p*-cocycles of K, $Z^p(K)$ is the kernel of δ^p ;
- the set $B^p(K)$ of *p*-coboundaries of K, $B^p(K)$ is the image of δ^{p-1} .

Fig. 1 depicts an annulus, with various cycles and cocyles, coloured in blue. In Fig. 1.a, we see a 1-cycle that is the 1-boundary of the two pink triangles. In Fig. 1.b, we have a 1-cycle that is not a 1-boundary. Such a cycle detects a "hole" by "contouring" it. In Fig. 1.c, we see a 1-cocycle which is the 1-coboundary of the four pink points. In Fig. 1.d, we have a 1-cocycle that is not a 1-coboundary. Such a cocycle detects a "hole" by "cutting" the annulus.

We also define the following quotient vector spaces:

 $-H_p(K) = Z_p(K) \setminus B_p(K), \text{ which is the } p^{\text{th}} \text{ homology vector space of } K; \\ -H^p(K) = Z^p(K) \setminus B^p(K), \text{ which is the } p^{\text{th}} \text{ cohomology vector space of } K.$



Fig. 1: An annulus, with various cycles and cocyles. See text for details.

An element h in $H_p(K)$ is such that $h = z + B_p(K)$ for some $z \in Z_p(K)$. We write $h = [z]_p$, which is the homology class of the cycle z.

Similarly, an element h in $H^p(K)$ is such that $h = z + B^p(K)$ for some $z \in Z^p(K)$. We write $h = [z]^p$, which is the cohomology class of the cocycle z.

Let $\beta_p(K) = \dim(H_p(K))$ and $\beta^p(K) = \dim(H^p(K))$. We have $\beta_p(K) = \beta^p(K)$ (See [7, Sec. V.1]). The number $\beta_p(K) = \beta^p(K)$ is the p^{th} Betti number (mod 2) of K.

3 Morse sequences and Morse frames

Let us first introduce the two following basic operators [15].

Let K, L be simplicial complexes. If $\sigma \in K$ is a facet of K, and if $L = K \setminus \{\sigma\}$, we say that L is an elementary perforation of K, and that K is an elementary filling of L.

The notion of a "Morse sequence" [2] may be defined by simply considering expansions and fillings of a simplicial complex.

Definition 1. Let K be a simplicial complex. A Morse sequence (on K) is a sequence $\vec{W} = \langle \emptyset = K_0, \dots, K_k = K \rangle$ of simplicial complexes such that, for each $i \in [1, k], K_i$ is either an elementary expansion or an elementary filling of K_{i-1} .

Let $\overrightarrow{W} = \langle K_0, \dots, K_k \rangle$ be a Morse sequence. For each $i \in [1, k]$:

- If K_i is an elementary filling of K_{i-1} , we write $\hat{\sigma}_i$ for the simplex σ such that $K_i = K_{i-1} \cup \{\sigma\}$. We say that the face σ is critical for \overrightarrow{W} .
- If K_i is an elementary expansion of K_{i-1} , we write $\hat{\sigma}_i$ for the free pair (σ, τ) such that $K_i = K_{i-1} \cup \{\sigma, \tau\}$. We say that $\hat{\sigma}_i, \sigma, \tau$, are regular for \overrightarrow{W} .

We write $\widehat{W} = \langle \hat{\sigma}_1, \ldots, \hat{\sigma}_k \rangle$, and we say that \widehat{W} is a *(simplex-wise) Morse sequence*. Clearly, \overrightarrow{W} and \widehat{W} are two equivalent forms. We shall pass from one of these forms to the other without notice.

There are several ways to obtain a Morse sequence \vec{W} from a given complex K. The two following schemes are basic ones to achieve this goal:

1. The increasing scheme. We build \overline{W} from the left to the right. Starting from \emptyset , we obtain K by iterative expansions and fillings. We say that this scheme is maximal if we make a filling only if no expansion can be made.

G. Bertrand and L. Najman

2. The decreasing scheme. We build \overrightarrow{W} from the right to the left. Starting from K, we obtain \emptyset by iterative collapses and perforations. We say that this

scheme is *maximal* if we make a perforation only if no collapse can be made. See [2, Section 7] for a discussion of the differences between these schemes.

Definition 2. The gradient vector field of a Morse sequence \overrightarrow{W} is the set of all regular pairs for \overrightarrow{W} . We say that two Morse sequences \overrightarrow{W} and \overrightarrow{V} on a given complex K are equivalent if they have the same gradient vector field.

It is worth mentioning that there is no loss of generality when using Morse sequences as a presentation of gradient vector fields. In fact, it may be proved that the gradient vector field of an arbitrary Morse function may be seen as the gradient vector field of a Morse sequence (see [2]).

Let \overrightarrow{W} be a Morse sequence on a simplicial complex K. We write $\ddot{W} = \{\sigma \in K \mid \sigma \text{ is critical for } \overrightarrow{W}\}$ and $\ddot{W}^{(p)} = \{\sigma \in K^{(p)} \mid \sigma \text{ is critical for } \overrightarrow{W}\}$. We write $\ddot{W}[p]$ for the set composed of all subsets of $\ddot{W}^{(p)}$, each $c \in \ddot{W}[p]$ is a *p*-chain of \ddot{W} . We have $\ddot{W}[p] \subseteq K[p]$.

A Morse frame is simply a map which assigns, to each p-simplex of K, a certain set of critical p-simplexes.

Definition 3. Let \overrightarrow{W} be a Morse sequence on a simplicial complex K. We say that Υ is a (Morse) frame on \overrightarrow{W} if Υ is a map such that: $\Upsilon : \sigma \in K^{(p)} \mapsto \Upsilon(\sigma) \in \overrightarrow{W}[p].$

If Υ is a Morse frame on \overline{W} , we also denote by Υ the map: $\Upsilon: c \in K[p] \mapsto \Upsilon(c) \in W[p]$, where $\Upsilon(c) = \sum_{\sigma \in c} \Upsilon(\sigma)$ and $\Upsilon(\emptyset) = 0$.

4 The Morse reference

A Morse complex is a basic tool for efficiently computing simplicial homology using discrete Morse theory. Since a Morse complex is built solely on critical complexes, its dimension is generally much smaller than the one of the original complex. In this section, we introduce two frames which allow simplifying the construction of a Morse complex. In particular, the boundary and co-boundary operators can easily be extracted from these two frames.

4.1 Reference and co-reference

Definition 4. Let \overrightarrow{W} be a Morse sequence and let Υ', Υ'' be two Morse frames on \overrightarrow{W} such that, for each critical simplex σ of \overrightarrow{W} , we have $\Upsilon'(\sigma) = \Upsilon''(\sigma) = \{\sigma\}$. We say that Υ' is the (Morse) reference of \overrightarrow{W} if, for each regular pair (σ, τ) of \overrightarrow{W} , we have $\Upsilon'(\tau) = 0$ and $\Upsilon'(\sigma) = \Upsilon'(\partial(\tau) \setminus \{\sigma\})$. We say that Υ'' is the (Morse) co-reference of \overrightarrow{W} if, for each regular pair (σ, τ) of \overrightarrow{W} , we have $\Upsilon''(\sigma) = 0$ and $\Upsilon''(\tau) = \Upsilon''(\delta(\sigma) \setminus \{\tau\})$.

If Υ is the reference of \overrightarrow{W} , we write Υ^* for the co-reference of \overrightarrow{W} .

 $\mathbf{4}$



Fig. 2: (a) A torus. Points with the same label are identified. (b) A Morse reference map. (c) A Morse co-reference map. See text for details.

Let \overrightarrow{W} be a Morse sequence on K and let $\widehat{W} = \langle \hat{\sigma}_1, \ldots, \hat{\sigma}_k \rangle$. We see that a Morse reference Υ of \overrightarrow{W} may be computed by scanning the sequence \widehat{W} from the left to the right. Also, a Morse co-reference Υ^* of \overrightarrow{W} may be computed by scanning \widehat{W} from the right to the left. The uniqueness of Υ and Υ^* is a consequence of these constructions. As a limit case, observe that:

- If τ is a facet of K, then we have $\Upsilon(\tau) = 0$ whenever τ is not critical.

– If σ is a 0-simplex of K, then we have $\Upsilon^*(\sigma) = 0$ whenever σ is not critical.

Fig. 2.a depicts a two-dimensional torus. We first illustrate, in Fig. 2.b, the Morse reference of a Morse sequence \overrightarrow{T} on this torus obtained by a maximal increasing scheme (depicted in detail in [2, Fig. 1]). In this figure, any simplex σ in grey is such that $\Upsilon(\sigma) = 0$. At the first step, the first critical simplex $\sigma_1 = a$ is coloured in pink, with $\Upsilon(\sigma_1) = a$. After all the possible expansions from a, we have $\Upsilon(\sigma) = a$ (in pink) for all σ of dimension 0. At the next stage, we introduce a first 1-critical simplex σ of dimension 1 highlighted in blue. We then introduce a second critical 1-simplex, c (in green), and we have $\Upsilon(c) = c$; this leads to $\Upsilon(\sigma) = c$ for all simplexes σ of dimension 1 in green. At the penultimate step of the Morse sequence, we have a free pair (σ, τ) , and $\Upsilon(\sigma) = x$, in purple, where $x = \Upsilon(\partial(\tau) \setminus \{\sigma\}) = b + c$. The ultimate step of the Morse sequence is the critical 2-simplex d, and we have $\Upsilon(d) = d$, highlighted in yellow.

Conversely, in Fig 2.c, by scanning \overrightarrow{T} from right to left, we obtain its Morse co-reference map. In this figure, any simplex σ in grey is such that $\Upsilon(\sigma) = 0$. Starting with the critical 2-simplex d, after several steps in the sequence, we have $\Upsilon^*(\sigma) = d$ for all simplexes σ of dimension 2, highlighted in yellow. We then have $\Upsilon^*(\sigma) = c$ for all simplexes σ of dimension 1 highlighted in green, and $\Upsilon^*(\sigma) = b$ for all simplexes σ of dimension 1 highlighted in blue. Finally, we have $\Upsilon^*(\sigma_1) = a$ for the last simplex of dimension 0, which is critical, and is coloured in pink.

It is straightforward to check that Morse references are invariant under the relation of equivalence given in Definition 2:

Proposition 1. Let \overrightarrow{V} and \overrightarrow{W} be two Morse sequences on K. Let Υ_V and Υ_W be the references of \overrightarrow{V} and \overrightarrow{W} , respectively.

1. The sequences \overrightarrow{V} and \overrightarrow{W} are equivalent if and only if $\Upsilon_V = \Upsilon_W$. 2. The sequences \overrightarrow{V} and \overrightarrow{W} are equivalent if and only if $\Upsilon_V^* = \Upsilon_W^*$.

4.2Gradient paths, co-gradient paths and gradient flows

References are closely related to the notion of a gradient path. In the following, we recall the classical definition of such a path. We also introduce the notion of a co-gradient path, which arises naturally from the definition of a co-reference.

Let \overline{W} be a Morse sequence on K.

- 1. Let $\pi = \langle \sigma_0, \tau_0, \dots, \sigma_{k-1}, \tau_{k-1}, \sigma_k \rangle$ be a sequence, with $\sigma_i \in K^{(p)}, \tau_i \in$ $K^{(p+1)}$. We say that π is a gradient path in \overrightarrow{W} (from σ_0 to σ_k) if, for any $i \in [0, k-1]$, the pair (σ_i, τ_i) is regular for \overrightarrow{W} and $\sigma_{i+1} \in \partial(\tau_i)$, with $\sigma_{i+1} \neq \sigma_i$.
- 2. Let $\pi = \langle \tau_0, \sigma_1, \tau_1, \dots, \sigma_k, \tau_k \rangle$ be a sequence, with $\tau_i \in K^{(p+1)}, \sigma_i \in K^{(p)}$. We say that π is a *co-gradient path in* \vec{W} (from τ_0 to τ_k) if, for any $i \in [1, k]$, the pair (σ_i, τ_i) is regular for \vec{W} and $\tau_{i-1} \in \delta(\sigma_i)$, with $\tau_i \neq \tau_{i-1}$.

The following result may be proved by induction, by considering the two scanning processes of \hat{W} that are mentioned above. Proposition 3 may be proved by changing the extremities of gradient and co-gradient paths.

Proposition 2. Let \overrightarrow{W} be a Morse sequence on K and let Υ the reference \overrightarrow{W} . Let σ be a p-simplex in K and ν be a critical p-simplex of \vec{W} .

- 1. We have $\nu \in \Upsilon(\sigma)$ if and only if the number of gradient paths from the simplex σ to the critical simplex ν is odd.
- 2. We have $\nu \in \Upsilon^*(\sigma)$ if and only if the number of co-gradient paths from the critical simplex ν to the simplex σ is odd.

Proposition 3. Let \overrightarrow{W} be a Morse sequence on K and Υ be the reference of \overrightarrow{W} . Let σ be a critical p-simplex of \vec{W} and τ be a critical (p+1)-simplex of \vec{W} . We have $\sigma \in \Upsilon(\partial(\tau))$ if and only if $\tau \in \Upsilon^*(\delta(\sigma))$.

An important concept in discrete Morse theory is the one of gradient flows [8], by which, using Forman's own words [9], loosely speaking, a simplex flows along the gradient paths for infinite time (see also [9] for the dual concept). Gradient flows are a basic ingredient for setting the fundamental property of a Morse complex, that is, the equality of homology between a complex and its Morse complex. In fact, there is a deep link between co-references of a Morse sequence and gradient flows. For the sake of space, we give only an informal presentation of this relation, which may be checked by the interested readers. If τ is a p-simplex of a complex K, the gradient flow which starts from τ is obtained:

1. By considering regular pairs (σ', τ') , with $\sigma' \in \partial(\tau)$. Such a pair may be seen as the beginning of a co-gradient path that starts at τ .

2. By considering some p-simplexes that are in the boundary of a (p + 1)-simplex ν , such that (τ, ν) is a free pair.

If τ is a critical simplex, then the case 2. cannot happen. Also, this case cannot happen for τ' , since τ' belongs to the regular pair (σ', τ') . By induction, we see the gradient flow starting at a critical simplex corresponds exactly to co-gradient paths. Thus, if σ is a simplex of K and τ is a critical simplex for \vec{W} , then:

The simplex σ is in the gradient flow starting at τ if and only if $\tau \in \Upsilon^*(\sigma)$.

It is interesting to compare Υ and Υ^* with the analogous constructions in smooth Morse theory. A gradient flow associates a critical simplex to a chain which is invariant under the flow. According to [9], this chain is the discrete analogue of the unstable (or descending) cell associated to a critical point of a smooth Morse function, and it is obtained with Υ^* . Forman [9] also studies the dual of the flow, the coflow. The coflow maps a critical simplex to a chain that is invariant under the coflow. This chain plays the role of the stable (or ascending) cell associated to a critical point of a smooth Morse function, and it is obtained thanks to Υ .

4.3 The Morse complex

Now, let us consider a boundary map that is restricted to the critical simplexes. This map may be easily built with a Morse reference.

Let Υ be the reference of \overrightarrow{W} . If $\sigma \in \overset{\circ}{W}^{(p)}$, we set $d(\sigma) = \Upsilon(\partial(\sigma))$. We denote by d_p the map:

 $\begin{array}{l} d_p: c \in \ddot{W}[p] \mapsto d_p(c) \in \ddot{W}[p-1], \text{ where } d_p(c) = \varUpsilon(\partial_p(c)). \\ \text{Thus, } d(\sigma) = \sum_{\tau \in \partial(\sigma)} \varUpsilon(\tau) \text{ and } d_p(c) = \sum_{\sigma \in c} d(\sigma). \end{array}$

Theorem 1. Let Υ be the reference of a Morse sequence \overrightarrow{W} on K. For each $c \in K[p]$, we have $d_p(\Upsilon(c)) = \Upsilon(\partial_p(c))$.

Proof. Let $\overrightarrow{W} = \langle \emptyset = K_0, ..., K_k = K \rangle$ be a Morse sequence on K, and let $\widehat{W} = \langle \hat{\sigma}_1, ..., \hat{\sigma}_k \rangle$. We consider the statement (S_i) : For each $c \in K_i[p]$, we have $d_p(\Upsilon(c)) = \Upsilon(\partial_p(c))$. We have $K_0[p] = \{\emptyset\}$. Thus (S_0) holds. Suppose (S_{i-1}) holds with $0 \leq i-1 \leq k-1$. Let $c \in K_i[p]$. 1) Suppose $\hat{\sigma}_i = \sigma$, with $\sigma \in W$. If $\sigma \notin c$, then we are done. Otherwise, we have $c = c' \cup \{\sigma\}$, with $c' \in K_{i-1}[p]$. We have $\partial_p(c) = \partial_p(c') + \partial(\sigma)$. Thus $\Upsilon(\partial_p(c)) = \Upsilon(\partial_p(c')) + \Upsilon(\partial(\sigma))$. By the induction hypothesis and by the definition of $d(\sigma)$, we obtain $\Upsilon(\partial_p(c)) = d_p(\Upsilon(c')) + d(\sigma)$. Therefore $\Upsilon(\partial_p(c)) = d_p(\Upsilon(c')) + d_p(\Upsilon(\{\sigma\})) = d_p(\Upsilon(c))$. 2) Suppose $\hat{\sigma}_i = (\sigma, \tau)$ is a free pair. If $\sigma \notin c$ and $\tau \notin c$, then we are done. 2.1) Suppose $\sigma \in c$. Let $c' = c + \partial_{p+1}(\tau)$. We have $\Upsilon(c') = \Upsilon(c) + \Upsilon(\partial_{p+1}(\tau)) =$ $\Upsilon(c)$. We also have $\partial_p(c') = \partial_p(c) + \partial_p(\partial_{p+1}(\tau)) = \partial_p(c)$. But $c' = (c \setminus \{\sigma\}) + d$, with $d = \{\eta \in \partial_{p+1}(\tau) \mid \eta \neq \sigma\}$. Thus $c' \in K_{i-1}[p]$. By the induction hypothesis, it follows that $d_p(\Upsilon(c')) = \Upsilon(\partial_p(c'))$. By the previous equalities, we obtain $d_p(\Upsilon(c)) = \Upsilon(\partial_p(c))$. 2.2) Suppose $\tau \in c$. Let $c = c' \cup \{\tau\}$, with $c' \in K_{i-1}[p]$. Since $\Upsilon(\tau) = 0$, we obtain $\Upsilon(c) = \Upsilon(c')$. Furthermore $\Upsilon(\partial_p(c)) = \Upsilon(\partial_p(c')) + \Upsilon(\partial(\tau)) = \Upsilon(\partial_p(c'))$. By the induction hypothesis, we have $d_p(\Upsilon(c')) = \Upsilon(\partial_p(c'))$. Therefore $d_p(\Upsilon(c)) = \Upsilon(\partial_p(c))$.

The two following results are direct consequences of Theorem 1.

Proposition 4. Let Υ be the reference of a Morse sequence \overrightarrow{W} on K. For any $c, c' \in K[p]$ we have $\Upsilon(\partial_p(c)) = \Upsilon(\partial_p(c'))$ whenever $\Upsilon(c) = \Upsilon(c')$.

Proof. Let $c, c' \in K[p]$ with $\Upsilon(c) = \Upsilon(c')$. Thus, $d_p(\Upsilon(c)) = d_p(\Upsilon(c'))$. By Theorem 1, we have $\Upsilon(\partial_p(c)) = \Upsilon(\partial_p(c))$.

Proposition 5. If \vec{W} is a Morse sequence, then the maps d_p are boundary operators. That is, we have $d_p \circ d_{p+1} = 0$.

Proof. Let $\sigma \in \ddot{W}^{(p+1)}$. We have $d_{p+1}(\{\sigma\}) = d(\sigma) = \Upsilon(\partial(\sigma)) = \Upsilon(\partial_{p+1}(\{\sigma\}))$. By Theorem 1, we have $d_p(\Upsilon(\partial_{p+1}(\{\sigma\}))) = \Upsilon(\partial_p(\partial_{p+1}(\{\sigma\}))) = \Upsilon(0) = 0$. Thus $d_p \circ d_{p+1}(\{\sigma\}) = 0$, which gives the result by linearity. \Box

Since $d_p \circ d_{p+1} = 0$, the couple $(\ddot{W}[p], d_p)$ satisfies the definition of a *chain* complex [12]. We say that $(\ddot{W}[p], d_p)$ is the Morse (chain) complex of \overrightarrow{W} . This notion of a Morse complex is equivalent to the classical one given in the context of discrete Morse theory. This fact may be verified using [14, Theorem 8.31], Proposition 2, and the very definition of the differential d_p .

Dual results for Th. 1, Prop. 4 and Prop. 5 can be written by considering Υ^{\star} instead of Υ .

In the following, we denote by $H_p(\ddot{W})$ (resp. $H^p(\ddot{W})$) the *pth* homology (resp. cohomology) vector space corresponding to the Morse complex of \overrightarrow{W} . By Theorem 1, the map Υ is a *chain map* [12] from the chain complex $(K[p], \partial_p)$ to the chain complex $(\ddot{W}[p], d_p)$. Hence, Υ induces a linear map between $H_p(K)$ and $H_p(\ddot{W})$; see [12]. Furthermore, we have the following.

Theorem 2 (from [8]). For all p, the vector spaces $H_p(K)$ and $H_p(\tilde{W})$ are isomorphic.

5 Annotations

If σ is a *p*-simplex in a complex *K*, an annotation for σ , as introduced in [4], is a length *g* binary vector, where *g* is the rank of the homology group $H_p(K)$. These annotations, when summed up for simplexes in a given cycle, provide a way to determine the homology class of this cycle. The following definition is an adaptation for Morse sequences of this notion. The main difference is that we annotate each simplex with a subset of the critical simplexes of the sequence, instead of a vector.

Let \overrightarrow{W} be a Morse sequence on K. We say that a Morse frame Υ on \overrightarrow{W} is an annotation on \overrightarrow{W} if Υ satisfies the three conditions:

8

- C1: For each $\sigma \in K^{(p)}$, we have $\Upsilon(\sigma) \subseteq \ddot{V}^{(p)}$ where $\ddot{V}^{(p)}$ is a subset of $\ddot{W}^{(p)}$;
- C2: For each p, we have $Card(\ddot{V}^{(p)}) = \beta_p(K);$
- C3: For any cycles $z, z' \in Z_p(K)$, we have $\Upsilon(z) = \Upsilon(z')$ if and only if their homology classes are such that $[z]_p = [z']_p$.

Let Υ be a frame on \overline{W} . If $\tau \in \overline{W}^{(p)}$, we set $\Upsilon^{\sharp}(\tau) = \{\sigma \in K^{(p)} \mid \tau \in \Upsilon(\sigma)\}$. The following proposition, derived from [6], indicates that an annotation may be seen as a way to determine a cohomology basis of the complex.

Proposition 6 (adapted from [6]). Let \overrightarrow{W} be a Morse sequence on K. A frame Υ on \overrightarrow{W} is an annotation on \overrightarrow{W} if and only if Υ satisfies the conditions C1, C2, and the following condition C4.

C4: The set of chains $\{\Upsilon^{\sharp}(\tau) \mid \tau \in \ddot{V}^{(p)}\}$ is a set of cocycles whose cohomology classes $\{[\Upsilon^{\sharp}(\tau)]^p \mid \tau \in \ddot{V}^{(p)}\}$ constitute a basis of $H^p(K)$.

We give a construction for obtaining an annotation. Again, it is an adaptation for a Morse sequence of the one given in [5] and [6]. Three cases are considered:

- 1. If a critical simplex is added, and if the annotation of the boundary of this simplex is trivial, then a new cycle is created. The label associated to this simplex is composed solely of the simplex itself.
- 2. If a critical simplex is added, and if the annotation of the boundary of this simplex is not trivial, then a cycle is removed. This is done by selecting one label in the annotation of the boundary of this simplex, and by removing this label from all the previous annotations.
- 3. If a free pair is added, we propagate the labels of the annotations to this pair, according to a simple rule.

See [5] and [6] for the validity of this construction for the cases 1 and 2 The validity for the case 3 is an easy consequence of the definition of a free pair.

Let $\overrightarrow{W} = \langle K_0, \dots, K_k \rangle$ be a Morse sequence and $\widehat{W} = \langle \hat{\sigma}_1, \dots, \hat{\sigma}_k \rangle$. We write $\overrightarrow{W}_i = \langle K_0, \dots, K_i \rangle$, $i \in [0, k]$. We consider the sequence $\langle \Upsilon_0, \dots, \Upsilon_i \rangle$, $i \in [0, k]$, such that Υ_i is a frame for \overrightarrow{W}_i , with $\Upsilon_0(\emptyset) = 0$ and:

- 1. If $\hat{\sigma}_i = \sigma_i$ and $\Upsilon_{i-1}(\partial(\sigma_i)) = 0$, then Υ_i is such that $\Upsilon_i(\sigma_i) = \{\sigma_i\}$ and $\Upsilon_i(\tau) = \Upsilon_{i-1}(\tau)$ otherwise.
- 2. If $\hat{\sigma}_i = \sigma_i$ and $\Upsilon_{i-1}(\partial(\sigma_i)) \neq 0$, then we select an arbitrary critical face $\nu \in \Upsilon_{i-1}(\partial(\sigma_i))$. The map Υ_i is such that $\Upsilon_i(\sigma_i) = 0$, $\Upsilon_i(\tau) = \Upsilon_{i-1}(\tau) + \Upsilon_{i-1}(\partial(\sigma_i))$ if $\nu \in \Upsilon_{i-1}(\tau)$, and $\Upsilon_i(\tau) = \Upsilon_{i-1}(\tau)$ otherwise.
- 3. If $\hat{\sigma}_i = (\sigma_i, \tau_i)$, then Υ_i such that $\Upsilon_i(\tau_i) = 0$, $\Upsilon_i(\sigma_i) = \Upsilon_{i-1}(\partial(\tau_i) + \{\sigma_i\})$, and $\Upsilon_i(\tau) = \Upsilon_{i-1}(\tau)$ otherwise.

Under the above construction, each frame Υ_i is an annotation on $\overrightarrow{W_i}$.

Let $\ddot{V}_i^{(p)} = \{\sigma \in K_i \mid \Upsilon_i(\sigma) = \{\sigma\}\}, \ \ddot{V}_i^{(p)}$ is composed of critical faces for $\overrightarrow{W_i}$. For each $\sigma \in K_i^{(p)}$, we have $\Upsilon_i(\sigma) \subseteq \ddot{V}_i^{(p)}$. Furthermore, for each p, we have $Card(\ddot{V}_i^{(p)}) = \beta_p(K_i)$.

The interested reader can check that the reference map of the torus, given in Fig. 2.b, is indeed an annotation. Here, the above case 2 does not happen. Due to space constraint, we do not describe in this paper a more meaningful example. G. Bertrand and L. Najman



Fig. 3: (a) A dunce hat. Points with the same label are identified. (b) A Morse reference map. (c) A Morse co-reference map. See text for details.

6 Computing Betti numbers with the Morse reference

In the construction described in Sec. 5, we have to remove a label from all previous annotations. We now present another construction that reduces the amount of operations required for this task. The basic idea is to use the information given by the reference of a Morse sequence, and to remove labels only for some faces which are in the boundary of critical simplexes. Thus, annotations are not computed for all simplexes, but this construction allows us to obtain the Betti numbers of the complex.

Let \overrightarrow{W} be a Morse sequence on a simplicial complex K, and let Υ be a Morse frame on \overrightarrow{W} . We say that Υ is *perfect* if each Betti number $\beta_p(K)$ is exactly equal to the number of critical *p*-simplexes σ in \overrightarrow{W} such that $\Upsilon(\sigma) = \{\sigma\}$.

A key observation is the following. The Morse reference Υ of \overrightarrow{W} is perfect if, and only if, for any critical simplex σ in \overrightarrow{W} , we have $\Upsilon(\partial(\sigma)) = 0$. In the next construction, we take advantage of this observation to iteratively remove suitable pairs of critical points from the image of Υ .

Let $\overrightarrow{W} = \langle \emptyset = K_0, ..., K_k = K \rangle$ be a Morse sequence on a complex K, and let Υ be the Morse reference of \overrightarrow{W} . We write $\widehat{W} = \langle \hat{\sigma}_1, ..., \hat{\sigma}_k \rangle$. We set:

 $\ddot{W}^+ = \ddot{W} \cup \{ \sigma \in K \mid \sigma \in \partial(\tau) \text{ for some } \tau \in \ddot{W} \}$

We consider the sequence of frames $\Upsilon_0, \ldots, \Upsilon_k$ such that $\Upsilon_0 = \Upsilon$ and:

- 1. If $\hat{\sigma}_i = \sigma_i$ and $\Upsilon_{i-1}(\partial(\sigma_i)) \neq 0$, then we select an arbitrary critical simplex $\nu \in \Upsilon_{i-1}(\partial(\sigma_i))$. The map Υ_i is such that $\Upsilon_i(\sigma_i) = 0$, $\Upsilon_i(\tau) = \Upsilon_{i-1}(\tau) +$
 - $\Upsilon_{i-1}(\partial(\sigma_i))$ if $\tau \in \ddot{W}^+$ and $\nu \in \Upsilon_{i-1}(\tau), \Upsilon_i(\tau) = \Upsilon_{i-1}(\tau)$ otherwise.

2. Otherwise, we have $\Upsilon_i = \Upsilon_{i-1}$.

We then have the following: the Morse frame Υ_k is perfect.

It is easy to check that the Morse reference Υ of the torus, given in Fig. 2.b, is such that $\Upsilon(\partial(\sigma)) = 0$ for all critical simplexes σ . Thus, this Morse reference is perfect, and directly gives the expected Betti numbers (1,2,1) for the torus.

Now, let us consider the Morse reference Υ of the dunce hat that is depicted in Fig. 3.b (see the corresponding Morse sequence in [2, Fig. 2]). The last critical simplex c in the sequence is such that $\Upsilon(\partial(c)) = b + b + b = b$. We thus need to remove both b and c from the set of annotations, in effect "killing" the blue cocycle (or dually, "killing" the blue cycle in Fig. 3.c). We then retrieve the expected Betti numbers (1,0,0) for the dunce hat. This result confirms that the dunce hat is acyclic, although the Morse sequence contains 3 critical simplexes, a, b and c.

7 Implementing Morse frames

In the literature, specific, independent algorithms are designed for computing the gradient vector field [13,1,11], the Morse complex [13,10], or the Betti numbers [6]. The framework of Morse frames shows that we can compute the gradient vector field and the Morse complex simultaneously, in only one pass, and the Betti numbers in two passes.

As long as we can check whether a pair is free in constant time (e.g., for example with cubical complexes and the use of a mask to check the neighborhood of a simplex), the complexity of a Morse sequence is linear in the number of simplexes. When we compute a Morse reference map, we need to maintain a list of labels for each simplex, each label corresponding to a critical simplex. This leads to a complexity in $\mathcal{O}(cn)$, where c is the number of critical simplexes, and n is the number of simplexes of the complex. The Morse reference also has a memory complexity of $\mathcal{O}(cn)$. In contrast, algorithms that compute a Morse complex, such as [13], claim a cubic-time complexity (because they have to run several times on each gradient path); furthermore, such algorithms can only be applied *after* the obtention of a gradient vector field.

The framework of Morse frames allow retrieving the concept of annotations [6]. Currently, annotations are implemented [3] using a Morse sequence with only fillings, *i.e.*, where all simplexes are critics. A key point for efficiency is the ordering of the simplexes: a heuristic is used to try preventing the creation of unnecessary cycles. Morse frames show that, with a simple modification (using, for example, a maximal increasing scheme), the annotation algorithm can take advantage of gradient fields. The ordering of the simplexes provided by such a scheme, avoids the creation of unnecessary cycles, by using expansions and fillings, instead of only fillings.

Section 6 provides an algorithm for computing the Betti numbers in mod 2 arithmetic, that is inspired by annotations. Using the reference map, and considering only the set of critical simplexes and their boundary, we obtain an efficient scheme for computing those Betti numbers.

8 Conclusion

Based on Morse sequences, a novel presentation of discrete Morse theory, we have introduced in this paper Morse frames. Morse frames allow for adding information to a Morse sequence, hence selecting specific simplexes depending on the task at hand. The main example of Morse frames, the Morse reference, is useful for homology. In particular, together with its dual, the Morse co-reference,

they provide the discrete analogue of ascending/stable and descending/unstable cell associated to a critical point of a smooth Morse function. The Morse reference also allows retrieving the Morse chain complex. Using Morse frame, we give a novel presentation of the Annotations. Inspired by these Annotations, we describe an efficient scheme for computing Betti numbers in mod 2 arithmetic.

On the theoretical side, for future work, we aim at providing a proof of Th. 2, that will rely only on the Morse reference. We also intend to generalize the Morse reference for other fields than the mod 2 arithmetic, and to compute persistence with the framework. On a more practical level, we also want to test the proposed algorithms, and to compare their efficiency with respect to the state-of-the-art.

References

- 1. Benedetti, B., Lutz, F.H.: Random discrete Morse theory and a new library of triangulations. Experimental Mathematics **23**(1), 66–94 (2014)
- 2. Bertrand, G.: Morse sequences. Tech. rep., University Gustave Eiffel (2023), https://hal.science/hal-04227281
- Boissonnat, J.D., Dey, T.K., Maria, C.: The Compressed Annotation Matrix: An Efficient Data Structure for Computing Persistent Cohomology. Algorithmica 73(3), 607–619 (2015)
- Busaryev, O., Cabello, S., Chen, C., Dey, T.K., Wang, Y.: Annotating simplices with a homology basis and its applications. In: Scandinavian workshop on algorithm theory. pp. 189–200. LNCS, Springer (2012)
- De Silva, V., Vejdemo-Johansson, M.: Persistent cohomology and circular coordinates. In: Proceedings of the twenty-fifth annual symposium on Computational geometry. pp. 227–236 (2009)
- Dey, T.K., Fan, F., Wang, Y.: Computing topological persistence for simplicial maps. In: Proceedings of the thirtieth annual symposium on Computational geometry. pp. 345–354 (2014)
- 7. Edelsbrunner, H., Harer, J.: Computational Topology an Introduction. American Mathematical Society (2010)
- Forman, R.: Witten-Morse theory for cell complexes. Topology 37(5), 945–979 (1998)
- Forman, R.: Discrete Morse theory and the cohomology ring. Transactions of the American Mathematical Society 354(12), 5063–5085 (2002)
- Fugacci, U., Iuricich, F., De Floriani, L.: Computing discrete Morse complexes from simplicial complexes. Graphical models 103, 101023 (2019)
- Harker, S., Mischaikow, K., Mrozek, M., Nanda, V.: Discrete Morse theoretic algorithms for computing homology of complexes and maps. Foundations of Computational Mathematics 14, 151–184 (2014)
- 12. Hatcher, A.: Algebraic topology. Cambridge University Press, Cambridge (2002)
- Robins, V., Wood, P.J., Sheppard, A.P.: Theory and algorithms for constructing discrete Morse complexes from grayscale digital images. IEEE Transactions on pattern analysis and machine intelligence 33(8), 1646–1658 (2011)
- 14. Scoville, N.A.: Discrete Morse Theory, vol. 90. American Mathematical Soc. (2019)
- 15. Whitehead, J.H.C.: Simplicial spaces, nuclei and m-groups. Proceedings of the London mathematical society 2(1), 243–327 (1939)