



Parametrization of the Coulomb interaction matrix with point-group symmetry

Coraline Letouzé, Guillaume Radtke, Benjamin Lenz, Christian Brouder

► To cite this version:

Coraline Letouzé, Guillaume Radtke, Benjamin Lenz, Christian Brouder. Parametrization of the Coulomb interaction matrix with point-group symmetry. TRIQS Summer School 2023, Aug 2023, Saint-Lambert-des-Bois, France. hal-04218994

HAL Id: hal-04218994

<https://hal.science/hal-04218994v1>

Submitted on 26 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Parametrization of the Coulomb interaction matrix with point-group symmetry

Coraline Letouzé¹ Guillaume Radtke¹ Benjamin Lenz¹ Christian Brouder¹

¹Institut de Minéralogie, de Physique des Matériaux et de Cosmochimie, Sorbonne Université, UMR CNRS 7590, Muséum National d'Histoire Naturelle, 75005 Paris, France



Motivation: beyond spherical symmetry

Coulomb integrals are relevant for spectroscopies and *ab initio* methods on strongly correlated materials

$$U_{ijtu} = \int d\vec{r} d\vec{r}' \phi_i(\vec{r})^* \phi_j(\vec{r}')^* U(\vec{r}, \vec{r}') \phi_t(\vec{r}) \phi_u(\vec{r}')$$

ϕ_i : a one-electron wave function

State of the art: Slater parametrization (1930), derived for $1/r$ potentials.

$$U_{ijtu} = \sum_k c_k(ij, tu) R^k(ij, tu)$$

↑
angular coefficient radial integral
→ but: exact **only** in spherical $SO(3)$ symmetry

Step 1: point-group symmetry

Let ϕ_i, \dots, ϕ_u transform as given irreducible representations (irreps) α, \dots, δ of the point group G . We diagonalize $U^{\alpha\beta\gamma\delta}$ in the basis of irreps $\{\eta\}$ of G (Schur's lemma).

$$U^{\alpha\beta\gamma\delta} = [[C^{\alpha\beta,\eta}]^{-1}] \begin{pmatrix} I^{\eta_1} & & & \\ & I^{\eta_2} & & \\ & & I^{\eta_3} & \\ & & & I^{\eta_3} \end{pmatrix} [C^{\gamma\delta,\eta}]$$

The Clebsch-Gordan coefficients $C^{\alpha\beta,\eta}$ couple the direct product of irreps $\alpha \otimes \beta \rightarrow \eta$


$$I^{\alpha\beta\gamma\delta,\eta} = \frac{1}{[\eta]} \sum_{e=1}^{[\eta]} C^{\alpha\beta,\eta} U^{\alpha\beta\gamma\delta} [C^{\gamma\delta,\eta}]^{-1}$$

→ exact parametrization of $U^{\alpha\beta\gamma\delta}$ for any point group


Step 2: permutation symmetry

The number of parameters can be reduced further by considering the invariance of U_{ijtu} under permutation of the wave functions.

The permutation group \mathcal{P} differs for real and complex wave functions.



real case:
 D_4 group



non-real case:
 $m'm'2'$ group

Projection of the invariants onto the fully-symmetric irrep A_1 of the permutation group:

$$\langle I^\eta \rangle \stackrel{\text{def}}{=} \frac{1}{|\mathcal{P}|} \sum_{\sigma_i \in \mathcal{P}} \sigma_i \triangleright I^\eta = M_{\eta\phi} I^\phi$$

↑ projection matrix
→ minimal set of independent parameters

Structural symmetry breaking


It corresponds to the subduction from the initial point group G (eg. $SO(3)$) to one of its subgroups G' . We relate the G invariants to the G' invariants using isoscalar factors.

$$C_{\alpha\beta,\eta e}^{\ell_1\ell_2,\ell} = \left(\begin{array}{c|c} \ell_1 \ell_2 & \ell \\ \hline \alpha & \beta \end{array} \right) C_{ab,e}^{\alpha\beta,\eta}$$



$$I^{(\ell_1\alpha\ell_2\beta\ell_3\gamma\ell_4\delta,\eta)} = \sum_{\ell} \left(\begin{array}{c|c} \ell_1 \ell_2 & \ell \\ \hline \alpha & \beta \end{array} \right) \left(\begin{array}{c|c} \ell_3 \ell_4 & \ell \\ \hline \gamma & \delta \end{array} \right)^* I^{(\ell_1\ell_2\ell_3\ell_4,\ell)}$$

NiO, a 3d shell in octahedral O_h symmetry


All values are in eV, for the bare Coulomb interaction. Also applicable to screened interactions (eg. via cRPA).



Wannier functions of the Ni 3d shell: a localized and symmetry-adapted basis set for the interacting subspace



O_h invariants



Conclusion and Outlook

- $U \approx \sum_k c_k R^k \rightarrow U = \sum_\eta c_\eta I^\eta$ for any point group
- in progress: DMFT calculations with point-group symmetrized interaction

Miscellaneous

- on arXiv: 2306.10186; accepted for publication in PRB
- C.L. acknowledges funding by a grant from the Fondation CFM pour la Recherche

