



HAL
open science

Review of the early Albian ammonites of the Montmell Formation near Marmellar (Salou-Garraf Basin, Tarragona, Catalonia, Spain)

Josep Anton Moreno-Bedmar, Emmanuel Robert, Rafel Matamales-Andreu, Telm Bover-Arnal

► To cite this version:

Josep Anton Moreno-Bedmar, Emmanuel Robert, Rafel Matamales-Andreu, Telm Bover-Arnal. Review of the early Albian ammonites of the Montmell Formation near Marmellar (Salou-Garraf Basin, Tarragona, Catalonia, Spain). *Carnets de Géologie / Notebooks on Geology*, 2017, 17 (1), pp.1-10. 10.4267/2042/62038 . hal-04220681

HAL Id: hal-04220681

<https://hal.science/hal-04220681>

Submitted on 28 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



**Review of the early Albian ammonites
of the Montmell Formation near Marmellar
(Salou-Garraf Basin, Tarragona, Catalonia, Spain)**

Josep Anton MORENO-BEDMAR¹

Emmanuel ROBERT²

Rafel MATAMALES-ANDREU³

Telm BOVER-ARNAL⁴

Abstract: In this work, we review the ammonites of the Montmell Formation in the Marmellar area housed in the collections of the Museo Geológico del Seminario de Barcelona and the Museu de Geologia de Barcelona. This taxonomic update allows the proper biostratigraphic analysis of the ammonite assemblage and assigns it to the early Albian, *Leymeriella tardefurcata* Zone. The taxonomic analysis of all the studied material allows us to recognize the presence of the following taxa: *Uhligella* sp., *Paregonoceras bassei*, *Hypacanthoplites plesiotypicus*, *Hypacanthoplites milletianus*, *Hypacanthoplites subelegans*, and *Hypacanthoplites* sp. The current work is a step forward in the chronostratigraphic knowledge of the Salou-Garraf Basin in the Catalan Coastal Ranges.

Key-words:

- Marmellar;
- Spain;
- Montmell Formation;
- ammonites;
- early Albian

Citation: MORENO-BEDMAR J.A., ROBERT E., MATAMALES-ANDREU R. & BOVER-ARNAL T. (2017).- Review of the early Albian ammonites of the Montmell Formation near Marmellar (Salou-Garraf Basin, Tarragona, Catalonia, Spain).- *Carnets Geol.*, Madrid, vol. 17, no. 1, p. 1-10.

Résumé : Révision des ammonites d'âge albien inférieur de la Formation de Montmell près de Marmellar (Bassin de Salou-Garraf, Tarragone, Catalogne, Espagne).- Dans cette étude, nous avons réexaminé les ammonites de la Formation de Montmell du secteur de Marmellar déposées dans les collections du Musée géologique du Séminaire de Barcelone et du Musée de Géologie de Barcelone. Cette mise à jour taxinomique a permis une analyse biostratigraphique précise de l'association ammonitique, qui est attribuée à la Zone à *Leymeriella tardefurcata* de l'Albien inférieur. L'analyse taxinomique de l'ensemble du matériel étudié révèle la présence des taxons suivants : *Uhligella* sp., *Paregonoceras bassei*, *Hypacanthoplites plesiotypicus*, *Hypacanthoplites milletianus*, *Hypacanthoplites subelegans* et *Hypacanthoplites* sp. Cette étude représente une étape importante dans la connaissance chronostratigraphique du Bassin de Salou-Garraf dans les Chaînes Côtières Catalanes.

¹ Corresponding author

Instituto de Geología, Universidad Nacional Autónoma de México, Ciudad Universitaria, Coyoacán, 04510, México, D.F. (Mexico)

josepamb@geologia.unam.mx

² Laboratoire de Géologie, UMR 5276 CNRS, Université Claude Bernard - Lyon 1, Campus de la Doua (bât. GEODE), 2, rue Raphaël Dubois, 69622 Villeurbanne Cedex (France)

emmanuel.robert@univ-lyon1.fr

³ Trafalgar 1, Manacor, 07500, Mallorca (Spain)

rafelmatamales@hotmail.com

⁴ Departament de Mineralogia, Petrologia i Geologia Aplicada, Facultat de Ciències de la Terra, Universitat de Barcelona, Martí i Franquès s/n, 08028 Barcelona (Spain)

telm.boverarnal@ub.edu

**Mots-clefs :**

- Marmellar ;
- Espagne ;
- Formation de Montmell ;
- ammonites ;
- Albién inférieur.

Resumen: Revisión de los ammonites del Albiense inferior de la Formación Montmell cerca de Marmellar (Cuenca de Salou-Garraf, Tarragona, Cataluña, España).- En este trabajo se revisan los ammonites de la Formación Montmell del área de Marmellar conservados en las colecciones del Museo Geológico del Seminario de Barcelona y del Museo de Geología de Barcelona. Esta actualización taxonómica permite realizar un preciso análisis bioestratigráfico de la asociación de ammonites asignándola a la Zona *Leymeriella tardefurcata*, del Albiense inferior. El análisis taxonómico del material estudiado permite reconocer la presencia de: *Uhligella* sp., *Paragonoceras bassei*, *Hypacanthoplites plesiotypicus*, *Hypacanthoplites milletianus*, *Hypacanthoplites subelegans* y *Hypacanthoplites* sp. Este trabajo constituye un paso adelante en el conocimiento cronoestratigráfico de la Cuenca de Salou-Garraf de las Cadenas Costeras Catalanas.

Palabras clave:

- Marmellar;
- España;
- Formación Montmell;
- ammonites;
- Albiense inferior.

Resum: Revisió dels ammonits de l'Albià inferior de la Formació Montmell a prop de Marmellar (Conca de Salou-Garraf, Tarragona, Catalunya, Espanya).- En aquest treball es revisen els ammonits de la Formació Montmell de l'àrea de Marmellar conservats en les col·leccions del Museu Geològic del Seminari de Barcelona i del Museu de Geologia de Barcelona. Aquesta actualització taxonòmica permet la correcta anàlisi bioestratigràfica de l'associació d'ammonits assignant-la a la Zona *Leymeriella tardefurcata* de l'Albià inferior. L'anàlisi taxonòmica del material estudiat permet reconèixer la presència d'*Uhligella* sp., *Paragonoceras bassei*, *Hypacanthoplites plesiotypicus*, *Hypacanthoplites milletianus*, *Hypacanthoplites subelegans* i *Hypacanthoplites* sp. Aquest treball constitueix un pas endavant en el coneixement cronoestratigràfic de la Conca de Salou-Garraf de les Cadenes Costaneres Catalanes.

Paraules clau:

- Marmellar;
- Espanya;
- Formació Montmell;
- ammonits;
- Albià inferior.

Introduction

The ammonite record of the Montmell Formation in the Marmellar area, Spain (Fig. 1), was first reported by ALMERA (1895), who concluded that it was Aptian Stage. He collected only one specimen that could be assigned to either the genus *Oppelia* or *Placentoceras*. There is no doubt that the best known and most abundant fossil record of the Marmellar area, and other nearby outcrops of the Montmell Formation, are the corals (e.g., DE ANGELIS D'OSSAT, 1905; BATALLER, 1937), currently under taxonomic review (LÖSER, 2012, 2013, 2014, 2015). The scarce ammonoid record is important in order to calibrate the age of the whole fossil assemblage. BATALLER (1954) described the best known (by the international ammonite-workers community) ammonite of Marmellar, an engonocerotid of the new genus and species *Platiknemiceras bassei*, and assigned it to the Albian. The subsequent work of MARTÍNEZ *et al.* (1994) was the first to provide a more exhaustive list of the ammonite assemblage of the Montmell Formation in the Marmellar area, men-

tioning the genera *Hypacanthoplites* and *Sonneratia*, as well as *Platiknemiceras*. However, in this latter work only one specimen was illustrated: *Platiknemiceras bassei*. MARTÍNEZ *et al.* (1994) assigned the ammonites of the Montmell Formation to the second zone of the lower Albian, *Douvilleiceras mamillatum*. MORENO-BEDMAR *et al.* (2009) also studied the ammonoids of the Montmell Formation, and reported the genus *Hypacanthoplites*, together with the engonocerotid *Platiknemiceras*, and assigned them to the first Albian ammonite zone, *Leymeriella tardefurcata*. However, no ammonoids of this formation were illustrated by MORENO-BEDMAR *et al.* (2009).

In this work, we review the ammonites of the Montmell Formation, Salou-Garraf Basin, collected in the area surrounding the currently abandoned town of Marmellar (Fig. 1). A field-work campaign in the area (2013) focused on ancient croplands where the ammonites were collected in the past, however, no ammonites were found because the croplands were abandoned and quite covered by vegetation. Therefore, in





Figure 1: A) Map of the Iberian Peninsula showing the location of the ammonoid sampling site in the Salou-Garraf Basin of the Catalan Coastal Ranges, as well as the nearby Maestrat Basin in the Iberian Chain. Geographic coordinates in degrees. B) Geological map of the area of Marmellar (Comarca of El Baix Penedès) sampled for ammonites (modified from Institut Cartogràfic de Catalunya, 2005). The sampling sites in the surroundings of Marmellar and Les Ventoses villages are indicated with red stars. Geographic coordinates in UTM.

this work we review the ammonites housed in two collections located at the Museo Geológico del Seminario de Barcelona (MGSB) and the Museu de Geologia de Barcelona (MGB). The detailed taxonomic identification of the ammonite assemblage of Marmellar allowed us to make a proper biostratigraphic analysis. In the literature, the only illustrated species of the Montmell Formation is *Platiknemiceras bassei*. In the current work, the complete ammonoid fauna is illustrated. Important improvements in the chronostratigraphic knowledge of the nearby Maestrat Basin were recently carried out (e.g., GARCIA *et al.*, 2014; BOVER-ARNAL *et al.*, 2016). Contrarily, the current state of knowledge of the chronostratigraphy of the Salou-Garraf Basin in the Catalan Coastal Ranges is considerably poorer. Factors that limit suitable data collection in the Salou-Garraf Basin include the very intense faulting caused by the Neogene extension and the opening of the Mediterranean Sea (GUIMERA & SANTANACH PRAT, 1978; BARTRINA *et al.*, 1992), which considerably complicates the study of stratigraphy, because the stratigraphic cover is intensely fragmented like a puzzle. The progressive abandonment of the farmlands resulting in the overgrowth of the outcrops and making them less accessible for stratigraphic study or fossil col-

lection. The combination of the aforementioned factors makes this basin less attractive for geologists. The current paper intends to update the chronostratigraphic knowledge of the Salou-Garraf Basin.

Stratigraphy

The ammonoids examined in this study come from the Montmell Formation and are of early Albian age. This lithostratigraphic unit is mainly characterized by an alternation of micaceous red clays, sandstones and calcarenites rich in *Mesorbitolina texana*, up to 90 m-thick (CANÉROT *et al.*, 1982). Other common skeletal components contained in those strata are miliolids, other foraminifera, echinoids, rudists, other molluscs, corals, and dasycladacean algae. Dolomitic levels, as well as ferruginous crusts, are locally found in the upper part of the formation (ESTEBAN, 1973). The lithologies and fossil content indicate a shallow subtidal to coastal depositional environment punctuated by continental conditions (ESTEBAN, 1973). The Montmell Formation is laterally equivalent with the lower part of the Escucha Formation in the Maestrat Basin (SALAS & MORENO, 2008; MORENO-BEDMAR *et al.*, 2009; BOVER-ARNAL *et al.*, 2016). The ammonite assemblage reported in Traiguera, from the basal part of the Escucha



◀ **Figure 2:** Chronostratigraphic chart for the upper Aptian, *pro parte*, to the upper Albian of the Salou-Garraf and Maestrat basins including the stratigraphic position of the ammonoids studied herein in the lower part of Montmell Formation and the equivalent ammonoid record from the lower part of Escucha Formation. Dashed in grey, the early Albian recognized ammonite zone *Leymeriella tardefurcata*.

Formation (MORENO-BEDMAR *et al.*, 2008; GARCIA *et al.*, 2014: p. 107), is quite similar to the herein studied ammonoids from the Montmell Formation.

The Montmell Formation is overlain and underlain by the Can Xuech Formation (CANÉROT *et al.*, 1982) and the Pujol Florit lithostratigraphic unit (SALAS & MORENO, 2008), respectively (Fig. 2). The Can Xuech Formation is upper Albian-Cenomanian (CANÉROT *et al.*, 1982), and is represented by a succession of marlstones, marly limestones, limestones and ferroan dolostones, up to 200 m-thick. The fossil content is characteristic of a shallow-water carbonate platform depositional environment and includes orbitolinids, miliolids, other foraminifera, brachiopods, oysters, rudists, other molluscs and corals (BATALLER, 1958; CANÉROT *et al.*, 1982).

The underlying Pujol Florit unit is upper Aptian (SALAS & MORENO, 2008; MORENO-BEDMAR *et al.*, 2009). This stratigraphic interval is a lateral equivalent of the upper part of the Benassal Formation from the Maestrat Basin (SALAS & MORENO, 2008; MORENO-BEDMAR *et al.*, 2009; BOVER-ARNAL *et al.*, 2016). The Pujol Florit unit is composed of a lower 20 m-thick succession of packstone to grainstone decimetric and metric beds of platform carbonates containing abundant orbitolinids and fragments of corals and rudist bivalves. Above, is a 40 m-thick alternation of marlstones with orbitolinids and brachiopods, and decimetric micritic limestone beds. The upper part of the Pujol Florit unit is characterized by a 50 m-thick alternation of marlstones with orbitolinids and micritic limestones with rudists bivalves (SALAS & MORENO, 2008).



Ammonite assemblage and biostratigraphic analysis

Review of the two museum collections allowed the identification of: *Uhligella* sp. (Fig. 3: A1-2, C), *Parengonoceras bassei* (BATALLER, 1954) (Fig. 3: D1-2, E1-2; Fig. 4: A1-2, B1-2), *Hypacanthoplites plesiotypicus* (FRITEL, 1906) (Fig. 3: B1-2, F1-2), *Hypacanthoplites milletianus* (ORBIGNY, 1841) (Fig. 4: D1-2), *Hypacanthoplites subelegans* (BREISTROFFER, 1936) (Fig. 4: E1-2) and *Hypacanthoplites* sp. (Fig. 4: C).


The taxon *Parengonoceras bassei* is well known to be restricted to the early Albian (MARTÍNEZ *et al.*, 1994; BULOT, 2010; BUTJOR, 2010). In the Traiguera and Marmellar areas of Spain, other recent works tend to assign this species to the *Leymeriella tardefurcata* Zone (MORENO-BEDMAR *et al.*, 2008, 2009; GARCIA *et al.*, 2014). This is the most common age assignment of this species, even though, at least in Tunisia, it is also present in the *Douvilleiceras mammillatum* Zone (LATIL, 2011). The species of the genus *Hypacanthoplites* range from the late Aptian to the early Albian, disappearing during the later part of the *Leymeriella tardefurcata* Zone. The *Hypacanthoplites* species, displaying flexuous ribs, belong to the late Aptian, whereas the taxa with straight ribs belong to the early Albian (e.g., AMÉDRO, 1992; ROBERT *et al.*, 2001; REY *et al.*, 2013). Most of the *Hypacanthoplites* specimens from the Marmellar area possess rigid costulation. In the particular case of *Hypacanthoplites milletianus*, it is well established that this is an Albian taxon from the *Leymeriella tardefurcata* Zone (e.g., PEYBERNÈS, 1976; LATIL, 1994; MATRION, 2010). The species *Hypacanthoplites subelegans* is present in Normandy, France, also in the *Leymeriella tardefurcata* Zone (DESTOMBES *et al.*, 1974; MARECHAL, 1994). Within the species of the genus *Hypacanthoplites* of the Montmell Formation, *Hypacanthoplites plesiotypicus* is the only one that possesses slightly flexuous costulation (Fig. 3, B2). The range of this species was reviewed in detail by KENNEDY *et al.* (2000: p. 601, Fig. 7), concluding that it belonged to the late Aptian. The aforementioned authors made a particular proposal for the boundary between the Aptian and Albian stages where some species of the genera *Leymeriella*, among which *L. germanica*, and *Preleymeriella* are included in the Aptian. These forms are usually included in the *Leymeriella tardefurcata* Zone and are considered Albian taxa. However, in the present work, we follow the boundary location as was proposed by REBOULET *et al.* (2011, 2014) which corresponds to the classical conception where the whole *Leymeriella tarde-*

furcata Zone is included in the Albian. When we apply REBOULET's definition, the range of *Hypacanthoplites plesiotypicus* is extended into the lowermost Albian. In summary, we conclude that the ammonite assemblage of the Montmell Formation belongs to the *Leymeriella tardefurcata* Zone of the standard Mediterranean ammonite zonation of REBOULET *et al.* (2014), and must be assigned without a doubt to the early Albian. This age-assignment was previously proposed by MORENO-BEDMAR *et al.* (2009) and it is confirmed in the present work.

Systematic notes

Parengonoceras bassei: CASEY (1961) reviewed the genus *Platknemiceras* and accepted its validity; however, he also remarked that the new genus was actually similar to *Parengonoceras*. Recent works (MORENO-BEDMAR *et al.*, 2008, 2009; BULOT, 2010) have begun to question the validity of the genus *Platknemiceras*, considering it a junior synonym of *Parengonoceras*. This question was analyzed in detail by Jean-Louis LATIL, who established that the genus *Platknemiceras* is indeed a synonym of *Parengonoceras* (LATIL, 2011: p. 348; LATIL & ALY, 2012: p. 161).

Hypacanthoplites subelegans and *Hypacanthoplites plesiotypicus*: these species of *Hypacanthoplites* possess a thick, wide whorl cross-section that differs from the common subrectangular compressed whorl section of the genus. *Hypacanthoplites subelegans*, *Hypacanthoplites plesiotypicus* and others, like *Hypacanthoplites ? buloti* LATIL, 2011, have a subrectangular to subquadrate whorl section, some of which are slightly depressed. This difference makes the identification of these species the most difficult taxonomic assignment of the current work. For this reason, in an initial version of the current paper we misidentified *Hypacanthoplites plesiotypicus* as *Archthoplites*. MARTÍNEZ *et al.* (1994), made a similar mistake for the same reason, which is the unusual wide whorl section of these *Hypacanthoplites*. Among specimens of the Marmellar ammonite assemblage that is housed in the collection of the Museo Geológico del Seminario de Barcelona, MARTÍNEZ *et al.* (1994) also incorrectly identified one specimen as *Sonneratia subglabra*, which probably corresponds to the one here illustrated in Figure 4: E1-2, now determined to be *Hypacanthoplites subelegans*. In the work of MARTÍNEZ *et al.* (1994), the biostratigraphic assignment of the whole assemblage to the *Douvilleiceras mammillatum* Zone was caused by this misidentification, since the majority of the *Sonneratia* species were restricted to this ammonite zone (CASEY, 1965).





◀ **Figure 3:** A1-2: *Uhligella* sp. lateral and ventral views of the specimen MGSB 10781-1, Les Ventoses, Marmellar. B1-2: *Hypacanthoplites plesiotypicus* ventral and lateral views of the specimen MGB 21739, Marmellar. C: *Uhligella* sp. lateral view of the specimen MGSB 10781-2, Les Ventoses, Marmellar. D1-2: *Parengonoceras bassei* lateral and ventral views of the specimen MGSB 1044, Marmellar. E1-2: *Parengonoceras bassei* lateral and ventral views of the specimen MGB 66888, Marmellar. F1-2: *Hypacanthoplites plesiotypicus* lateral and ventral views of the specimen MGB 20319, Marmellar. Scale bar = 1 cm.




Figure 4: A1-2: *Parengonoceras bassei* lateral and ventral views of the holotype MGSB 11478, Marmellar. B1-2: *Parengonoceras bassei* lateral and ventral views of the specimen MGSB 8689, Les Ventoses, Marmellar. C: *Hypacanthoplites* sp. lateral view of the specimen MGSB 10762, Marmellar. D1-2: *Hypacanthoplites milletianus* lateral and ventral views of the specimen MGSB 11032, Marmellar. E1-2: *Hypacanthoplites subelegans* ventral and lateral views of the specimen MGSB 10571, Marmellar. Scale bar = 1 cm.



◀ **Figure 5:** *Hypacanthoplites plesiotypicus* lateral and ventral views of the specimen AC 5 (Andrés CUMBA collection, La Jana, Province of Castelló, E Spain). Scale bar = 1 cm.

Hypacanthoplites plesiotypicus in Traiguera, Maestrat Basin: in the work of GARCIA *et al.* (2014), one of the depicted ammonites was misidentified as *Douvilleiceras* gr. *leightonense*. We reviewed this specimen in the current work and we reassign it to *Hypacanthoplites plesiotypicus* (Fig. 5) because of the similar whorl section and ornamentation pattern found in the Marmellar specimens. This misidentification also happened because of the unusual wide whorl section of this species of *Hypacanthoplites*.

Conclusions

The ammonite assemblage of the Montmell Formation near Marmellar, Salou-Garraf Basin, must be assigned to an early Albian age, the *Leymeriella tardefurcata* Zone.

The detailed taxonomic analysis of all the studied material allows the identification of the following taxa: *Uhligella* sp., *Paregonoceras bassei*, *Hypacanthoplites plesiotypicus*, *Hypacanthoplites milletianus*, *Hypacanthoplites subelegans*, and *Hypacanthoplites* sp. The complete ammonoid assemblage is illustrated for the first time.

Identification of *Hypacanthoplites plesiotypicus* and *Hypacanthoplites subelegans*, constitutes the most challenging taxonomic issue, because these species possess a characteristic wide whorl section, an unusual trait among the species of the genus *Hypacanthoplites*.

The current work is a step forward in the chronostratigraphic knowledge of the Salou-Garraf Basin by means of age-calibration of the lower part of the Montmell Formation.

Acknowledgements

This study was supported by the projects PAPIIT IA104715 (DGAPA, Universidad Nacional Autónoma de México), the Grup de Recerca Reconegut per la Generalitat de Catalunya 2014 SGR 251 "Geologia Sedimentària" and CGL 2015-69805-P (MINECO, FEDER, EU). We appreciate the help of Dr. Jaume GALLEMÍ and Dr. Vicent VICEDO for providing access to the ammonites of the Marmellar area that are housed in the Museu de Geologia de Barcelona, Catalonia, Spain. We appreciate the assistance and access to facilities to review the ammonites of the Marmellar area by Dr. Sebastián CALZADA, director of the Museo Geológico del Seminario de Barcelona, Catalonia, Spain. We are also very grateful to Robert W. SCOTT for linguistic corrections.

Bibliographic references

- ALMERA J. (1895).- Étude stratigraphique du massif crétacé du littoral de la province de Barcelone.- *Bulletin de la Société Géologique de France*, Paris, (3e Série), t. XXIII, p. 564-571.
- AMÉDRO F. (1992).- L'Albien du bassin anglo-parisien : Ammonites, zonation phylétique, séquences.- *Bulletin des Centres de Recherches Exploration-Production elf-Aquitaine*, Pau, vol. 16, no. 1, p. 187-233.
- BARTRINA M.T., CABRERA L., JURADO M.J., GUIMERA J., ROCA E. (1992).- Evolution of the central Catalan margin of the Valencia trough (western Mediterranean).- *Tectonophysics*, vol. 203, p. 219-247.



- BARTRINA M.T., CABRERA L., JURADO M.J., GUIMERA J., ROCA E. (1992).- Evolution of the central Catalan margin of the Valencia trough (western Mediterranean).- *Tectonophysics*, vol. 203, p. 219-247.
- BATALLER J.R. (1937).- La fauna coral·lina del Cretàcic de Catalunya i regions limítrofes.- *Arxius de l'Escola Superior d'Agricultura*, Barcelona, (n. s.), vol. 3, no. 1, p. 1-299.
- BATALLER J.R. (1954).- Los Engonoceratidos en España. In: Homenaje póstumo al Dr. D. FRANCISCO PARDILLO VAQUER.- Facultad de Ciencias de la Universidad de Barcelona, p. 173-178.
- BATALLER J.R. (1958).- El Cretácico del Macizo del Montmell.- *Memorias de la Real Academia de Ciencias y Artes de Barcelona*, (3.a época), vol. 32, no. 16, p. 493-549.
- BOVER-ARNAL T., MORENO-BEDMAR J.A., FRIJIA G., PASCUAL-CEBRIAN E. & SALAS R. (2016).- Chronostratigraphy of the Barremian-Early Albian of the Maestrat Basin (E Iberian Peninsula): Integrating strontium-isotope stratigraphy and ammonoid biostratigraphy.- *Newsletters on Stratigraphy*, Stuttgart, vol. 49, p. 41-68.
- BULOT L.G. (2010).- Appendix. Systematic paleontology of Aptian and Albian ammonites from Southwest Iran. In: VINCENT B., BUCHEM F.S.P. van, BULOT L.G., IMMENHAUSER A., CARON M., BAGHBANI D. & HUC A.Y., Carbon-isotope stratigraphy, biostratigraphy and organic matter distribution in the Aptian-Lower Albian successions of southwest Iran (Dariyan and Kazhdumi formations).- *GeoArabia*, Manama, *Special Publication*, vol. 4, p. 167-195.
- BUJTOR L. (2010).- Systematics, phylogeny and homeomorphy of the Engonoceratidae HYATT, 1900 (Ammonoidea, Cretaceous) and revision of *Engonoceras duboisi* LATIL, 1989.- *Carnets Geol.*, Madrid, vol. 10, no. A08 (CG2010_A08), p. 1-31.
- CANÉROT J., CUNY P., PARDO G., SALAS R. & VILLENA J. (1982).- Ibérica Central-Maestrazgo. In: El Cretácico de España.- Universidad Complutense de Madrid, p. 273-344.
- CASEY R. (1961).- The Cretaceous (Albian) ammonite genus *Platinknemicer* BATALLER.- *The Annals and Magazine of Natural History*, London, (Thirteenth Series), vol. 4, p. 353-357.
- CASEY R. (1965).- The Ammonoidea of the Lower Greensand, Part VI.- *Monograph of the Paleontographical Society*, London, vol. 118 (1964), p. 399-546.
- DE ANGELIS D'OSSAT G. (1905).- Coralli del Cretácico inferiore della Catalogna.- *Palaeontographia Italica*, Pisa, vol. 9, p. 169-251.
- DESTOMBES P., JUIGNET P. & RIOULT M. (1974).- Ammonites de l'Aptien-Albien du Bec de Caux, Normandie (NW France).- *Bulletin de la Société Géologique de Normandie et Amis du Muséum du Havre*, Le Havre, vol. 61 (1973), p. 49-106.
- ESTEBAN M. (1973, unpublished).- Petrología de las calizas cretácicas del sector central de los Catalánides (Provincia de Tarragona y Barcelona).- Ph.D. Thesis, Universitat de Barcelona, 425 p.
- GARCIA R., MORENO-BEDMAR J.A., BOVER-ARNAL T., COMPANY M., SALAS R., LATIL J-L, MARTÍN-MARTÍN J.D., GOMEZ-RIVAS E., BULOT L-G., DELANOY G., MARTÍNEZ R. & GRAUGES A. (2014).- Lower Cretaceous (Hauterivian-Albian) ammonite biostratigraphy in the Maestrat Basin (E Spain).- *Journal of Iberian Geology*, Madrid, vol. 40, no. 1, p. 99-112.
- GUIMERA J.J. & SANTANACH PRAT P.F. (1978).- Sobre la compresión alpina en el sector central de las Cadenas Costeras Catalanas.- *Acta geológica hispánica*, Barcelona, vol. 13, no. 2, p. 33-42.
- Institut Cartogràfic de Catalunya (2005).- Baix Penedès.- Mapa geològic Comarcal de Catalunya 1:50 000.
- KENNEDY W.J., GALE A.S., BOWN P.R., CARON M., DAVEY R.J., GRÖCKE D. & WRAY D.S. (2000).- Integrated stratigraphy across the Aptian-Albian boundary in the Marnes Bleues, at the Col Pré-Guittard, Arnayon (Drôme), and at Tartonne (Alpes-de-Haute-Provence), France: a candidate Global Boundary Stratotype Section and Boundary Point for the base of the Albian Stage.- *Cretaceous Research*, vol. 21, p. 591-720.
- LATIL J.-L. (1994).- Les Lyelliceratinae SPATH, 1921 (Ammonitina, Ammonoidea) de l'Albien inférieur et moyen dans le bassin de Paris et sur les bordures du bassin vocontien : Stratigraphie, paléobiogéographie et taxonomie.- *Géologie Alpine*, Grenoble, no. 20, p. 327-381.
- LATIL J.-L. (2011).- Early Albian ammonites from Central Tunisia and adjacent areas of Algeria.- *Revue de Paléobiologie*, Genève, vol. 30, p. 321-429.
- LATIL J.-L. & ALY M.F. (2012).- *Knemiceras gracile* DOUVILLÉ, 1916: A misunderstood Early Albian ammonite from north Sinai (Egypt), and considerations on the genus *Platinknemicer* BATALLER, 1954.- *Revue de Paléobiologie*, Genève, vol. 31, no. 1, p. 159-170.
- LÖSER H. (2012).- The type of *Aplosmilium vidali* ANGELIS D'OSSAT, 1905 (Scleractinia; Early Cretaceous).- *Treballs del Museu de Geologia de Barcelona*, vol. 18 (2011), p. 5-7.
- LÖSER H. (2013).- Taxonomy and distribution of the Early Cretaceous coral genus *Actinastraeopsis*.- *Neues Jahrbuch für Geologie und Paläontologie Monatshefte*, Stuttgart, vol. 269, no. 2, p. 189-202.
- LÖSER H. (2014).- The genus *Brachyseris* ALLOI-TEAU 1946/47 and remarks on *Latimaeandra raea felixi* ANGELIS D'OSSAT, 1905 (Scleractinia; Cretaceous).- *Treballs del Museu de Geologia de Barcelona*, vol. 20, p. 15-20.



- LÖSER H. (2015).- Remarks on the genus *Angelismilia* REIG, 1988 (Scleractinia, Early Cretaceous).- *Treballs del Museu de Geologia de Barcelona*, vol. 21, p. 29-33.
- MARÉCHAL M. (1994).- Les *Hypacanthoplites* SPATH, 1923 des Argiles à *Bucaillella cayeuxi* de Cauville (76) et leur position stratigraphique.- *Bulletin trimestriel de la Société Géologique de Normandie et des Amis du Muséum du Havre*, Le Havre, vol. 81, no. 2, p. 83-96.
- MARTÍNEZ R., GRAUGES A. & SALAS R. (1994).- Distribución de los ammonites del Cretácico inferior de la Cordillera Costera Catalana e Ibérica Oriental.- *Cuadernos de Geología ibérica*, Madrid, vol. 18, p. 337-354.
- MATRION B. (2010).- Les ammonites. In: COLLETÉ C. (éd.), *Stratotype de l'Albien*.- Publications Scientifiques du Muséum, Paris, Collection "Patrimoine géologique"; Biotope Editions, Mèze; BRGM Editions, Orléans, p. 99-193.
- MORENO-BEDMAR J.A., BULOT L., LATIL J.-L., MARTÍNEZ R., FERRER O., BOVER-ARNAL T. & SALAS R. (2008).- Precisiones sobre la edad de la base de la Fm. Escucha, mediante ammonioideos, en la subcuena de la Salzedella, Cuenca del Maestrat (E Cordillera Ibérica).- *Geo-Temas*, Madrid, vol. 10, p. 1269-1272.
- MORENO-BEDMAR J.A., RAMÍREZ L., COMPANY M., DELANOY G., BOVER-ARNAL T., BULOT L.-G., LATIL J.-L. & SALAS R. (2009).- Biostratigrafía de los ammonites de Can Casanyes. Macizo Del Garraf (Barcelona, España).- *Batalleria*, Barcelona, vol. 14, p. 91-98.
- PEYBERNÈS B. (1976).- Le Jurassique et le Crétacé inférieur des Pyrénées franco-espagnoles, entre la Garonne et la Méditerranée.- Thèse de Doctorat ès-Sciences Naturelles, Université Paul-Sabatier, Toulouse, 459 p.
- REBOULET S., RAWSON P.F. & MORENO-BEDMAR J.A. (reporters), AGUIRRE-URRETA M.B., BARRAGÁN R., BOGOMOLOV Y., COMPANY M., GONZÁLEZ-ARREOLA C., IDAKIEVA STOYANOVA V., LUKENEDER A., MATRION B., MITTA V., RANDRIANALY H., VAŠIČEK Z., BARABOSHKIN E.J., BERT D., BERSAC S., BOGDANOVA T.N., BULOT L.G., LATIL J.-L., MIKHAILOVA I.A., ROPOLO P. & SZIVES O. (2011).- Report on the 4th International Meeting of the IUGS Lower Cretaceous Ammonite Working Group, the "KILIAN Group" (Dijon, France, 30th August 2010).- *Cretaceous Research*, vol. 32, p. 786-793.
- REBOULET S. & SZIVES O. (reporters), AGUIRRE-URRETA B., BARAGÁN R., COMPANY M., IDAKIEVA V., IVANOV M., KAKABADZE M.V., MORENO-BEDMAR J.A., SANDOVAL J. BARABOSHKIN E., ÇAĞLAR M.K., FÖZY I., GONZÁLEZ-ARREOLA C., KENJO S., LUKENEDER A., RAISOSSADAT S.N., RAWSON P.F. & TAVERA J.M. (2014).- Report on the 5th International meeting of the IUGS Lower Cretaceous ammonite working group, the KILIAN Group (Ankara, Turkey, 31st August, 2013).- *Cretaceous Research*, vol. 50, p. 126-137.
- REY J., MORENO-BEDMAR J.-A., BILOTTE M. & MARTÍNEZ R. (2013).- Nouvelles données paléontologiques et stratigraphiques au passage Aptien - Albien dans les Pyrénées ariégeoises (France).- *Carnets Geol.*, Madrid, vol. 13, no. A01 (CG2013_A01), p. 67-81.
- ROBERT E., PEYBERNÈS B. & BULOT L.G. (2001).- Caractérisation d'une nouvelle sous-zone d'ammonites au passage Aptien-Albien dans les "Marnes noires à *Hypacanthoplites*" des Pyrénées espagnoles.- *Géobios*, Villeurbanne, vol. 34, fasc. 1, p. 53-62.
- SALAS R. & MORENO J.A. (2008, unpublished).- Revisió de la geologia i altres treballs complementaris de quatre fulls del MGC25M a l'àrea del massís de garraf fulls Nos.: 420-1-1, 420-1-2, 448-1-1 i 448-1-2. Part I: Estratigrafia dels Materials Aptians.- Institut Geològic de Catalunya, Barcelona, Internal report, 41 p.