

# The consumption of ice-derived resources is associated with higher mercury contamination in an Arctic seabird

Fanny Cusset, Julie Charrier, Guillaume Massé, Mark Mallory, Birgit Braune,

Jennifer Provencher, Gaël Guillou, Philippe Massicotte, Jérôme Fort

## ▶ To cite this version:

Fanny Cusset, Julie Charrier, Guillaume Massé, Mark Mallory, Birgit Braune, et al.. The consumption of ice-derived resources is associated with higher mercury contamination in an Arctic seabird. Environmental Research, 2023, 238, pp.117066. 10.1016/j.envres.2023.117066 . hal-04223391

# HAL Id: hal-04223391 https://hal.science/hal-04223391v1

Submitted on 21 Nov 2023  $\,$ 

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# 1 The consumption of ice-derived resources is associated with

# 2 higher mercury contamination in an Arctic seabird

3 Fanny Cusset<sup>1,2,a\*</sup>, Julie Charrier<sup>2,a\*</sup>, Guillaume Massé<sup>1,3</sup>, Mark Mallory<sup>4</sup>, Birgit Braune<sup>5</sup>, Jennifer

4 Provencher<sup>5</sup>, Gaël Guillou<sup>2</sup>, Philippe Massicotte<sup>1</sup> and Jérôme Fort<sup>2</sup>

5

- 6 <sup>1</sup> Takuvik International Research Laboratory (IRL 3376) ULaval-CNRS, Biolgy Department,
- 7 Laval University, 1045 Avenue de la Médecine, Québec, QC, Canada G1V 0A6
- 8 <sup>2</sup> LIENSs, UMR 7266, CNRS-La Rochelle Université, 2 Rue Olympe de Gouges, 17000 La
  9 Rochelle, France
- <sup>3</sup> LOCEAN, UMR 7159, CNRS, MNHN, IRD, Sorbonne-Université, Station Marine de
   Concarneau, BP225, 29900, Concarneau, France
- <sup>4</sup>Biology Department, Acadia University, 15 University Avenue, Wolfville, NS, B4P 2R6, Canada
   <sup>5</sup> Environment and Climate Change Canada, National Wildlife Research Centre, Carleton
   University, Raven Road, Ottawa, ON, K1A 0H3, Canada
- 15
- 16 \* Co-first author
- 17 <u>aCorresponding author</u>: Fanny Cusset (cussetfanny@gmail.com), Julie Charrier
  18 (julie.charrier@univ-lr.fr)

19

21 Sea ice plays a fundamental role in Arctic marine environments, by driving primary productivity 22 and sustaining ice-associated ecosystems. Simultaneously, sea ice influences the contamination of 23 Arctic marine organisms, by modifying contaminant cycles or their bioavailability. Changes in sea 24 ice conditions could therefore profoundly impact the functioning of Arctic marine food webs and 25 their contamination. Top predators such as seabirds, which are subject to bioaccumulation and 26 biomagnification of contaminants, are particularly exposed. In this context, the present study aims 27 to investigate the influence of sea ice and of the use of ice-derived resources on the contamination 28 of seabirds by mercury (Hg). To this end, eggs of thick-billed murres (Brünnich's guillemots, Uria 29 *lomvia*; n = 60) were collected on Prince Leopold Island (Canadian High Arctic) during four years 30 of varying ice conditions (2010–2013). Trophic tracers (*i.e.*, Highly Branched Isoprenoids, HBIs 31 - an indicator of the use of ice-derived resources; carbon and nitrogen stable isotopes - indicators 32 of foraging habitats and trophic status), as well as total Hg concentrations were quantified. Results 33 showed that feeding on ice-derived resources (as indicated by HBI concentrations) was positively 34 correlated to sea ice cover, and both positively influenced Hg concentrations in murre eggs. 35 However, when testing for the best predictor with model selection, sea ice concentration only 36 drove Hg contamination in murres. This work provides new insights into the role of sea ice and 37 ice-derived resources in the contamination by Hg of Arctic wildlife. Further research is now 38 needed to better understand the relationship between sea ice and Hg contamination in Arctic biota 39 and its underlying mechanisms, but also to identify Hg sources in rapidly changing environmental 40 conditions in the Arctic.

#### 41 KEYWORDS

42 Brünnich's guillemot – Hg – Highly Branched Isoprenoids – Sea ice – Sympagic food web –
43 Thick-billed murre

44

#### 45 FUNDING SOURCES

Funding for sample collections was provided by Environment and Climate Change Canada, the Northern Contaminants Program of Crown-Indigenous Relations and Northern Affairs Canada and Acadia University. FC benefited from scholarships from Natural Sciences and Engineering Research Council (Discovery grant to GM). JC is funded by the project ARCTIC-STRESSORS (French National Research Agency; ANR- 20- CE34- 0006 to JF). The preparation of this paper was supported by the Sentinel North program of Laval University, made possible, in part thanks to funding from the Canada First Research Excellence Fund.

#### 53 1. INTRODUCTION

54 Sea ice plays a fundamental role in shaping Arctic ecosystems, by driving the phenology of marine 55 primary production. During spring, melting sea-ice drives several consecutive blooms of primary 56 producers. In early spring, while ocean surface waters are still covered by ice, a bloom of ice algae 57 represents a major contributor to the total primary production. This can represent up to 57% of the 58 entire primary production in the water column and sea ice in the central Arctic (Gosselin et al., 59 1997), which constitutes the first levels of sympagic (i.e., ice-associated) food webs. Later, a 60 bloom of pelagic phytoplankton occurs in ice-free water, supporting pelagic food webs (Horner 61 and Schrader, 1982; Tremblay et al., 2012). These two consecutive blooms and their phenology 62 are crucial for secondary production and the functioning of entire marine ecosystems in the Arctic, 63 highlighting the potential impacts of melting sea ice under the influence of global warming 64 (Wassmann and Reigstad, 2011).

65 In addition, sea ice plays a central role in the biogeochemical cycle of chemical contaminants, in 66 their bioavailability to marine wildlife, and thus in the contamination of Arctic marine ecosystems. 67 This is the case for mercury (Hg), a non-essential and toxic metal that represents a major threat to 68 Arctic wildlife (AMAP, 2021). Mercury is naturally present in the environment, yet anthropogenic 69 activities release considerable amounts of Hg into the environment (Zhang et al., 2014). Once in 70 the atmosphere, this highly volatile contaminant travels via the global atmospheric circulation and 71 reaches even the most remote oceanic regions, such as the Arctic Ocean (Dastoor et al., 2022). As 72 Hg is highly soluble, it reacts with snow, rain and is easily absorbed by particles (Chételat et al., 73 2022). Therefore, it may be deposited in all ecosystems, including sea ice, where it can be stored 74 for long periods of time (Beattie et al., 2014). Melting sea ice thus represents a major route of Hg 75 deposition to the ocean, where it is assimilated by primary producers and magnifies along the food

76 webs (Campbell et al., 2005). Furthermore, sea ice can influence the Hg cycle by regulating air-77 sea exchanges of Hg species (DiMento et al., 2019) and controlling the photochemical 78 transformation processes of Hg (Zheng et al. 2021). These processes further modify Hg 79 bioavailability and the contamination of Arctic marine organisms. Sea ice can also drive Hg levels 80 in marine predators through changes in their trophic ecology and diet. In marine mammals, 81 different contaminant patterns were observed in different sea ice conditions, which were related to 82 prev shifts (Gaden and Stern, 2010; McKinney et al., 2009). In seabirds, changing ice conditions 83 in northern Hudson Bay has resulted in prey shifts (Gaston et al., 2003) toward a higher 84 consumption of less contaminated prey (Braune et al., 2014; Gaston et al., 2012b). Further, weather 85 patterns over multiple years influence exposure of Arctic seabirds to Hg (Foster et al. 2019), 86 presumably due both to deposition but also release or access to Hg already stored in Arctic archives 87 (e.g., glaciers and permafrost). Understanding how Arctic predators rely on sympagic versus 88 pelagic food webs is crucial, not only to foresee future impacts of climate change and associated 89 sea ice melt on ecosystem structure and dynamics, but also because it could largely alter, directly 90 or indirectly, their exposure to contaminants. This is particularly important for species that are 91 used in long-term contaminant monitoring programs, so that changes in contaminant burdens can 92 be interpreted within the context of climate change (Braune et al., 2014).

Multi-tracer approaches enable to track and distinguish the fingerprint of primary producers (*i.e.*, ice algae and pelagic phytoplankton) throughout the entire food web, revealing the dependency of Arctic consumers to sympagic and pelagic resources (Amiraux et al., 2021; Kohlbach et al., 2016; Yurkowski et al., 2020). Highly branched isoprenoids (HBIs) are lipid biomarkers which are specifically synthesized by a restricted number of diatom species (Belt and Müller, 2013). In the Arctic, a restricted number of ice-associated diatoms belonging to genera *Haslea* and *Navicula* 

99 synthesize IP<sub>25</sub> and diene (a mono- and a di-unsaturated HBI, respectively), which allow the 100 tracking of a sympagic fingerprint through a food web. In contrast, triene (a tri-unsaturated HBI) 101 is mainly synthesized by open-water diatom species, allowing for the tracking of a pelagic 102 fingerprint through a food web (Belt et al., 2013; Brown et al., 2011). In the last decade, these 103 biomarkers have provided key insights into the association with sympagic and pelagic food webs 104 in various consumers (e.g. Amiraux et al., 2023; Brown et al., 2013; Brown and Belt, 2012). 105 Recently, HBIs were combined with carbon ( $\delta^{13}$ C) and nitrogen stable isotopes ( $\delta^{15}$  N) – which 106 are commonly used as proxies of predator trophic ecology: the feeding habitat and the diet, 107 respectively (Kelly, 2000) - to establish an index of sea ice use (i.e., Ice Use Index, Cusset et al., 108 2019).

109 In this context, the present study investigated the relationship between sea ice, the use of sympagic 110 resources and Hg contamination of an Arctic seabird: the thick-billed murre (or Brünnich's 111 guillemot, Uria lomvia, hereafter «murres») breeding in the Canadian high Arctic. Canadian 112 murres, like other marine top predators, show high levels of Hg (Braune et al., 2015, Bond et al., 113 2015; Chastel et al., 2022; Mallory et al., 2015), and rely on sea ice and ice-derived resources when 114 ice conditions are favorable (Cusset et al., 2019). More specifically, we (i) investigated differences 115 in Hg concentrations measured in bird eggs (an indicator of female short-term contamination, 116 Bond and Diamond, 2010) during four years of contrasting ice conditions (between 2010 and 117 2013), and (ii) evaluated the drivers of egg Hg contamination (*i.e.*, ice association and trophic 118 ecology of birds). According to previous studies, we expected an increase of egg Hg concentrations 119 during icier years and hence a positive association between Hg contamination and the reliance on 120 sympagic resources.

#### 121 2. MATERIAL AND METHODS

#### 122 **2.1. Study area, species and egg collection**

The present study was carried out on Prince Leopold Island (Lancaster Sound, hereafter "PLI"; 74°N, 90°W; Figure 1). This site represents one of the largest seabird colonies in the Canadian Arctic, supporting approximately 100,000 pairs of thick-billed murres during the breeding season (Gaston et al., 2012a). On PLI, the breeding season runs as follows: colony attendance in mid-June, egg formation (one single egg per nest) in the vicinity of the colony (*i.e.*, for 15 days on average; Gaston and Nettleship, 1981) until laying in early July, hatching in end-July and chick fledging from late August to September.

130 A total of 60 eggs (15 eggs per year) were collected at the beginning of the breeding season (late 131 June – early July) each year between 2010 and 2013, under appropriate permits as part of a long-132 term monitoring program supported by Environment and Climate Change Canada and the 133 Northern Contaminants Program (e.g., Braune et al., 2016). Eggs were collected from nests located 134 on the eastern and southern cliffs of PLI. Eggs were kept cool in the field and shipped to the 135 National Wildlife Research Centre (NWRC, Ottawa) for processing. Egg contents were 136 homogenized for 10-20 s using either a stainless steel Sorvall Omni Mixer or a Brinkmann 137 Polytron homogenizer. Egg homogenates (1-2 g aliquots) were then stored frozen at -40 °C in 138 acid-rinsed polyethylene vials for chemical analyses.

In seabirds, Hg acquired through feeding accumulates over time in the whole body (including liver, muscle, kidney). In female seabirds, Hg is excreted *via* two pathways: feather moult and egg formation/laying. Egg Hg could thus result from Hg acquired locally during egg formation in the Arctic, but also more globally, during the previous wintering in subarctic regions where Arctic 143 seabirds, including murres, can accumulate large concentrations of Hg (Albert et al., 2021). 144 However, murres are income breeders (*i.e.*, allocation of local nutrient to egg formation; Bond and 145 Diamond, 2010; Gaston and Hipfner, 2020). Thus, murre eggs from PLI were assumed to reflect 146 mainly local diet and Hg contamination of females during egg formation (~two weeks).

147

#### 148 **2.2. Ice conditions**

149 As a high Arctic colony, PLI is surrounded by a dense sea-ice cover in winter that progressively 150 breaks up from west to east in spring and summer. Across years, accessibility to open waters for 151 birds to forage is highly variable, as the ice edge position can range from the colony as far east as 152 the Lancaster Sound Polynya in northern Baffin Bay (> 250 km) (Gaston et al., 2005). Sea-ice 153 conditions were considered during four sampling years (from 2010 to 2013). Since weather 154 patterns over multiple years influence Hg exposure of Arctic birds (Foster et al., 2019), ice 155 conditions during the previous years were also considered to investigate potential time-lags. 156 Because murres exhibit a local feeding range (0-150 km; Gaston et al., 2013), Sea Ice 157 Concentrations (SIC) were calculated daily for the overall breeding season (between May 1 and 158 August 31), over a 9-year period (2006–2013), in a 100 km radius around PLI. SIC were expressed 159 as proportion of ice versus open water pixels, using satellite data from the National Oceanic and 160 Atmospheric Administration dataset 161 (ftp://sidads.colorado.edu/DATASETS/NOAA/G02186/shapefiles/4km/); 4 km resolution raster 162 images). Since egg formation was the critical phase investigated in this study, SIC were compared

163 for June 15 over the years. The 4-year target period exhibited an appreciable sea ice gradient during

164 murre egg formation, from highest SIC in 2012 (latest ice break-up) to the lowest SIC in 2011 165 (earliest ice break-up) as follows: 2012 > 2010 > 2013 > 2011 (Table 1).

166

#### 167 **2.3. Chemical analyses**

#### 168 2.3.1. Trophic tracers

169 Highly Branched Isoprenoid (HBI) analyses were carried out at Laval University (Québec), using 170 established techniques (Belt et al., 2012), and were published in Cusset et al. (2019). Briefly, egg 171 aliquots were freeze-dried for 48 h prior to lipid extraction. Internal standards (7-hexylnonadecane 172 and 9-octylheptadec-8-ene, 10 µL, 9.68 µg/ml each) were added to egg homogenates to quantify 173  $IP_{2,5}$  and blanks were analysed with each sample set (n=14) to guarantee analytical accuracy. 174 Total lipids were extracted four consecutive times with dichloromethane:methanol mixture (2:1, 3 175 min vortex, 15 min sonication). Once dried under a gentle stream of N<sub>2</sub>, total lipid extracts were saponified (4 mL, 5% MeOH/H2 O, 90/10, 90°C, 1h). Non-saponifiable lipids (NSL) were then 176 177 extracted from the saponification mixture by liquid-liquid extraction (hexane, 3 x 2 mL), 178 evaporated (nitrogen stream  $< 35^{\circ}$ C) and purified using open column chromatography (SiO<sub>2</sub> 179 50g/g NSL, hexane), to yield an apolar lipid fraction containing HBIs. HBIs were then analyzed 180 by gas chromatography-mass spectrometry (GC-MS) and quantified by measuring mass spectral 181 intensities for each HBI (IP<sub>2 5</sub>: m/z 350.3, diene: m/z 348.3, triene:m/z 346.3). Finally, HBI 182 relative concentrations were combined into the H-Print index (Brown et al., 2014) using Eq. 1:

184 where H–Print indicates a range of association to sympagic (low H-Print), mixed (intermediate)
185 and pelagic (high H-Print) resources.

186 Stable isotope analyses were conducted at the Littoral, Environment and Societies (LIENSs) Joint 187 Research Unit and its stable isotope facility (La Rochelle) on delipidized samples loaded into tin 188 cups (0.2 to 0.8 mg dry weight). Results were previously published in Cusset et al. (2019). Briefly, 189 stable isotope abundances were determined using an elemental analyzer (Flash EA 1112, Thermo Scientific, Milan, Italy) coupled in continuous flow mode to an isotope ratio mass spectrometer 190 191 (Delta V Plus with a Conflo IV interface, Thermo Scientific, Bremen, Germany). Results are 192 expressed in the  $\delta$  notation as deviations from standards (Vienna Pee Dee Belemnite for  $\delta^{13}$ C and N<sub>2</sub> in air for  $\delta^{15}$ N) following the formula:  $\delta^{13}$ C or  $\delta^{15}$ N = [(R<sub>sample</sub>/R<sub>standard</sub>) - 1] x 10<sup>3</sup>, where R is 193 <sup>13</sup>C/<sup>12</sup>C or <sup>15</sup>N/<sup>14</sup>N, respectively. Replicate measurements of laboratory standards (USGS-61 and 194 195 USGS-62) indicated that the measurement uncertainty was < 0.15 % for both  $\delta^{15}$  N and  $\delta^{13}$ C 196 values. Based on the distinct  $\delta^{13}$ C signatures of ice algae and phytoplankton in the Arctic (Hobson 197 et al. 1995),  $\delta^{13}$ C isotopic values and H-Print were combined to create an index of sea-ice use, the 198 Ice Use Index (see Cusset et al., 2019 for detailed methodology).

#### 199 2.3.2. Mercury analyses

Analyses of total Hg (thereafter 'Hg') in egg homogenates were carried out at the NWRC (Ottawa) and were published in Braune et al. (2016). Briefly, samples were homogenized, freeze-dried, homogenized again and weighed into nickel combustion boats. They were then analysed for Hg by direct combustion of the solid sample in an oxygen-rich atmosphere (Salvato and Pirola, 1996), using an Advanced Mercury Analyser (AMA-254, Altec) equipped with an autosampler. Moisture content was determined by weight loss during freeze-drying. One or two blank samples were analysed with each sample set to determine analytical accuracy, as well as standard reference materials (SRMSs). Practical detection limits were 0.006 and 0.004  $\mu$ g/g dry weight for the 2010-2012 samples and the 2013 samples, respectively. All Hg concentrations are expressed in  $\mu$ g/g dry weight (dw). For further analytical details, including reference materials and recoveries, the reader is referred to Braune et al. (2016).

#### 211 **2.4. Statistical analyses**

All statistical analyses and figures were computed using R version 4.0.3 (R Core Team, 2020). First, inter-annual differences in egg Hg concentrations were tested using one-way ANOVA. Data homogeneity and residuals normality were checked prior interpretation, using Breush-Pagan (package «lmtest»; Zeileis and Hothorn, 2002) and Shapiro tests, respectively. In addition, differences in ice regimes (extensive- *versus* reduced-ice years) were tested using Welch twosample t-tests.

218 Then, drivers of egg Hg concentrations were investigated following two steps. First, simple linear 219 regressions were used to investigate independently the relationships between egg Hg 220 concentrations and (i) ice concentrations, (ii) ice association (Ice Use Index) and (iii)  $\delta^{15}$  N values. 221 Second, we used general linear models (GLMs) to investigate the role of these explanatory 222 variables and their combination on Hg concentrations in murre eggs («nlme» package; Pinheiro et 223 al., 2020). Correlations between variables were checked prior to model construction («cor» 224 function). Explanatory variables were standardized (mean = 0, SD = 1) to facilitate comparison of 225 effect sizes. Model assumptions were checked visually with diagnostic functions («plot» function). 226 All potential combinations of variables (Table 2) were subjected to model selection using the 227 «AICcmodavg» package (Mazerolle, 2020). Models were compared using Akaike's Information 228 Criterion corrected for small sample sizes (AIC<sub>c</sub>). The model with the lowest AIC<sub>c</sub> value was 229 selected as the best model (Burnham and Anderson, 2002). Model support was assessed using 230 Akaike weights (w<sub>i</sub>), following Johnson and Omland (2004) and model fit was checked by residual 231 analysis.

During data exploration, time-lags were investigated following previous findings (Foster et al., 2019). Available ice data for the present dataset only dated back to 2006, implying a maximum time-lag of four years for sea ice (instead of seven years as showed by Foster et al., 2019). Given the lack of statistical improvement when added in the models, time-lags were excluded from final analyses.

237

#### 238 **3. RESULTS**

#### 239 **3.1.** Hg concentrations in changing ice conditions

Analyses of Hg concentrations in eggs varied significantly among years (p<0.0001,  $F_{3,56} = 7.4$ , ANOVA) : 2012 ( $1.2 \pm 0.2 \mu g/g$ ; heavy dense ice) differed from 2011 ( $0.9 \pm 0.2 \mu g/g$ ; ice-free) and 2013 ( $1.0 \pm 0.2 \mu g/g$ ; low ice), and 2010 ( $1.1 \pm 0.2 \mu g/g$ ; heavy patchy ice) differed from 2011 (Figure S1). Hg concentrations were lower during reduced-ice years (2011 and 2013;  $0.92 \pm 0.19$  $\mu g/g$ , n=30) than during extensive-ice years (2010 and 2012;  $1.16 \pm 0.23 \mu g/g$ , n=30) (p<0.0001,  $t_{56.7}=4.6$ , Welch two-sample t-test; Figure S2). Hence, Hg concentrations increased by 30% between the minimum and maximum ice years (Table 1).

#### 247 **3.2.** Drivers of egg Hg contamination in changing ice conditions

Unifactorial analyses showed that Hg concentrations in murre eggs increased linearly with increasing ice concentrations (R<sup>2</sup>=0.25, p<0.0001, y=0.0037x + 0.78), and with increasing Ice Use Index (R<sup>2</sup>=0.19, p=0.001, y=0.082 x + 1.0; Figure 2), but not with  $\delta^{15}$  N values (p=0.08) (Figure 3). Multifactorial analyses revealed that the best model only included ice concentrations as an explanatory variable (Table 2; Figure 4). Effect size for this model was -0.14 ± 0.03 (-0.2, -0.08; Gamma inverse).

254

#### 255 4. DISCUSSION

Climate change has a strong impact on sea ice in the Arctic, for instance its distribution, extent, thickness and phenology (Garcia-Soto et al., 2021). Yet, melting sea ice could in turn play a significant role in the contamination by Hg of Arctic marine biota and human populations, through changes in Hg biogeochemistry or modifications of the food web structure (AMAP, 2022; Chételat et al., 2022). By using a new approach combining multiple trophic tracers, Hg analyses and sea ice data, we highlighted the influence of sea ice and sympagic food webs on Hg contamination in murres during the early breeding season.

In years with more extensive sea ice, murre eggs contained higher concentrations of Hg (Figure 4). When considering two distinct ice regimes with relatively "warm" (in 2011 and 2013) and "cold" (in 2010 and 2012) years characterized by reduced and extensive ice cover, respectively, Hg concentrations were higher during cold years compared to warm ones (Figure S2). These results are in agreement with previous investigations using different ice metrics (Mallory and Braune, 2018). Contrasting Hg levels could result from different foraging ecologies and consumed prey according to ice conditions. The main route of Hg uptake in predators is through their diet

270 (Atwell et al., 1998; Campbell et al., 2005; Power et al., 2002; Thompson et al., 1998). As pursuit-271 divers, murres at PLI feed extensively on fish, such as Arctic cod (Boreogadus saida), capelin 272 (Mallotus villosus) and sandlance (Ammodytes sp.), as well as on macrozooplankton, including 273 amphipods (Hyperiidae and Gammaridae) and copepods (*Calanus spp.*; Gaston and Nettleship, 274 1981). Higher Hg contamination during extensive ice years could therefore result from predation 275 on (i) larger prev species which have bioaccumulated more Hg in their tissues and/or (ii) prev with higher  $\delta^{15}$  N values resulting in higher Hg contamination (biomagnification; Dietz et al., 1996). 276 277 Following this hypothesis, murres'  $\delta^{15}$  N values should directly influence their Hg contamination. However, there was a weak relationship between egg Hg concentrations and  $\delta^{15}$  N isotopic values 278 279 (Figure 3), suggesting that temporal changes in trophic ecology alone could not explain higher Hg 280 contamination during icier years, a result consistent with previous findings (Braune et al. 2016, 281 Moody et al. 2012, Provencher et al. 2012).

282 Alternatively, higher Hg concentrations in murres during icy years could be the consequence of a 283 higher contamination of sympagic food webs, compared to the more pelagic ones. Indeed, our 284 results showed that egg Hg contamination was positively related to the Ice Use Index (Figure 2), 285 which reflects bird association with sympagic resources. Hence, murres relying more heavily on 286 ice-derived resources during the early breeding season exhibited higher Hg concentrations. Similar 287 findings were observed in polar bears (Ursus maritimus), which exhibited higher Hg 288 concentrations when feeding on food webs dominated by sea-ice associated diatoms (Routti et al. 289 2012). Higher Hg concentrations were also documented in Arctic cod, an emblematic Arctic fish 290 species that is closely tied to sea ice (Kohlbach et al. 2017). Arctic cod, the main prey of murres 291 in the High Arctic, especially at PLI (Provencher et al., 2012), use under-ice habitats for part of 292 their life cycle (David et al. 2016, Gradinger & Bluhm 2004), for instance as a refuge to escape visual predators (LeBlanc et al., 2019). Marginal ice zones (MIZ) thus represent an important foraging ground for murres (Bradstreet and Cross, 1982). Yet, high Hg concentrations were observed in MIZ, below the productive surface layer (Heimbürger et al. 2015), enhancing Hg uptake and bioavailability in ice-associated food webs up to top predators and seabirds such as murres. Consequently, foraging in these productive habitats may also expose murres to higher Hg in their diet.

299 Sea ice is composed of different semi-solid matrices, such as brines, gases, particles, and biota 300 (e.g., bacteria, viruses, algae). These matrices are subject to physical, biological, and chemical 301 processes which could influence Hg contamination in Arctic marine ecosystems during years of 302 extensive ice (Steffen et al., 2008; Thomas and Dieckmann, 2009). Sea-ice permeability, shaped 303 by temperatures and the creation of brine pockets and drainage channels, considerably influences 304 Hg processes through changing redox conditions and the presence of sea-ice biota (Thomas and 305 Dieckmann, 2009). Indeed, Hg methylation has been documented in sea ice (Beattie et al., 2014) 306 and Hg concentrations were the highest in brines (Chaulk et al., 2011), which represent the primary 307 habitat for microbial communities that sustain sympagic food webs (Thomas and Dieckmann, 308 2002). Brines provide more Hg-enriched environments for primary producers (Schartup et al. 309 2020), including ice algae, potentially increasing food web contamination (Stern et al. 2012). Sea 310 ice could act as an important entry point for Hg at the base of sympagic food webs, additionally 311 enhancing murre Hg contamination when relying on sympagic resources. Thus, sources of Hg 312 found in murre eggs could either originate from melting sea ice and snow where Hg can be stored 313 (Zdanowicz et al., 2013) or from air-ocean exchanges of Hg species (Lalonde et al., 2002; Poulain 314 et al., 2007).

315 Despite the existing relationship between Hg contamination in murre eggs and their use of 316 sympagic resources, model selection revealed that the best predictor for egg Hg concentrations was sea ice concentration only, and did not include the Ice Use Index nor the  $\delta^{15}$  N. This suggests 317 318 that the processes involved are more complex than the reliance on sympagic resources alone. Hg 319 contamination in murres can result from long-term processes with a timescale of years, decades 320 and even centuries (UNEP, 2018). For instance, environmental factors such as the North Atlantic 321 Oscillation (NAO), snow cover or air temperatures were previously identified as drivers of Hg 322 contamination in murre eggs, with a lag time of several years (Foster et al., 2019). Mercury can 323 indeed remain up to one year in the atmosphere (Streets et al., 2019), before deposition and its 324 resulting methylation, assimilation and final transfer along the food chain. Mercury dynamics are 325 driven by a large number of interacting post-depositional processes, which control the transport, 326 methylation and biological uptake of Hg stored in the Arctic Ocean (Dastoor et al., 2022; Outridge 327 et al., 2008). Thus, reduced sea ice cover or earlier ice break-up may expose murres to lower Hg 328 contamination if this was the only change underway. However, air and ocean warming actually 329 increase local export of Hg to Arctic marine areas from environmental archives found largely on 330 land (Chételat et al., 2022; Dastoor et al., 2022). For this reason, it is difficult to disentangle all the 331 processes involved here and one might expect a period of flux and unpredictability, at least until 332 Arctic Hg archives decrease their contribution to marine environments.

Considering its long half-life, Hg is highly persistent in the body of living organisms. Although eggs mainly reflect local diet and Hg contamination in income-breeders like murres (i.e., in the Arctic, during egg formation), they could also integrate, to a low extent, Hg accumulated outside the Arctic (i.e., during the non-breeding season) and remobilized from storing tissues during egg synthesis. Previous studies have reported carry-over effects of adult winter contamination on the subsequent summer Hg concentrations in blood of little auks (Carravieri et al., 2023), as well as higher Hg contamination during the non-breeding season relative to the breeding season in many alcids, including thick-billed murres (Albert et al., 2021). Future research should now further investigate how the reliance of Arctic seabirds to ice-derived resources during the non-breeding season can expose them to Hg and other chemical contaminants.

343

#### 344 **5. CONCLUSION**

345 Sea ice is a key component of Arctic marine environments, where it drives primary productivity 346 and shape the phenology of ice algae and pelagic phytoplankton. At the base of Arctic food webs, 347 these primary producers provide the vital energy pulses for the entire marine food web to thrive. 348 Simultaneously, they assimilate contaminants, such as Hg, which are then transferred to all 349 consumers and predators along the food web. In this study, association with sea ice, and more 350 particularly the reliance on sympagic resources, was positively related to Hg concentrations in 351 eggs of thick-billed murres from the Canadian High Arctic. This work provides new insights into 352 the role of sea ice and ice-derived resources in Hg contamination for Arctic wildlife and human 353 populations that heavily depend on marine resources for subsistence. Further research is now 354 required to fully understand the role of ice-derived resources and sea-ice melt in future Hg 355 contamination of Arctic biota under climate change, as well as the ecological and biogeochemical 356 interactions involved.

357

#### 359 ACKNOWLEDGMENTS

The authors are thankful to the numerous people involved in field work over the years. Logistical support for the fieldwork out of Resolute Bay was provided by the Polar Continental Shelf Program of Natural Resources Canada. Special thanks are due to Caroline Guilmette for her support during HBI analyses, and to Céline Albert and Alice Carravieri for statistical guidance during data analyses. We acknowledge the use of the NOAA satellite dataset. This study also represents a contribution to the French Polar Institute ADACLIM research program (IPEV Pgr 388).

366

#### 367 ADDITIONAL INFORMATION

368 Competing of interest. Authors have no conflict of interest to declare.

#### 369 Contributions.

- 370 Conception and design: FC, JF, GM, MM
- 371 Laboratory work and data acquisition: FC (trophic tracers, ice parameters), BB (mercury), GG
- 372 (stable isotopes) and PM (ice analyses)
- 373 Data analysis: FC
- 374 Interpretation: FC, JC, JF, GM, MM, JP
- 375 First draft: FC, JC
- 376 Supervision: GM, JF
- 377 Revision and final approval: FC, JC, GM, MM, BB, JP, GG, PM, JF

#### **379 REFERENCES**

380 Albert, C., Helgason, H.H., Brault-Favrou, M., Robertson, G.J., Descamps, S., Amélineau, F., 381 Danielsen, J., Dietz, R., Elliott, K., Erikstad, K.E., Eulaers, I., Ezhov, A., Fitzsimmons, 382 M.G., Gavrilo, M., Golubova, E., Grémillet, D., Hatch, S., Huffeldt, N.P., Jakubas, D., 383 Kitaysky, A., Kolbeinsson, Y., Krasnov, Y., Lorentsen, S.-H., Lorentzen, E., Mallory, 384 M.L., Merkel, B., Merkel, F.R., Montevecchi, W., Mosbech, A., Olsen, B., Orben, R.A., 385 Patterson, A., Provencher, J., Plumejeaud, C., Pratte, I., Reiertsen, T.K., Renner, H., Rojek, 386 N., Romano, M., Strøm, H., Systad, G.H., Takahashi, A., Thiebot, J.-B., Thórarinsson, 387 T.L., Will, A.P., Wojczulanis-Jakubas, K., Bustamante, P., Fort, J., 2021. Seasonal 388 variation of mercury contamination in Arctic seabirds: A pan-Arctic assessment. Sci. Total 389 Environ. 750, 142201. https://doi.org/10.1016/j.scitotenv.2020.142201

- AMAP, 2021. AMAP Assessment 2021: Mercury in the Arctic. Arctic Monitoring and Assessment
   Programme (AMAP), Tromsø, Norway. viii + 324pp
- Amiraux, R., Archambault, P., Moriceau, B., Lemire, M., Babin, M., Memery, L., Massé, G.,
   Tremblay, J.-E., 2021. Efficiency of sympagic-benthic coupling revealed by analyses of n 3 fatty acids, IP<sub>25</sub> and other highly branched isoprenoids in two filter-feeding Arctic benthic
   molluscs: *Mya truncata* and *Serripes groenlandicus*. Org. Geochem. 151, 104160.
   https://doi.org/10.1016/j.orggeochem.2020.104160
- Amiraux, R., Mundy, C.J., Pierrejean, M., Niemi, A., Hedges, K.J., Brown, T.A., Ehn, J.K., Elliott,
  K.H., Ferguson, S.H., Fisk, A.T., Gilchrist, G., Harris, L.N., Iken, K., Jacobs, K.B.,
  Johnson, K.F., Kuzyk, Z.A., Limoges, A., Loewen, T.N., Love, O.P., Matthews, C.J.D.,
  Ogloff, W.R., Rosenberg, B., Søreide, J.E., Watt, C.A., Yurkowski, D.J., 2023. Tracing
  carbon flow and trophic structure of a coastal Arctic marine food web using highly
  branched isoprenoids and carbon, nitrogen and sulfur stable isotopes. Ecol. Indic. 147,
  109938. https://doi.org/10.1016/j.ecolind.2023.109938
- 404 Atwell, L., Hobson, K.A., Welch, H.E., 1998. Biomagnification and bioaccumulation of mercury 405 in an Arctic marine food web: Insights from stable nitrogen isotope analysis 55, 8.
- Beattie, S.A., Armstrong, D., Chaulk, A., Comte, J., Gosselin, M., Wang, F., 2014. Total and
  methylated mercury in Arctic multiyear sea ice. Environ. Sci. Technol. 48, 5575–5582.
  <u>https://doi.org/10.1021/es5008033</u>
- Belt, S.T., Brown, T.A., Ringrose, A.E., Cabedo-Sanz, P., Mundy, C.J., Gosselin, M., Poulin, M.,
  2013. Quantitative measurement of the sea ice diatom biomarker IP<sub>25</sub> and sterols in Arctic
  sea ice and underlying sediments: Further considerations for palaeo sea ice reconstruction.
  Org. Geochem. 62, 33–45. <u>https://doi.org/10.1016/j.orggeochem.2013.07.002</u>
- Belt, S.T., Brown, T.A., Rodriguez, A.N., Sanz, P.C., Tonkin, A., Ingle, R., 2012. A reproducible
  method for the extraction, identification and quantification of the Arctic sea ice proxy IP<sub>25</sub>
  from marine sediments. Anal. Methods 4, 705–713. <u>https://doi.org/10.1039/C2AY05728J</u>

- Belt, S.T., Müller, J., 2013. The Arctic sea ice biomarker IP<sub>25</sub>: A review of current understanding,
   recommendations for future research and applications in palaeo sea ice reconstructions.
   Quat. Sci. Rev. 79, 9–25. https://doi.org/10.1016/j.quascirev.2012.12.001
- Bond, A.L., Diamond, A.W., 2010. Nutrient allocation for egg production in six Atlantic seabirds.
  Can. J. Zool. 88, 1095–1102. <u>https://doi.org/10.1139/Z10-082</u>
- Bond, A.L., Hobson, K.A., Branfireun, B.A., 2015. Rapidly increasing methyl mercury in endangered ivory gull (*Pagophila eburnea*) feathers over a 130 year record. Proc. R. Soc.
  B. 282:20150032. http://doi.org/10.1098/rspb.2015.0032
- Bradstreet, M.S.W., Cross, W.E., 1982. Trophic relationships at High Arctic ice edges. Arctic 35,
  1–12.
- Braune, B., Chételat, J., Amyot, M., Brown, T., Clayden, M., Evans, M., Fisk, A., Gaden, A.,
  Girard, C., Hare, A., Kirk, J., Lehnherr, I., Letcher, R., Loseto, L., Macdonald, R., Mann,
  E., McMeans, B., Muir, D., O'Driscoll, N., Poulain, A., Reimer, K., Stern, G., 2015.
  Mercury in the marine environment of the Canadian Arctic: Review of recent findings. Sci.
  Total Environ., Special Issue: Mercury in Canada's North 509–510, 67–90.
  https://doi.org/10.1016/j.scitotenv.2014.05.133
- Braune, B.M., Gaston, A.J., Elliott, K.H., Provencher, J.F., Woo, K.J., Chambellant, M., Ferguson,
  S.H., Letcher, R.J., 2014. Organohalogen contaminants and total mercury in forage fish
  preyed upon by thick-billed murres in northern Hudson Bay. Mar. Pollut. Bull. 78, 258–
  266. https://doi.org/10.1016/j.marpolbul.2013.11.003
- Braune, B.M., Gaston, A.J., Mallory, M.L., 2016. Temporal trends of mercury in eggs of five
  sympatrically breeding seabird species in the Canadian Arctic. Environ. Pollut. 214, 124–
  131. <u>https://doi.org/10.1016/j.envpol.2016.04.006</u>
- Brown, T.A., Belt, S.T., 2012. Identification of the sea ice diatom biomarker IP<sub>25</sub> in Arctic benthic
  macrofauna: Direct evidence for a sea ice diatom diet in Arctic heterotrophs. Polar Biol.
  35, 131–137. https://doi.org/10.1007/s00300-011-1045-7
- Brown, T.A., Belt, S.T., Philippe, B., Mundy, C.J., Massé, G., Poulin, M., Gosselin, M., 2011.
  Temporal and vertical variations of lipid biomarkers during a bottom ice diatom bloom in the Canadian Beaufort Sea: Further evidence for the use of the IP<sub>25</sub> biomarker as a proxy for spring Arctic sea ice. Polar Biol. 34, 1857–1868. <u>https://doi.org/10.1007/s00300-010-</u> 0942-5
- Brown, T.A., Bicknell, A.W.J., Votier, S.C., Belt, S.T., 2013. Novel molecular fingerprinting of
  marine avian diet provides a tool for gaining insights into feeding ecology. Environ. Chem.
  Lett. 11, 283–288. <u>https://doi.org/10.1007/s10311-013-0402-x</u>
- Burnham, K.P., Anderson, D.R., 2002. Model selection and multimodel inference: A practical
   information 616 Theoretic approach, 2nd ed. Springer US.

- 452 Campbell, L.M., Norstrom, R.J., Hobson, K.A., Muir, D.C.G., Backus, S., Fisk, A.T., 2005a.
  453 Mercury and other trace elements in a pelagic Arctic marine food web (Northwater
  454 Polynya, Baffin Bay). Sci. Total Environ., 351–352, 247–263.
  455 <u>https://doi.org/10.1016/j.scitotenv.2005.02.043</u>
- Carravieri, A., Lorioux, S., Angelier, F., Chastel, O., Albert, C., Bråthen, V.S., Brisson-Curadeau, 456 457 É., Clairbaux, M., Delord, K., Giraudeau, M., Perret, S., Poupart, T., Ribout, C., Viricel-458 Pante, A., Grémillet, D., Bustamante, P., Fort, J. 2023. Carryover effects of winter mercury 459 contamination on summer concentrations and reproductive performance in little auks, 460 Pollut. Environ. 318. 120774. ISSN 0269-7491. 461 https://doi.org/10.1016/j.envpol.2022.120774.
- 462 Chastel, O., Fort, J., Ackerman, J.T., Albert, C., Angelier, F., Basu, N., Blévin, P., Brault-Favrou, 463 M., Bustnes, J.O., Bustamante, P., Danielsen, J., Descamps, S., Dietz, R., Erikstad, K.E., 464 Eulaers, I., Ezhov, A., Fleishman, A.B., Gabrielsen, G.W., Gavrilo, M., Gilchrist, G., Gilg, 465 O., Gíslason, S., Golubova, E., Goutte, A., Grémillet, D., Hallgrimsson, G.T., Hansen, E.S., 466 Hanssen, S.A., Hatch, S., Huffeldt, N.P., Jakubas, D., Jónsson, J.E., Kitaysky, A.S., 467 Kolbeinsson, Y., Krasnov, Y., Letcher, R.J., Linnebjerg, J.F., Mallory, M., Merkel, F.R., Moe, B., Montevecchi, W.J., Mosbech, A., Olsen, B., Orben, R.A., Provencher, J.F., 468 Ragnarsdottir, S.B., Reiertsen, T.K., Rojek, N., Romano, M., Søndergaard, J., Strøm, H., 469 470 Takahashi, A., Tartu, S., Thórarinsson, T.L., Thiebot, J.-B., Will, A.P., Wilson, S., 471 Wojczulanis-Jakubas, K., Yannic, G., 2022. Mercury contamination and potential health 472 risks to Arctic seabirds and shorebirds. Sci. Total Environ. 156944. 473 https://doi.org/10.1016/j.scitotenv.2022.156944
- Chaulk, A., Stern, G.A., Armstrong, D., Barber, D.G., Wang, F., 2011. Mercury distribution and transport across the ocean-sea-ice-atmosphere interface in the Arctic Ocean. Environ. Sci.
  Technol. 45, 1866–1872. https://doi.org/10.1021/es103434c
- Chételat, J., McKinney, M.A., Amyot, M., Dastoor, A., Douglas, T.A., Heimbürger-Boavida, L.E., Kirk, J., Kahilainen, K.K., Outridge, P.M., Pelletier, N., Skov, H., St. Pierre, K.,
  Vuorenmaa, J., Wang, F., 2022. Climate change and mercury in the Arctic: Abiotic
  interactions. Sci. Total Environ. 824, 153715.
  https://doi.org/10.1016/j.scitotenv.2022.153715
- 482 Cusset, F., Fort, J., Mallory, M., Braune, B., Massicotte, P., Massé, G., 2019. Arctic seabirds and
  483 shrinking sea-ice: Egg analyses reveal the importance of ice-derived resources. Sci. Rep.
  484 9, 1–15. https://doi.org/10.1038/s41598-019-51788-4
- Dastoor, A., Angot, H., Bieser, J., Christensen, J.H., Douglas, T.A., Heimbürger-Boavida, L.-E.,
  Jiskra, M., Mason, R.P., McLagan, D.S., Obrist, D., Outridge, P.M., Petrova, M.V.,
  Ryjkov, A., St. Pierre, K.A., Schartup, A.T., Soerensen, A.L., Toyota, K., Travnikov, O.,
  Wilson, S.J., Zdanowicz, C., 2022a. Arctic mercury cycling. Nat. Rev. Earth Environ. 3,
  270–286. <a href="https://doi.org/10.1038/s43017-022-00269-w">https://doi.org/10.1038/s43017-022-00269-w</a>
- 490 Dastoor, A., Angot, H., Bieser, J., Christensen, J.H., Douglas, T.A., Heimbürger-Boavida, L.-E.,
  491 Jiskra, M., Mason, R.P., McLagan, D.S., Obrist, D., Outridge, P.M., Petrova, M.V.,

- 492 Ryjkov, A., St. Pierre, K.A., Schartup, A.T., Soerensen, A.L., Toyota, K., Travnikov, O.,
  493 Wilson, S.J., Zdanowicz, C., 2022b. Arctic mercury cycling. Nat. Rev. Earth Environ. 3,
  494 270–286. <u>https://doi.org/10.1038/s43017-022-00269-w</u>
- 495 David, C., Lange, B., Krumpen, T., Schaafsma, F., van Franeker, J.A., Flores, H., 2016. Under-ice
  496 distribution of polar cod *Boreogadus saida* in the central Arctic Ocean and their association
  497 with sea-ice habitat properties. Polar Biol. 39, 981–994. <u>https://doi.org/10.1007/s00300-</u>
  498 015-1774-0
- 499 Dietz, R., Letcher, R.J., Aars, J., Andersen, M., Boltunov, A., Born, E.W., Ciesielski, T.M., Das, 500 K., Dastnai, S., Derocher, A.E., Desforges, J.-P., Eulaers, I., Ferguson, S., Hallanger, I.G., 501 Heide-Jørgensen, M.P., Heimbürger-Boavida, L.-E., Hoekstra, P.F., Jenssen, B.M., 502 Kohler, S.G., Larsen, M.M., Lindstrøm, U., Lippold, A., Morris, A., Nabe-Nielsen, J., 503 Nielsen, N.H., Peacock, E., Pinzone, M., Rigét, F.F., Rosing-Asvid, A., Routti, H., Siebert, 504 U., Stenson, G., Stern, G., Strand, J., Søndergaard, J., Treu, G., Vikingsson, G.A., Wang, 505 F., Welker, J.M., Wiig, Ø., Wilson, S.J., Sonne, C., 2022. A risk assessment review of 506 mercury exposure in Arctic marine and terrestrial mammals. Sci. Total Environ. 829, 507 154445. https://doi.org/10.1016/j.scitotenv.2022.154445
- 508Dietz, R., Riget, F., Johansen, P., 1996. Lead, cadmium, mercury and selenium in Greenland509marine animals. Sci. Total Environ. 186, 67–93. <a href="https://doi.org/10.1016/0048-9697(96)05086-3">https://doi.org/10.1016/0048-9697(96)05086-3</a>
- 511 DiMento, B.P., Mason, R.P., Brooks, S., Moore, C., 2019. The impact of sea ice on the air-sea
   512 exchange of mercury in the Arctic Ocean. Deep Sea Res. Part Oceanogr. Res. Pap. 144,
   513 28-38. https://doi.org/10.1016/j.dsr.2018.12.001
- 514Foster, K.L., Braune, B.M., Gaston, A.J., Mallory, M.L., 2019. Climate influence on mercury in515Arctic seabirds.Sci.TotalEnviron.693,133569.516https://doi.org/10.1016/j.scitotenv.2019.07.375
- Garcia-Soto, C., Cheng, L., Caesar, L., Schmidtko, S., Jewett, E.B., Cheripka, A., Rigor, I.,
  Caballero, A., Chiba, S., Báez, J.C., Zielinski, T., Abraham, J.P., 2021. An overview of
  ocean climate change indicators: Sea surface temperature, ocean heat content, ocean pH,
  dissolved oxygen concentration, Arctic sea ice extent, thickness and volume, sea level and
  strength of the AMOC (Atlantic Meridional Overturning Circulation). Front. Mar. Sci. 8,
  642372. https://doi.org/10.3389/fmars.2021.642372
- Gaston, A.J., Elliott, K.H., Ropert-Coudert, Y., Kato, A., Macdonald, C.A., Mallory, M.L.,
  Gilchrist, H.G., 2013. Modeling foraging range for breeding colonies of thick-billed murres *Uria lomvia* in the Eastern Canadian Arctic and potential overlap with industrial
  development. Biol. Conserv. 168, 134–143. <a href="https://doi.org/10.1016/j.biocon.2013.09.018">https://doi.org/10.1016/j.biocon.2013.09.018</a>
- Gaston, A.J., Gilchrist, H.G., Mallory, M.L., 2005. Variation in ice conditions has strong effects
   on the breeding of marine birds at Prince Leopold Island, Nunavut. Ecography 28, 331–
   344. <u>https://doi.org/10.1111/j.0906-7590.2005.04179.x</u>

- Gaston, A. J., & Hipfner, J. M. (2020). Thick-billed Murre (*Uria lomvia*), version 1.0. In Birds of
   the World (S. M. Billerman, Editor). Cornell Lab of Ornithology, Ithaca, NY, USA.
   https://doi-org/10.2173/bow.thbmur.01
- Gaston, A.J., Mallory, M.L., Gilchrist, H.G., 2012a. Populations and trends of Canadian Arctic
  seabirds. Polar Biol. 35, 1221–1232. <u>https://doi.org/10.1007/s00300-012-1168-5</u>
- Gaston, A.J., Nettleship, D.N., 1981. The thick-billed murre of Prince Leopold Island, Monograph
   Series. Environment Canada Canadian Wildlife Service, Ottawa.
- Gaston, A.J., Smith, P.A., Provencher, J.F., 2012b. Discontinuous change in ice cover in Hudson
   Bay in the 1990s and some consequences for marine birds and their prey. ICES J. Mar. Sci.
   69, 1218–1225. <u>https://doi.org/10.1093/icesjms/fss040</u>
- Gaston, A.J., Woo, K., Hipfner, J.M., 2003. Trends in forage fish populations in Northern Hudson
   Bay since 1981, as determined from the diet of nestling thick-billed murres, *Uria lomvia*.
   ARCTIC 56, 227–233. https://doi.org/10.14430/arctic618
- Gosselin, M., Levasseur, M., Wheeler, P.A., Horner, R.A., Booth, B.C., 1997. New measurements
  of phytoplankton and ice algal production in the Arctic Ocean. Deep Sea Res. Part II Top.
  Stud. Oceanogr. 44, 1623–1644. https://doi.org/10.1016/S0967-0645(97)00054-4
- Gradinger, R.R., Bluhm, B.A., 2004. In-situ observations on the distribution and behavior of
  amphipods and Arctic cod (Boreogadus saida) under the sea ice of the High Arctic Canada
  Basin. Polar Biol. 27, 595–603. https://doi.org/10.1007/s00300-004-0630-4
- Heimbürger, L.-E., Sonke, J.E., Cossa, D., Point, D., Lagane, C., Laffont, L., Galfond, B.T.,
  Nicolaus, M., Rabe, B., van der Loeff, M.R., 2015. Shallow methylmercury production in
  the marginal sea ice zone of the central Arctic Ocean. Sci. Rep. 5, 10318.
  <a href="https://doi.org/10.1038/srep10318">https://doi.org/10.1038/srep10318</a>
- Horner, R., Schrader, G.C., 1982. Relative contributions of ice algae, phytoplankton, and benthic
   microalgae to primary production in nearshore regions of the Beaufort Sea. Arctic 35, 485–
   503.
- 556 Johnson, J.B., Omland, K.S., 2004. Model selection in ecology and evolution. Trends Ecol. Evol. 557 19, 101744 108. https://doi.org/10.1016/j.tree.2003.10.013
- Kelly, J.F., 2000. Stable isotopes of carbon and nitrogen in the study of avian and mammalian
   trophic ecology. Can. J. Zool. 78, 1–27. <u>https://doi.org/10.1139/z99-165</u>
- Kohlbach, D., Graeve, M., Lange, B.A., David, C., Peeken, I., Flores, H., 2016. The importance
  of ice algae-produced carbon in the central Arctic Ocean ecosystem: Food web
  relationships revealed by lipid and stable isotope analyses. Limnol. Oceanogr. 61, 2027–
  2044. https://doi.org/10.1002/lno.10351
- Kohlbach, D., Schaafsma, F.L., Graeve, M., Lebreton, B., Lange, B.A., David, C., Vortkamp, M.,
   Flores, H., 2017. Strong linkage of polar cod (*Boreogadus saida*) to sea ice algae-produced

- 566 carbon: Evidence from stomach content, fatty acid and stable isotope analyses. Prog. 567 Oceanogr. 152, 62–74. https://doi.org/10.1016/j.pocean.2017.02.003
- Lalonde, J.D., Poulain, A.J., Amyot, M., 2002. The role of mercury redox reactions in snow on
   snow-to-air mercury transfer. Environ. Sci. Technol. 36, 174–178.
   <a href="https://doi.org/10.1021/es010786g">https://doi.org/10.1021/es010786g</a>
- LeBlanc, M., Gauthier, S., Garbus, S.E., Mosbech, A., Fortier, L., 2019. The co-distribution of
   Arctic cod and its seabird predators across the marginal ice zone in Baffin Bay. Elem Sci
   Anth 7, 1. https://doi.org/10.1525/elementa.339
- Mallory, M.L., Braune, B.M., 2018. Do concentrations in eggs and liver tissue tell the same story
   of temporal trends of mercury in high Arctic seabirds? J. Environ. Sci. 68, 65–72.
   <u>https://doi.org/10.1016/j.jes.2017.10.017</u>
- Mallory, M.L., Braune, B.M., Provencher, J.F., Callaghan, D.B., Gilchrist, H.G., Edmonds, S.T.,
  Allard, K., O'Driscoll, N.J., 2015. Mercury concentrations in feathers of marine birds in
  Arctic Canada. Mar. Pollut. Bull. 98, 308–313.
  https://doi.org/10.1016/j.marpolbul.2015.06.043
- 581 Mazerolle, M.J., 2020. AICcmodavg: Model selection and multimodel inference based on 582 (Q)AIC(c). R package version 2.3-1. <u>https://cran.r-project.org/package=AICcmodavg</u>.
- McKinney, M.A., Peacock, E., Letcher, R.J., 2009. Sea ice-associated diet change increases the
   levels of chlorinated and brominated contaminants in polar bears. Environ. Sci. Technol.
   43, 4334–4339. https://doi.org/10.1021/es900471g
- 586Outridge, P.M., Macdonald, R.W., Wang, F., Stern, G.A., Dastoor, A.P., 2008. A mass balance587inventory of mercury in the Arctic Ocean. Environ. Chem. 5, 89.588https://doi.org/10.1071/EN08002
- Perrette, M., Yool, A., Quartly, G.D., Popova, E.E., 2011. Near-ubiquity of ice-edge blooms in the
   Arctic. Biogeosciences 8, 515–524. <u>https://doi.org/10.5194/bg-8-515-2011</u>
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., R Core Team, 2020. nlme: Linear and nonlinear
   mixed effects models. R package version 3.1-149, <u>https://CRAN.R-</u>
   project.org/package=nlme
- 594 Poulain, A.J., Garcia, E., Amyot, M., Campbell, P.G.C., Raofie, F., Ariya, P.A., 2007. Biological 595 and chemical redox transformations of mercury in fresh and salt waters of the High Arctic 596 during spring and summer. Environ. Sci. Technol. 41, 1883-1888. 597 https://doi.org/10.1021/es061980b
- 598 Power, M., Klein, G.M., Guiguer, K.R.R.A., Kwan, M.K.H., 2002. Mercury accumulation in the 599 fish community of a sub-Arctic lake in relation to trophic position and carbon sources: Hg 600 in а sub-Arctic lacustrine fish community. J. Appl. Ecol. 39. 819-830. 601 https://doi.org/10.1046/j.1365-2664.2002.00758.x

- Provencher, J.F., Gaston, A.J., O'Hara, P.D., Gilchrist, H.G., 2012. Seabird diet indicates changing
   Arctic marine communities in eastern Canada. Mar. Ecol. Prog.Ser. 454, 171–182.
   https://doi.org/10.3354/meps09299
- 605 R Core Team, 2020. R: A language an environment for statistical computing. R Foundation for 606 statistical computing, Vienna, Austria. URL: <u>https://www.R-project.org/</u>
- Routti, H., Letcher, R.J., Born, E.W., Branigan, M., Dietz, R., Evans, T.J., McKinney, M.A.,
  Peacock, E., Sonne, C., 2012. Influence of carbon and lipid sources on variation of mercury
  and other trace elements in polar bears (*Ursus maritimus*). Environ. Toxicol. Chem. 31,
  2739–2747. <u>https://doi.org/10.1002/etc.2005</u>
- Salvato, N., Pirola, C., 1996. Analysis of mercury traces by means of solid sample atomic
   absorption spectrometry. Microchim. Acta 123, 63–71.
   https://doi.org/10.1007/BF01244379
- Schartup, A.T., Soerensen, A.L., Heimbürger-Boavida, L.-E., 2020. Influence of the Arctic sea ice regime shift on sea-ice methylated mercury trends. Environ. Sci. Technol. Lett. 7, 708–
   713. <u>https://doi.org/10.1021/acs.estlett.0c00465</u>
- Steffen, A., Douglas, T., Amyot, M., Ariya, P., Aspmo, K., Berg, T., Bottenheim, J., Brooks, S.,
  Cobbett, F., Dastoor, A., Dommergue, A., Ebinghaus, R., Ferrari, C., Gardfeldt, K.,
  Goodsite, M.E., Lean, D., Poulain, A.J., Scherz, C., Skov, H., Sommar, J., Temme, C.,
  2008. A synthesis of atmospheric mercury depletion event chemistry in the atmosphere and
  snow. Atmospheric Chem. Phys. 8, 1445–1482. https://doi.org/10.5194/acp-8-1445-2008
- Stern, G.A., Macdonald, R.W., Outridge, P.M., Wilson, S., Chételat, J., Cole, A., Hintelmann, H.,
  Loseto, L.L., Steffen, A., Wang, F., Zdanowicz, C., 2012. How does climate change
  influence Arctic mercury? Sci. Total Environ. 414, 22–42.
  https://doi.org/10.1016/j.scitotenv.2011.10.039
- Streets, D.G., Horowitz, H.M., Lu, Z., Levin, L., Thackray, C.P., Sunderland, E.M., 2019. Five
  hundred years of anthropogenic mercury: spatial and temporal release profiles\*. Environ
  Res Lett 14:084004. <u>https://doi.org/10.1088/1748-9326/ab281f</u>
- 629 Thomas, D.N., Dieckmann, G.S., 2009. Sea ice. John Wiley & Sons.
- Thomas, D.N., Dieckmann, G.S., 2002. Antarctic sea ice: A Habitat for extremophiles. Science
  295, 641–644. https://doi.org/10.1126/science.1063391
- Thompson, D.R., Furness, R.W., Monteiro, L.R., 1998. Seabirds as biomonitors of mercury inputs
   to epipelagic and mesopelagic marine food chains. Sci. Total Environ. 213, 299–305.
   <a href="https://doi.org/10.1016/S0048-9697(98)00103-X">https://doi.org/10.1016/S0048-9697(98)00103-X</a>
- Tremblay, J.-É., Robert, D., Varela, D.E., Lovejoy, C., Darnis, G., Nelson, R.J., Sastri, A.R., 2012.
  Current state and trends in Canadian Arctic marine ecosystems: I. Primary production.
  Clim. Change 115, 161–178. <a href="https://doi.org/10.1007/s10584-012-0496-3">https://doi.org/10.1007/s10584-012-0496-3</a>

- Wassmann, P., Reigstad, M., 2011. Future Arctic Ocean seasonal ice zones and implications for
   pelagic-benthic coupling. Oceanography 24, 220–231.
   https://doi.org/10.5670/oceanog.2011.74
- Yurkowski, D.J., Brown, T.A., Blanchfield, P.J., Ferguson, S.H., 2020. Atlantic walrus signal 641 642 latitudinal differences in the long-term decline of sea ice-derived carbon to benthic fauna 643 Canadian Arctic. Proc. R. Soc. B Biol. Sci. in the 287, 20202126. 644 https://doi.org/10.1098/rspb.2020.2126
- Zeileis, A., Hothorn, T., 2002. Diagnostic checking in regression relationships. R News 2(3), 7 <u>https://CRAN.R-project.org/doc/Rnews</u>
- Zdanowicz, C., Krümmel, E.M., Lean, D., Poulain, A.J., Yumvihoze, E., Chen, J., Hintelmann,
  H., 2013. Accumulation, storage and release of atmospheric mercury in a glaciated Arctic
  catchment, Baffin Island, Canada. Geochim. Cosmochim. Acta 107, 316–335.
  <u>https://doi.org/10.1016/j.gca.2012.11.028</u>
- Zhang, Y., Jaeglé, L., Thompson, L., Streets, D.G., 2014. Six centuries of changing oceanic
   mercury: Anthropogenic mercury in ocean. Glob. Biogeochem. Cycles 28, 1251–1261.
   <a href="https://doi.org/10.1002/2014GB004939">https://doi.org/10.1002/2014GB004939</a>
- 654 Zheng, W., Chandan, P., Steffen, A., Stupple, G., De Vera, J., Mitchell, C.P.J., Wania, F., 655 Bergquist, B.A., 2021. Mercury stable isotopes reveal the sources and transformations of 656 atmospheric in the high Arctic. Geochem. 105002. Hg Appl. 131. 657 https://doi.org/10.1016/j.apgeochem.2021.105002

## 659 FIGURES







Figure 1. Location of Prince Leopold Island in the Canadian High Arctic



Figure 2. Relationship between total Hg concentrations measured in eggs of thick-billed murres (*Uria lomvia*) from Prince Leopold Island (Nunavut, Canada) and Ice Use Index. Lower and higher values of Ice Use Index correspond to the most pelagic and sympagic feeding association, respectively.

667



Figure 3. Relationship between total Hg concentrations ( $\mu g/g \, dw$ ) measured in eggs of thick-billed murres (*Uria lomvia*) from Prince Leopold Island (Nunavut, Canada) and  $\delta^{15}$  N values (proxy for trophic position).



**Figure 4.** Egg total Hg concentrations of thick-billed murres (*Uria lomvia*) from Prince Leopold Island (Nunavut, Canada) with varying ice concentrations. Ice concentrations were scaled for statistical analyses (raw concentrations range from 17.02 to 97.96 %). Solid and dotted lines represent the best-fitting model detected from construction of Generalized Linear Models (GLMs) and the confidence interval, respectively (see "Material and Methods" for further details).

#### TABLES

**Table 1.** Ice conditions and egg parameters from thick-billed murres collected between 2010 and 2013 on Prince Leopold Island (Nunavut, Canada): trophic tracers include the Ice Use Index, carbon- ( $\delta^{13}$ C) and nitrogen- ( $\delta^{15}$  N) stable isotopes; total mercury (THg) concentrations ( $\mu g/g \, dw$ ).  $\delta^{13}$ C and  $\delta^{15}$  N are proxies for bird feeding habitat and trophic position, respectively. Values represent means  $\pm$  SD (n).

	2010	2011	2012	2013
Ice concentrations <sup>1</sup> (%)	99.0	33.4	99.0	79.8
Ice characteristics <sup>2</sup>	Patchy, large floes, leads with water access	Ice-free	Dense pack-ice, no leads, ice edge > 250 km away	Dense pack ice, ice edge 50 km away
Ice Use Index <sup>3</sup>	$1.0 \pm 1.1$ (15)	-1.1 ± 1.1 (15)	0.7 ± 1.0 (15)	0.7 ± 0.7 (15)
H-Print (%) <sup>4</sup>	28.1 ± 20.8 (15)	82.5 ± 18.8 (15)	56.9 ± 27.3 (15)	63.4 ± 18.1 (15)
δ <sup>13</sup> C (‰)	$-19.7 \pm 0.4$ (11)	$-20.3 \pm 0.5$ (11)	$-19.3 \pm 0.4$ (15)	$-20.3 \pm 0.3$ (15)
δ <sup>15</sup> N (‰)	$16.0 \pm 0.5 (11)$	$15.3 \pm 0.7$ (11)	15.8 ± 0.5 (15)	15.4 ± 0.8 (15)
THg (µg/g dw)	$1.1 \pm 0.2$ (15)	0.9 ± 0.2 (15)	$1.2 \pm 0.2$ (15)	$1.0 \pm 0.2$ (15)
Moisture (%)	73.3 ± 0.9 (15)	69.1 ± 2.5 (15)	73.7 ± 1.0 (15)	73.6 ± 1.0 (15)

<sup>1</sup>Ice concentrations represent the proportion of sea ice *versus* open water in a 100 km radius around the colony at the time of egg-laying (on June 15). Please refer to the "Material and Methods" for further details.

<sup>2</sup>Despite similar ice concentrations, 2010 and 2012 differed in their ice characteristics. Complete description of ice conditions is available in Cusset et al. (2019).

<sup>3</sup>The Ice Use Index indicates the degree of ice feeding association (values < and > 0 indicates low and high reliance on sympagic resources, respectively).

<sup>4</sup>H-Print (i.e., HBI-fingerprint) indicates the degree of association to ice-derived resources (from 0 to 100% for sympagic and pelagic association, respectively).

**Table 2.** AIC<sub>c</sub> model ranking from statistical analyses of egg THg concentrations from thick-billed murres breeding on Prince Leopold Island (2010-2013). Models are Generalized Linear Models (GLMs) with a Gamma distribution and an inverse link function. Explanatory variables include: ice concentrations within 100 km around the colony («Ice»), the type of association to sea-ice and its derived resources (pelagic *versus* sympagic; «IceUse»), the trophic position (« $\delta^{15}$  N») and their relevant interactions. Abbreviations: k, the number of parameters; AIC<sub>c</sub>, Akaike's Information Criteria adjusted for small sample-size values; w<sub>i</sub>, AICc weights (weights across all models sum to 1.00); LL, Log-Likelihood.

Models	k	AIC <sub>c</sub>	ΔAIC <sub>c</sub>	Wi	LL
THg ~ Ice*	3	-16.43	0.00	0.83	11.46
$THg \sim \delta^{I^{5}} N + IceUse + Ice$	5	-12.79	3.64	0.13	12.05
$THg \sim \delta^{15} N + IceUse + \delta^{15} N$ : $IceUse + Ice + Ice$ : $IceUse$	7	-9.44	6.99	0.03	12.99
THg ~ IceUse	3	-7.04	9.38	0.01	6.77
$THg \sim \delta^{I^{5}} N + IceUse$	4	-5.02	11.40	0.00	6.94
$THg \sim \delta^{I^{5}} N + IceUse + \delta^{I^{5}} N: IceUse$	5	-4.13	12.30	0.00	7.72
$THg \sim \delta^{I^{5}} N + IceUse + Ice:IceUse$	5	-2.96	13.46	0.00	7.13
$THg \sim \delta^{I^5} N$	3	0.18	16.60	0.00	3.16
$THg \sim NULL$	2	1.14	17.56	0.00	1.55

\*The best model (in bold) is selected with the lowest AICcvalue.

### **CREDIT AUTHOR STATEMENT**

### Contributions.

- Conception and design: FC, JF, GM, MM
- Laboratory work and data acquisition: FC (trophic tracers, ice parameters), BB (mercury), GG
- (stable isotopes) and PM (ice analyses)
- Data analysis: FC
- Interpretation: FC, JC, JF, GM, MM, JP
- First draft: FC, JC
- Supervision: GM, JF
- Revision and final approval: FC, JC, GM, MM, BB, JP, GG, PM, JF

