



HAL
open science

Dietary methylmercury and selenium interactions in rainbow trout juveniles (*Oncorhynchus mykiss*)

Marius Bidon, Khoulood El Hanafi, Silvia Queipo-Abad, Claudia Marchán Moreno, Jérôme Roy, David Amouroux, Maïté Bueno, Zoyne Pedrero, Stéphanie Fontagné-Dicharry

► To cite this version:

Marius Bidon, Khoulood El Hanafi, Silvia Queipo-Abad, Claudia Marchán Moreno, Jérôme Roy, et al.. Dietary methylmercury and selenium interactions in rainbow trout juveniles (*Oncorhynchus mykiss*). Aquaculture Europe 2021, EAS, Oct 2021, Funchal, Portugal. hal-04226080

HAL Id: hal-04226080

<https://hal.science/hal-04226080v1>

Submitted on 3 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DIETARY METHYLMERCURY AND SELENIUM INTERACTIONS IN RAINBOW TROUT JUVENILES (*Oncorhynchus mykiss*)

M. Bidon^{1*}, C. Marchan Moreno², K. El Hanafi², S. Queipo Abad², J. Roy¹, D. Amouroux², M. Bueno², Z. Pedrero-Zayas², S. Fontagné-Dicharry¹

¹INRAE, Univ. Pau & Pays Adour, E2S UPPA, NUMEA, 64310 Saint-Pée-sur-Nivelle, France

²Université de Pau et des Pays de l'Adour, E2S UPPA, CNRS, IPREM, Institut des Sciences Analytiques et de Physico-chimie pour l'Environnement et les matériaux, Pau, France.

E-mail: marius.bidon@inrae.fr

Introduction

Mercury (Hg), especially the organic, methylated form (MeHg), is a toxic heavy metal that can accumulate in large amounts in fish through time in various organs, like liver and brain. Brain, with neurons, is the principal target of MeHg toxicity (Ni et al. 2012). MeHg induces its toxicity through induction of oxidative stress and inflammation (Morcillo et al. 2017). Selenium (Se), an essential micronutrient mainly involved in antioxidant protection, has been shown to reduce MeHg toxicity. However, Se and MeHg interactions are far to be totally understood. As relatively high levels of these two elements can be found in marine ingredients, a better understanding of their interactions is important to improve aquafeed formulations.

Material and Methods

Six plant protein-rich diets were formulated to contain two different MeHg levels (0 and 2.5 mg Hg/kg in HgC and HgH diets) combined with 3 different Se levels and forms (0.3 mg Se/kg in SeC diets and 2.6 mg Se/kg supplied either as sodium selenite in SeI diets or as L-selenomethionine from Orffa in SeO diets). Each diet was distributed to 3 replicate tanks of 50 fish (initial mean body weight: 26±1 g) to apparent satiation over a 3-week feeding trial at a water temperature of 17.5°C. Fish growth performance (n=3 tanks/diet) was analysed as well as total Hg and Se contents (n=3 individuals/diet) and relative gene expression levels (RT-PCR, n=9 individuals/diet) in liver and brain.

Results

Reduced growth performance of fish fed MeHg was recorded only within the non-Se supplemented SeC-group (Table 1). Feeding MeHg significantly increased total Hg concentrations in both liver and brain but to a significant smaller extent in liver when fish were supplemented with Se. Along with the hepatic increase of total Hg, dietary MeHg supplementation decreased total Se content in liver. Dietary inorganic Se supplementation increased total Se concentrations in liver to a larger extent than dietary organic Se supplementation but inorganic Se intake did not increase significantly Se content in brain contrary to organic Se intake.

In liver, MeHg intake increased transcript levels of enzymes involved in cellular stress response and antioxidant defences such as GST π and MsrB3 (Table 1). Both dietary organic and inorganic Se supplementation increased hepatic transcript levels of the selenoprotein GPX1a and also SelPa1 but only in HgC groups. In brain, dietary MeHg supplementation decreased transcript levels of RBFOX3, a neural marker, and up-regulated expression of IL-1 β , a pro-inflammatory cytokine.

Discussion and conclusion

The growth inhibition by MeHg after only 21 days of feeding could be due to the prooxidant and proinflammatory properties highlighted by gene expression analysis in brain and liver. Fighting against MeHg-induced oxidative stress and inflammation consumes energy that is thus less available for growth (Morcillo et al. 2017). The mitigation by dietary Se supplementation that promoted antioxidant genes lends further support to this hypothesis.

Table 1. Effects of dietary MeHg and Se supplementation and their interaction on growth performance, total Hg, total Se and gene expression (normalised with luciferase and EF1 α to the control group HgC-SeC) in liver and brain of rainbow trout juveniles fed for 3 weeks using two-way ANOVA. GST π , glutathione S-transferase; MsrB3, methionine sulfoxide reductase B3; GPX1a, glutathione peroxidase 1a; SelPa1, selenoprotein Pa1; IL-1 β , interleukin 1 β ; RBOX3, RNA binding fox-1 homolog 3.

	Dietary MeHg group		Dietary Se group			Interaction
	HgC	HgH	SeC	SeI	SeO	
Final mean body weight (g)	42.7	41.9	41.7	42.6	42.6	0.025
Liver total Hg content ($\mu\text{g/g}$)	0.3 ^b	5.9 ^a	4.1 ^y	2.5 ^z	2.6 ^z	0.008
Liver total Se content ($\mu\text{g/g}$)	18.8 ^a	16.0 ^b	2.6 ^z	33.4 ^x	16.1 ^y	0.002
Brain total Hg content ($\mu\text{g/g}$)	0.4 ^b	3.6 ^a	1.9	1.8	2.2	0.624
Brain total Se content ($\mu\text{g/g}$)	1.2	1.1	0.8 ^y	1.0 ^y	1.7 ^z	0.920
Liver GST π transcript level	1.4 ^b	2.1 ^a	1.4	2.1	1.8	0.858
Liver MsrB3 transcript level	1.9 ^b	2.4 ^a	1.8	2.3	2.3	0.040
Liver GPX1a transcript level	1.3	1.3	1.1 ^y	1.4 ^z	1.4 ^z	0.230
Liver SelPa1 transcript level	1.5	1.6	1.5	1.6	1.7	0.002
Brain IL-1 β transcript level	1.6 ^b	2.5 ^a	1.7	2.2	2.3	0.190
Brain RBFOX3 transcript level	1.1 ^a	1.0 ^b	1.0	1.1	1.0	0.740

Indeed, at the transcript level we found that MeHg induced inflammation in brain and antioxidant defences in liver to fight against oxidative stress. The antioxidant support by dietary Se supplementation in plant-based diets at the transcript level, in agreement with previous studies (Fontagné-Dicharry et al., 2020), could provide better protection against xenobiotics like MeHg in rainbow trout juveniles, even if the controlled genes by dietary MeHg and Se are not systematically similar and deserved further investigation for a better understanding.

The present results provide evidence of interactions between dietary MeHg and Se on tissue Hg and Se concentrations, especially in liver. The difference of Hg and Se storage between the two tissues suggest post-absorptive interacting effects between Hg and Se maybe at the depuration level due to their strong affinity for each other as suggested by Amlund et al. (2015) in zebrafish but do not exclude interactive effects at the digestive or absorption level that deserve further investigation. As muscle, the edible part, has been shown to readily accumulate MeHg (Berntssen et al. 2004), this tissue should be included in the study to ascertain this finding with different dietary Se levels and forms.

Our data showed that the dietary level of 2 mg Hg/kg diet supplied as organic mercury induced adverse effects as early as 21 days of feeding, especially in Se-deficient diets, in agreement with the threshold levels of dietary MeHg set as 5 mg Hg/kg by Berntssen et al. (2004) for salmonid diets. In brain, our RBFOX3 results confirm the high sensitivity of neurons to MeHg toxicity (Ni et al. 2014). Longer exposure, for example up to 6 months, should be assessed in rainbow trout juveniles for a better understanding on the adverse effects of MeHg and the interacting effects with dietary Se.

Acknowledgements

This project was supported by the French National Research Agency (ANR-18-CE34-0004), the Region Nouvelle-Aquitaine (BENESEL) and I-site E2S UPPA (contract 2020-32) and has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 101007962.

References

- Amlund, H., Lundebye, A.-K., Boyle, D., Ellingsen, S., 2015. Dietary selenomethionine influences the accumulation and depuration of dietary methylmercury in zebrafish (*Danio rerio*). *Aquat. Toxicol.* 158: 211-217.
- Berntssen, M.H.G., Hylland, K., Julshamn, K., Lundebye, A.-K., Waagbø, R., 2004. Maximum limits of organic and inorganic mercury in fish feed. *Aquac. Nutr.* 10: 83-97.
- Fontagné-Dicharry, S., Véron, V., Larroquet, L., Godin, S., Wischhusen, P., Aguirre, P., Terrier, F., Richard, N., Bueno, M., Bouyssi re, B., Antony Jesu Prabhu, P., Tacon, P., Kaushik, S.J., 2020. Effect of selenium sources in plant-based diets on antioxidant status and oxidative stress-related parameters in rainbow trout juveniles under chronic stress exposure. *Aquaculture* 529: 735684.
- Morcillo, P., Esteban, M.A., Cuesta, A., 2017. Mercury and its toxic effects on fish. *AIMS Environ. Sci.* 4: 386-402.
- Ni, M., Li, X., Rocha, J.B.T., Farina, M., Aschner, M., 2012. Glia and methylmercury neurotoxicity. *J. Toxicol. Environ. Health A* 75: 1091-1101.