



**HAL**  
open science

## Comment on “Fickian Non-Gaussian Diffusion in Glass-Forming Liquids”

L. Berthier, E. Flenner, G. Szamel

► **To cite this version:**

L. Berthier, E. Flenner, G. Szamel. Comment on “Fickian Non-Gaussian Diffusion in Glass-Forming Liquids”. *Physical Review Letters*, 2023, 131 (11), pp.119801. 10.1103/PhysRevLett.131.119801 . hal-04228262

**HAL Id: hal-04228262**

**<https://hal.science/hal-04228262v1>**

Submitted on 4 Oct 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L’archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d’enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Comment on ‘Fickian Non-Gaussian Diffusion in Glass-Forming Liquids’.

A recent Letter [1] examined the statistics of individual particles displacements  $\Delta x(t)$  over time  $t$  in two-dimensional glass-formers and concluded that the corresponding probability distribution  $G_s(\Delta x, t)$ , called the van-Hove distribution, is non-Gaussian in a time regime where the mean-squared displacement (MSD) is Fickian,  $\langle \Delta x^2(t) \rangle \propto D_s t$ , where  $D_s$  is the self-diffusion constant. If this analysis were correct, glass-formers would be ‘Fickian non-Gaussian’ materials [2, 3].

Here, we clarify that the multiple lengthscales and timescales reported in [1] have either been characterized before, or are not well-defined. This leads us to dispute the conclusions that glass-formers display Fickian non-Gaussian behaviour and that this analogy fruitfully addresses the central questions regarding the nature of dynamic heterogeneity in these systems.

The main features of self-diffusion in supercooled liquids are explained by invoking two characteristic timescales [4–7]. The self-diffusion coefficient  $D_s$  controls the first one,  $\tau_D = \sigma^2/D_s$ , where  $\sigma$  is the particle size. The second one is the structural relaxation time  $\tau_\alpha$  determined from usual time correlations, such as the self-intermediate scattering function. The adimensional ratio  $X = \tau_\alpha/\tau_D$  plays a special role. It controls both the amount of decoupling  $X(T) \sim D_s \tau_\alpha$  (akin to violations of the Stokes-Einstein relation [8]), and the Fickian length-scale  $\ell_F \propto \sqrt{X}$  [4]. Known results [4–10] paint a picture that is inconsistent with several conclusions reported in [1, 3] as we now show.

Let us start with the van Hove distribution. It was found in [9] that  $G_s(\Delta x, t)$  approaches a Gaussian distribution only for times much longer than  $\tau_D$ , a result rediscovered in [1] with equivalent tools. However, the non-Gaussian parameter  $\alpha_2(t)$  used in [1] to reveal Gaussianity decays as a power law at large times. Hence, the gradual emergence of Gaussian behaviour from  $\alpha_2(t)$  is a scale-free process and there is no characteristic timescale after which self-diffusion is Gaussian, although of course empirically  $\alpha_2(t)$  will be small when  $\langle \Delta x^2 \rangle \gg \ell_F^2$ . It was found numerically [8] and explained theoretically [4] that  $\ell_F$  in fact controls the crossover in the wavevector dependence of the self-intermediate scattering function, a result ignored in [1].

For times  $t < \tau_\alpha$ ,  $G_s(\Delta x, t)$  is characterized by a Gaussian core at small  $\Delta x$  and a nearly exponential tail  $G_s \propto \exp(-|\Delta x|/\lambda)$  at large  $\Delta x$  [5]. Refs. [1, 3] discuss the existence and possible universality of a power law description of the time evolution of the exponential tail,  $\lambda(t) \sim t^\alpha$ . As noticed in [5, 6, 11], the exponential tail is generically explained by a large deviation argument, but asymptotic convergence is so slow that the actual value of  $\lambda$  depends on the fitted range (see [6] for an explicit test and [11] for quantitative statements suggesting that  $\alpha = 0$ ), which may explain reported discrepancies [1, 3].

More fundamentally  $\lambda(t)$  is difficult to measure, and  $\alpha$  is not a novel characteristic exponent.

Linear behaviour of the MSD is visually detected [1] in log-log representations after a time  $\tau_D$  which grows more slowly than  $\tau_\alpha$  at low temperature, but the approach to linearity is again algebraic [12]. This power law approach to Fickian behaviour is again scale free and no characteristic timescale controls the emergence of Fickian behaviour in  $\langle \Delta x^2(t) \rangle$ .

Even though glass-formers may appear empirically close to Fickian non-Gaussian materials, there is no characteristic timescales or lengthscales controlling the approach to either Fickian and Gaussian dynamics, and the existence of a Fickian non-Gaussian regime cannot be decided. Instead, the salient features of self-diffusion, including algebraic approach to Fickian and Gaussian behaviours as well as nearly exponential van Hove distributions, are analytically captured by (effective) non-interacting continuous time random walk models [4–7, 10] based on the only two important and well-defined characteristic timescales  $\tau_D$  and  $\tau_\alpha$ . The multiple time and length scales determined empirically in [1, 3] are either related to those, or conceptually ill-defined.

The complexity of glass-formers is that the timescales  $\tau_D$  and  $\tau_\alpha$  emerge from many-body interactions (disorder is self-induced) and have non-trivial temperature dependencies which are not fully understood, but from which the very rich statistics of single particle displacement naturally follows. The behaviour of supercooled liquids is very different from several Fickian non Gaussian materials, which are described by interesting, but quite different, models [15].

We end by noting that the use of two-dimensional simulations to study the statistics of particle displacements in glass-formers is profoundly influenced at all timescales by Mermin-Wagner fluctuations [13, 14], which presumably adds to the profusion of timescales reported in [1].

L. Berthier<sup>1,2</sup>, E. Flenner<sup>3</sup>, G. Szamel<sup>3</sup>

<sup>1</sup>Laboratoire Charles Coulomb (L2C), Université de Montpellier, CNRS, 34095 Montpellier, France

<sup>2</sup>Yusuf Hamied Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge CB2 1EW, United Kingdom

<sup>3</sup>Department of Chemistry, Colorado State University, Fort Collins Colorado 80523, USA

- 
- [1] F. Rusciano, R. Pastore, and F. Greco, Phys. Rev. Lett. **128**, 168001 (2022).
  - [2] B. Wang, S. M. Anthony, S. C. Bae, and S. Granick, Proc. Natl. Acad. Sci. USA **106**, 15160 (2009).
  - [3] J. M. Miotto, S. Pigolotti, A. V. Chechkin, and S. Roldan-Vargas, Phys. Rev. X **11**, 031002 (2021).
  - [4] L. Berthier, D. Chandler, and J.P. Garrahan, Europhys. Lett. **69**, 320 (2005).

- [5] P. Chaudhuri, L. Berthier, and W. Kob, Phys. Rev. Lett. **99**, 060604 (2007).
- [6] P. Chaudhuri, Y. Gao, L. Berthier, M. Kilfoil, and W. Kob, J. Phys.: Condens. Matter **20**, 244126 (2008).
- [7] L. O. Hedges, L. Maibaum, D. Chandler, and J. P. Garahan, J. Chem. Phys. **127**, 211101 (2007).
- [8] L. Berthier, Phys. Rev. E **69**, 020201(R) (2004).
- [9] G. Szamel and E. Flenner, Phys. Rev. E **73**, 011504 (2006).
- [10] J. Helfferich, F. Ziebert, S. Frey, H. Meyer, J. Farago, A. Blumen, and J. Baschnagel, Phys. Rev. E **89**, 042604 (2014).
- [11] E. Barkai and S. Burov, Phys. Rev. Lett. **124**, 060603 (2020).
- [12] T. B. Schroder and J. C. Dyre, J. Chem. Phys. **152**, 141101 (2020).
- [13] E. Flenner and G. Szamel, Nature Comm. **6**, 7392 (2015).
- [14] E. Flenner and G. Szamel, Proc. Natl. Acad. Sci. U.S.A. **116**, 2015 (2019).
- [15] A. V. Chechkin, F. Seno, R. Metzler, and I. M. Sokolov, Phys. Rev. X **7**, 021002 (2017).