



Plants response to estuarine constraints: a study of controlled salinity and inundation

Markus Neupert, Michaël Aubert, Estelle Langlois

► To cite this version:

Markus Neupert, Michaël Aubert, Estelle Langlois. Plants response to estuarine constraints: a study of controlled salinity and inundation. 65th Annual IAVS Symposium, Sep 2023, Coffs Harbour, Australia. hal-04228323

HAL Id: hal-04228323

<https://hal.science/hal-04228323>

Submitted on 4 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PLANTS RESPONSE TO ESTUARINE CONSTRAINTS: A STUDY OF CONTROLLED SALINITY AND INUNDATION





Markus Neupert¹, Michaël Aubert¹, Estelle Langlois¹

¹ Ecodiv Lab USC INRAE 1499, University of Rouen Normandy, France
Contact: markus.neupert1@univ-rouen.fr



- Inbetween terrestrial, marine and freshwater ecosystems, estuarine wetlands are shaped by strong environmental constraints, including tidal cycles, varying salt concentrations, and erosion/sedimentation dynamics (Perillo et al., 2019).
- Vegetation community composition is strongly driven by the salinity and inundation gradients (Silvestri et al., 2005), which impact growth performance and select species with fitting functional traits (Keddy et al., 1992).



The salt gradient of the Seine estuary as a reference.


- In a context of human-made climate change, the current salinities and inundation frequencies are bound to change (IPCC, 2022). How do plant growth and resource allocation of estuarine wetland plants respond to them?

- Working hypotheses:**
 - Disturbance gradients of salinity and inundation impact plant survival and functional traits at the species level.
 - Productivity is related to disturbance intensity and frequency.

LINEAR MODEL COEFFICIENTS OF LEAF TRAIT RESPONSE

Formula: Trait ~ Inundation * Salinity
Only most significant coefficients shown ($p < 0.07$)


Species: ● *Alopecurus geniculatus* ● *Festuca arundinacea* ● *Holcus lanatus*



- Across the gradients of inundation and salinity, there is evidence for different responses of traits between species, related to their life strategies.

- With approx. 5000 photographs taken over the course of the experiment, we found an important synergy of the effects of inundation and salinity on plant survival and growth performance.
- Given the linear increase of leaf overlap, the photographic data can be corrected to estimate total leaf area.

PHOTOGRAPHED LEAF AREA OVER THE COURSE OF THE EXPERIMENT (not adjusted for leaf overlap)



CONTEXT

1

FIRST RESULTS: TRAITS

6


FIRST RESULTS: GROWTH

5

PROOF OF CONCEPT

- A preliminary experiment with 4 different plant species was used to develop the photoanalysis algorithm and test its use case under benign growth conditions.
- The average leaf area values from 4 photographs per individual allowed for the most accurate estimations of leaf area and plant height.
- Compared to destructive leaf area scans, photography underestimated the total leaf area due to increasing leaf overlap with plant size.
- Photographed leaf area is a good estimator of dry biomass ($R^2=0.93$), except for *J. gerardii*, whose thin leaves were not detected as well ($R^2=0.46$).

PHOTOGRAPHED LEAF AREA COMPARED TO SCANNED LEAF AREA



EXPERIMENTAL SETUP

2


PHOTOANALYSIS

- Photographic growth monitoring is cost-effective, time saving and allows for repeated measures on the same individuals.
- Various applications: vegetation cover estimation, N content estimation, phenological analyses, biomass estimation.
- The custom algorithm calculates leaf area in few seconds, simplifying batch analyses.




A scanned individual.

DRY SHOOT BIOMASS AS A LINEAR FUNCTION OF PHOTOGRAPHED LEAF AREA




Input Photograph



Color Adjustment



Thresholding



Pixel to surface conversion

