

Towards a better characterization of heat stress recurrence: a case study in oilseed rape

Sophie Brunel-Muguet, Lethicia Magno Massuia de Almeida, Jérémy Delamare, Lorane Le Franc, Benoît Bernay, Julien Pontin, Jessica L. Soyer

▶ To cite this version:

Sophie Brunel-Muguet, Lethicia Magno Massuia de Almeida, Jérémy Delamare, Lorane Le Franc, Benoît Bernay, et al.. Towards a better characterization of heat stress recurrence: a case study in oilseed rape. International Society for Seed Science, Jul 2023, Paris, France. hal-04232194

HAL Id: hal-04232194 https://hal.science/hal-04232194

Submitted on 7 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Towards a better characterization of heat stress recurrence: A case study in oilseed rape

Brunel-Muguet S¹, Magno Massuia de Almeida L¹, Delamare J¹, Le Franc L¹, Bernay B², Pontin J², Fremont J¹, Avice JC¹, Mollier A³, Personeni E¹, Soyer J⁴ 1: UMR 950 Ecophysiologie Végétale, Agronomie et Nutritions NCS (UCBN, INRAE) 2 : Plateforme Proteogen, US Emerode (UCBN) 3: UMR 1391 Interactions Sol Plante Atmosphère (Bordeaux science agro, INRAE) 4 : UMR 1290 BIOlogie et GEstion des Risques en agriculture, (Univ. Paris-Saclay, INRAE)



Our objectives were to (i) induce a thermopriming effect to alleviate the negative impacts of later intense heat stress on seed yield and quality criteria and (ii) to identify physiological pathways and epigenetic markers that account for positive memory (priming) or negative memory (cumulated effects). **Two sets of experiments** were designed such that the effects of an early mild stress event followed by a recovery phase (EMS+R) or a gradual increase (GI) in temperature prior to 5 day-intense heat (IH) were tested.

Seed yield and quality criteria were measured to assess any priming effect of EMS+R or GI prior to the IH. In order to consider the age of pods at the time of stress application, measurements were also performed at the cohort level.

For the GI experiment, global and targeted proteomics analyses were performed by Mass Spectrometry following technical adjustments (Le Franc, in prep.) to quantify the abundance of histone post translational modifications (histones PTMs). Clustering and Gene ontology (GO) analyses were carried out to highlight the physiological processes involved in response to Giinduced priming.

Plant performances: seed yield and quality criteria



Priming effects were mostly observed in the Gradual Increase, modality while negative memory effects (cumulated) were observed in the Early Mild Stress + Recovery modality (Table 1).

Most differences among modalities observed at the whole plant scale are accounted for by a given cohort of pods: the **intermediate cohort** of pods at the time of pre stress application for the **EMS+R experiment and** the **oldest cohort** for the **GI** experiment (Fig. 4)

These observations lead to conclude that (i) there is a narrow phenological window (in relation with the age of the seeds) for the pre stress to be priming which reflects the need to synchronize the "priming" event and the building processes of the seed storage compounds and (ii) there is a specific duration of the recovery phase over which the mild stress induced memory mechanisms are reset (Lamke and Baurle, 2017).



Glo<mark>bal and</mark> Targeted (histone PTM) proteomics



Clustering (Fig. 5) and 60 (Fig. 6) analyses reveal 4 clusters of proteins that were differentially represented according to our modalities.

- Cluster 1 proteins are responsive to the duration of stress : CS vs. [control, GI, IH]
- Cluster 2 proteins are responsive to GI (alone or combined with IH). They involved in stress tolerance acquisition and are involved in S metabolism
- Cluster 3 proteins are more abundant in modalities that include IH (alone or combined with GI).
- Cluster 4 is a mirror of Cluster 3. It gathers proteins that are less abundant under intense heat stress (CS and IHS) and that are involved in oxidative stress.

In addition, significant enrichment of three histone PTMs by MS was highlighted : H3K27me1, H3K9me1 and H3K4me1, the later being more abundant under IH that is, in proteins of cluster 4 (not shown).

Conclusions and Perspectives

vui, nicmalis DUII

Thermo-sensitization protocols are difficult to picture in indeterminate growth species due to mixed age organs (and hence different extent of stress sensitivity) and compensation effects (plasticity). Designing thermo-sensitization protocols to maintain seed quality are likely to be criteria specific as a consequence of a required synchrony between the priming event and the quality-related building processes. S metabolism is involved in thermo-sensitization acquisition which raises the question of methylation mediated regulation with S-Adenosylmethionine (-CH3 donor) Several PTMs of histones were associated to heat stress responses and more specifically H3K4me1 which was found to be more abundant upon intense heat.

Chromatin Immuno Precipitation sequencing (ChIP seq) analyses are in progress which will allows heat stress (memory) responsive genes to be associated with the over/under abundance of targeted histone PTMs.

References:

IPCC (2021.) doi:10.1017/9781009157896. Lamke J and Baurle I (2017) doi: 10.1186/s13059-017-1263-6. Magno Massuia de Almeida L et al. (2021) doi: 10.1016/j.envexpbot.2021.104400