



On two dimensional Schrödinger operators

Pierre Gaillard

► To cite this version:

| Pierre Gaillard. On two dimensional Schrödinger operators. 2023. hal-04232392

HAL Id: hal-04232392

<https://hal.science/hal-04232392>

Preprint submitted on 8 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On two dimensional Schrödinger operators

Pierre Gaillard,

Université de Bourgogne-Franche Comté, Dijon, France,

Abstract

Following the works of Berest [4], we compute explicitly Hadamard's coefficients for two dimensional Schrödinger operators.

For this, we use some specific generating functions and compute their relative Wronskians.

1 Introduction

In 2008 Berest, Cramer and Eshmatov [4] studied the Schrödinger operator $\mathcal{L} = -\Delta + V$, with a C^∞ potential V in some open domain Ω , following several works on similar themes [2, 3]. The heat kernel of \mathcal{L} is the solution $\Phi_+(x, \zeta, t)$ in $\Omega \times \Omega \times \mathbf{R}^+$ of the initial value problem

$$(\partial_t + \mathcal{L})\Phi_+(x, \xi, t) = 0, \quad \lim_{t \rightarrow 0^+} \Phi_+(x, \xi, t) = \delta(x - \xi), \quad (1)$$

where $\delta(x - \xi)$ is the Dirac delta function on Ω with support at ξ .

The function $\Phi_+(x, \xi, t)$ has an asymptotic expansion of the form [71]

$$\Phi_+(x, \xi, t) \sim \frac{e^{|x-\xi|^2/4t}}{(4\pi t)^{n/2}} \left(1 + \sum_{\nu=1}^{\infty} U_\nu(x, \xi) t^\nu \right) \text{ as } t \rightarrow 0^+. \quad (2)$$

The functions $U_\nu(x, \xi)$ are called Hadamard's coefficients of the Schrödinger operator $\mathcal{L} = -\Delta + V$.

In their paper, they consider the potential V defined by

$$V(x) = -\frac{2}{r^2} \frac{\partial^2}{\partial \varphi^2} \ln W(F_0, \dots, F_m), \quad (3)$$

where $W(F_0, \dots, F_m)$ is the wronskian of the functions F_0, \dots, F_m , and $k = (k_0, k_1, \dots, k_m)$ is a strictly increasing sequence of integers.

Considering the functions F_i defined by $F_i(x) = \cos(k_i x + x_i)$, where x is equal to φ or ϕ and x_i equal to φ_i or ϕ_i , they proved that the Hadamard's coefficients of $\mathcal{L} = -\Delta + V_k$ are given in terms of polar coordinates $x = (r \cos \varphi; r \sin \varphi)$, $\xi = (\rho \cos \phi; \rho \sin \phi)$ by

$$U_\nu(x, \xi) = \frac{(-2)^\nu}{(r\rho)^\nu} \sum_{i=0}^m c_i \Psi_i(\varphi) \Psi_i(\phi) T_{k_i}^{(\nu)}(\cos(\varphi - \phi)), \text{ for } \nu \geq 0 \quad (4)$$

where

$$\Psi_i := \frac{W(F_0, \dots, F_{i-1}, F_{i+1}, \dots, F_m)}{W(F_0, \dots, F_m)}, \quad (5)$$

$$c_i := \prod_{j=0, j \neq i}^m k_i^2 - k_j^2, \quad (6)$$

$T_N(z) := \cos N(\arccos z)$, with $T_N^{(\nu)}(z)$ being its derivative of order ν with respect to z .

In this statement we recognize the Chebychev polynomials defined by $T_N(z) = \cos N(\arccos z)$.

We set some classic notations for the text. We will denote the columns of order m of a matrix by C_m ; the classical binomial coefficient $\frac{n!}{p!(n-p)!}$ will be denoted by C_n^p .

In the following, we will compute $W(f_1, \dots, f_{i-1}, f_{i+1}, \dots, f_m)$ and $W(f_1, \dots, f_m)$ for a certain type of functions f_i , following the ideas of [10, 7, 16]. Then we will be able to evaluate explicitly the Hadamard's coefficients and solutions to the two-dimensional Schrödinger equations with Huygens' potentials.

2 Computation of Wronskians

2.1 First type of formulas

Let m be some non negative integer, α and x_0 be some real parameters, and f_j the functions defined as follows :

$$f_j = \sin \alpha j(x - x_0), \quad 1 \leq j \leq m. \quad (7)$$

We use below the standard notation,

$$W(f_1, f_2, \dots, f_m) := \det \partial_x^{j-1} f_k(x); \quad j, k = 1, \dots, m, \quad (8)$$

for the Wronskian determinant of any m functions and $[x]$ the largest integer less than or equal to x .

We will denote W_{m-1}^i the determinant $W(f_1, \dots, f_{i-1}, f_{i+1}, \dots, f_m)$.

We will use the following result [5, 6] :

Proposition 2.1 *For any smooth functions $f, \varphi_1, \dots, \varphi_n$, the Wronskian $W(f \cdot \varphi_1, \dots, f \cdot \varphi_n)$ verify*

$$W(f \cdot \varphi_1, \dots, f \cdot \varphi_n) = (f(x))^n W(\varphi_1, \dots, \varphi_n), \quad (9)$$

and the formula of change of independent variable in a Wronskian :

Proposition 2.2 *For $t = \varphi(x)$ a new variable so that $f_j(x) = \tilde{f}_j(t)$,*

$$W(f_1, \dots, f_n) = (\varphi'(x))^{\frac{n(n-1)}{2}} \cdot W(\tilde{f}_1, \tilde{f}_2, \dots, \tilde{f}_n). \quad (10)$$

With these notations we have the following formula

Theorem 2.1

$$\begin{aligned} W_{m-1}^i &= i(2)^{\frac{m(m-1)}{2}-i+1} (-\alpha)^{\frac{(m-1)(m-2)}{2}} \prod_{k=1}^{m-1} k! \times \sin^{\frac{m(m-1)}{2}} \alpha(x - x_0) \\ &\times \left(\sum_{k=0}^{[(m-i)/2]} \frac{2^{-2k}}{(i+k)!k!} \frac{\cos^{m-i-2k} \alpha(x - x_0)}{(m-i-2k)!} \right), \quad 1 \leq i \leq m. \end{aligned} \quad (11)$$

Proof:

We start with the cases where $i = m$ and $i = m - 1$.

The details of the justifications will be given in the general case.

1. For W_{m-1}^m , all functions involved in the determinant are of the form $\sin(px)$. We use the expansion

$$\begin{aligned} \sin px &= \sin x \sum_{k=0}^{\lfloor \frac{p-1}{2} \rfloor} (-1)^k C_{p-k-1}^k \cdot (2 \cos x)^{p-1-2k} = \\ &\sin x [(2 \cos x)^{p-1} - (p-2)(2 \cos x)^{p-3} + \dots]. \end{aligned} \quad (12)$$

the combinations on the columns and the previous properties (9), (10) to get the series of equalities as already used in [10, 7, 16]

$$\begin{aligned}
W_{m-1}^m &= W(\sin \alpha(x - x_0), \sin 2\alpha(x - x_0), \dots, \sin(m-1)\alpha(x - x_0)) \\
&= 2^{\frac{(m-1)(m-2)}{2}} W(\sin \alpha(x - x_0), \sin \alpha(x - x_0) \cos \alpha(x - x_0), \dots, \\
&\quad \dots, \sin \alpha(x - x_0) \cos^{m-2} \alpha(x - x_0)) \\
&= 2^{\frac{(m-1)(m-2)}{2}} (\sin \alpha(x - x_0))^{m-1} W(1, \cos \alpha(x - x_0), \dots, \cos^{m-2} \alpha(x - x_0)) \\
&= 2^{\frac{(m-1)(m-2)}{2}} (-\alpha)^{\frac{(m-1)(m-2)}{2}} \sin^{\frac{m(m-1)}{2}} \alpha(x - x_0) W(1, t, \dots, t^{m-2}) \\
&\quad (\text{by the change } \cos \alpha(x - x_0) = t); \text{ thus} \\
W_{m-1}^m &= 2^{\frac{(m-1)(m-2)}{2}} (-\alpha)^{\frac{(m-1)(m-2)}{2}} \prod_{j=1}^{m-2} j! \sin^{\frac{m(m-1)}{2}} \alpha(x - x_0).
\end{aligned}$$

So we get (11) for the case $i = m$.

2. For W_{m-1}^{m-1} , we use the same ideas that these given in the case $i = m$ and previous works [10, 7, 16]. We use the expansion 12, the combinations on the columns and the previous properties (9), (10) to get the following equalities :

$$\begin{aligned}
W_{m-1}^{m-1} &= W(\sin \alpha(x - x_0), \sin 2\alpha(x - x_0), \dots, \sin(m-2)\alpha(x - x_0), \sin m\alpha(x - x_0)), \\
&= 2^{\frac{m(m-1)}{2}-(m-2)} W(\sin \alpha(x - x_0), \sin \alpha(x - x_0) \cos \alpha(x - x_0), \dots, \\
&\quad \dots, \sin \alpha(x - x_0) \cos^{m-3} \alpha(x - x_0), \sin \alpha(x - x_0) \cos^{m-1} \alpha(x - x_0)) \\
&= 2^{\frac{m(m-1)}{2}-(m-2)} \sin^{m-1} \alpha(x - x_0) W(1, \cos \alpha(x - x_0), \dots, \\
&\quad \dots, \cos^{m-3} \alpha(x - x_0), \cos^{m-1} \alpha(x - x_0)). \\
&= 2^{\frac{m(m-1)}{2}-(m-2)} (-\alpha)^{\frac{(m-1)(m-2)}{2}} \sin^{\frac{m(m-1)}{2}} (\alpha(x - x_0) W(1, t, \dots, t^{m-3}, t^{m-1}) \\
&\quad (\text{by the change } \cos \alpha(x - x_0) = t); \text{ thus} \\
W_{m-1}^{m-1} &= 2^{\frac{m(m-1)}{2}-(m-2)} (-\alpha)^{\frac{(m-1)(m-2)}{2}} \prod_{j=1, j \neq m-2}^{m-1} j! \sin^{\frac{m(m-1)}{2}} \alpha(x - x_0) \cos \alpha(x - x_0).
\end{aligned}$$

This gives exactly formula (11) for $i = m - 1$.

3. Let i be an integer such that $1 \leq i \leq m - 2$.

First we use the expansion

$$\begin{aligned}
\sin px &= \sin x \sum_{k=0}^{\lfloor \frac{p-1}{2} \rfloor} (-1)^k C_{p-k-1}^k \cdot (2 \cos x)^{p-1-2k} = \\
&\quad \sin x [(2 \cos x)^{p-1} - (p-2)(2 \cos x)^{p-3} + \dots].
\end{aligned} \tag{13}$$

Therefore

$$\begin{aligned}
W_{m-1}^i &= W(\sin \alpha(x - x_0), \sin 2\alpha(x - x_0), \dots, \\
&\quad \sin(i-1)\alpha(x - x_0), \sin(i+1)\alpha(x - x_0), \dots, \sin m\alpha(x - x_0)),
\end{aligned}$$

can be reduced by replacing each term in the expansion of $\sin p\alpha(x - x_0)$, formula (13) by their first terms for $1 \leq p \leq i$, and p of the form $i + 2k$,

$1 \leq k \leq [(m-i)/2]$.

We can factorize in each column p the term 2^{p-1} for $1 \leq p \leq i-1$.

So we get

$$\begin{aligned} W_{m-1}^i &= 2^{\frac{(i-1)(i-2)}{2}} W(\sin \alpha(x - x_0), \sin \alpha(x - x_0) \cos \alpha(x - x_0), \dots, \\ &\quad \sin \alpha(x - x_0) \cos^{i-2} \alpha(x - x_0), 2^i \sin \alpha(x - x_0) \cos^i \alpha(x - x_0), \\ &\quad 2^{i+1} \sin \alpha(x - x_0) \cos^{i+1} \alpha(x - x_0) - i 2^{i-1} \sin \alpha(x - x_0) \cos^{i-1} \alpha(x - x_0), \\ &\quad \dots, 2^{m-1} \sin \alpha(x - x_0) \cos^{m-1} \alpha(x - x_0) + \dots). \end{aligned} \tag{14}$$

Using the previous propositions 10, 10, then W_{m-1}^i can be written

$$\begin{aligned} W_{m-1}^i &= 2^{\frac{(i-1)(i-2)}{2}} (\sin \alpha(x - x_0))^{m-1} W(1, \cos \alpha(x - x_0), \dots, \\ &\quad \cos^{i-2} \alpha(x - x_0), 2^i \cos^i \alpha(x - x_0), 2^{i+1} \cos^{i+1} \alpha(x - x_0) - i 2^{i-1} \cos^{i-1} \alpha(x - x_0), \\ &\quad \dots, 2^{m-1} \cos^{m-1} \alpha(x - x_0) + \dots). \end{aligned} \tag{15}$$

Then, we make the following variable change defined by $t = \cos \alpha(x - x_0)$ is made and we can factorize $(-\alpha \sin \alpha(x - x_0))^{\frac{(m-1)(m-2)}{2}}$.

Therefore the Wronskian W_{m-1}^i can be written as

$$\begin{aligned} W_{m-1}^i &= 2^{\frac{(i-1)(i-2)}{2}} (-\alpha)^{\frac{(m-1)(m-2)}{2}} (\sin \alpha x)^{\frac{m(m-1)}{2}} \\ &\quad \cdot \tilde{W}_{m-1}^i(1, t, \dots, t^{i-2}, 2^i t^i, 2^{i+1} t^{i+1} - i 2^{i-1} t^{i-1}, 2^{i+2} t^{i+2}, \\ &\quad 2^{i+3} t^{i+3} - (i+2) 2^{i+1} t^{i+1} + \frac{i(i+1)}{2} 2^{i-1} t^{i-1}, \\ &\quad \dots, 2^{m-1} t^{m-1} - (m-2) 2^{m-3} t^{m-3} + \dots + (-1)^{\frac{m-1}{2}} 2^{i-1} C_{(m+i)/2-1}^{(m-i)/2} t^{i-1}). \end{aligned} \tag{16}$$

If $m - i - 1 = 2p$, we can factorize $2^i 2^{i+2} \dots 2^{m-1} (2^{i-1})^{\frac{(m-i-1)}{2}}$ which is equal to $2^{\frac{(m-i+1)(m+i-1)}{4} + \frac{(m-i-1)(i-1)}{2}}$.

If $m - i - 1 = 2p + 1$, we can factorize $2^i 2^{i+2} \dots 2^{m-2} (2^{i-1})^{\frac{(m-i)}{2}}$ which is equal to $2^{\frac{(m-i)(m+i-2)}{4} + \frac{(m-i)(i-1)}{2}}$.

So, the Wronskian $\tilde{W}_{m-1}^i := W(1, t, \dots, t^{i-2}, 2^i t^i, 2^{i+1} t^{i+1} - i 2^{i-1} t^{i-1}, \dots, 2^{m-1} t^{m-1} - (m-2) 2^{m-3} t^{m-3} + \dots)$ can be written as

$$\begin{aligned} \tilde{W}_{m-1}^i &= \prod_{j=1}^{i-2} j! 2^{\frac{(m-i+1)(m+i-1)}{4} + \frac{(m-i-1)(i-1)}{2}} \tilde{w}_{m-1,e}^i, \text{ if } m - i - 1 = 2p \text{ is even ,} \\ \tilde{W}_{m-1}^i &= \prod_{j=1}^{i-2} j! 2^{\frac{(m-i)(m+i-2)}{4} + \frac{(m-i)(i-1)}{2}} \tilde{w}_{m-1,o}^i, \text{ if } m - i - 1 = 2p + 1 \text{ is odd} \end{aligned} \tag{17}$$

where $\tilde{w}_{m-1,o}^i$ is the determinant if $m - i - 1 = 2p + 1$ is odd defined by,

$$\left| \begin{array}{cccccc} i!t & 2^2(i+1)!t^2/2! - i(i-1)! & \dots & (m-2)!\frac{t^{m-i-1}}{(m-i-1)!} & 2^{m-i}(m-1)!\frac{t^{m-i}}{(m-i)!} + \dots + (-1)^{(m-i)/2}C_{(m+i)/2-1}^{(m-i)/2}2^{i-1}(i-1)! \\ i! & 2^2(i+1)!t & \dots & (m-2)!\frac{t^{m-i-2}}{(m-i-2)!} & 2^{m-i}(m-1)!\frac{t^{m-i-1}}{(m-i-1)!} + \dots + (-1)^{(m-i)/2+1}C_{(m+i)/2}^{(m-i)/2+1}2^{i+1}(i+1)t \\ 0 & 2^2(i+1)! & \dots & (m-2)!\frac{t^{m-i-3}}{(m-i-3)!} & 2^{m-i}(m-1)!\frac{t^{m-i-2}}{(m-i-2)!} + \dots + (-1)^{(m-i)/2+1}C_{(m+i)/2}^{(m-i)/2+1}2^{i+1}(i+1)! \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \dots & (m-2)! & 2^{m-i}(m-1)!t \end{array} \right| \quad (18)$$

and $\tilde{w}_{m-1,e}^i$ is the determinant if $m - i - 1 = 2p$ is even, defined by,

$$\left| \begin{array}{cccccc} i!t & 2^2(i+1)!t^2/2! - i(i-1)! & \dots & 2^{m-i-1}(m-2)!\frac{t^{m-i-1}}{(m-i-1)!} + \dots + (-1)^{(m-i-1)/2}C_{(m+i-1)/2}^{(m-i-1)/2}2^{i-1}(i-1)! & (m-1)!\frac{t^{m-i}}{(m-i)!} \\ i! & 2^2(i+1)!t & \dots & 2^{m-i-1}(m-2)!\frac{t^{m-i-2}}{(m-i-2)!} + \dots + (-1)^{(m-i-1)/2+1}C_{(m+i-1)/2+1}^{(m-i-1)/2+1}2^{i+1}(i+1)t & (m-1)!\frac{t^{m-i-1}}{(m-i-1)!} \\ 0 & 2^2(i+1)! & \dots & 2^{m-i-1}(m-2)!\frac{t^{m-i-3}}{(m-i-3)!} + \dots + (-1)^{(m-i-1)/2+1}C_{(m+i-1)/2+1}^{(m-i-1)/2+1}2^{i+1}(i+1)! & (m-1)!\frac{t^{m-i-2}}{(m-i-2)!} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \dots & 2^{m-i-1}(m-2)! & (m-1)!t \end{array} \right| \quad (19)$$

In the following, we calculate $\tilde{w}_{m-1,o}^i$ and $\tilde{w}_{m-1,e}^i$.

All the columns of the determinant $\tilde{w}_{m-1,o}^i$ and $\tilde{w}_{m-1,e}^i$ of the form C_{i+2k} can be written as

$$C_{i+2k}^* = {}^t ((i+2k)!\frac{t^{2k+1}}{(2k+1)!}; (i+2k)!\frac{t^{2k}}{(2k)!}; \dots; (i+2k)!; 0; \dots; 0) \quad (20)$$

It remains to reduce the columns of the determinant $\tilde{w}_{m-1,o}^i$ and $\tilde{w}_{m-1,e}^i$ of the form C_{i+2k+1} .

a. We begin with the case $m - i - 1 = 2p$.

We consider the column

$$C_{i+1}^* = {}^t (2^2(i+1)!\frac{t^2}{2!} - i!; 2^2(i+1)!t; 2^2(i+1)!; 0; \dots; 0)$$

On each column C_{i+2k+1} of \tilde{w}_{m-1}^i , $1 \leq k \leq (m - i - 1)/2$, we make the following transformations

$$\begin{aligned} C_{i+3} &\rightarrow C_{i+3} + (i+2)C_{i+1}^* = C_{i+3}^* \\ &= {}^t (2^4(i+3)!\frac{t^4}{4!} - \frac{i!(i+3)}{2!}; 2^4(i+3)!\frac{t^3}{3!}; 2^4(i+3)!\frac{t^2}{2!}; 2^4(i+3)!t; 2^4(i+3)!; 0; \dots; 0) \end{aligned} \quad (21)$$

$$\begin{aligned} C_{i+5} &\rightarrow C_{i+5} + (i+4)C_{i+3}^* - \frac{(i+3)(i+2)}{2}C_{i+1}^* = C_{i+5}^* \\ &= {}^t (2^6(i+5)! \frac{t^6}{6!} - \frac{i!(i+4)(i+5)}{3!}; 2^6(i+5)! \frac{t^5}{5!}; 2^6(i+5)! \frac{t^4}{4!}; 2^6(i+5)! \frac{t^3}{3!} \\ &\quad ; 2^6(i+5)! \frac{t^2}{2!}; 2^6(i+5)!t; 2^6(i+5)!; 0; \dots; 0) \end{aligned} \quad (22)$$

until

$$\begin{aligned} C_{i+2k+1} &\rightarrow C_{i+2k+1} + (i+2k)C_{i+2k-1}^* - \frac{(i+2k-1)(i+2k-2)}{2!}C_{i+2k-3}^* + \dots \\ &\dots + (-1)^{k+1} \frac{(i+k+1)\dots(i+2)}{k!}C_{i+1}^* = C_{i+2k+1}^* \end{aligned} \quad (23)$$

We claim that

$$\begin{aligned} C_{i+2k+1}^* &= {}^t (2^{2k+2}(i+2k+1)! \frac{t^{2k+2}}{(2k+2)!} - \frac{i!(i+2k+1)\dots(i+k+2)}{(k+1)!}; \\ &2^{2k+2}(i+2k+1)! \frac{t^{2k+1}}{(2k+1)!}; \dots; 2^{2k+2}(i+2k+1)!; 0; \dots; 0). \end{aligned} \quad (24)$$

Suppose that the column C_{i+2i+1}^* can be written as in (24) for $i \leq k$. We have clearly the relation for $k = 1$ (21) and $k = 2$ (22).

Then we make the following transformation on the column C_{i+2k+3} :

$$\begin{aligned} C_{i+2k+3} &\rightarrow C_{i+2k+3} + (i+2k+2)C_{i+2k+1}^* - \frac{(i+2k+1)(i+2k)}{2!}C_{i+2k-1}^* \\ &+ \dots + (-1)^{k+2} \frac{(i+k+2)\dots(i+2)}{(k+1)!}C_{i+1}^* = C_{i+2k+3}^*. \end{aligned}$$

Clearly, all the rows from 2 to $m-1$ of the column C_{i+2k+3}^* take the form

$$(2^{k+4}(i+2k+3)! \frac{t^{2k+3}}{(2k+3)!}, 2^{k+4}(i+2k+3)! \frac{t^{2k+2}}{(2k+2)!}, \dots, 2^{k+4}(i+2k+3)!, 0, \dots, 0). \quad (25)$$

It remains to evaluate the first row of the column C_{i+2k+3} .

In a shortened way, the first row can be written as

$$(C_{i+2k+3}^*)_1 = -i! \sum_{j=1}^{k+2} \frac{(-1)^{j+1}}{i+k-2-j} C_{i+2k+3-j}^j C_{i+2k+3-2j}^{k+2-j}.$$

By reduction, it can be written as

$$(C_{i+2k+3}^*)_1 = -i! \frac{C_{i+2k+3}^{k+2}}{i+k+2} = -i! \frac{C_{i+2k+3}^{k+1}}{k+2} = -i! \frac{\prod_{j=0}^k (i+2k+3-j)}{(k+2)!}. \quad (26)$$

The relations (25) and (26) gives exactly the relation (24) at the rank $k+1$.

Then we can write the determinant $\tilde{w}_{m-1,e}^i$ defined in (18) as the following sum $\tilde{w}_{m-1,e}^{i,*} + \tilde{w}_{m-1,e}^{i,**}$ where $\tilde{w}_{m-1,e}^{i,*}$ is defined by

$$\tilde{w}_{m-1,e}^{i,*} = \begin{vmatrix} i!t & 2^2(i+1)!\frac{t^2}{2!} & \dots & 2^{m-i-1}(m-2)!\frac{t^{m-i-1}}{(m-i-1)!} & (m-1)!\frac{t^{m-i}}{(m-i)!} \\ i! & 2^2(i+1)!t & \dots & 2^{m-i-1}(m-2)!\frac{t^{m-i-2}}{(m-i-2)!} & (m-1)!\frac{t^{m-i-1}}{(m-i-1)!} \\ 0 & 2^2(i+1)! & \dots & 2^{m-i-1}(m-2)!\frac{t^{m-i-3}}{(m-i-3)!} & (m-1)!\frac{t^{m-i-2}}{(m-i-2)!} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \dots & 2^{m-i-1}(m-2)! & (m-1)!t \end{vmatrix}.$$

$\tilde{w}_{m-1,e}^{i,**}$ is equal to

(27)

$$\tilde{w}_{m-1,e}^{i,**} = \begin{vmatrix} 0 & -(i)! & 0 & \dots & 0 & -\frac{C_{i+2k+1}^k}{(k+1)} & 0 & -\frac{(i)!\frac{C_{m-2}^{(m-i-3)/2}}{(m-i-1)/2}}{(m-i-1)!} & 0 \\ i! & 2^2(i+1)!t & (i+2)!\frac{t^2}{2!} & \dots & \dots & 2^{2k+2}(i+2k+1)!\frac{t^{2k+1}}{(2k+1)!} & \dots & 2^{m-i-1}(m-2)!\frac{t^{m-i-2}}{(m-i-2)!} & (m-1)!\frac{t^{m-i-1}}{(m-i-1)!} \\ 0 & 2^2(i+1)! & (i+2)!t & \dots & \dots & 2^{2k+2}(i+2k+1)!\frac{t^{2k}}{(2k)!} & \dots & 2^{m-i-1}(m-2)!\frac{t^{m-i-3}}{(m-i-3)!} & (m-1)!\frac{t^{m-i-2}}{(m-i-2)!} \\ \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & 0 & \dots & 2^{m-i-1}(m-2)! & (m-1)!t \end{vmatrix}.$$

(28)

Clearly, we have

$$\tilde{w}_{m-1,e}^{i,*} = \prod_{j=1}^{m-1} j! \frac{t^{m-i}}{(m-i)!} 2^{\frac{(m-i-1)(m-i+1)}{4}}.$$

For $\tilde{w}_{m-1,e}^{i,**}$, we have

$$\tilde{w}_{m-1,e}^{i,**} = i! \sum_{k=0}^{(m-i-3)/2} \frac{C_{i+2k+1}^k}{(k+i)} \delta_{i+2k+1}.$$

where δ_{i+2k+1} is the determinant defined by

$$\delta_{i+2k+1} = \begin{vmatrix} i! & 2^2(i+1)!t & \dots & (i+2k)!\frac{t^{2k}}{(2k)!} & (i+2k+2)!\frac{t^{2k+2}}{(2k+2)!} & \dots & 2^{m-i-1}(m-2)!\frac{t^{m-i-2}}{(m-i-2)!} & (m-1)!\frac{t^{m-i-1}}{(m-i-1)!} \\ 0 & 2^2(i+1)! & \dots & (i+2k)!\frac{t^{2k-1}}{(2k-1)!} & (i+2k+2)!\frac{t^{2k+1}}{(2k+1)!} & \dots & 2^{m-i-1}(m-2)!\frac{t^{m-i-3}}{(m-i-3)!} & (m-1)!\frac{t^{m-i-2}}{(m-i-2)!} \\ \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & 0 & \dots & 2^{m-i-1}(m-2)! & (m-1)!t \end{vmatrix}.$$

δ_{i+2k+1} is equal to

$$\delta_{i+2k+1} = 2^{\frac{(m-i+1)(m-i-1)}{4} - (2k+2)} \prod_{j=i, j \neq i+2k+1}^{m-1} j! \frac{t^{m-i-2k-2}}{(m-i-2k-2)!}.$$

So $\tilde{w}_{m-1,e}^{i,**}$ can be written as

$$\tilde{w}_{m-1,e}^{i,**} = i! \sum_{k=0}^{(m-i-3)/2} \frac{C_{i+2k+1}^k}{(k+1)} 2^{\frac{(m-i+1)(m-i-1)}{4} - (2k+2)} \prod_{j=i, j \neq i+2k+1}^{m-1} j! \frac{t^{m-i-2k-2}}{(m-i-2k-2)!}. \quad (30)$$

Taking into account (29) and (30), with (17) we have for W

$$\begin{aligned} \tilde{W}_{m-1}^i &= \prod_{j=1}^{i-2} j! 2^{\frac{(m-i+1)(m+i-1)}{4} + \frac{(m-i-1)(i-1)}{2}} \left(\prod_{j=i}^{m-1} j! \frac{t^{m-i}}{(m-i)!} 2^{\frac{(m-i-1)(m-i+1)}{4}} \right. \\ &\quad \left. + i! \sum_{k=0}^{(m-i-3)/2} \frac{C_{i+2k+1}^k}{(k+1)} 2^{\frac{(m-i+1)(m-i-1)}{4} - (2k+2)} \prod_{j=i, j \neq i+2k+1}^{m-1} j! \frac{t^{m-i-2k-2}}{(m-i-2k-2)!} \right) \end{aligned} \quad (31)$$

Thus, we have

$$\begin{aligned} \text{if } m-i-1 &= 2p, \\ W_{m-1}^i &= (2)^{\frac{(m-1)(m-i+1)}{2} + \frac{(i-1)(m-3)}{2}} (-\alpha)^{\frac{(m-1)(m-2)}{2}} \sin^{\frac{m(m-1)}{2}} \alpha(x-x_0) \prod_{j=0}^{i-2} j! \\ &\cdot \left(\prod_{j=i}^{m-1} j! \frac{\cos^{m-i} \alpha(x-x_0)}{(m-i)!} + i! \cdot \sum_{k=0}^{(m-i-3)/2} \frac{C_{i+2k+1}^k}{k+1} 2^{-2k-2} \cdot \prod_{j=i, j \neq i+2k+1}^{m-1} j! \frac{\cos^{m-i-2k-2} \alpha(x-x_0)}{(m-i-2k-2)!} \right), \end{aligned} \quad (32)$$

Taking into account that $\frac{m(m-i)}{2} + \frac{(i-1)(m-2)}{2} = \frac{m(m-1)}{2} - (i-1)$, and $\frac{m-i-3}{2} = [\frac{m-i-2}{2}]$, we have :

$$W_{m-1}^i = (2)^{\frac{m(m-1)}{2} - (i-1)} (-\alpha)^{\frac{(m-1)(m-2)}{2}} \sin^{\frac{m(m-1)}{2}} \alpha(x-x_0) \cdot i \prod_{j=1}^{m-1} j! \left(\frac{\cos^{m-i} \alpha(x-x_0)}{i!(m-i)!} + \sum_{k=0}^{[m-i-2]/2} \frac{2^{-2(k+1)}}{(i+k+1)!(k+1)!} \frac{\cos^{m-i-2k-2} \alpha(x-x_0)}{(m-i-2k-2)!} \right),$$

which can be rewritten as

$$W_{m-1}^i = (2)^{\frac{m(m-1)}{2} - (i-1)} (-\alpha)^{\frac{(m-1)(m-2)}{2}} \sin^{\frac{m(m-1)}{2}} \alpha(x-x_0) \cdot i \prod_{j=1}^{m-1} j! \left(\frac{\cos^{m-i} \alpha(x-x_0)}{i!(m-i)!} + \sum_{k=1}^{[m-i]/2} \frac{2^{-2k}}{(i+k)!(k)!} \frac{\cos^{m-i-2k} \alpha(x-x_0)}{(m-i-2k)!} \right).$$

It is easy to see that we get exactly (11).

b. We consider now the case $m-i-1 = 2p+1$.

The ideas are the same, but the structure of $\tilde{w}_{m-1,o}^i$ is different (the two last columns are different).

On each column C_{i+2k+1} of $\tilde{w}_{m-1,o}^i$, $1 \leq k \leq m-i-1$, we make the same transformations as defined in the case $m-i-1 = 2p+1$, i.e. (21), (22) and (23) for the column C_{i+2k+1} .

We have the same relation (24) whose proof is the same. Clearly, all the rows

from 1 to $m - 1$ of the column C_{i+2k+3}^* take the same form as in (25) and (26). Then we can write the determinant $\tilde{w}_{m-1,o}^i$ defined in (19) as the following sum $\tilde{w}_{m-1,o}^{i,*} + \tilde{w}_{m-1,o}^{i,**}$ where $\tilde{w}_{m-1,o}^{i,*}$ is defined by

$$\tilde{w}_{m-1,o}^{i,*} = \begin{vmatrix} i!t & 2^2(i+1)!\frac{t^2}{2!} & \dots & (m-2)!\frac{t^{m-i-1}}{(m-i-1)!} & 2^{m-i}(m-1)!\frac{t^{m-i}}{(m-i)!} \\ i! & 2^2(i+1)!t & \dots & (m-2)!\frac{t^{m-i-2}}{(m-i-2)!} & 2^{m-i}(m-1)!\frac{t^{m-i-1}}{(m-i-1)!} \\ 0 & 2^2(i+1)! & \dots & 2^{m-i-1}(m-2)!\frac{t^{m-i-3}}{(m-i-3)!} & (m-1)!\frac{t^{m-i-2}}{(m-i-2)!} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \dots & (m-2)! & 2^{m-i}(m-1)!t \end{vmatrix}.$$

$\tilde{w}_{m-1,o}^{i,**}$ is equal to

$$\tilde{w}_{m-1,o}^{i,**} = \begin{vmatrix} 0 & -(i)! & 0 & \dots & 0 & -(i)!\frac{C_{i+2k+1}^k}{(k+1)} & 0 & 0 & -(i)!\frac{C_{m-1}^{(m-i-2)/2}}{(m-i)/2} \\ i! & 2^2(i+1)!t & (i+2)!\frac{t^2}{2!} & \dots & \dots & 2^{2k+2}(i+2k+1)!\frac{t^{2k+1}}{(2k+1)!} & \dots & (m-2)!\frac{t^{m-i-2}}{(m-i-2)!} & 2^{m-i}(m-1)!\frac{t^{m-i-1}}{(m-i-1)!} \\ 0 & 2^2(i+1)! & (i+2)!t & \dots & \dots & 2^{2k+2}(i+2k+1)!\frac{t^{2k}}{(2k)!} & \dots & (m-2)!\frac{t^{m-i-3}}{(m-i-3)!} & 2^{m-i}(m-1)!\frac{t^{m-i-2}}{(m-i-2)!} \\ \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & 0 & \dots & (m-2)! & 2^{m-i}(m-1)!t \end{vmatrix} \quad (34)$$

We have

$$\tilde{w}_{m-1,o}^{i,*} = \prod_{j=1}^{m-1} j! \frac{t^{m-i}}{(m-i)!} 2^{\frac{(m-i)(m-i+2)}{4}}. \quad (35)$$

$\tilde{w}_{m-1,o}^{i,**}$ can be written as

$$\tilde{w}_{m-1,o}^{i,**} = i! \sum_{k=0}^{(m-i-2)/2} \frac{C_{i+2k+1}^k}{(k+1)} 2^{\frac{(m-i+2)(m-i)}{4} - (2k+2)} \prod_{j=i, j \neq i+2k+1}^{m-1} j! \frac{t^{m-i-2k-2}}{(m-i-2k-2)!}. \quad (36)$$

Taking into account (35) and (36), with (17) we have for W

$$W = \prod_{j=1}^{i-2} j! 2^{\frac{(m-i)(m+i-2)}{4} + \frac{(m-i)(i-1)}{2}} \left(\prod_{j=i}^{m-1} j! \frac{t^{m-i}}{(m-i)!} 2^{\frac{(m-i)(m-i+2)}{4}} \right. \\ \left. + i! \sum_{k=0}^{(m-i-2)/2} \frac{C_{i+2k+1}^k}{(k+1)} 2^{\frac{(m-i+2)(m-i)}{4} - (2k+2)} \prod_{j=i, j \neq i+2k+1}^{m-1} j! \frac{t^{m-i-2k-2}}{(m-i-2k-2)!} \right) \quad (37)$$

Thus we are able to give the expression of W_i^{m-1}

$$\begin{aligned} \text{if } m - i - 1 = 2p + 1, \\ W_{m-1}^i = (2)^{\frac{m(m-i)}{2} + \frac{(i-1)(m-2)}{2}} (-\alpha)^{\frac{(m-1)(m-2)}{2}} \sin^{\frac{m(m-1)}{2}} \alpha(x - x_0) \prod_{j=1}^{i-2} j! \\ \cdot \left(\prod_{j=i}^{m-1} j! \frac{\cos^{m-i} \alpha(x - x_0)}{(m-i)!} + i! \cdot \sum_{k=0}^{(m-i-2)/2} \frac{C_{i+2k+1}^k}{k+1} 2^{-2k-2} \cdot \prod_{j=i, j \neq i+2k+1}^{m-1} j! \frac{\cos^{m-i-2k-2} \alpha(x - x_0)}{(m-i-2k-2)!} \right), \end{aligned} \quad (38)$$

Taking into account that $\frac{m-i-2}{2} = [\frac{m-i-2}{2}]$, and after some simplifications like these done in the part a., we have the expression of W_i^{m-1} according (11).

□

Remark 2.1 The possible extension of the same calculations to the sequence of the hyperbolic functions can be easily made by the change $\alpha \rightarrow i\alpha$.

2.2 Second type of formulas

Let m some non negative integer, α and x_0 be some real parameters, and g_j the functions defined as follows :

$$g_j = \cos \alpha j(x - x_0), \quad 1 \leq j \leq m. \quad (39)$$

We use the standard notation (8) for the Wronskian determinant of any m functions.

We will denote \overline{W} the determinant $W(g_1, \dots, g_m)$.

With these notations we have the following formulas

Theorem 2.2

$$\overline{W} = 2^{\frac{m(m-1)}{2}} \prod_{k=1}^m k! (-\alpha)^{\frac{m(m-1)}{2}} \sin^{\frac{m(m-1)}{2}} \alpha(x - x_0) \times \left(\sum_{k=0}^{[m/2]} \frac{2^{-2k}}{k!^2} \frac{\cos^{m-2k} \alpha(x - x_0)}{(m-2k)!} \right). \quad (40)$$

Proof

First we use the expansion

$$\begin{aligned} \cos px &= 2^{p-1} \cos^p x + \sum_{k=1}^{[\frac{p}{2}]} \frac{p}{k} (-1)^k C_{p-1-k}^{k-1} \cdot 2^{p-1-2k} \cos^{p-1-2k} x = \\ &2^{p-1} \cos^p x - p 2^{p-3} \cos^{p-2} x + \dots \end{aligned} \quad (41)$$

We replace each term $\cos p\alpha(x - x_0)$ by its expansion (formula 41).

Then we use the formula of change of independent variable (10) defined by

$t = \cos \alpha(x - x_0)$ and we can factorize $(-\alpha \sin \alpha(x - x_0))^{\frac{(m)(m-1)}{2}}$. Then \bar{W} can be written

$$\bar{W} = (-\alpha \sin \alpha(x - x_0))^{\frac{m(m-1)}{2}} W(t, 2t^2 - 1, \dots, 2^{m-1}t^m - m 2^{m-3}t^{m-2} + \dots). \quad (42)$$

We denote in the following w the Wronskian

$$w := W(t, 2t^2 - 1, \dots, 2^{m-1}t^m - m 2^{m-3}t^{m-2} + \dots). \quad (43)$$

In the following, we calculate w .

We are going to define the succession of transformations on the columns in the following .

All the odd columns of the form C_{2k+1} can be easily reduced to

$$C_{2k+1}^* = {}^t (2^{2k}(2k+1)! \frac{t^{2k+1}}{(2k+1)!}; 2^{2k}(2k+1)! \frac{t^{2k}}{(2k)!}; \dots; 2^{2k}(2k+1)!; 0; \dots; 0) \quad (44)$$

It remains to reduce the even columns of the form C_{2k+2} .

We consider the second column

$$C_2^* = {}^t (2 \cdot 2! \frac{t^2}{2!} - 1; 2 \cdot 2!t; 2 \cdot 2!; 0; \dots; 0). \quad (45)$$

On each column C_{2k} of w , $k \geq 2$, we make the following transformations

$$\begin{aligned} C_4 &\rightarrow C_4 + 4C_2^* = C_4^* \\ &= {}^t (2^3 \cdot 4! \frac{t^4}{4!} - 3; 2^3 \cdot 4! \frac{t^3}{3!}; 2^3 \cdot 4! \frac{t^2}{2!} 2^3 \cdot 4!t; 2^3 \cdot 4!; 0; \dots; 0) \end{aligned} \quad (46)$$

$$\begin{aligned} C_6 &\rightarrow C_6 + 6C_4^* - \frac{6 \times 3}{2} C_2^* = C_6^* \\ &= {}^t (2^5 \cdot 6! \frac{t^6}{6!} - 10; 2^5 \cdot 6! \frac{t^5}{5!}; 2^5 \cdot 6! \frac{t^4}{4!} 2^5 \cdot 6! \frac{t^3}{3!} \\ &\quad ; 2^5 \cdot 6! \frac{t^2}{2!}; 2^5 \cdot 6!t; 2^5 \cdot 6!; 0; \dots; 0) \end{aligned} \quad (47)$$

until

$$\begin{aligned} C_{2k} &\rightarrow C_{2k} + (2k)C_{2k-2}^* - \frac{(2k)(2k-3)}{2!} C_{2k-4}^* + \dots \\ &\quad \dots + (-1)^k \frac{2k}{k-1} \frac{(2k-1-(k-1)) \dots (2k-1-(k-1+k-1-2))}{(k-2)!} C_2^* = C_{2k}^* \end{aligned} \quad (48)$$

which can be rewritten as

$$\begin{aligned} C_{2k} &\rightarrow C_{2k} + (2k)C_{2k-2}^* - \frac{(2k)(2k-3)}{2!} C_{2k-4}^* + \dots \\ &\quad \dots + (-1)^k k^2 C_2^* = C_{2k}^*. \end{aligned} \quad (49)$$

We claim that

$$\begin{aligned} C_{2k}^* &= {}^t (2^{2k-1}(2k)!\frac{t^{2k}}{(2k)!} - C_{2k-1}^{k-1}; \\ &\quad 2^{2k-1}(2k)!\frac{t^{2k-1}}{(2k-1)!}; \dots; 2^{2k-1}(2k)!; 0; \dots; 0). \end{aligned} \quad (50)$$

Suppose that the column C_{2i}^* can be written as in (24) for $i \leq k$.

We have clearly the relation for $k = 1$ (45), $k = 2$ (46) and $k = 3$ (47). Then we make the following transformation on the column C_{2k+2} :

$$\begin{aligned} C_{2k+2} &\rightarrow C_{2k+2} + (2k+2)C_{2k}^* - \frac{(2k+2)(2k-1)}{2!}C_{2k-2}^* \\ &+ \dots + (-1)^{k+1}(k+1)^2C_2^* = C_{2k+2}^*. \end{aligned}$$

Clearly, all the rows from 2 to m of the column C_{2k+2}^* take the form

$$(2^{2k+1}(2k+2)!\frac{t^{2k+1}}{(2k+1)!}, (2^{2k+1}(2k+2)!\frac{t^{2k}}{(2k)!}, \dots, (2^{2k+1}(2k+2)!, 0, \dots, 0)). \quad (51)$$

It remains to evaluate the first row of the column C_{2k+2} .

In a shortened way, the first row can be written as

$$(C_{2k+2}^*)_1 = (-1)^{k+1} + (2k+2) \sum_{j=1}^k \frac{(-1)^j}{j} C_{2k+1-j}^{j-1} C_{2k+1-2j}^{k-j}.$$

By reduction, it can be written as

$$(C_{2k+2}^*)_1 = -C_{2k+1}^k = \prod_{j=0}^{k-1} \frac{(2k+1-l)}{(k-l)}. \quad (52)$$

The relations (51) and (52) gives exactly the relation (50) at the rank $k+1$.

Then we can write the determinant w defined in (43) as the following sum $w_1 + w_2$ where w_1 is defined by depending on m :

a. If $m = 2q$ we have for w_1

$$w_1 = \begin{vmatrix} t & 2 \cdot 2! \frac{t^2}{2!} & \dots & 2^{m-1} m! \frac{t^m}{m!} \\ 1 & 2 \cdot t & \dots & 2^{m-1} m! \frac{t^{m-1}}{(m-1)!} \\ 1 & 2 & \dots & 2^{m-1} m! \frac{t^{m-2}}{(m-2)!} \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \dots & 2^{m-1} m! t \end{vmatrix}.$$

w_2 is equal to

(53)

$$w_2 = \begin{vmatrix} 0 & -1 & 0 & \dots & 0 & -C_{2k-1}^k t^{2j-1} & 0 & \dots & -C_{m-1}^{m/2} \\ 1 & 2 \cdot t & 2^2(2)! \frac{t^2}{2!} & \dots & \dots & 2^{2j-1}(2j)! \frac{t^{2j-1}}{(2j-1)!} & \dots & \dots & 2^{m-1}(m)! \frac{t^{m-1}}{(m-1)!} \\ 0 & 2 & 2^2 t & \dots & \dots & 2^{2k+2}(i+2k+1)! \frac{t^{2k}}{(2k)!} & \dots & 2^{m-i-1}(m-2)! \frac{t^{m-i-3}}{(m-i-3)!} & (m-1)! \frac{t^{m-i-2}}{(m-i-2)!} \\ \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & 0 & \dots & 2^{m-2}(m-1)! & 2^{m-1}m!t \end{vmatrix} \quad (54)$$

Then we have

$$w_1 = \prod_{j=1}^m j! \frac{t^m}{m!} 2^{\frac{m(m-1)}{2}}. \quad (55)$$

For w_2 , we have

$$w_2 = \sum_{j=1}^{m/2} C_{2j-1}^{j-1} \delta_{2j},$$

where δ_{2j} is the determinant defined by

$$\delta_{2j} = \begin{vmatrix} 1 & 2t & 2^2 \cdot 3! \frac{t^2}{2!} & \dots & 2^{2j-2}(2j-1)! \frac{t^{2j-2}}{(2j-2)!} & 2^{2j}(2j+1)! \frac{t^{2j}}{(2j)!} & \dots & 2^{m-1}m! \frac{t^{m-1}}{(m-1)!} \\ 0 & 2 & 2^2 \cdot 3!t & \dots & 2^{2j-2}(2j-1)! \frac{t^{2j-3}}{(2j-3)!} & 2^{2j}(2j+1)! \frac{t^{2j-1}}{(2j-1)!} & \dots & 2^{m-1}m! \frac{t^{m-2}}{(m-2)!} \\ \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & \dots & 2^{m-2}(m-1)! & 2^{m-1}m!t \end{vmatrix}.$$

δ_{2j} is equal to

$$\delta_{2j} = 2^{\frac{m(m-1)}{2}-2j+1} \prod_{k=1, k \neq 2j}^m k! \frac{t^{m-2j}}{(m-2j)!}.$$

So w_2 can be written as

$$w_2 = \sum_{j=1}^{m/2} C_{2j-1}^{j-1} 2^{\frac{m(m-1)}{2}-2j+1} \prod_{k=1, k \neq 2j}^m k! \frac{t^{m-2j}}{(m-2j)!}. \quad (56)$$

Taking into account (55), (56) and $[m/2] = m/2$, with (42) we have :

$$\overline{W} = 2^{\frac{m(m-1)}{2}} \prod_{k=i}^m k! (-\alpha)^{\frac{m(m-1)}{2}} \sin^{\frac{m(m-1)}{2}} \alpha(x-x_0) \\ \cdot \left(\frac{\cos^m \alpha(x-x_0)}{m!} + \sum_{j=1}^{[m/2]} \frac{(2j-1)! 2^{-2j+1}}{(j-1)! j! (2j)!} \frac{\cos^{m-2j} \alpha(x-x_0)}{(m-2j)!} \right),$$

which can be rewritten as

$$\overline{W} = 2^{\frac{m(m-1)}{2}} \prod_{k=i}^m k! (-\alpha)^{\frac{m(m-1)}{2}} \sin^{\frac{m(m-1)}{2}} \alpha(x-x_0) \\ \cdot \left(\frac{\cos^m \alpha(x-x_0)}{m!} + \sum_{j=1}^{[m/2]} \frac{2^{-2j}}{j!^2} \frac{\cos^{m-2j} \alpha(x-x_0)}{(m-2j)!} \right)$$

This gives for \bar{W} exactly the formula (40).

b. We consider now the case $m = 2q + 1$.

The ideas are the same, but the structure of w is different (the last column is different).

We can write the determinant w defined in (43) as the following sum $w_1 + w_2$ where w_1 is defined by

$$w_1 = \begin{vmatrix} t & 2 \cdot 2! \frac{t^2}{2!} & \dots & 2^{m-1} m! \frac{t^m}{m!} \\ 1 & 2 \cdot t & \dots & 2^{m-1} m! \frac{t^{m-1}}{(m-1)!} \\ 1 & 2 & \dots & 2^{m-1} m! \frac{t^{m-2}}{(m-2)!} \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \dots & 2^{m-1} m! t \end{vmatrix}.$$

w_2 is equal to

$$w_2 = \begin{vmatrix} 0 & -1 & 0 & \dots & 0 & -C_{2k-1}^k & \dots & -C_{m-2}^{(m-1)/2} & 0 \\ 1 & 2 \cdot t & 2^2 (2!) \frac{t^2}{2!} & \dots & \dots & 2^{2j-1} (2j)! \frac{t^{2j-1}}{(2j-1)!} & \dots & \dots & 2^{m-1} (m)! \frac{t^{m-1}}{(m-1)!} \\ 0 & 2 & 2^2 t & \dots & \dots & 2^{2k+2} (i+2k+1)! \frac{t^{2k}}{(2k)!} & \dots & 2^{m-i-1} (m-2)! \frac{t^{m-i-3}}{(m-i-3)!} & (m-1)! \frac{t^{m-i-2}}{(m-i-2)!} \\ \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & 0 & \dots & 2^{m-2} (m-1)! & 2^{m-1} m! t \end{vmatrix} \quad (57)$$

(58)

Then we have

$$w_1 = \prod_{j=1}^m j! \frac{t^m}{m!} 2^{\frac{m(m-1)}{2}}. \quad (59)$$

For w_2 , we have

$$w_2 = \sum_{j=1}^{(m-1)/2} C_{2j-1}^{j-1} \delta_{2j},$$

where δ_{2j} is the determinant defined by

$$\delta_{2j} = \begin{vmatrix} 1 & 2t & 2^2 \cdot 3! \frac{t^2}{2!} & \dots & 2^{2j-2} (2j-1)! \frac{t^{2j-2}}{(2j-2)!} & 2^{2j} (2j+1)! \frac{t^{2j}}{(2j)!} & \dots & 2^{m-1} m! \frac{t^{m-1}}{(m-1)!} \\ 0 & 2 & 2^2 \cdot 3! t & \dots & 2^{2j-2} (2j-1)! \frac{t^{2j-3}}{(2j-3)!} & 2^{2j} (2j+1)! \frac{t^{2j-1}}{(2j-1)!} & \dots & 2^{m-1} m! \frac{t^{m-2}}{(m-2)!} \\ \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & \dots & 2^{m-2} (m-1)! & 2^{m-1} m! t \end{vmatrix}.$$

δ_{2j} is equal to

$$\delta_{2j} = 2^{\frac{m(m-1)}{2}-2j+1} \prod_{k=1, k \neq 2j}^m k! \frac{t^{m-2j}}{(m-2j)!}.$$

So w_2 can be written as

$$w_2 = \sum_{j=1}^{(m-1)/2} C_{2j-1}^{j-1} 2^{\frac{m(m-1)}{2}-2j+1} \prod_{k=1, k \neq 2j}^m k! \frac{t^{m-2j}}{(m-2j)!}. \quad (60)$$

Taking into account (59), (60), $[m/2] = (m-1)/2$, with (42) and some simplifications as these done in the part a., we have for \bar{W} exactly the formula (40).

So we get the result.

□

Remark 2.2 As in the previous section, the possible extension of the same calculations to the sequence of the hyperbolic functions can be easily made by the change $\alpha \rightarrow i\alpha$.

3 Hadamard's coefficients and solutions to the 2D-Schrödinger equation with Huygens' potentials

3.1 Explicit expressions of Hadamard's coefficients

We want to give a reduction of Hadamard's coefficients, given in terms of polar coordinates $x = (r \cos \varphi; r \sin \varphi)$, $\xi = (\rho \cos \phi; \rho \sin \phi)$ by (4)

$$U_\nu(x, \xi) = \frac{(-2)^\nu}{(r\rho)^\nu} \sum_{i=0}^m c_i \Psi_i(\varphi) \Psi_i(\phi) T_{k_i}^{(\nu)}(\cos(\varphi - \phi)), \text{ for } \nu \geq 0,$$

where

$$\Psi_i = \frac{W(F_0, \dots, F_{i-1}, F_{i+1}, \dots, F_m)}{W(F_0, \dots, F_m)},$$

$$c_i = \prod_{j=0, j \neq i}^m k_i^2 - k_j^2,$$

and $T_N(x) := \cos N(\arccos x)$.

Let F_i the functions defined by

$$\text{for } x = \varphi, \text{ or } x = \phi, \quad F_i(x) = \cos k_i(x), \text{ with } k_i = i, \text{ for } 0 \leq i \leq n \quad (61)$$

Then, using the same notations as before, we have the obvious relations, where the functions f_i are defined in (7) with $\alpha = 1$ and $x_0 = 0$.

$$\begin{aligned} W(F_0, F_1, \dots, F_{i-1}, F_{i+1}, \dots, F_m) &= (-1)^{m-1} \frac{m!}{i} W(f_1, \dots, f_{i-1}, f_{i+1}, \dots, f_m) \\ &= (-1)^{\frac{(m)(m-1)}{2}} \times 2^{\frac{m(m-1)}{2}-i+1} \prod_{k=1}^m k! \times \sin^{\frac{m(m-1)}{2}} x \times \left(\sum_{k=0}^{[(m-i)/2]} \frac{2^{-2k}}{(i+k)!k!} \frac{\cos^{m-i-2k} x}{(m-i-2k)!} \right). \end{aligned}$$

We have also,

$$\begin{aligned} W(F_1, \dots, F_m) &= W(g_1, \dots, g_m) = \overline{W} \\ &= 2^{\frac{m(m-1)}{2}} \prod_{k=1}^m k! (-1)^{\frac{m(m-1)}{2}} \sin^{\frac{m(m-1)}{2}} x \times \left(\sum_{k=0}^{[m/2]} \frac{2^{-2k}}{k!^2} \frac{\cos^{m-2k} x}{(m-2k)!} \right) \end{aligned}$$

Moreover,

$$W(F_0, \dots, F_m) = (-1)^m m! W(f_1, \dots, f_m) = (-1)^m m! W_m.$$

But, we have from [16],

$$W_m = W(f_1, \dots, f_m) = (-2)^{\frac{m(m-1)}{2}} \prod_{k=1}^{m-1} k! \sin^{\frac{m(m+1)}{2}} x.$$

This gives the equality

$$W(F_0, \dots, F_m) = (-1)^m (-2)^{\frac{m(m-1)}{2}} \prod_{k=1}^m k! \sin^{\frac{m(m+1)}{2}} x.$$

Then for $1 \leq i \leq m$, we have for Ψ_i :

$$\Psi_i = -\frac{W_{m-1}^i}{iWm} = (-1)^m (2)^{-i+1} \sin^{-m} x \cdot \left(\sum_{k=0}^{[(m-i)/2]} \frac{2^{-2k}}{(i+k)!k!} \frac{\cos^{m-i-2k} x}{(m-i-2k)!} \right) \quad (62)$$

For Ψ_0 we get :

$$\Psi_0 = -\frac{\overline{W}}{(-1)^m m! W m} = (-1)^m \sin^{-m} x \cdot \left(\sum_{k=0}^{[m/2]} \frac{2^{-2k}}{k!^2} \frac{\cos^{m-2k} x}{(m-2k)!} \right). \quad (63)$$

We use also the following relations:

$$c_i = \frac{(-1)^{m-i}(m-i)!(m+i)!}{2},$$

It is obvious that for $0 \leq \nu \leq i$

$$\begin{aligned} & T_{k_i}^{(\nu)}(\cos(\varphi - \phi)) = T_i^{(\nu)}(\cos(\varphi - \phi)) \\ &= \sum_{p=0}^{[i/2]} \sum_{j=0, i-2p+2j-\nu \geq 0}^p (-1)^{p+j} \binom{i}{2p} \binom{p}{j} \prod_{l=0}^{\nu-1} (i-2p+2j-l) \cos^{i-2p+2j-\nu}(\varphi - \phi). \\ & T_{k_i}^{(\nu)} = 0 \text{ for } \nu > i, \text{ and } T_{k_0}^{(\nu)}(\cos(\varphi - \phi)) = T_0^{(\nu)}(\cos(\varphi - \phi)) = 1. \end{aligned}$$

At this step, we can evaluate Hadamard's coefficients given in (4) :

Proposition 3.1 *With $x = (x_1, x_2) = (r \cos(\varphi), r \cos(\varphi))$, $\xi = (\xi_1, \xi_2) = (\rho \cos(\phi), \rho \cos(\phi))$, the Hadamard's coefficients can be written as*

$$\begin{aligned} U_0(x, \xi) &= \left\{ (-1)^m 2^{-1} (m!)^2 \sin^{-m} \varphi \sin^{-m} \phi \times \left(\sum_{k=0}^{[m/2]} \frac{2^{-2k}}{k!^2} \frac{\cos^{m-2k} \varphi}{(m-2k)!} \right) \right. \\ &\times \left(\sum_{k=0}^{[m/2]} \frac{2^{-2k}}{k!^2} \frac{\cos^{m-2k} \phi}{(m-2k)!} \right) + \sum_{i=1}^m (-1)^{m-i} 2^{-2i+1} (m-i)!(m+i)! \sin^{-m} \varphi \sin^{-m} \phi \\ &\times \sum_{p=0}^{[i/2]} \sum_{j=0, i-2p+2j-\nu \geq 0}^p (-1)^{p+j} \binom{i}{2p} \binom{p}{j} \cos^{i-2p+2j}(\varphi - \phi) \\ &\times \left. \left(\sum_{k=0}^{[(m-i)/2]} \frac{2^{-2k}}{(i+k)!k!} \frac{\cos^{m-i-2k} \varphi}{(m-i-2k)!} \right) \left(\sum_{k=0}^{[(m-i)/2]} \frac{2^{-2k}}{(i+k)!k!} \frac{\cos^{m-i-2k} \phi}{(m-i-2k)!} \right) \right\}. \end{aligned}$$

$$\begin{aligned}
U_\nu(x, \xi) = & \frac{(-2)^\nu}{(r\rho)^\nu} \left\{ \sum_{i=1}^m (-1)^{m-i} 2^{-2i+1} (m-i)! (m+i)! \sin^{-m} \varphi \sin^{-m} \phi \right. \\
& \times \sum_{p=0}^{[i/2]} \sum_{j=0, i-2p+2j-\nu \geq 0}^p (-1)^{p+j} \binom{i}{2p} \binom{p}{j} \prod_{l=0}^{\nu-1} (i-2p+2j-l) \cos^{i-2p+2j-\nu} (\varphi - \phi) \\
& \times \left. \left(\sum_{k=0}^{[(m-i)/2]} \frac{2^{-2k}}{(i+k)! k!} \frac{\cos^{m-i-2k} \varphi}{(m-i-2k)!} \right) \left(\sum_{k=0}^{[(m-i)/2]} \frac{2^{-2k}}{(i+k)! k!} \frac{\cos^{m-i-2k} \phi}{(m-i-2k)!} \right) \right\}, \text{ for } \nu \geq 1.
\end{aligned}$$

3.2 Expression of the eigenfunctions of the 2-D Schrödinger operator with Huygens' potentials.

Following the results of Berest [4], the expressions of the eigenfunctions of the two dimensional Schrödinger operator with $x = (x_1, x_2) = (r \cos(\varphi), r \cos(\varphi))$, $\xi = (\xi_1, \xi_2) = (\rho \cos(\phi), \rho \cos(\phi))$ are given by the following result

Proposition 3.2 *The eigenfunctions to the two dimensional Schrödinger operator*

$$-\Delta + V$$

with

$$V = -\frac{2}{r^2} \partial_x^2 \ln(W(\psi_0, \dots, \psi_m))$$

and ψ defined by (61) can be written as

$$\Psi(x, \xi) = \left(\sum_{\nu=0}^m \frac{1}{2^\nu} U_\nu(x, \xi) \right) e^{r\rho \cos \varphi \cos \phi + r\rho \sin \varphi \sin \phi}$$

$$\begin{aligned}
&= e^{r\rho \cos \varphi \cos \phi + r\rho \sin \varphi \sin \phi} \left\{ (-1)^m 2^{-1} (m!)^2 \sin^{-m} \varphi \sin^{-m} \phi \times \left(\sum_{k=0}^{[m/2]} \frac{2^{-2k}}{k!^2} \frac{\cos^{m-2k} \varphi}{(m-2k)!} \right) \right. \\
&\quad \times \left(\sum_{k=0}^{[m/2]} \frac{2^{-2k}}{k!^2} \frac{\cos^{m-2k} \phi}{(m-2k)!} \right) + \sum_{i=1}^m (-1)^{m-i} 2^{-2i+1} (m-i)! (m+i)! \sin^{-m} \varphi \sin^{-m} \phi \\
&\quad \times \sum_{p=0}^{[i/2]} \sum_{j=0, i-2p+2j-\nu \geq 0}^p (-1)^{p+j} \binom{i}{2p} \binom{p}{j} \cos^{i-2p+2j} (\varphi - \phi) \\
&\quad \times \left(\sum_{k=0}^{[(m-i)/2]} \frac{2^{-2k}}{(i+k)!k!} \frac{\cos^{m-i-2k} \varphi}{(m-i-2k)!} \right) \left(\sum_{k=0}^{[(m-i)/2]} \frac{2^{-2k}}{(i+k)!k!} \frac{\cos^{m-i-2k} \phi}{(m-i-2k)!} \right) \\
&\quad + \sum_{\nu=1}^m \frac{(-1)^\nu}{(r\rho)^\nu} \sum_{i=1}^m (-1)^{m-i} 2^{-2i+1} (m-i)! (m+i)! \sin^{-m} \varphi \sin^{-m} \phi \\
&\quad \times \sum_{p=0}^{[i/2]} \sum_{j=0, i-2p+2j-\nu \geq 0}^p (-1)^{p+j} \binom{i}{2p} \binom{p}{j} \prod_{l=0}^{\nu-1} (i-2p+2j-l) \cos^{i-2p+2j-\nu} (\varphi - \phi) \\
&\quad \times \left(\sum_{k=0}^{[(m-i)/2]} \frac{2^{-2k}}{(i+k)k!} \frac{\cos^{m-i-2k} \varphi}{(m-i-2k)!} \right) \left(\sum_{k=0}^{[(m-i)/2]} \frac{2^{-2k}}{(i+k)k!} \frac{\cos^{m-i-2k} \phi}{(m-i-2k)!} \right) \Big\}
\end{aligned}$$

4 Conclusion

We have constructed in this paper explicit expressions of Hadamard's coefficients by means of explicit expressions of some Wronskians.

Expressions of the Hadamard's coefficients for the Schrödinger operator with a matrix potential were given by algorithms in [1].

Other expressions of these coefficients were given in [70] in function of the tau functions of the KdV hierarchy. But these expressions are not explicit and practically unusable because they depend on a product of the sum of Schur polynomials by tau functions of the KdV equation which are more difficult to calculate.

As a byproduct, we give explicit expressions of the eigenfunctions of the two dimensional Schrödinger operator with Huygens' potentials.

References

- [1] I.G.Avramidi, R. Schiming, Heat kernel coefficients for the matrix Schrödinger operator, *J. Math. Phys.*, V. **36**, 5042, (1995)
- [2] Y.Berest, Solution of a restricted Hadamard problem on Minkowski space, *Comm. Pure Appl. Math.*, V. **50**, 1019-1052, (1997)
- [3] Y. Berest, I. Loutsenko, Huygens principle in Minkowski space and soliton solutions to the Korteweg de Vries equation, *CMP Phys.* V. **190**, 113-132, (1997)
- [4] Y. Berest, T. Cramer, F. Eshmatov Heat Kernel Coefficients for Two-Dimensional Schrödinger equation, *Commun. Math. Phys.*, V. **283**, 853-860 (2008).
- [5] G .Frobenius, L. Stickelberg, Über die addition und Multiplication der elliptischen function, *J. Reine und Angew. Math.* V. **88**, 146-184, (1880)
- [6] G. Frobenius, Über die Determinant mehrer Functionen einer Variablen, *ibid.*, V. **77**, 245-257, (1874)
- [7] P. Gaillard, V.B. Matveev, Wronskian addition formula and its applications, Max-Planck-Institut für Mathematik, MPI 02-31, V. **161**, 2002
- [8] P. Gaillard, A new family of deformations of Darboux-Pöschl-Teller potentials, *Lett. Math. Phys.*, V. **68**, 77-90, 2004
- [9] P. Gaillard, V.B. Matveev, New formulas for the eigenfunctions of the two-particle Calogero-Moser system, *Lett. Math. Phys.*, V. **89**, 1-12, 2009
- [10] P. Gaillard, V.B. Matveev, Wronskian and Casorai determinant representations for Darboux-Pöschl-Teller potentials and their difference extensions, *J. Phys A : Math. Theor.*, V. **42**, 404409-1-16, 2009
- [11] P. Dubard, P. Gaillard, C. Klein, V. B. Matveev, On multi-rogue wave solutions of the NLS equation and positon solutions of the KdV equation, *Eur. Phys. J. Spe. Top.*, V. **185**, 247-258, 2010
- [12] P. Gaillard, Families of quasi-rational solutions of the NLS equation and multi-rogue waves, *J. Phys. A : Meth. Theor.*, V. **44**, 435204-1-15, 2011

- [13] P. Gaillard, Wronskian representation of solutions of the NLS equation and higher Peregrine breathers, *J. Math. Sciences : Adv. Appl.*, V. **13**, N. 2, 71-153, 2012
- [14] P. Gaillard, Degenerate determinant representation of solution of the NLS equation, higher Peregrine breathers and multi-rogue waves, *J. Math. Phys.*, V. **54**, 013504-1-32, 2013
- [15] P. Gaillard, Wronskian representation of solutions of NLS equation and seventh order rogue waves, *J. Mod. Phys.*, V. **4**, N. 4, 246-266, 2013
- [16] P. Gaillard, V.B. Matveev, Wronskian addition formula and Darboux-Pöschl-Teller potentials, *J. Math.*, V. **2013**, ID 645752, 1-10, 2013
- [17] P. Gaillard, Two-parameters determinant representation of seventh order rogue waves solutions of the NLS equation, *J. Theor. Appl. Phys.*, V. **7**, N. **45**, 1-6, 2013
- [18] P. Gaillard, Deformations of third order Peregrine breather solutions of the NLS equation with four parameters, *Phys. Rev. E*, V. **88**, 042903-1-9, 2013
- [19] P. Gaillard, Two parameters deformations of ninth Peregrine breather solution of the NLS equation and multi rogue waves, *J. Math.*, V. **2013**, 1-111, 2013
- [20] P. Gaillard, Six-parameters deformations of fourth order Peregrine breather solutions of the NLS equation, *J. Math. Phys.*, V. **54**, 073519-1-22, 2013
- [21] P. Gaillard, The fifth order Peregrine breather and its eight-parameters deformations solutions of the NLS equation, *Commun. Theor. Phys.*, V. **61**, 365-369, 2014
- [22] P. Gaillard, Ten parameters deformations of the sixth order Peregrine breather solutions of the NLS equation, *Phys. Scripta*, V. **89**, 015004-1-7, 2014
- [23] P. Gaillard, Higher order Peregrine breathers, their deformations and multi-rogue waves, *J. Of Phys. : Conf. Ser.*, V. **482**, 012016-1-7, 2014

- [24] P. Gaillard, M. Gastineau, Eighteen parameter deformations of the Peregrine breather of order ten solutions of the NLS equation, *Int. J. Mod. Phys. C*, V. **26**, N. 2, 1550016-1-14, 2014
- [25] P. Gaillard, Two parameters wronskian representation of solutions of nonlinear Schrödinger equation, eight Peregrine breather and multi-rogue waves, *J. Math. Phys.*, V. **5**, 093506-1-12, 2014
- [26] P. Gaillard, Hierarchy of solutions to the NLS equation and multi-rogue waves, *J. Phys. : Conf. Ser.*, V. **574**, 012031-1-5, 2015
- [27] P. Gaillard, Tenth Peregrine breather solution of the NLS, *Ann. Phys.*, V. **355**, 293-298, 2015
- [28] P. Gaillard, M. Gastineau, The Peregrine breather of order nine and its deformations with sixteen parameters solutions of the NLS equation *Phys. Lett. A*, V. **379**, 1309-1313, 2015
- [29] P. Gaillard, Other 2N-2 parameters solutions to the NLS equation and 2N+1 highest amplitude of the modulus of the N-th order AP breather, *J. Phys. A: Math. Theor.*, V. **48**, 145203-1-23, 2015
- [30] P. Gaillard, Multi-parametric deformations of the Peregrine breather of order N solutions to the NLS equation and multi-rogue waves, *Adv. Res.*, V. **4**, N. **5**, 346-364, 2015
- [31] P. Gaillard, Higher order Peregrine breathers solutions to the NLS equation, *Jour. Phys. : Conf. Ser.*, V. **633**, 012106-1-6, 2016
- [32] P. Gaillard, M. Gastineau Patterns of deformations of Peregrine breather of order 3 and 4, solutions to the NLS equation with multi-parameters, *Journal of Theoretical and Applied Physics*, V. **10**, 1-7, 2016
- [33] P. Gaillard, M. Gastineau Twenty parameters families of solutions to the NLS equation and the eleventh Peregrine breather, *Commun. Theor. Phys.*, V. **65**, N. **2**, 136-144, 2016
- [34] P. Gaillard, Rational solutions to the KPI equation and multi rogue waves, *Annals Of Physics*, V. **367**, 1-5, 2016

- [35] P. Gaillard, M. Gastineau Twenty two parameters deformations of the twelfth Peregrine breather solutions to the NLS equation, Adv. Res., V. **10**, 83-89, 2016
- [36] P. Gaillard, Towards a classification of the quasi rational solutions to the NLS equation, Theor. And Math. Phys., V. **189**, N. **1**, 1440-1449, 2016
- [37] P. Gaillard, Fredholm and Wronskian representations of solutions to the KPI equation and multi-rogue waves, Jour. of Math. Phys., V. **57**, 063505-1-13, doi: 10.1063/1.4953383, 2016
- [38] P. Gaillard, M. Gastineau Families of deformations of the thirteenth Peregrine breather solutions to the NLS equation depending on twenty four parameters, Jour. Of Bas. And Appl. Res. Int., V. **21**, N. 3, 130-139, 2017
- [39] P. Gaillard, From Fredholm and Wronskian representations to rational solutions to the KPI equation depending on $2N2$ parameters, Int. Jour. of Appl. Sci. And Math., V. **4**, N. 3, 60-70, 2017
- [40] P. Gaillard, Families of rational solutions of order 5 to the KPI equation depending on 8 parameters, New Hor. in Math. Phys., V. 1, N. 1, 26-31, 2017
- [41] P. Gaillard, 6-th order rational solutions to the KPI Equation depending on 10 parameters, Jour. Of Bas. And Appl. Res. Int., V. **21**, N. 2, 92-98, 2017
- [42] P. Gaillard, N -Order rational solutions to the Johnson equation depending on $2N - 2$ parameters, Int. Jour. of Adv. Res. in Phys. Sci., V. **4**, N. 9, 19-37, 2017
- [43] P. Gaillard, Families of rational solutions to the KPI equation of order 7 depending on 12 parameters, Int. Jour. of Adv. Res. in Phys. Sci., V. **4**, N. 11, 24-30, 2017
- [44] P. Gaillard, Rational solutions to the Johnson equation and rogue waves, Int. Jour. of Inn. In Sci. and Math., V. **6**, N. 1, 14-19, 2018

- [45] P. Gaillard, Multiparametric families of solutions of the KPI equation, the structure of their rational representations and multi-rogue waves, *Theo. And Mat. Phys.*, V. **196**, N. 2, 1174-1199, 2018
- [46] P. Gaillard, The Johnson equation, Fredholm and wronskian representations of solutions and the case of order three, *Adv. In Math. Phys.*, V. **2018**, 1-18, 2018
- [47] P. Gaillard, Families of Solutions of Order 5 to the Johnson Equation Depending on 8 Parameters, *NHIMP*, V. **2**, N. 4, 53-61, 2018
- [48] P. Gaillard, Multiparametric families of solutions to the Johnson equation, *J. Phys. : Conf. Series*, V. **1141**, 012102-1-10, 2018
- [49] P. Gaillard, Rational solutions to the Boussinesq equation, *Fund. Jour. Of Math. And Appl.*, V. ., 109-112, 2019
- [50] P. Gaillard, Differential relations for the solutions to the NLS equation and their different representations, *Comm. In Adv. Math. Sci.*, V. **2**, N. 4, 1-4, 2019
- [51] P. Gaillard, Multi-parametric families of solutions of order N to the Boussinesq and KP equations and the degenerate rational case, *UJMA*, V. **3**, N. 2, 44-52, 2020
- [52] P. Gaillard, The mKdV equation and multi-parameters rational solutions, *Wave Motion*, V. **100**, 102667-1-9, 2021
- [53] P. Gaillard, Degenerate Riemann theta functions, Fredholm and wronskian representations of the solutions to the KdV equation and the degenerate rational case, *Jour. Of Geom. And Phys.*, V. **161**, 104059-1-12, 2021
- [54] P. Gaillard, 2N parameter solutions to the Burgers equation, *Jour. Of Appl. Nonlin. Dyn.*, V. **1**, N. **1**, 1-6, 2021
- [55] P. Gaillard, Multi-Parametric rational Solutions to the KdV equation, *Asi. Jour. Of Res. and Rev. In Phys.*, V. **3**, N. 3, 14-21, 2021
- [56] P. Gaillard, Multiparametric solutions to the Gardner equation and the degenerate rational case, *Jour. Of Appl. Ana. And Appl.*, V. **11**, N. 4, 2102-2113, 2021

- [57] P. Gaillard, Other families of rational solutions to the KPI equation, *Asi. Res. Jour Of Math.*, V. **17**, N. 6, 27-34, 2021
- [58] P. Gaillard, Rational solutions to the mKdV equation associated to particular polynomials, *Wave Motion*, V. **107**, 102824-1-11, 2021
- [59] P. Gaillard, Multiparametric Rational Solutions of Order N to the KPI Equation and the Explicit Case of Order 3, *Arch. Of cur. Res. Int.*, V. **21**, N . 6, 58-71, 2021
- [60] P. Gaillard, Rational solutions to the KPI equation from particular polynomials, *Wave Motion*, V. **108**, 102828-1-9, 2022
- [61] P. Gaillard, Rogue Waves of the Lakshmanan Porsezian Daniel Equation Depending on Multi-parameters, *As. Jour. Of Adv. Res. And Rep.*, V 16, N. 3, 32-40, 2022
- [62] P. Gaillard, Quasi-rational and rational solutions to the defocusing NLS equation, *Jour. Of Math. Sci. And Mod. Res. And Rep.*, V 5, N. 1, 24-34, 2022
- [63] P. Gaillard, Rational solutions to the KdV equation in terms of particular polynomials, *Jour. Of Nonlin. Mod. and Ana.*, V. 4, N. 4, 615-626, 2022
- [64] P. Gaillard, Rational solutions to the Painlev II equation from particular polynomials, *Int. J. Of Appl. Nonlin. Sci.*, V. 3, N. 3, 189-196, 2022
- [65] P. Gaillard, Multi-lump solutions to the KPI equation with a zero degree of derivation, *Jour. Of Theor. And Appl. Phys.*, V. 16, N. 4, 1-8, 2022
- [66] P. Gaillard, Families of solutions to the KPI equation given by an extended Darboux transformation, *Part. Diff. Equ. And Appl.*, V. 3, N. 80, 1-14, 2022
- [67] P. Gaillard, Rogue waves of the Hirota equation in terms of quasi-rational solutions depending on multi-parameters, *WSEAS Trans. On Math.*, DOI: 10.37394/23206.2023.22.24, V. 22, 190-203, 2023
- [68] P. Gaillard, Different representations of the solutions to the cylindrical nonlinear Schrödinger equation, *Dis. Nonlin. And Compl.*, V. 12, N. 3, 539-553, 2023

- [69] P. Gaillard, Rational solutions to the KdV equation from Riemann theta functions, Jour. Of Theor. And Appl. Phys., V. 17, N. 3, 1-5, 2023
- [70] P. Iliev, On the heat kernel and the Kortweg de Vries hierarchy, Ann. Inst. Fourier,, V. **55**, 2117-2127, 2005
- [71] S. Rosenberg The Laplacian on a riemannian manifold, Lon. Math. Soc., Stu. Texts, **31**, Camb. Uni. Press, Cambridge, (1987)