



HAL
open science

Hecke cycles on moduli of vector bundles and orbital degeneracy loci

Vladimiro Benedetti, Michele Bolognesi, Daniele Faenzi, Laurent Manivel

► **To cite this version:**

Vladimiro Benedetti, Michele Bolognesi, Daniele Faenzi, Laurent Manivel. Hecke cycles on moduli of vector bundles and orbital degeneracy loci. 2023. hal-04233167

HAL Id: hal-04233167

<https://hal.science/hal-04233167>

Preprint submitted on 9 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HECKE CYCLES ON MODULI OF VECTOR BUNDLES AND ORBITAL DEGENERACY LOCI

V. BENEDETTI, M. BOLOGNESI, D. FAENZI, L. MANIVEL

ABSTRACT. Given a smooth genus two curve C , the moduli space $SU_C(3)$ of rank three semi-stable vector bundles on C with trivial determinant is a double cover in \mathbb{P}^8 branched over a sextic hypersurface, whose projective dual is the famous Coble cubic, the unique cubic hypersurface that is singular along the Jacobian of C . In this paper we continue our exploration of the connections of such moduli spaces with the representation theory of GL_9 , initiated in [GSW13] and pursued in [GS15, RS18, RS19, BMT21]. Starting from a general trivector v in $\wedge^3 \mathbb{C}^9$, we construct a Fano manifold $D_{Z_{10}}(v)$ in $G(3, 9)$ as a so-called orbital degeneracy locus, and we prove that it defines a family of Hecke lines in $SU_C(3)$. We deduce that $D_{Z_{10}}(v)$ is isomorphic to the odd moduli space $SU_C(3, \mathcal{O}_C(c))$ of rank three stable vector bundles on C with fixed effective determinant of degree one. We deduce that the intersection of $D_{Z_{10}}(v)$ with a general translate of $G(3, 7)$ in $G(3, 9)$ is a K3 surface of genus 19.

1. INTRODUCTION

The Coble hypersurfaces are among the most fascinating objects in algebraic geometry, being connected with an unusual amount of different topics such as abelian varieties and their moduli spaces, moduli spaces of points in \mathbb{P}^2 , theta-groups and theta-representations à la Vinberg, or the arithmetic invariant theory of Bhargava-Gross. Long after their discovery in the first quarter of 20th century, they were realized to be directly connected to certain moduli spaces of vector bundles on curves. Indeed, a Coble quartic in \mathbb{P}^7 is isomorphic to the moduli space $SU_C(2)$ of rank two semistable bundles with trivial determinant on a genus three curve C . For a curve C of genus two, the moduli space $SU_C(3)$ of rank three semistable bundles with trivial determinant is a double cover of \mathbb{P}^8 , branched over a sextic hypersurface whose projective dual is a Coble cubic [Ort05, Ngu07, BB12]. The geometry of the Coble hypersurfaces reflects in many ways the geometry of these curves.

More generally, the moduli spaces $SU_C(r, L)$ of semi-stable vector bundles of rank r on C with fixed determinant L have been thoroughly investigated, from many different perspectives, including theta maps, conformal blocks and the Verlinde formula. Up to isomorphism, they do not depend on L but only on its degree d modulo r , and up to sign. Moreover they are smooth when d and r are coprime. So the moduli spaces that are connected to the Coble hypersurfaces have each one smooth companion, and one can wonder how these Fano manifolds fit into the picture.

For genus three, we provided an answer to this question in [BBFM23], where we proved that the moduli space $SU_C(2, L)$ of stable vector bundles with fixed determinant L of degree one, can be embedded in the Grassmannian $G(2, 8)$ as the singular locus of a special quadratic section of the Grassmannian. This quadratic section is uniquely determined by this property, exactly as the Coble quartic is uniquely determined by the fact that it is singular along the

2020 *Mathematics Subject Classification.* 14H60; 22E46.

Key words and phrases. Moduli spaces of stable bundles, Coble hypersurfaces, degeneracy loci, Hecke lines, self-dual hypersurfaces, subvarieties of Grassmannians.

Kummer threefold of C ; for this reason we coined it the *Coble quadric*. Our main purpose in this paper is to obtain a similar description of the odd moduli space in genus two.

As in [BBFM23], our study will be based on the beautiful relationships with certain theta-representations, in the sense of Vinberg. To be more specific, it was observed in [GSW13] that a Coble quartic can easily be constructed from a general element in $\wedge^4 V_8$, if we denote by V_8 a complex vector space of dimension eight. Similarly, a Coble cubic can be constructed from a general element in $\wedge^3 V_9$, if V_9 is now a complex vector space of dimension nine. In [BMT21] we used the approach of *orbital degeneracy loci*, whose general theory was developed in [BFMT20a, BFMT20b], to explore this connection further and construct the Coble sextic and its singular locus, and even the generalized Kummer fourfold associated to the Jacobian of the curve, which is a famous example of hyperKähler manifold. Here we further apply these ideas to linear spaces in $\mathrm{SU}_C(3)$.

Lines in the moduli spaces $\mathrm{SU}_C(r, L)$ have been investigated by many authors, where by lines we mean rational curves that have degree one with respect to the ample generator of the Picard group. In $\mathrm{SU}_C(3)$ there are three families of lines passing through the general point: one family of Hecke lines, and two families of lines contained in the \mathbb{P}^4 's of the two natural rulings of the moduli space. Starting from a general element v in $\wedge^3 V_9$, we will be able to describe these two rulings in terms of orbital degeneracy loci (Proposition 3.2). In order to describe the Hecke lines, we will first construct an orbital degeneracy locus in $G(3, V_9)$, that we denote by $D_{Z_{10}}(v)$, and we will prove that each point of $D_{Z_{10}}(v)$ defines two Hecke planes in $\mathrm{SU}_C(3)$ (where we define a Hecke plane as a plane whose lines are all Hecke lines). It is probably known to the experts that although the family of Hecke lines is irreducible, the family of Hecke planes has two different components, and this is one of the key properties that will allow us to prove our main result (Theorem 4.9):

Theorem. *For v general in $\wedge^3 V_9$, the Fano eightfold $D_{Z_{10}}(v) \subset G(3, V_9)$ is isomorphic to the odd moduli space $\mathrm{SU}_C(3, \mathcal{O}_C(c))$, $c \in C$.*

An important advantage of this approach by orbital degeneracy loci is that it provides lots of informations on the embedding of $\mathrm{SU}_C(3, \mathcal{O}_C(c))$ into the Grassmannian. For example we get a free resolution of its ideal sheaf, from which we can confirm an expectation recently expressed by Mukai and Kanemitsu [KM23], that the intersection of the moduli space with a general translate of $G(3, 7)$ in $G(3, 9)$ is a K3 surface of genus 19 (Proposition 4.14). In fact, there is a natural embedding of $\mathrm{SU}_C(3, \mathcal{O}_C(p))$ in $G(3, V_9)$ for any choice of a point p on C , and playing with these one expects to be able to describe a locally complete family of K3 surfaces of genus 19. Such locally complete families have been constructed so far for most values of the genus up to 20, but genus 19 would be new.

The plan of the paper is the following. In section 2, we recall some useful facts about the lines and planes inside the moduli space $\mathrm{SU}_C(3)$, for C a curve of genus 2. In section 3 we recall the connections with representation theory and use them to describe explicitly the rulings of $\mathrm{SU}_C(3)$ by \mathbb{P}^4 's, starting from a general element v of $\wedge^3 V_9$. In section 4 we apply the same approach to Hecke lines and planes. We define the orbital degeneracy locus $D_{Z_{10}}(v)$, show that it parametrizes pairs of Hecke planes and deduce our main Theorem. Finally we describe the minimal resolution of its ideal sheaf and the application to K3 surfaces of genus 19.

Acknowledgements. All authors partially supported by FanoHK ANR-20-CE40-0023. D.F. and V.B. partially supported by SupToPhAG/EIPHI ANR-17-EURE-0002, Région Bourgogne-Franche-Comté, Feder Bourgogne and Bridges ANR-21-CE40-0017.

We warmly thank Christian Pauly for useful discussions, and Shigeru Mukai and Akihiro Kanemitsu for sharing the results of [KM23].

2. MODULI SPACE OF RANK 3 VECTOR BUNDLES ON GENUS 2 CURVES

2.1. Moduli spaces. We will use the same notations as in [BBFM23]: we denote by $U_C(r, d)$ the moduli space of semi-stable vector bundles on a curve C of rank r and determinant of degree d . If L is a degree d line bundle on C , we will denote by $SU_C(r, L)$ the subvariety of $U_C(r, d)$ parametrizing vector bundles of fixed determinant L ; moreover $SU_C(r) := SU_C(r, \mathcal{O}_C)$. Since all the moduli spaces $SU(r, L)$ are (non canonically) isomorphic when the degree of L is fixed, we will also denote their isomorphism class by $SU_C(r, d)$; it does depend on d only modulo r .

2.2. Theta maps. In this paper we focus on the case where C has genus two and the rank $r = 3$. The moduli space $SU_C(3)$ has Picard rank one, and the positive generator of the Picard group defines the Theta map

$$\theta : SU_3(C) \longrightarrow |3\Theta| = \mathbb{P}(V_9)$$

to the linear system $|3\Theta|$ on C . This map is a double cover branched over a sextic hypersurface \mathcal{S} , whose projective dual is the Coble cubic, characterized as the unique cubic hypersurface in $\mathbb{P}(V_9^\vee)$ that is singular along the Jacobian of C [Ort05, Ngu07, Bea03].

The sextic \mathcal{S} is also singular, in fact its singular locus identifies with that of $SU_C(3)$, which has dimension five and can be described set theoretically as the locus of properly semi-stable vector bundles, equivalent to a direct sum $E = F \oplus \det(F)^\vee$ for some rank two vector bundle F of degree zero on C . As a deeper stratum, we get the fourfold \mathcal{K} parametrizing vector bundles $E = L_1 \oplus L_2 \oplus L_3$, for three degree zero line bundles L_1, L_2, L_3 with trivial tensor product. When two among the three bundles L_1, L_2, L_3 coincide, we get a surface that is a singular model of the Jacobian of C . Its singular locus, where the three bundles L_1, L_2, L_3 all coincide, is in bijection with the 81 three-torsion points of the Jacobian.

2.3. Lines in the moduli space. Rational curves in the moduli spaces $SU_C(r, d)$ were extensively studied, see e.g. [NR75, NR78, Sun05, MS09, Pal16, MTiB20]. Restricting to $g = 2$, $r = 3$ and $d = 0$, the results of [MTiB20] show that there exist three different families of covering lines, i.e. families of rational curves of degree one with respect to the Theta embedding and passing through a general point of the moduli space. We will denote these three families, as subvarieties of the Hilbert scheme, by $\mathcal{F}_H, \mathcal{F}_{R_1}, \mathcal{F}_{R_2}$, and we will denote by $\overline{\mathcal{F}}_H, \overline{\mathcal{F}}_{R_1}, \overline{\mathcal{F}}_{R_2}$ their bijective images inside the Grassmannian $G(2, V_9)$. They all have dimension eleven, which is the expected dimension.

2.3.1. Hecke lines. The family \mathcal{F}_H of Hecke lines is most naturally defined as parametrizing lines in planes that are constructed as follows. Start with a point $p \in C$ and a stable rank three vector bundle E on C with $\det(E) = \mathcal{O}_C(p)$. Any non zero linear form on the fiber E_p defines a rank three vector bundle F on C as the kernel of the composition $E \rightarrow E_p \rightarrow \mathcal{O}_p$. Since E is stable of degree one, by [NR78, Remark 5.2 (v)] it is $(0, 1)$ -semistable, and then by [NR78, Lemma 5.5] the vector bundle F is automatically semistable.

We thus get a linear embedding of $\mathbb{P}(E_p^\vee)$ inside $SU_C(3)$, whose image we will denote by $\Pi(E, p)$ (this is a *good Hecke cycle* in the terminology of [NR78]; the fact that the embedding is linear is e.g. in [MTiB20]). By the first paragraph of the proof of [NR78, Theorem 5.13], the resulting morphism

$$\phi_p : SU_C(3, \mathcal{O}_C(p)) \longrightarrow G(3, V_9)$$

is injective.

Note also that the point p can be uniquely recovered from any good Hecke plane. Indeed, the image of $\mathbb{P}(E_p^\vee)$ in $SU_C(3)$ is defined by the rank three bundle M over $C \times \mathbb{P}(E_p^\vee)$ fitting

into the exact sequence

$$0 \longrightarrow M \longrightarrow p_1^* E \longrightarrow p_1^* \mathcal{O}_p \otimes p_2^* \mathcal{O}_{\mathbb{P}(E_p^\vee)}(1) \longrightarrow 0$$

[NR78, page 362]. For a point $q \neq p$ of C , the restriction $M(q)$ of M to $\{q\} \times \mathbb{P}(E_p^\vee)$ is a trivial bundle on \mathbb{P}^2 , while $M(p)$ is the direct sum of the hyperplane bundle and the dual tautological bundle, in particular it is non trivial. This allows to distinguish p among the points of C .

The lines in the planes $\Pi(E, p)$ are the Hecke lines. They are specified by fixing a one-dimensional subspace L of E_p , and considering the planes $P \subset E_p$ that contain L ; we will denote the corresponding Hecke line by $\delta(E, p, L)$. In fact the previous argument works over such a line:

Lemma 2.1. *A Hecke line $\delta(E, p, L)$ uniquely determines the vector bundle E and also the point p in general.*

As a result, the projectivization $\hat{\mathcal{F}}_H$ of the dual universal bundle on what is denoted $U_C(3, C)$ in [NR78] (the moduli space of stable bundles with effective determinant of degree one) maps birationally to \mathcal{F}_H , and fibers over the curve C .

2.3.2. Lines in the rulings. Another way to construct lines in $SU_C(3)$ is to use extensions between bundles of smaller ranks. Consider stable vector bundles E_1, E_2 with $\text{rank}(E_1) = r_1$ and $\text{rank}(E_2) = r_2 = 3 - r_1$, r_1 being either 1 or 2. Suppose that E_1 has degree -1 and E_2 degree 1, with $\det(E_1) = \det(E_2)^{-1}$. So if $r_1 = 1$, the line bundle E_1 is determined by E_2 , and vice versa: if $r_1 = 2$, the line bundle E_2 is determined by E_1 . The space of extensions between E_1 and E_2 is five dimensional, and we get a linear embedding of $\mathbb{P}(\text{Ext}^1(E_2, E_1)) \simeq \mathbb{P}^4$ inside $SU_C(3)$. Hence we obtain two rulings R_1 and R_2 in four-dimensional projective spaces. The families \mathcal{F}_{R_1} and \mathcal{F}_{R_2} consist in the lines contained in these two rulings.

Lemma 2.2. *A line in \mathcal{F}_{R_1} (or \mathcal{F}_{R_2}) that is contained in the stable locus of the moduli space is contained in a unique linear space of the ruling.*

Proof. Consider a line δ in $SU_C(3)$ which is contained in two different members of the ruling \mathcal{R}_1 , corresponding to pairs of vector bundles (E_1, E_2) and (E'_1, E'_2) , and two planes $A \subset \text{Ext}^1(E_2, E_1)$, $A' \subset \text{Ext}^1(E'_2, E'_1)$. Each point of $\mathbb{P}(A)$ and $\mathbb{P}(A')$ can be identified to a point of $\delta = \mathbb{P}(B)$, and we can choose isomorphisms $A \simeq B \simeq A'$ compatible with these identifications. Hence an element e of $\text{Hom}(B, \text{Ext}^1(E_2, E_1)) \simeq \text{Ext}^1(p_1^* E_2, p_1^* E_1 \otimes p_2^* \mathcal{O}_{\mathbb{P}(B)}(1))$, where the latter Ext group is taken between vector bundles on $C \times \mathbb{P}(B) = C \times \delta$, with p_1 and p_2 the two projections. This defines on this surface an extension (see [MTiB20, Proposition 2.5])

$$0 \longrightarrow p_1^* E_1 \otimes p_2^* \mathcal{O}_{\mathbb{P}(B)}(1) \longrightarrow \mathcal{E} \longrightarrow p_1^* E_2 \longrightarrow 0.$$

Similarly, we get an element e' of $\text{Hom}(B, \text{Ext}^1(E'_2, E'_1)) \simeq \text{Ext}^1(p_1^* E'_2, p_1^* E'_1 \otimes p_2^* \mathcal{O}_{\mathbb{P}(B)}(1))$, and an extension

$$0 \longrightarrow p_1^* E'_1 \otimes p_2^* \mathcal{O}_{\mathbb{P}(B)}(1) \longrightarrow \mathcal{E}' \longrightarrow p_1^* E'_2 \longrightarrow 0.$$

We claim that $\mathcal{E} \simeq \mathcal{E}'$. Since δ is supposed not to meet the locus of properly semistable bundles (the singular locus of the moduli space), the isomorphism class of $\mathcal{E}_{C \times \{t\}}$ is uniquely determined at every point $t \in \delta$, and has to coincide with that of $\mathcal{E}'_{C \times \{t\}}$. Since this is true for any $t \in \delta$, we deduce from [Ram73, Lemma 2.5] that $\mathcal{E}' = \mathcal{E} \otimes p_2^* N$ for some line bundle N on δ ; computing determinants we see that N is in fact trivial.

But then the composition

$$p_1^* E_1 \otimes p_2^* \mathcal{O}_{\mathbb{P}(B)}(1) \longrightarrow \mathcal{E} \simeq \mathcal{E}' \longrightarrow p_1^* E'_2$$

clearly has to vanish, hence there is an induced morphism from $p_1^*E_1 \otimes p_2^*\mathcal{O}_{\mathbb{P}(B)}(1)$ to $p_1^*E'_1 \otimes p_2^*\mathcal{O}_{\mathbb{P}(B)}(1)$, and then from E_1 to E'_1 . Exchanging e and e' we deduce that in fact $E_1 \simeq E'_1$ and $E_2 \simeq E'_2$. \square

In the construction of the rulings, the rank two bundle (or its dual) is parametrized by $U_C(2, 1)$ and determines the rank one bundle. So the rulings are parametrized by $U_C(2, 1)$. By the previous Lemma, the corresponding families of lines \mathcal{F}_{R_1} and \mathcal{F}_{R_2} are the birational images of $\hat{\mathcal{F}}_{R_1}$ and $\hat{\mathcal{F}}_{R_2}$, which are $G(2, 5)$ -fibrations over $U_C(2, 1)$. Notice that since $U_C(2, 1)$ is a fiber bundle over the abelian surface $\text{Pic}^1(C)$, this is also the case for $\hat{\mathcal{F}}_{R_1}$ and $\hat{\mathcal{F}}_{R_2}$.

2.3.3. Action of the involution. According to [Ort05], the covering involution τ of the double cover θ of $\text{SU}_C(3)$ is given by $E \mapsto \tau(E) := \iota^*E^\vee$, where ι is the hyperelliptic involution of C .

Lemma 2.3. *The two rulings R_1 and R_2 are exchanged by τ .*

Proof. If the bundle E fits into an exact sequence

$$0 \rightarrow E_1 \rightarrow E \rightarrow E_2 \rightarrow 0,$$

with $\text{rank}(E_1) = 2$, $\text{deg}(E_1) = -1$, $\text{rank}(E_2) = 1$, $\text{deg}(E_2) = 1$, then ι^*E^\vee fits into the exact sequence

$$0 \rightarrow \iota^*E_2^\vee \rightarrow \iota^*E^\vee \rightarrow \iota^*E_1^\vee \rightarrow 0,$$

with $\text{rank}(\iota^*E_2^\vee) = 1$, $\text{deg}(\iota^*E_2^\vee) = -1$, $\text{rank}(\iota^*E_1^\vee) = 2$, $\text{deg}(\iota^*E_1^\vee) = 1$. This implies the claim. \square

The upshot is that the images of \mathcal{F}_{R_1} and \mathcal{F}_{R_2} inside $G(2, V_9)$ form a single family $\overline{\mathcal{F}}_R := \overline{\mathcal{F}}_{R_1} = \overline{\mathcal{F}}_{R_2}$, and their fibrations over $\text{Pic}^1(C)$ descend to $\overline{\mathcal{F}}_R$.

Lemma 2.4. *The family \mathcal{F}_H of Hecke lines is preserved by τ .*

Proof. This is clear from the previous Lemma since the lines are preserved by τ . But let us be more precise. Let E be a stable rank three vector bundle such that $\det(E) = \mathcal{O}_C(p)$ for a point $p \in C$. Consider a complete flag $L \subset P \subset E_p$. There is a commutative diagram

$$\begin{array}{ccccccc}
 & & & 0 & & & \\
 & & & \uparrow & & & \\
 & & & \mathcal{O}_p \otimes P/L & & & 0 \\
 & & & \uparrow & & & \uparrow \\
 0 & \longrightarrow & F_P & \longrightarrow & E & \longrightarrow & \mathcal{O}_p \otimes E_p/P \longrightarrow 0 \\
 & & \uparrow & & \parallel & & \uparrow \\
 0 & \longrightarrow & G_L^\vee & \xrightarrow{\gamma_L} & E & \longrightarrow & \mathcal{O}_p \otimes E_p/L \longrightarrow 0 \\
 & & \uparrow & & & & \uparrow \\
 & & 0 & & & & \mathcal{O}_p \otimes P/L \\
 & & & & & & \uparrow \\
 & & & & & & 0
 \end{array}$$

The bottom horizontal exact sequence defines a vector bundle G_L^\vee , whose dual G_L has again rank three and determinant $\mathcal{O}_C(p)$. Moreover G_L is also stable. Indeed, F_P is (semi)stable of degree 0, hence $(0, 1)$ -(semi)stable. By [NR78, Lemma 5.5] G_L^\vee is then $(0, 0)$ -(semi)stable, hence (semi)stable as well as its dual.

The plane $\Pi(E, p)$ in $\text{SU}_C(3)$ parametrized by $\mathbb{P}(E_p^\vee)$ consists in the vector bundles F_P , for $P \subset E_p$. We get a line in this plane, that is, a Hecke line, by imposing to the planes P to

contain a common line L . Then dualizing the first vertical exact sequence in the previous diagram, and pulling-back by the hyperelliptic involution, we get

$$0 \longrightarrow \iota^* F_P^\vee \longrightarrow \iota^* G_L \longrightarrow \mathcal{O}_{\iota(p)} \longrightarrow 0,$$

confirming that the image of our Hecke line is again a Hecke line. \square

Nevertheless, observe that the bundle $\iota^* G_L$ that defines this Hecke line does really depend on L , not only on E . This will be a direct consequence of the next

Lemma 2.5. *The map from $\mathbb{P}(E_p)$ to $\mathrm{SU}_C(3, \mathcal{O}_C(p))$, sending L to G_L , is non constant.*

Proof. The proof mimicks the proof of [NR78, Lemma 5.9], where it is proved that the differential of the morphism $L \mapsto G_L$ is injective in a context that needs a little adaptation, as follows. The differential is computed in [NR78, Lemma 5.10] in a rather general setting.

To deduce it is injective in our situation as well, let us consider the exact sequence

$$0 \rightarrow G_L^\vee \rightarrow E \rightarrow \mathcal{O}_p \otimes E_p/L \rightarrow 0.$$

By letting L vary in $\mathbb{P}(E_p)$, we get a family of extensions

$$0 \rightarrow \mathcal{G}^\vee \rightarrow p_1^* \mathcal{E} \rightarrow p_2^* \mathcal{Q} \otimes \mathcal{O}_{\{p\} \times \mathbb{P}(E_p)} \rightarrow 0$$

on $C \times \mathbb{P}(E_p)$, where \mathcal{Q} is the rank two tautological bundle on $\mathbb{P}(E_p)$. By [NR78, Lemma 5.10], the differential of $L \mapsto G_L$ is given by the connecting homomorphism

$$\mathcal{H}om((\mathcal{G}^\vee|_{\{p\} \times \mathbb{P}(E_p)}), \mathcal{Q}) \rightarrow R^1 p_{2*} \mathcal{E}nd(\mathcal{G}).$$

But this fits in the exact sequence

$$0 \rightarrow p_{2*} \mathcal{E}nd(\mathcal{G}) \rightarrow p_{2*} \mathcal{H}om(\mathcal{G}^\vee, p_1^* \mathcal{E}) \rightarrow \mathcal{H}om((\mathcal{G}^\vee|_{\{p\} \times \mathbb{P}(E_p)}), \mathcal{Q}) \rightarrow R^1 p_{2*} \mathcal{E}nd(\mathcal{G}).$$

Since \mathcal{G}^\vee is a family of stable bundles, $p_{2*} \mathcal{E}nd(\mathcal{G}) = \mathcal{O}_{\mathbb{P}(E_p)}$. Similarly to what happens in the proof of [NR78, Lemma 5.9], the result will follow if we can show that $\mathrm{Hom}(G_L^\vee, E)$ is one dimensional, which is essentially a slight modification of [NR78, Lemma 5.6].

In order to check this, the main observation is the following. The morphism $\gamma_L : G_L^\vee \rightarrow E$ in the diagram above has full rank outside p . Suppose $\gamma : G_L^\vee \rightarrow E$ is another a morphism, generically of full rank. Then $\det(\gamma)$ defines a section of $\det(G_L) \otimes \det(E) = \mathcal{O}_C(2p)$. But this line bundle has a unique global section, vanishing only (with multiplicity two) at p . So γ also has to be an isomorphism outside p . But if the dimension of $\mathrm{Hom}(G_L^\vee, E)$ was bigger than one, we could construct morphisms $s\gamma_L + t\gamma$ whose rank would drop at any chosen point of C , and this would yield a contradiction. \square

Proposition 2.6. *The image of the plane $\Pi(E, p)$ by the involution τ is*

$$\tau(\Pi(E, p)) = \Pi^*(\iota^* E, \iota(p)),$$

where $\Pi^*(F, q)$ is the plane that parametrizes the duals of the bundles parametrized by $\Pi(F, q)$. In particular, it is not a plane of the same type.

Proof. Indeed, if we had $\tau(\Pi(E, p)) = \Pi(E', p')$ for some bundle E' and some point p' , then by the proof of Lemma 2.4 we would deduce that $G_L = \iota^* E'$ does not depend on L . \square

Let us define a *Hecke plane* in $\mathrm{SU}_C(3)$ as a plane whose lines are all Hecke lines. We have defined two distinct families of Hecke planes parametrized by $U_C(3, C)$ (which is nine dimensional), and we want prove that there is no other Hecke plane through the general point of $\mathrm{SU}_C(3)$. The key point to prove this claim is the following result [Hwa02, Theorem 3]:

Proposition 2.7. *The VMRT at a general point $[F]$ of the family of Hecke lines is the image of the relative flag manifold $Fl(F) \xrightarrow{\psi} C$ by the linear system $|\mathcal{O}_{Fl(F)}(1, 1) \otimes \psi^* K_C|$, into the projectivized tangent space of $SU_C(3)$ at F , that is, $\mathbb{P}(H^1(C, \mathcal{E}nd_0(F))) \simeq \mathbb{P}(H^0(C, S_{2,1}(F) \otimes K_C)^\vee)$.*

Corollary 2.8. *A general Hecke line is contained in exactly two Hecke planes.*

Proof. Consider a general Hecke line $\delta(E, p, L)$ in $SU_C(3)$, and let Π be a Hecke plane containing this line. Let $[F]$ be a general point on $\delta(E, p, L)$, and let us apply the previous Proposition. In the VMRT at $[F]$, $\delta(E, p, L)$ defines a point δ , and Π defines a projective line π through δ . Observe that any line in Π is a Hecke line, hence of the form $\delta(E', p', L')$; since the point p' is uniquely determined in general, we get a rational map from the dual plane of lines in Π , to C , and such a map must be constant, equal to p . This means that π is contained in the fiber over p of the relative flag bundle $Fl(F)$. So we just need to consider the ordinary three-dimensional flag manifold $Fl(1, 2, \mathbb{C}^3)$. And then it is clear that any point $(L_0 \subset P_0)$ is contained in exactly two lines, parametrizing flags of the form $(L_0 \subset P)$ or $(L \subset P_0)$. \square

It would be interesting to describe the VMRT at a general point of the families \mathcal{F}_{R_1} and \mathcal{F}_{R_2} . They must be \mathbb{P}^3 -bundles over the families of \mathbb{P}^4 's in the rulings passing through the general point $[E] \in SU_C(3)$, which are parametrized by curves C_1 and C_2 . Note that like the rulings R_1 and R_2 , the curves C_1 and C_2 map to $\text{Pic}^1(C)$, and in fact these curves have a simple description in terms of theta divisors. Each of them parametrizes line bundles L of degree -1 such that $h^0(L^\vee \otimes E) \neq 0$. The locus of line bundles $L^\vee \in \text{Pic}^1(C)$ that verify this condition is just the *theta divisor* associated with E , which in this case is a divisor in $\text{Pic}^1(C)$ belonging to the linear system $|3\Theta|$. For general E , the associated theta divisor is smooth and irreducible. An easy computation shows that the generic element of $|3\Theta|$ is a curve of genus 10.

Let us summarize the results we will use in the sequel.

Proposition 2.9. *The moduli space $SU_C(3)$ admits three covering families of lines:*

- *The family \mathcal{F}_H of Hecke lines, which is birationally fibered over C .*
- *Two families of lines \mathcal{F}_{R_1} and \mathcal{F}_{R_2} contained in the rulings, which are birationally fibered over $\text{Pic}^1(C)$ and exchanged by the involution τ ; they both descend to the same family $\overline{\mathcal{F}}_R$ inside $G(2, V_9)$, which is still birationally fibered over the abelian surface $\text{Pic}^1(C)$.*

The moduli space $SU_C(3)$ also admits four covering families of planes:

- *Two families $\mathcal{P}_H, \mathcal{P}_{H^*}$ of Hecke planes, both birationally fibered over C .*
- *Two families \mathcal{P}_{R_1} and \mathcal{P}_{R_2} of planes contained in the rulings, both birationally fibered over $\text{Pic}^1(C)$.*

Proof. The only point that does not follow from the previous discussion is that there is not other plane Π through the general point of $SU_C(3)$, which is made of lines contained in one of the rulings. What we have to exclude is that the lines in Π belong to \mathbb{P}^4 's, say from R_1 , that move with the line. In this situation, we would get a map from the dual plane Π^\vee to R_1 , that we could compose with the projection to $\text{Pic}^1(C)$. The composition would need to be constant, which means that the line bundle we denoted E_1 must be constant.

But then, consider the pencil of lines in Π passing through a fixed general point $[E]$ of $SU_C(3)$. For each of these lines we get a morphism $E_1 \rightarrow E$, and the quotient bundle has to change with the line. So we deduce that $F = \text{Hom}(E_1, E)$ has at least two global sections. This means that F belongs to the Brill-Noether stratum $\mathcal{W}_{3,3}^1$, which according to [BPGN97] has the expected dimension given by the Brill-Noether number $\rho_{3,3}^1 = 6$. Since F determines E_1 up to 3-torsion, this leaves only six parameters also for E , which cannot be general. \square

3. ODL INTERPRETATIONS

In this section we summarize the results of [GSW13] and [BMT21] about the relationships between the moduli space $SU_C(3)$, the Coble cubic and a skew-symmetric cubic tensor in nine variables.

3.1. On the Coble side. The starting point is a general $v \in \wedge^3 V_9$. By the Borel-Weil theorem, there is a natural identification

$$\wedge^3 V_9 \simeq H^0(\mathbb{P}(V_9^\vee), \wedge^2 \mathcal{U}(1)),$$

where \mathcal{U} denotes the rank eight tautological bundle. This allows to consider the skew-symmetric cubic tensor v as a family of skew-symmetric bilinear forms, and then the loci where these forms drop rank: first a Pfaffian cubic hypersurface, and then its singular locus where the rank drops to six. This turns out to be an abelian surface, and the Pfaffian hypersurface is the corresponding Coble cubic, being singular exactly on the abelian surface.

3.2. On the dual side. The Borel-Weil theorem gives another identification

$$\wedge^3 V_9 \simeq H^0(\mathbb{P}(V_9), \wedge^3 \mathcal{Q}),$$

where \mathcal{Q} denotes the rank eight tautological quotient bundle. This allows to consider the skew-symmetric cubic tensor v in nine variables as a family of cubic tensors in eight variables, which are considerably simpler. In particular, $\wedge^3 V_8$ admits finitely many $GL(V_8)$ -orbits, and for each orbit closure $Y \subset \wedge^3 V_8$ there is an orbital degeneracy locus (ODL) $D_Y(v) \subset \mathbb{P}(V_9)$. Set theoretically, this is simply the set of points x for which v defines an element of $\wedge^3 \mathcal{Q}_x \simeq \wedge^3 V_8$ that falls into Y . For v generic, we need only consider the orbits Y_i of codimension $i \leq 8$, since this will also be the codimension of $D_{Y_i}(v) \subset \mathbb{P}(V_9)$. Moreover, this locus will be almost always singular, with singular locus $D_{\text{Sing}(Y_i)}(v)$, and it will come with some natural desingularisations, induced by desingularisations of Y of a special type, called Kempf collapsings. Concretely, in this case we get the following loci:

- The Coble sextic

$$\mathcal{S} := \{[U_1] \in \mathbb{P}(V_9) \mid \exists U_6 \supset U_1, v \in U_1 \wedge (\wedge^2 V_9) + \wedge^2 U_6 \wedge V_9\} \subset \mathbb{P}(V_9).$$

- Its singular locus (five dimensional, corresponding to strictly semistable bundles)

$$\text{Sing}(\mathcal{S}) := \{[U_1] \in \mathbb{P}(V_9) \mid \exists U_5 \supset U_2 \supset U_1, v \in U_2 \wedge (\wedge^2 V_9) + \wedge^2 U_5 \wedge V_9\}.$$

- The fourfold \mathcal{K} parametrizing rank three vector bundles with trivial determinant, splitting as a sum of three line bundles:

$$\mathcal{K} := \{[U_1] \in \mathbb{P}(V_9) \mid \exists U_6 \supset U_3 \supset U_1, v \in U_1 \wedge (\wedge^2 V_9) + U_3 \wedge U_6 \wedge V_9 + \wedge^3 U_6\}.$$

As proved in [BMT21], \mathcal{K} is a singular model of the generalized Kummer fourfold associated to the Jacobian of C . Moreover, its singular locus (when two of the three line bundles are equal) is a surface which is birational to the Jacobian itself, but with 81 singular points in bijection with the 3-torsion points (when the three line bundles are equal).

3.3. The rulings. We have seen that the two rulings of $SU_C(3)$ by \mathbb{P}^4 's are parametrized by pairs of bundles (E_1, E_2) , of ranks $(1, 2)$ or $(2, 1)$, such that $\det(E_1) \otimes \det(E_2) = \mathcal{O}_C$. Up to a shift in the degrees, this is very similar to the data that defines a strictly semistable rank three vector bundle on C , hence a point in $\text{Sing}(\mathcal{S})$. So we expect the latter to be the parameter space for some natural families of five-dimensional subspaces of V_9 . This is exactly the content of the next two statements.

Since $SU_C(3)$ is a double cover of $\mathbb{P}(V_9)$ branched over the sextic \mathcal{S} , a $\mathbb{P}^4 = \mathbb{P}(U_5)$ in $\mathbb{P}(V_9)$ will lift to a pair of \mathbb{P}^4 's in $SU_C(3)$, meeting in codimension one, exactly when the sextic hypersurface restricts on $\mathbb{P}(U_5)$ to a double cubic. We will check this phenomenon does happen by rather indirect arguments. Define

$$\overline{\text{Sing}}(\mathcal{S}) := \{(U_1 \subset U_2 \subset U_5) \in Fl(1, 2, 5, V_9), v \in U_2 \wedge (\wedge^2 V_9) + \wedge^2 U_5 \wedge V_9\}.$$

Lemma 3.1. *The projection $\overline{\text{Sing}}(\mathcal{S}) \rightarrow \text{Sing}(\mathcal{S})$ is generically finite of degree two.*

Proof. This is contained in [BMT21, Remark 3.1]. □

Proposition 3.2. *Let $(U_1 \subset U_2 \subset U_5)$ be general in $\overline{\text{Sing}}(\mathcal{S})$. Then $\mathbb{P}(U_5) \cap \mathcal{S}$ is a double cubic in $\mathbb{P}(U_5) \simeq \mathbb{P}^4$, and $\theta^{-1}(\mathbb{P}(U_5))$ is the union of two copies of \mathbb{P}^4 in $SU_C(3)$.*

Proof. Let us fix a general $(U_1 \subset U_2 \subset U_5)$ in $\overline{\text{Sing}}(\mathcal{S})$. This means that

$$v \in U_2 \wedge (\wedge^2 V_9) + \wedge^2 U_5 \wedge V_9.$$

The intersection $\mathbb{P}(U_5) \cap \mathcal{S}$ parametrizes the flags $(W_1 \subset W_6)$, with $W_1 \subset U_5$, such that

$$v \in W_1 \wedge (\wedge^2 V_9) + \wedge^2 W_6 \wedge V_9.$$

We need to understand the relative position of the flags $(U_1 \subset U_2 \subset U_5)$ and $(W_1 \subset W_6)$.

Lemma 3.3. *$\dim(W_6 \cap U_2) = 2$ and $\dim(W_6 \cap U_5) = 4$.*

Proof of the Lemma. We will argue as follows. Let

$$\mathcal{A} = U_2 \wedge (\wedge^2 V_9) + \wedge^2 U_5 \wedge V_9, \quad \mathcal{B} = W_1 \wedge \wedge^2 V_9 + \wedge^2 W_6 \wedge V_9.$$

The dimension of \mathcal{A} is 62. We can stratify the space of flags $(W_1 \subset W_6)$, with $W_1 \subset U_5$, by the possible values of $\dim(W_1 \cap U_2)$, $\dim(W_6 \cap U_2)$ and $\dim(W_6 \cap U_5)$. On each stratum S , the quotient $\mathcal{P} := \mathcal{A}/\mathcal{A} \cap \mathcal{B}$ has a fixed dimension, hence defines a vector bundle of a certain rank. Moreover this bundle is generated by global sections and v defines such a global section. Although S is in general not compact, we can apply the general Bertini type arguments to the space $\Gamma = U_2 \wedge (\wedge^2 V_9) + \wedge^2 U_5 \wedge V_9$ of global sections, which is finite dimensional and generates \mathcal{P} everywhere on S .

Remark that we consider the same space $U_2 \wedge (\wedge^2 V_9) + \wedge^2 U_5 \wedge V_9$ in two different ways: first as a trivial bundle \mathcal{A} on S , then as a space of global sections of the quotient bundle \mathcal{P} . We will be able to conclude that when $\text{rank}_S(\mathcal{P}) > \dim(S)$, the general $v \in U_2 \wedge (\wedge^2 V_9) + \wedge^2 U_5 \wedge V_9$ defines a section that vanishes nowhere, which means that there exists no flag $(W_1 \subset W_6)$ satisfying the corresponding conditions. If $\text{rank}_S(\mathcal{P}) \leq \dim(S)$, the general section either vanishes nowhere, or on a locus of dimension $\dim(S) - \text{rank}_S(\mathcal{P})$, and we need this difference to be equal to three (at least) since we are looking for a hypersurface in \mathbb{P}^4 . Note in particular that we can, and will always suppose in the sequel that $\dim(W_1 \cap U_2) = 0$, since $\mathbb{P}(U_2)$ is only one dimensional.

When convenient, we will occasionally replace \mathcal{P} by some quotient $\overline{\mathcal{P}}$, to which the same arguments will a fortiori apply. In some cases we will even need that this quotient is in fact obtained by pull-back from another parameter space $\overline{\mathcal{S}}$; under the condition that $\text{rank}(\overline{\mathcal{P}}) > \dim \overline{\mathcal{S}} - 3$, this will allow as well to eliminate the stratum S . In the end, a unique stratum will survive and we will have proved the Lemma. More precisely, we will prove that for all but one couple of values $(\dim(W_6 \cap U_2), \dim(W_6 \cap U_5))$ the section defined by v vanishes on a locus of dimension at most two. So there will be a unique couple of values $(\dim(W_6 \cap U_2), \dim(W_6 \cap U_5))$ describing the intersection dimensions of a general flag $W_1 \subset W_6$ inside $\mathbb{P}(U_5) \cap \mathcal{S}$. The conclusion will hold for any flag $(U_2 \subset U_5)$ and for the general $v \in U_2 \wedge (\wedge^2 V_9) + \wedge^2 U_5 \wedge V_9$, hence conversely for the general v and the general flag $(U_2 \subset U_5)$ such that $v \in U_2 \wedge (\wedge^2 V_9) + \wedge^2 U_5 \wedge V_9$.

Let us discuss the different possibilities for $(\dim(W_6 \cap U_2), \dim(W_6 \cap U_5))$, which can a priori be $(0, 2), (1, 2), (2, 2), (0, 3), (1, 3), (2, 3), (1, 4), (2, 4), (2, 5)$.

(0,2) This is the open stratum. We compute that

$$\mathcal{A} \cap \mathcal{B} = W_1 \wedge U_5 \wedge V_9 + U_2 \wedge (\wedge^2 W_6) + (W_6 \cap U_5) \wedge W_6 \wedge U_5$$

has dimension 46. Since this is the first time we make such a computation, let us explain once and for all how we proceed; the bravest readers should be able to check the next computations by themselves. Let v_1, \dots, v_9 be a basis of V_9 such that $U_2 = \langle v_1, v_2 \rangle$ and $U_5 = \langle v_1, \dots, v_5 \rangle$. Since we are on the stratum indexed by the intersection dimensions $(0, 2)$, we can suppose that $W_1 = \langle v_3 \rangle$ and $W_6 = \langle v_3, v_4, v_6, v_7, v_8, v_9 \rangle$. Then a brute force computation shows that a basis of $\mathcal{P} := \mathcal{A}/\mathcal{B}$ is given by the classes, if we denote $v_{ijk} := v_i \wedge v_j \wedge v_k$, of the following 16 trivectors: $v_{145}, v_{156}, v_{157}, v_{158}, v_{159}, v_{245}, v_{256}, v_{257}, v_{258}, v_{259}, v_{124}, v_{125}, v_{126}, v_{127}, v_{128}, v_{129}$. So \mathcal{P} has rank 16 and $\mathcal{A} \cap \mathcal{B}$ has dimension $62 - 16 = 46$, as claimed.

Since the open stratum has dimension $19 = 16 + 3$, at first sight there is no obstruction. But let $W_2 := W_6 \cap U_5$, $W_4 := W_2 + U_2$ and $W_8 := W_6 + U_2$. The parameter space for such flags $(W_1 \subset W_2 \subset W_4 \subset W_8)$, with $U_2 \subset W_4 \subset U_5$ has dimension 11 (six dimensions for the choice of W_2 in U_5 , transverse to U_2 ; then one extra dimension for the choice of W_1 ; W_4 is uniquely determined; finally four extra dimensions for the choice of W_8 containing W_4). Let

$$\mathcal{C} := W_2 \wedge U_5 \wedge U_9 + U_2 \wedge (\wedge^2 W_8).$$

We check that $\mathcal{A} \cap \mathcal{B} \subset \mathcal{C} \subset \mathcal{A}$ and that \mathcal{C} has dimension 53, hence codimension 9 in \mathcal{A} . Therefore the zero locus of the section of the bundle \mathcal{A}/\mathcal{C} induced by our general v must have dimension at most $11 - 9 = 2 < 3$. This means that a general point in the intersection $\mathbb{P}(U_5) \cap \mathcal{S}$ does not belong to this stratum.

(1,2) Let $T_1 := W_6 \cap U_2$ and $W_2 := W_6 \cap U_5$. The parameter space for such flags has dimension 17: one dimension for the choice of T_1 in U_2 , four dimensions for the choice of W_1 in U_5 , then twelve dimensions for the choice of W_6 containing $W_2 = T_1 + W_1$. We compute that

$$\mathcal{A} \cap \mathcal{B} = W_1 \wedge U_5 \wedge V_9 + U_2 \wedge (\wedge^2 W_6) + T_1 \wedge W_6 \wedge V_9$$

has dimension 46, so the rank of \mathcal{P} is $16 = \dim(\mathcal{S}) - 1$, and we are safe.

(2,2) In this case $W_6 \cap U_5 = W_6 \cap U_2 = U_2$. But this implies that $W_1 \subset U_2$, which we excluded right from the beginning.

(0,3) Let $W_3 := W_6 \cap U_5$. The parameter space for such flags $(W_1 \subset W_3 \subset W_6)$ has dimension 17 (6 dimensions for the choice of W_3 in U_5 , transverse to U_2 ; then two extra dimensions for the choice of W_1 in W_3 ; finally nine extra dimensions for the choice of W_6 containing W_3). In this case $\mathcal{A} \cap \mathcal{B}$ has dimension 45, so the rank of \mathcal{P} is 17 and we are safe.

(1,3) Let $T_1 := W_6 \cap U_2$ and $W_3 := W_6 \cap U_5$. Here the parameter space has dimension 16 and the dimension of $\mathcal{A} \cap \mathcal{B}$ jumps to 49, so \mathcal{P} has rank 13 on this stratum. So we need to dig a little bit more. Let $W_8 := W_6 + U_5$. In this case we check that

$$\begin{aligned} \mathcal{A} \cap \mathcal{B} &= W_1 \wedge U_5 \wedge V_9 + T_1 \wedge W_6 \wedge V_9 + U_2 \wedge (\wedge^2 W_6) + W_3 \wedge U_5 \wedge W_6 \subset \\ &\subset \mathcal{C} := ((W_1 + T_1) \wedge W_8 \wedge V_9 + \wedge^3 W_8) \cap \mathcal{A} \subset \mathcal{A}. \end{aligned}$$

The dimension of \mathcal{C} is 56, so the rank of \mathcal{A}/\mathcal{C} is 6. But now \mathcal{C} and \mathcal{A}/\mathcal{C} only depend on W_1, T_1 and W_8 . The parameter space for (W_1, T_1, W_8) has dimension eight (one dimension for T_1 in U_2 ; four dimensions for W_1 in U_5 ; and three extra dimensions for

W_8 , that contains U_5). So the zero locus of the section induced by v has dimension at most 2 inside this stratum.

(2,3) Here $W_3 := W_6 \cap U_5$ has dimension 3 and $W_6 \supset U_2$. In this case we check that

$$\mathcal{A} \cap \mathcal{B} = W_1 \wedge U_5 \wedge V_9 + U_2 \wedge W_6 \wedge V_9 \subset \mathcal{C} := W_1 \wedge U_5 \wedge V_9 + U_2 \wedge (\wedge^2 V_9) \subset \mathcal{A}.$$

The bundle \mathcal{A}/\mathcal{C} has rank four and depends only on W_1 for which we have four parameters; so we are safe.

(1,4) Let $T_1 := W_6 \cap U_2$ and $W_4 := W_6 \cap U_5$. The parameter space for these flags has dimension 13 (one dimension for T_1 in U_2 ; four dimensions for W_1 in U_5 ; then two dimensions for W_4 in U_5 and containing $T_1 + W_1$; finally six extra dimensions for W_6 containing W_4). We check that

$$\mathcal{A} \cap \mathcal{B} = W_1 \wedge U_5 \wedge V_9 + T_1 \wedge W_6 \wedge V_9 + U_2 \wedge (\wedge^2 W_6) + W_4 \wedge U_5 \wedge W_6$$

has dimension 50. So \mathcal{P} has rank 12 on this stratum, and since $\dim(S) - \text{rank}_S(\mathcal{P}) = 1 < 3$, we are safe.

(2,5) This means that $U_5 \subset W_6$. But then $\mathcal{A} \cap \mathcal{B} = \wedge^2 U_5 \wedge V_9$. A dimension count shows that the general v does not belong to any such space. \square

Now, let us go back to the proof of the proposition. The only possibility for $[W_1]$ to be a general point of the hypersurface $\mathbb{P}(U_5) \cap \mathcal{S}$ is that $\dim(W_6 \cap U_2) = 2$ and $\dim(W_6 \cap U_5) = 4$. The parameter space for such pairs $[W_1 \subset W_6]$ has dimension 11 while the quotient bundle \mathcal{P} has rank 8 on this stratum. So the set of points $[W_1] \in \mathbb{P}(V_5) \cap \mathcal{S}$ such that $\dim(U_6 \cap V_2) = 2$ and $\dim(U_6 \cap V_5) = 4$ has expected dimension 3, i.e. it is a hypersurface inside $\mathbb{P}(U_5)$.

With the help of [GS] we can construct the parameter space for the stratum (2,4) and the quotient bundle \mathcal{P} , and check that the degree of $\mathcal{O}_{\mathbb{P}(U_5)}(1)$ on the zero locus of a general section of \mathcal{P} inside the parameter space is equal to three. So this zero locus projects to a cubic hypersurface in $\mathbb{P}(U_5)$. Set theoretically this cubic coincides with $\mathbb{P}(V_5) \cap \mathcal{S}$. Since \mathcal{S} is a sextic, this means that \mathcal{S} cuts $\mathbb{P}(U_5)$ along a double cubic hypersurface. The result follows. \square

In order to ensure we really get \mathbb{P}^4 's in a ruling of $\text{SU}_C(3)$, there remains to check that our family of linear spaces covers the whole moduli space. To be precise, we expect to get codimension one families of the two rulings. Indeed recall that the base of the rulings is five dimensional, exactly as $\overline{\text{Sing}}(\mathcal{S})$. But the latter is actually, by its very definition, a \mathbb{P}^1 -fibration over its image $\pi_{2,5}(\overline{\text{Sing}}(\mathcal{S}))$ through the projection $\pi_{2,5} : Fl(1, 2, 5, V_9) \rightarrow Fl(2, 5, V_9)$, whose birational image in $G(5, V_9)$ is the basis of our family. A similar phenomenon will happen for Hecke lines, as we will see in the next section. We have therefore a four dimensional family of \mathbb{P}^4 's and we want to show that these cover $\mathbb{P}(V_9)$.

Proposition 3.4. *There are 18 \mathbb{P}^4 's of the family $\pi_{2,5}(\overline{\text{Sing}}(\mathcal{S}))$ passing through the general point of $\mathbb{P}(V_9)$. In particular, the \mathbb{P}^4 's parametrized by $\pi_{2,5}(\overline{\text{Sing}}(\mathcal{S}))$ cover $\mathbb{P}(V_9)$.*

Proof. This is a degree computation that we carried out with [GS]. In order to do so, construct $\pi_{2,5}(\overline{\text{Sing}}(\mathcal{S}))$ inside $Fl(2, 5, V_9)$ as the zero locus of a general section of the quotient of $\wedge^3 V_9$ by $\mathcal{U}_2 \wedge (\wedge^2 V_9) + \wedge^2 \mathcal{U}_5 \wedge V_9$. Then consider the projective bundle $\mathbb{P}(\mathcal{U}_5)$ over $\pi_{2,5}(\overline{\text{Sing}}(\mathcal{S}))$ and its relative tautological bundle $\mathcal{O}_{\mathbb{P}(\mathcal{U}_5)}(-1)$. We get

$$\int_{\mathbb{P}(\mathcal{U}_5)} c_1(\mathcal{O}_{\mathbb{P}(\mathcal{U}_5)}(1))^8 = 18$$

and the result follows. \square

In fact we get the two rulings of $\mathrm{SU}_C(3)$, since by Remark 2.3.3 they are exchanged by the covering involution of $\mathrm{SU}_C(3)$. Indeed the two copies of \mathbb{P}^4 constituting $\theta^{-1}(\mathbb{P}(U_5))$, being exchanged by the involution, must belong to different rulings.

4. THE ODD MODULI SPACE AS AN ODL

4.1. Orbital degeneracy loci in $G(3, V_9)$. Over the Grassmannian, the Borel-Weil theorem provides still another identification

$$\wedge^3 V_9 \simeq H^0(G(3, V_9), \wedge^3 \mathcal{Q}),$$

where \mathcal{Q} now denotes the rank six tautological quotient bundle. This allows to consider the skew-symmetric cubic tensor v as a family of skew-symmetric cubic tensors in six variables. The $GL(V_6)$ -orbit closures in $\wedge^3 V_6$ are easy to describe: by increasing dimension (and forgetting the extremal ones) we have

- the cone Z_{10} over the Grassmannian $G(3, 6)$,
- the variety Z_5 of partially decomposable tensors,
- the tangent variety Z_1 to the Grassmannian, which is a quartic hypersurface.

Correspondingly, we get inside $G(3, V_9)$

- a hypersurface $D_{Z_1}(v)$ of the Grassmannian, which is a singular quadratic section,
- its singular locus $D_{Z_5}(v)$, whose singular locus is
- $D_{Z_{10}}(v)$, a smooth eightfold since $D_{\mathrm{Sing}(Z_{10})}(v) = \emptyset$ for dimensional reasons.

Remark 4.1. The hypersurface $D_{Z_1}(v)$ of $G(3, V_9)$ has obvious similarities with the Coble quadric discovered in [BBFM23]. As the latter, it is a special quadratic section of the Grassmannian. Nevertheless its relationship with the moduli space is less direct. The Coble quadric can be characterized as the unique quadric section of the Grassmannian $G(2, V_8)$ which is singular exactly on a copy of the odd moduli space of rank two vector bundles on a (non hyper-elliptic) genus three curve. Here, it is the singular locus of the quadric section, for which we are not aware of any modular interpretation, which is itself singular along a copy of the odd moduli space of rank three vector bundles on the genus two curve (indeed, we will show in Theorem 4.9 that $D_{Z_{10}}(v)$ identifies with such a moduli space). The quadric section is presumably uniquely determined by this property, but we did not prove this.

4.2. A Fano eightfold. By definition, $D_{Z_{10}}(v) \subset G(3, V_9)$ is the projection from $Fl(3, 6, V_9)$ of the locus

$$\tilde{D}_{Z_{10}}(v) := \{(U_3 \subset U_6) \in Fl(3, 6, V_9), \quad v \in U_3 \wedge (\wedge^2 V_9) + \wedge^3 U_6\}.$$

Proposition 4.2. $\tilde{D}_{Z_{10}}(v) \simeq D_{Z_{10}}(v)$ is a smooth Fano eightfold of even index.

Proof. First observe that $\tilde{D}_{Z_{10}}(v)$ is defined as the zero locus of a general section of a globally generated vector bundle of rank 19 on the flag manifold $Fl(3, 6, V_9)$, namely

$$\mathcal{E} = \wedge^3 V_9 / (\mathcal{U}_3 \wedge (\wedge^2 V_9) + \wedge^3 \mathcal{U}_6).$$

We deduce that $\tilde{D}_{Z_{10}}(v)$ is smooth of dimension eight. Moreover its canonical bundle can be computed by the adjunction formula: since the canonical bundle of $Fl(3, 6, V_9)$ is $\det(\mathcal{U}_3)^6 \otimes \det(\mathcal{U}_6)^6$, we find that

$$K_{\tilde{D}_{Z_{10}}(v)} = \det(\mathcal{U}_3)^{-3} \otimes \det(\mathcal{U}_6)^5|_{\tilde{D}_{Z_{10}}(v)}.$$

This does not seem to be negative, but observe that the quotient of $\mathcal{U}_3 \wedge (\wedge^2 V_9) + \wedge^3 \mathcal{U}_6$ by $\mathcal{U}_3 \wedge (\wedge^2 V_9)$ is just the line bundle $\wedge^3(\mathcal{U}_6/\mathcal{U}_3)$. Over $\tilde{D}_{Z_{10}}(v)$, our v defines a section of this line bundle, and this section is nowhere vanishing for v general. Indeed $U_3 \wedge (\wedge^2 V_9)$ has

codimension 30 in $\wedge^3 V_9$, and this is much bigger than the dimension of $G(3, V_9)$. We conclude that on $\tilde{D}_{Z_{10}}(v)$, the line bundle $\wedge^3(\mathcal{U}_6/\mathcal{U}_3)$ is trivial, which means that $\det(\mathcal{U}_3) \simeq \det(\mathcal{U}_6)$. As a consequence, we can rewrite the canonical bundle as

$$K_{\tilde{D}_{Z_{10}}(v)} = \det(\mathcal{U}_3) \otimes \det(\mathcal{U}_6)|_{\tilde{D}_{Z_{10}}(v)} \simeq \det(\mathcal{U}_6)^2|_{\tilde{D}_{Z_{10}}(v)}.$$

Since $\det(\mathcal{U}_3) \otimes \det(\mathcal{U}_6)$ is dual to the minimal very ample line bundle on the flag manifold $Fl(3, 6, V_9)$, the first identity shows that $\tilde{D}_{Z_{10}}(v)$ is Fano. The second identity implies it has even index.

We now prove that $\tilde{D}_{Z_{10}}(v)$ and $D_{Z_{10}}(v)$ are isomorphic. Let π be the projection from $Fl(3, 6, V_9)$ to $G(3, V_9)$. We denote in the same way its restriction from $\tilde{D}_{Z_{10}}(v)$ to $D_{Z_{10}}(v)$. Since $U_3 \wedge (\wedge^2 V_9)$ has codimension one inside $U_3 \wedge (\wedge^2 V_9) + \wedge^3 U_6$, the fact that $[U_3] \in D_{Z_{10}}(v)$ has two pre-images in $\tilde{D}_{Z_{10}}(v)$ would imply that v actually belongs to $U_3 \wedge (\wedge^2 V_9)$, and we already noticed this cannot be the case for v general. So $\pi : \tilde{D}_{Z_{10}}(v) \rightarrow D_{Z_{10}}(v)$ is a bijection. There remains to check it is everywhere immersive. For this we use the normal exact sequence

$$0 \rightarrow T\tilde{D}_{Z_{10}}(v) \rightarrow TFl(3, 6, V_9)|_{\tilde{D}_{Z_{10}}(v)} \rightarrow \mathcal{E}|_{\tilde{D}_{Z_{10}}(v)} \rightarrow 0.$$

Since the flag manifold is homogeneous, its tangent space at some point $(U_3 \subset U_6)$ can be seen as a quotient of $\mathfrak{sl}(V_9)$, by the space of morphisms that preserve the flag. The projection to $G(3, V_9)$ induces a surjective map between tangent spaces, whose kernel, the vertical tangent space, is just the tangent space $\text{Hom}(U_3, U_6/U_3)$ to the fiber $G(3, U_6)$. This vertical tangent space should be seen as the quotient of the subspace of $\mathfrak{sl}(V_9)$ consisting of morphisms that preserve U_3 . So the fact that π is immersive at $(U_3 \subset U_6)$ amounts to the following claim: if $X \in \mathfrak{sl}(V_9)$ preserves U_3 and sends v to $U_3 \wedge (\wedge^2 V_9) + \wedge^3 U_6$, then it preserves the full flag $(U_3 \subset U_6)$.

In order to check this, note that we can decompose $v = v' + e_4 \wedge e_5 \wedge e_6$, where v' belongs to $U_3 \wedge (\wedge^2 V_9)$ (which does not contain v , as we already stressed), and e_4, e_5, e_6 are a basis of U_6 modulo U_3 . Since X preserves U_3 , $X(v')$ belongs to $U_3 \wedge (\wedge^2 V_9)$. On the other hand, since $X(e_4 \wedge e_5 \wedge e_6) = X(e_4) \wedge e_5 \wedge e_6 + e_4 \wedge X(e_5) \wedge e_6 + e_4 \wedge e_5 \wedge X(e_6)$, it is clear that it belongs to $U_3 \wedge (\wedge^2 V_9) + \wedge^3 U_6$ if and only if X preserves U_6 , and the proof is complete. \square

4.3. A family of planes in the moduli space. We want to prove that the projective planes in $\mathbb{P}(V_9)$ parametrized by $D_{Z_{10}}(v)$ lift to planes in the moduli space $\text{SU}_C(3)$. As we already noticed for the \mathbb{P}^4 's of the rulings, this happens if the sextic \mathcal{S} restricted to such a plane restricts to a double cubic.

Proposition 4.3. *Let $[U_3] \in D_{Z_{10}}(v)$, then $\mathbb{P}(U_3) \cap \mathcal{S}$ is a double cubic, and $\theta^{-1}(\mathbb{P}(U_3))$ is the union of two projective planes meeting along a cubic curve.*

Proof. Since $[U_3] \in D_{Z_{10}}(v)$, there exists $U_6 \supset U_3$ such that $v \in U_3 \wedge (\wedge^2 V_9) + \wedge^3 U_6$. We are looking for flags $(W_1 \subset W_6)$ such that $W_1 \subset U_3$ and $v \in W_1 \wedge (\wedge^2 V_9) + (\wedge^2 W_6) \wedge V_9$. These flags correspond to points $[W_1] \in \mathbb{P}(U_3) \cap \mathcal{S}$. For a generic element $[W_1]$ in this intersection, we want to compute the values of $(\dim(W_6 \cap U_3), \dim(W_6 \cap U_6))$. We will prove:

Lemma 4.4. *$W_6 \supset U_3$ and $\dim(W_6 \cap U_6) = 5$.*

Proof. For convenience let us denote $\mathcal{A} = U_3 \wedge (\wedge^2 V_9) + \wedge^3 U_6$ and $\mathcal{B} = W_1 \wedge (\wedge^2 V_9) + (\wedge^2 W_6) \wedge V_9$. According to the relative position of the flags $(U_3 \subset U_6)$ and $(W_1 \subset W_6)$, the intersection of \mathcal{A} and \mathcal{B} will have a specific dimension, and will be constrained by the fact that it has to contain the general tensor v . Let us discuss these constraints more precisely, for each possible values of $(\dim(W_6 \cap U_3), \dim(W_6 \cap U_6))$; this couple of integers can a priori be either $(3, 6)$, $(3, 5)$, $(3, 4)$,

(3, 3), (2, 5), (2, 4), (2, 3), (1, 4), or (1, 3). We will discuss these cases roughly by increasing complexity, and exclude all of them except (3, 5).

(3,3) In this case $\mathcal{A} \cap \mathcal{B} = W_1 \wedge (\wedge^2 V_9) + U_3 \wedge W_6 \wedge V_9$, and we would deduce that $v \in \mathcal{A} \cap \mathcal{B} \subset U_3 \wedge (\wedge^2 V_9)$. But this is a contradiction, since we have already observed that the general tensor v is contained in no such space.

(2,3) Here we compute that

$$\mathcal{A} \cap \mathcal{B} = W_1 \wedge (\wedge^2 V_9) + (U_3 \cap W_6) \wedge W_6 \wedge V_9 + U_3 \wedge (\wedge^2 W_6),$$

So again $\mathcal{A} \cap \mathcal{B} \subset U_3 \wedge (\wedge^2 V_9)$ and we get the same contradiction as before.

(3,4) Same argument.

(1,4) In this case the parameter space for the flag $(U_1 \subset U_6)$ has dimension 14. Moreover

$$\mathcal{A} \cap \mathcal{B} = W_1 \wedge (\wedge^2 V_9) + U_3 \wedge (\wedge^2 W_6) + \wedge^3 U_6$$

has dimension 49. So v defines a general section of the vector bundle $\mathcal{A}/\mathcal{A} \cap \mathcal{B}$, of rank 16 over the 14-dimensional parameter space, and will therefore vanish nowhere in general.

(2,5) Here the parameter space for the flag $(U_1 \subset U_6)$ has dimension 9, while

$$\mathcal{A} \cap \mathcal{B} = W_1 \wedge (\wedge^2 V_9) + (W_6 \cap U_3) \wedge W_6 \wedge V_9 + U_3 \wedge (\wedge^2 W_6) + \wedge^3 U_6$$

has dimension 53. So the vector bundle $\mathcal{A}/\mathcal{A} \cap \mathcal{B}$ has rank 12 and we conclude as in the previous case.

(3,6) In this case $U_6 = V_6$ and the parameter space is a projective plane. Moreover

$$\mathcal{A} \cap \mathcal{B} = W_1 \wedge (\wedge^2 V_9) + U_3 \wedge U_6 \wedge V_9 + \wedge^3 U_6$$

has dimension 59. So the vector bundle $\mathcal{A}/\mathcal{A} \cap \mathcal{B}$ has rank 6 and we conclude as before.

(1,3) This is the biggest stratum, of dimension 17. Here

$$\mathcal{A} \cap \mathcal{B} = W_1 \wedge (\wedge^2 V_9) + U_3 \wedge (\wedge^2 W_6) + \wedge^2(U_6 \cap W_6) \wedge U_6$$

has dimension 49, so the quotient $\mathcal{A}/\mathcal{A} \cap \mathcal{B}$ has rank 16 which is smaller than the dimension of the parameter space. So we need to modify the argument a little bit. For this we observe that

$$\mathcal{A} \cap \mathcal{B} \subset \mathcal{C} := W_1 \wedge (\wedge^2 V_9) + \wedge^2(W_6 + U_3) \wedge U_3 + \wedge^3 U_6,$$

which has dimension 54 but only depends on W_1 and $W_8 := W_6 + U_3$. And now the parameter space for the flag $(W_1 \subset U_3 \subset W_8)$ is only 7-dimensional, while the bundle \mathcal{A}/\mathcal{C} has rank 11. So the previous argument applies.

(2,4) This case is somewhat similar to the previous case. Indeed the parameter space for the flag $(U_1 \subset U_6)$ has dimension 13, and we compute that

$$\mathcal{A} \cap \mathcal{B} = W_1 \wedge (\wedge^2 V_9) + (W_6 \cap U_3) \wedge W_6 \wedge V_9 + U_3 \wedge (\wedge^2 W_6) + \wedge^2(W_6 \cap U_6) \wedge U_6$$

has dimension 53, so the quotient $\mathcal{A}/\mathcal{A} \cap \mathcal{B}$ has rank 12 which is smaller than the dimension of the parameter space. But we observe that

$$\mathcal{A} \cap \mathcal{B} \subset \mathcal{D} := (W_6 \cap U_3) \wedge (\wedge^2 V_9) + \wedge^2(W_6 + U_6) \wedge U_3 + \wedge^3 U_6,$$

which has dimension 60 but only depends on $W_2 := W_6 \cap U_3$ and $W_8 := W_6 + U_6$. And now the parameter space for the flag $(W_2 \subset U_3 \subset U_6 \subset W_8)$ is only 4-dimensional, while the bundle \mathcal{A}/\mathcal{D} has rank 5. So the previous argument applies.

So we are only left with the case $(\dim(U_6 \cap V_3), \dim(U_6 \cap V_6)) = (3, 5)$, which concludes the proof of the Lemma. \square

In order to conclude the proof of the Proposition, we proceed as follows. The parameter space for the flags $W_1 \subset U_3 \subset W_6$ with $\dim(U_6 \cap W_6) \geq 5$ has dimension equal to 7. Over this space we have

$$\mathcal{A} \cap \mathcal{B} = W_1 \wedge (\wedge^2 V_9) + U_3 \wedge W_6 \wedge V_9 + \wedge^3 U_6,$$

which has constant dimension 59, even when W_6 specializes to U_6 . So $\mathcal{A}/\mathcal{A} \cap \mathcal{B}$ defines a rank six vector bundle \mathcal{W} on the parameter space.

Then with the help of [GS] we construct the parameter space and the vector bundle \mathcal{W} , and we check that the degree of $\mathcal{O}_{\mathbb{P}(V_3)}(1)$ on the zero locus of a section of \mathcal{W} inside the parameter space is equal to three. So this zero locus projects to a cubic curve E in $\mathbb{P}(V_3)$, and set-theoretically this cubic coincides with $\mathbb{P}(V_3) \cap \mathcal{S}$. But \mathcal{S} is a sextic, so it has to cut E doubly in $\mathbb{P}(V_3)$. The result follows. \square

So we get a family of planes in $SU_C(3)$. Let us denote by \mathcal{F} the family of lines contained in these planes. We need to check that the lines from \mathcal{F} cover the whole moduli space. This follows from the next statement:

Proposition 4.5. *The family of planes parametrized by $D_{Z_{10}}(v)$ covers $\mathbb{P}(V_9)$.*

Proof. It is sufficient to check that the eighth Chern class of the bundle $(\mathcal{O}_{\mathbb{P}(V_9)}(1))^{\oplus 8}$ over the projective bundle $p : \mathbb{P}(\mathcal{U}_3) \rightarrow D_{Z_{10}}(v)$ is different from zero. This can be done with the help of [GS]. Let \mathcal{L} be the pullback to $\mathbb{P}(\mathcal{U}_3)$ of the restriction of the bundle $\mathcal{O}_{G(3, V_9)}(1)$ to $D_{Z_{10}}(v)$. By constructing $D_{Z_{10}}(v)$ as a zero locus inside $Fl(3, 6, V_9)$, and $\mathbb{P}(\mathcal{U}_3)$ as a projective bundle over it, we checked that

$$\int_{\mathbb{P}(\mathcal{U}_3)} c_1(\mathcal{L})^2 c_1(\mathcal{O}_{\mathbb{P}(\mathcal{U}_3)}(1))^8 = 18,$$

which implies the claim. Similarly, in order to check this non-vanishing, we could have used the fact that $\int_{\mathbb{P}(\mathcal{U}_3)} c_2(\mathcal{M}) c_1(\mathcal{O}_{\mathbb{P}(\mathcal{U}_3)}(1))^8 = 6 \neq 0$, where $\mathcal{M} := (p^* \mathcal{U}_3 / \mathcal{O}_{\mathbb{P}(\mathcal{U}_3)}(-1))^\vee$, which can again be checked with [GS]. \square

Remark 4.6. The 18 appearing in the formula here above has a nice interpretation. Recall that $\mathbb{P}(\mathcal{U}_3)$ has dimension 10, hence - given a general point $p \in \mathbb{P}(V_9)$ - there is a surface inside $D_{Z_{10}}(v)$ that parametrizes planes through p . Moreover each point p defines two points $[E_1], [E_2] = \tau^*[E_1]$ in $SU_C(3)$. It turns out that the surface is a disjoint union of two \mathbb{P}^2 's embedded via their anticanonical class since the degree 18 is twice the degree of each \mathbb{P}^2 inside $D_{Z_{10}}(v) \subset G(3, V_9)$ in the Plücker embedding. In view of Theorem 4.9, these two planes correspond to the two Hecke planes inside $SU_C(3, \mathcal{O}_C(c))$ defined as extensions of E_1 by \mathcal{O}_c and E_2 by \mathcal{O}_c .

We will also need the following technical result.

Lemma 4.7. *The general line in \mathcal{F} does not meet $\text{Sing}(\mathcal{S})$.*

Proof. It is enough to prove that the general plane in $D_{Z_{10}}(v)$ meets $\text{Sing}(\mathcal{S})$ at most in a finite set. In order to prove this, we proceed as follows. A plane in $D_{Z_{10}}(v)$ corresponds to a flag $(U_3 \subset U_6)$, a point in $\text{Sing}(\mathcal{S})$ (or rather $\overline{\text{Sing}}(\mathcal{S})$) to a flag $(U_1 \subset U_2 \subset U_5)$. The point is contained in the plane if $U_1 \subset U_3$ and we need that $v \in \mathcal{C} \cap \mathcal{D}$ where

$$\mathcal{C} = U_3 \wedge (\wedge^2 V_9) + \wedge^3 U_6, \quad \mathcal{D} = U_2 \wedge (\wedge^2 V_9) + (\wedge^2 U_5) \wedge V_9.$$

We will stratify the parameter space by the relative position of the two flags $(U_1 \subset U_3 \subset U_6)$ and $(U_1 \subset U_2 \subset U_5)$. There are 23 possibilities. We can eliminate 9 of them right away by

observing that if $U_2 \cap U_3 = U_2 \cap U_6$, and $U_5 \cap U_3$ has codimension at most one in $U_5 \cap U_6$, then $\mathcal{C} \cap \mathcal{D} \subset U_3 \wedge (\wedge^2 V_9)$, which cannot happen in general.

For the 14 remaining strata S , we will compute the dimension of S and the rank of $\mathcal{C} \cap \mathcal{D}$ on S . If $\dim S + \text{rank}_S(\mathcal{C} \cap \mathcal{D}) \leq 92$, then the set of flags compatible with the general v will have dimension at most $92 - 84 = 8$, and will project to $D_{Z_{10}}(v)$ with finite (possibly empty) general fibers.

The data we computed are summarized in the following table, where the intersection dimensions in the first four columns specify each stratum S .

$U_2 \cap U_3$	$U_2 \cap U_6$	$U_5 \cap U_3$	$U_5 \cap U_6$	$\dim S$	$\text{rank}_S(\mathcal{C} \cap \mathcal{D})$
1	1	1	3	46	46
1	2	1	2	45	46
1	2	1	3	44	46
1	1	1	4	42	46
1	2	1	4	41	46
1	2	2	3	42	50
1	1	2	4	41	50
1	2	2	4	40	50
1	2	2	5	36	50
2	2	2	4	38	53
1	2	3	4	37	53
1	2	3	5	34	53
2	2	2	5	33	53
2	2	3	5	32	59

The sum of the two numbers in the righthmost columns does never exceed 92, and this implies our claim. \square

Proposition 4.8. *The family of lines in the planes parametrized by $D_{Z_{10}}(v)$ is a codimension one family of Hecke lines.*

Proof. The family \mathcal{F} of lines covers $\text{SU}_C(3)$ by the previous statement, so it must consist of Hecke lines or lines contained in the rulings. Let $\overline{\mathcal{F}}$ denote its image in $G(2, V_9)$, which is the image of a projective plane bundle over $D_{Z_{10}}(v)$. Since $D_{Z_{10}}(v)$ is Fano, it is rationally connected, so $\overline{\mathcal{F}}$ is rationally connected as well. If it was contained in $\overline{\mathcal{F}}_R$, we could consider its preimage inside \mathcal{F}_{R_1} and then $\hat{\mathcal{F}}_{R_1}$. By Lemmas 2.2 and 4.7, the preimage $\hat{\mathcal{F}}$ would still be rationally connected, and would therefore be contained in a fiber of the fibration of $\hat{\mathcal{F}}_{R_1}$ to the abelian surface $\text{Pic}^1(C)$. But these fibers have codimension two, while $\hat{\mathcal{F}}$ is a divisor: indeed $\overline{\mathcal{F}}$ is a projective plane bundle over $D_{Z_{10}}(v)$, so it has dimension 10, hence codimension one in \mathcal{F}_H . This is a contradiction. \square

4.4. The odd moduli space. We finally prove our main result.

Theorem 4.9. *For generic $v \in \wedge^3 V_9$, the orbital degeneracy locus $D_{Z_{10}}(v)$ is isomorphic to the moduli space $\text{SU}_C(3, \mathcal{O}_C(c))$ of semistable rank three vector bundles on C with fixed determinant $\mathcal{O}_C(c)$, for a certain point $c \in C$.*

Proof. By Proposition 4.3, each point of $D_{Z_{10}}(v)$ defines a pair of planes in $\text{SU}_C(3)$, exchanged by the cover involution. By Proposition 4.8 these planes are Hecke planes, that cover the whole moduli space by Proposition 4.5. According to Proposition 2.9, there are only two such covering families of Hecke planes, and they are exchanged by the cover involution. This implies that each point of $D_{Z_{10}}(v)$ defines a plane in each of the two families \mathcal{P}_H and \mathcal{P}_{H^*} . We




FIGURE 1. Minimal resolution of $I_{Z_{10}}$

can therefore lift $D_{Z_{10}}(v)$, at least birationally, to a hypersurface in either \mathcal{P}_H and \mathcal{P}_{H^*} . But recall from Proposition 2.9 that these two families fiber over C . Since $D_{Z_{10}}(v)$ is Fano, hence rationally connected, its pre-image in \mathcal{P}_H is also rationally connected, hence contained in a fiber of the projection to C , over some point c . Recall that this fiber is a birational image of $\mathrm{SU}_C(3, \mathcal{O}_C(c))$, and therefore the latter maps birationally to $D_{Z_{10}}(v)$. Since $\mathrm{SU}_C(3, \mathcal{O}_C(c))$ is smooth and its Picard group is cyclic, this birational morphism has to be an isomorphism. \square

We know that the moduli space $\mathrm{SU}_C(3, \mathcal{O}_C(c))$ has index two, so comparing with Proposition 4.2 we see that its canonical bundle must be the restriction of $\det(\mathcal{U}_3)^2$ on the Grassmannian, or also $\det(\mathcal{U}_3) \otimes \det(\mathcal{V}_6)$ on the flag manifold $Fl(3, 6, V_9)$. We deduce:

Corollary 4.10. $\tilde{D}_{Z_{10}}(v)$ is projectively equivalent to $\mathrm{SU}_C(3)$ in its anticanonical embedding.

4.5. Resolving the structure sheaf. An immediate consequence of our approach is that we can provide a resolution of the structure sheaf of $\mathrm{SU}_C(3, \mathcal{O}_C(c))$ inside $G(3, V_9)$. Indeed, by construction $D_{Z_{10}}(v)$ is modeled on the cone Z_{10} over $G(3, V_6) \subset \mathbb{P}(\wedge^3 V_6)$. The minimal resolution of its ideal $I_{Z_{10}}$ has been computed in [PW86]. Letting $S = \mathbb{C}[\wedge^3 V_6]$, this minimal resolution is reproduced in Figure 1.

Since this is an equivariant resolution, we can relativize it to the total space of $\wedge^3 Q$ over $G(3, V_9)$. This yields the resolution of the ideal sheaf of $D_{Z_{10}}(v)$ in Figure 2.

$$\begin{array}{c}
0 \\
\uparrow \\
I_{D_{Z_{10}}(v)} \\
\uparrow \\
S_{211111}Q^\vee \\
\uparrow \\
S_{21}Q^\vee(-1) \oplus S_{22221}Q^\vee(1) \\
\uparrow \\
S_{2211}Q^\vee(-1) \quad \longleftarrow \quad S_3Q^\vee(-2) \oplus S_{33333}Q^\vee \\
\uparrow \quad \longleftarrow \quad S_{3111}Q^\vee(-2) \oplus S_{222}Q^\vee(-2) \oplus S_{33222}Q^\vee(-1) \\
\uparrow \quad \longleftarrow \quad S_{32211}Q^\vee(-2) \oplus S_{32211}Q^\vee(-2) \\
\uparrow \quad \longleftarrow \quad S_{33222}Q^\vee(-2) \oplus S_{3111}Q^\vee(-3) \oplus S_{222}Q^\vee(-3) \\
\uparrow \quad \longleftarrow \quad S_{2211}Q^\vee(-4) \quad \longleftarrow \quad S_{33333}Q^\vee(-2) \oplus S_3Q^\vee(-4) \\
\uparrow \quad \longleftarrow \quad S_{22221}Q^\vee(-4) \oplus S_{21}Q^\vee(-5) \\
\uparrow \\
S_{211111}Q^\vee(-5) \\
\uparrow \\
\mathcal{O}(-7) \\
\uparrow \\
0
\end{array}$$

FIGURE 2. Minimal resolution of the ideal of $D_{Z_{10}}(v)$

Remark 4.11. Notice the remarkable fact that this corresponds to a Cohen-Macaulay (length equal to the codimension) and even Gorenstein (last term of rank one, which implies that the resolution is self-dual) locally free resolution of $\mathcal{O}_{D_{Z_{10}}(v)}$. In general, if \mathcal{F}_\bullet is a Gorenstein \mathcal{O}_X -locally free resolution of \mathcal{O}_Z , with $Z \subset X$, then the canonical bundle K_Z of Z is given by the formula:

$$K_Z = (K_X \otimes F_{-\text{codim}_X(Z)}^\vee)|_Z.$$

In our situation, we deduce that $K_{D_{Z_{10}}(v)}$ is the restriction of $K_{G(3, V_9)} \otimes \mathcal{O}(7) = \mathcal{O}(-2)$, which confirms what we proved in Proposition 4.2.

We can derive the following facts from the previous resolution.

Corollary 4.12. *The Hilbert polynomial of $(D_{Z_{10}}(v), \mathcal{O}_{D_{Z_{10}}(v)}(1))$ is*

$$\chi(\mathcal{O}_{D_{Z_{10}}(v)}(m-1)) = \frac{477}{2240} m^8 + \frac{63}{160} m^6 + \frac{99}{320} m^4 + \frac{47}{560} m^2.$$

In full generality, the Hilbert polynomial of the moduli space of vector bundles of fixed rank and determinant on a curve of given genus is given by the Verlinde formula, but in a form which is far from being obviously polynomial, see e.g. [Bea97, Proposition 3.1].

Proposition 4.13. $D_{Z_{10}}(v)$ has the following properties.

- (1) It is not linearly normal, but k -normal for any $k \geq 2$.
- (2) It is contained in 810 quadratic sections of the Grassmannian.
- (3) It is scheme theoretically cut-out by quadrics.

Proof. The Hilbert polynomial (recall that $D_{Z_{10}}(v)$ is Fano) gives $h^0(\mathcal{O}_{D_{Z_{10}}(v)}(1)) = 85$, which is bigger by one than the dimension of $\wedge^3 V_9$. The difference comes from the term $S^3 \mathcal{Q}^\vee(-2)$ in the resolution of $I_{D_{Z_{10}}(v)}$: by Bott's theorem $h^3(S^3 \mathcal{Q}^\vee(-1)) = 1$. From the same theorem we readily reduce that $h^1(I_{D_{Z_{10}}(v)}(k)) = 0$ for any $k \geq 2$, which proves the first assertion. The second one is just a computation. The last one follows from the fact that $S_{21111} \mathcal{Q}^\vee(2) = S_{21111} \mathcal{Q}$ is generated by global sections, hence also $I_{D_{Z_{10}}(v)}(2)$. \square

It would be nice to find an explanation of this failure of linear normality.

4.6. Application to K3 surfaces of genus 19. Consider again the embedding

$$D_{Z_{10}}(v) \simeq \mathrm{SU}_C(3, \mathcal{O}_C(c)) \subset G(3, V_9).$$

Since the moduli space is a Fano eightfold of index two, we will obtain a surface S with trivial canonical bundle by considering the zero locus of two general sections of the dual tautological bundle (or equivalently, the intersection of $D_{Z_{10}}(v)$ with a general translate of $G(3, 7)$ inside $G(3, V_9)$).

Proposition 4.14. *The surface S is a K3 surface of genus 19.*

Proof. In order to simplify the notations we let $D := D_{Z_{10}}(v)$ and $G := G(3, V_9)$. We need to check that S is connected and that $h^1(\mathcal{O}_S) = 0$. This is a standard computation: the Koszul complex of the section of $\mathcal{G} = \mathcal{U}^\vee \oplus \mathcal{U}^\vee$ that defines S in D resolves its structure sheaf as a \mathcal{O}_D -module, and we will be able to conclude if we can check that

$$h^i(D, \wedge^i \mathcal{G}^\vee) = h^i(D, \wedge^{i-1} \mathcal{G}^\vee) = 0 \quad \forall i > 0.$$

Since the bundle \mathcal{G} on D is the restriction of a bundle defined on the whole Grassmannian, we can check these vanishing by twisting the complex \mathcal{J}^\bullet of vector bundles on the Grassmannian that resolves \mathcal{O}_D , and verify that

$$(1) \quad h^{i+j}(G, \wedge^i \mathcal{G}^\vee \otimes \mathcal{J}^j) = h^{i+j}(D, \wedge^{i-1} \mathcal{G}^\vee \otimes \mathcal{J}^j) = 0 \quad \forall i > 0, \forall j \geq 0.$$

This is just an application of Bott's theorem on the Grassmannian. All bundles in play are homogeneous and even completely reducible: they come from certain semisimple representations of the parabolic subgroup P defining $G(3, V_9) = \mathrm{GL}(9)/P$. The exterior powers and tensor products appearing in (1) can thus be computed with [vLCL92] as exterior powers and tensor products of representations of the Levi factor of P , which is just $GL(3) \times GL(6)$. Then, it is possible to implement Bott's Theorem with a Python script to compute the cohomology of the corresponding bundles. In this way we were able to obtain all cohomology groups $H^k(G, \wedge^i \mathcal{G}^\vee \otimes \mathcal{J}^j)$ for all $i \geq 0, j \geq 0, k \geq 0$. In fact they all vanish except $H^0(G, \wedge^0 \mathcal{G}^\vee \otimes \mathcal{J}^0)$ and $H^{18}(G, \wedge^6 \mathcal{G}^\vee \otimes \mathcal{J}^{10})$, which are both one dimensional. This implies that $h^0(\mathcal{O}_S) = 1, h^1(\mathcal{O}_S) = 0$ and $h^2(\mathcal{O}_S) = 1$, proving that S is a K3 surface. The genus is 19 because the degree of $\mathcal{O}_S(1) := \mathcal{O}_G(1)|_S$ is equal to 36, as computed with [GS]. \square

Remark 4.15. Similar computations lead to the following observations and questions:

- (1) The Euler characteristic $\chi(S, \mathcal{U}_{|S}^\vee) = 9$, suggesting that $\mathcal{U}_{|S}^\vee$ has nine global sections, while we would expect seven. What is the explanation?
- (2) Since the normal bundle of S in $D = \mathrm{SU}_C(3, \mathcal{O}_C(p))$ is $N_{S/D} = \mathcal{U}_{|S}^\vee \oplus \mathcal{U}_{|S}^\vee$, this would imply that $h^0(S, N_{S/D}) = 9 + 9 = 18$. Do the intersection of the moduli space with translates of $G(3, 7)$ really form a family of K3 surfaces of dimension 18?
- (3) The Euler characteristic $\chi(S, \mathcal{E}nd_0(\mathcal{U}_{|S}^\vee)) = -2$. If we could prove that $\mathcal{U}_{|S}^\vee$ is stable, hence simple, we would deduce that his moduli space is a K3 surface. It is conjectured in [KM23, Section 1.2] that this moduli space should be a quartic K3 surface.

Remark 4.16. The construction of Hecke lines and planes in $\mathrm{SU}_C(3)$ shows that the odd moduli space $\mathrm{SU}_C(3, \mathcal{O}_C(p))$ has a different embedding inside $G(3, V_9)$ for each point $p \in C$. Equivalently, since this moduli space does not depend on p , it admits a rank three vector bundle \mathcal{E}_p for each p , generated by global sections. Considering a global section of $\mathcal{E}_p \oplus \mathcal{E}_q$, for p and q any two points in C , it follows from Proposition 4.14 that we will in general still get a K3 surface. Mukai and Kanemitsu recently suggested that this construction should yield a locally complete family of K3 surfaces of genus 19, a genus for which no such projective model was previously known [KM23].

REFERENCES

- [Bea97] Arnaud Beauville, *The Verlinde formula for PGL_p* , The mathematical beauty of physics (Saclay, 1996), Adv. Ser. Math. Phys., vol. 24, World Sci. Publ., River Edge, NJ, 1997, pp. 141–151.
- [Bea03] ———, *The Coble hypersurfaces*, C. R. Math. Acad. Sci. Paris **337** (2003), no. 3, 189–194.
- [BBFM23] Vladimiro Benedetti, Michele Bolognesi, Daniele Faenzi, and Laurent Manivel, *The Coble quadric*, arXiv e-print math.AG/2307.05993, 2023.
- [BFMT20a] Vladimiro Benedetti, Sara Angela Filippini, Laurent Manivel, and Fabio Tanturri, *Orbital degeneracy loci and applications*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **21** (2020), 169–206.
- [BFMT20b] ———, *Orbital degeneracy loci II: Gorenstein orbits*, Int. Math. Res. Not. IMRN (2020), no. 24, 9887–9932.
- [BMT21] Vladimiro Benedetti, Laurent Manivel, and Fabio Tanturri, *The geometry of the Coble cubic and orbital degeneracy loci*, Math. Ann. **379** (2021), no. 1-2, 415–440.
- [BB12] Michele Bolognesi and Sonia Brivio, *Coherent systems and modular subvarieties of $\mathrm{SU}_C(r)$* , Internat. J. Math. **23** (2012), no. 4, 1250037, 23.
- [BPGN97] Leticia Brambila-Paz, Iwona Grzegorzczak, and Peter E. Newstead, *Geography of Brill-Noether loci for small slopes*, J. Algebraic Geom. **6** (1997), no. 4, 645–669.
- [GS] Daniel R. Grayson and Michael E. Stillman, *Macaulay2, a software system for research in algebraic geometry*, Available at <http://www.math.uiuc.edu/Macaulay2/>.
- [GS15] Laurent Gruson and Steven V. Sam, *Alternating trilinear forms on a nine-dimensional space and degenerations of $(3, 3)$ -polarized Abelian surfaces*, Proc. Lond. Math. Soc. (3) **110** (2015), no. 3, 755–785.
- [GSW13] Laurent Gruson, Steven V. Sam, and Jerzy Weyman, *Moduli of abelian varieties, Vinberg θ -groups, and free resolutions*, Commutative algebra, Springer, New York, 2013, pp. 419–469.
- [Hwa02] Jun-Muk Hwang, *Hecke curves on the moduli space of vector bundles over an algebraic curve*, Algebraic geometry in East Asia (Kyoto, 2001), World Sci. Publ., River Edge, NJ, 2002, pp. 155–164.
- [KM23] Akihiro Kanemitsu and Shigeru Mukai, *Polarized k3 surfaces of genus thirteen and genus three*, arXiv e-print math.AG/2310.02078, 2023.
- [MS09] Ngaiming Mok and Xiao Tao Sun, *Remarks on lines and minimal rational curves*, Sci. China Ser. A **52** (2009), no. 4, 617–630.
- [MTiB20] Yusuf Mustopa and Montserrat Teixidor i Bigas, *Rational curves on moduli spaces of vector bundles*, arXiv e-print math.AG/2007.10511, 2020.
- [NR75] Mudumbai S. Narasimhan and Sundararaman Ramanan, *Deformations of the moduli space of vector bundles over an algebraic curve*, Ann. of Math. (2) **101** (1975), 391–417.
- [NR78] ———, *Geometry of Hecke cycles. I.*, C. P. Ramanujam—a tribute, Tata Inst. Fund. Res. Studies in Math., 8, ,, 1978, pp. 291–345.

- [Ngu07] Quang Minh Nguyen, *Vector bundles, dualities and classical geometry on a curve of genus two*, Internat. J. Math. **18** (2007), no. 5, 535–558.
- [Ort05] Angela Ortega, *On the moduli space of rank 3 vector bundles on a genus 2 curve and the Coble cubic*, J. Algebraic Geom. **14** (2005), no. 2, 327–356.
- [Pal16] Sarbeswar Pal, *Moduli of rank 2 stable bundles and Hecke curves*, Canad. Math. Bull. **59** (2016), no. 4, 865–877.
- [PW86] Piotr Pragacz and Jerzy Weyman, *On the construction of resolutions of determinantal ideals: a survey*, Séminaire d’algèbre Paul Dubreil et Marie-Paule Malliavin, Lecture Notes in Math., vol. 1220, Springer, 1986, pp. 73–92.
- [RS18] Eric M. Rains and Steven V. Sam, *Invariant theory of $\wedge^3(9)$ and genus-2 curves*, Algebra Number Theory **12** (2018), no. 4, 935–957.
- [RS19] ———, *Vector bundles on genus 2 curves and trivectors*, Algebr. Geom. **6** (2019), no. 3, 328–345.
- [Ram73] Sundararaman Ramanan, *The moduli spaces of vector bundles over an algebraic curve*, Math. Ann. **200** (1973), 69–84.
- [Sun05] Xiaotao Sun, *Minimal rational curves on moduli spaces of stable bundles*, Math. Ann. **331** (2005), no. 4, 925–937.
- [vLCL92] Marc A.A. van Leeuwen, Arjeh M. Cohen, and Bert Lisser, *LiE, a package for Lie group computations*, Computer Algebra Nederland, Amsterdam, 1992.

Université Côte d’Azur, CNRS – Laboratoire J.-A. Dieudonné, Parc Valrose, F-06108 Nice Cedex 2, FRANCE.
Email address: vladimiro.benedetti@univ-cotedazur.fr

Institut Montpellierain Alexander Grothendieck, Université de Montpellier, Place Eugène Bataillon, 34095 Montpellier Cedex 5, FRANCE.

Email address: michele.bolognesi@umontpellier.fr

Institut de Mathématiques de Bourgogne, Université de Bourgogne et Franche-Comté, 9 Avenue Alain Savary, 21078 Dijon Cedex, FRANCE.

Email address: daniele.faenzi@u-bourgogne.fr

Institut de Mathématiques de Toulouse, Paul Sabatier University, 118 route de Narbonne, 31062 Toulouse Cedex 9, FRANCE.

Email address: manivel@math.cnrs.fr