



HAL
open science

Stereoselective Preparation of CF 3 -Containing Cyclopropanes

Veronika Myronova, Dominique Cahard, Ilan Marek

► **To cite this version:**

Veronika Myronova, Dominique Cahard, Ilan Marek. Stereoselective Preparation of CF 3 -Containing Cyclopropanes. *Organic Letters*, 2022, 24 (49), pp.9076-9080. 10.1021/acs.orglett.2c03714 . hal-04234818

HAL Id: hal-04234818

<https://hal.science/hal-04234818v1>

Submitted on 10 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


Stereoselective Preparation of CF₃-Containing Cyclopropanes

Veronika Myronova^{†‡}, Dominique Cahard^{†*} and Ilan Marek^{‡*}

[‡] The Mallat Family Laboratory of Organic Chemistry, Schulich Faculty of Chemistry, Technion - Israel Institute of Technology, Haifa 3200009, Israel.

[†] UMR 6014 CNRS COBRA, Normandie Université, 76821 Mont Saint Aignan, France


Supporting Information Placeholder



ABSTRACT: A regio- and diastereoselective carbometalation of easily accessible CF₃-substituted cyclopropenes is developed with a diastereoselectivity of the addition opposite to the CF₃ group. This simple strategy allows the preparation of polysubstituted (up to penta-) cyclopropyl rings possessing two adjacent quaternary carbon stereocenters with excellent diastereoselectivities.


The unique properties of fluorine atoms incorporated into organic molecules have created a lot of interests from the synthetic, physical and biological communities.^{1,2} Similarly, cyclopropyl ring is a basic structural subunit, widespread both in natural and synthetic bioactive molecules.³ It was therefore natural that combining cyclopropane and fluorine moieties (CH₂F, CHF₂ or CF₃) in the same molecular structure would attract the curiosity of chemists.⁴ Indeed, extensive studies have appeared in the last few years,⁵ leading to several bioactive approved drug candidates such as Glecaprevir,⁶ Patesicab⁷ and VX-659⁸ (Scheme 1a). It became rapidly clear that CF₃-substituted cyclopropanes as synthetic scaffold is of a constant interest giving rise to several powerful stereoselective approaches.⁹ Among the most prominent strategies reported in the literature (Scheme 1a),¹⁰ the metal-catalyzed (Fe, Ru, Rh, Co)^{9a, 11} and the bioengineered myoglobin-catalyzed¹² addition of diazo compounds to variously substituted alkenes represent certainly the best routes to these important potential drugs subunits. In recent years, we have developed several approaches to the synthesis of polysubstituted cyclopropanes by diastereoselective carbometalation reaction of cyclopropenyl ester and achiral cyclopropenes.¹³ In this vein, we questioned if we could use this strategy for the diastereoselective preparation of these important polysubstituted trifluoromethyl cyclopropyl cores. Although the synthetic scheme looks straightforward, several important questions arise such as the effect of the CF₃ group as potential chemical handle to control (or not) the diastereoselectivity of the addition¹⁴ as well as the stability of a cyclopropyl metal species towards ring-fragmentation containing a CF₃ group (Scheme 1b). Trifluoromethyl-substituted cyclopropenes **1** have already been described in the literature¹⁵ and are therefore easily accessible, even in enantiomerically enriched form (see supporting information for all details).¹⁶ Having in hand various cyclopropenes **1**, we started our study by performing

the copper-catalyzed carbomagnesiation on our model compound **1a** (R = Bu, R¹ = Ph).



Scheme 1. CF₃-containing cyclopropanes: state-of-the-art and research proposal.


When the reaction was performed in Et₂O at low temperature, we were gratified to realize that the addition proceeded very rapidly but an instantaneous ring-fragmentation occurred to provide the diene **3** (Scheme 2).¹⁷ Interestingly, from the stereochemistry of **3**, it appeared that the carbometalation reaction proceeded from the opposite face of the CF₃ moiety, *syn* to the aryl group.



Scheme 2. Copper-catalyzed carbomagnesiation of trifluoromethyl cyclopropene 1a

To avoid the ring-fragmentation, we surmised that a more covalent cyclopropyl metal species should be less prone towards this β -elimination reaction. We therefore considered that the addition of an organocopper, easily prepared by addition of alkyllithium or alkylmagnesium bromide to a stoichiometric amount of copper salt, would lead to a stable cyclopropyl copper species.¹⁸ Indeed, when 1 equiv. of alkylmagnesium bromide (or alkyllithium) was added to 1 equiv. of CuBr·Me₂S in THF at -40 °C, the resulting organocopper reacted smoothly with **1a** to provide after hydrolysis, the expected trifluoromethyl cyclopropane **2a** in 73% yield with an excellent diastereoselectivity (*dr* 97:03, Scheme 3). The opposite diastereomer can easily be prepared by simply permuting the nature of the alkyl groups on the organocopper and cyclopropene. In this case, the opposite diastereomer **2b** was obtained with a similar diastereomeric ratio (Scheme 3). The organocopper could similarly be generated from an alkyllithium instead of the alkylmagnesium halide with similar diastereoselectivity; a subscript (**Li**) is indicated in those cases after the compound numbers to indicate the nature of the initial organocopper. The formation of a discrete organometallic species was proved by reaction with DCI (**2c**, Scheme 3). The reaction proceeds with various R² group on the organocopper as well as various R group on the cyclopropene (**2d-l**, Scheme 3). The electronic nature of the aromatic group can also be varied as shown for the examples **2m** and **2n**. The resulting configurationally stable cyclopropyl copper could react with various carbon electrophiles such as allyl and propargyl bromide or acyl chlorides to provide pentasubstituted cyclopropanes (**2o-2ac** respectively) possessing two adjacent quaternary carbon stereocenters including the desired trifluoromethyl group with excellent diastereomeric ratios (Scheme 3). The relative configuration was established by X-ray analysis of **2ab** and the configurations of all other products were assigned by analogy. It is interesting to note that the CF₃ group is not coordinating to the copper complex, and therefore the latter reacts *anti* to the trifluoromethyl group. The diastereoselectivity of the addition diverges completely from all the additions on cyclopropene that has a methyl group instead of the CF₃.¹⁹ Following the positive results for the diastereocontrol of the carbometalation reaction of sp³-disubstituted **1** (possessing an aryl group geminated to the CF₃ unit), we then further continued our investigation on the diastereoselectivity of sp³-monosubstituted trifluoromethyl cyclopropenes (R¹ = H, Scheme 1). The syn-


thesis of the starting material was again performed by using reported procedures (see supporting information for all details)^{15a} and two sp²-monosubstituted cyclopropenes **1g,h** were easily prepared (Scheme 4). Having in hand, these two sp²-substituted cyclopropenes **1g,h**, the same experimental conditions as above were tested for the carbometalation reaction.



Scheme 3. Diastereoselective carbocupration of sp³-disubstituted CF₃-cyclopropenes 1a-f

We were again pleased to observe a smooth addition of the organocopper to the strained double bond of trifluoromethyl cyclopropene but surprisingly, final acidic hydrolysis needed to be performed with 1 M HCl to get reproducible results (most probably MeOH or H₂O were not acidic enough to quantitatively quench the copper salt leading to potential side reactions). We then generalized this reaction to our two substrates and in all cases, the carbometalation proceeds with an *anti* diastereofacial preference towards the CF₃ group (Scheme 4). The relative configurations were assigned by analysis of the coupling constants (observed value *J* = 5.8 - 6 Hz, litera-


ture values for $J_{\text{trans}} = 4 - 7.5$ Hz and for $J_{\text{cis}} = 7.5 - 9$ Hz).²⁰ It is interesting to note that although one of the two diastereotopic faces of the cyclopropenes **1g,h** is now unshielded as compared to **1a-f**, the level of diastereoselectivity of the carbometalation reaction for the formers are intriguingly lower than that the latter (compare **2f**(Li) with **2ae**(Li), **2l** with **2ah** or **2o** with **2ai**, Schemes 3 and 4 respectively), underlining the positive influence of an aryl group on the diastereoselectivity of the carbometalation reaction.



Scheme 4. Diastereoselective carbocupration of sp^3 -monosubstituted CF_3 -cyclopropenes **1g,h**


An alternative group that would exert a strong diastereocontrol for a *syn*-addition reaction should be an ester. Indeed, when the easily accessible cyclopropenyl ester **1i** possessing a CF_3 group,²¹ was treated with a methyl Grignard reagent in the presence of a substoichiometric amount of CuI, the carbometalation reaction proceeded at low temperature to give a thermally stable cyclopropyl metal species that could further react with carbon electrophile to provide the corresponding products **2aq-2as** in excellent yields and diastereoselectivities (Scheme 5). It should be noted that although the stereochemistry at the carbinol center of **2as** is formed in a 1:1 ratio, the cyclopropyl magnesium species is stable towards β -

fragmentation, most probably due to the intramolecular chelation from the ester.



Scheme 5. Diastereoselective carbocupration of CF_3 -cyclopropenyl ester **1i**

Finally, the use of chiral ligand in the cyclopropenation of alkynes provided the cyclopropenes **1a,b** with high enantiomeric ratios (*er* 95:05, Scheme 6, see Supporting Information).¹⁶ After carbocupration reaction and eventually reaction with an electrophile, the corresponding tetra- and penta-substituted cyclopropanes **2b** and **2r**, respectively, were obtained with excellent diastereomeric and enantiomeric ratios (Scheme 6).



Scheme 6. Enantio- and diastereoselective preparation of trifluoromethyl cyclopropanes

In conclusion, we have developed a regio- and diastereoselective carbometalation of easily accessible CF_3 -substituted cyclopropenes. The diastereoselectivity of the addition is opposite to the CF_3 group and the exact reasons still need to be fully clarified. This simple strategy allows the preparation of polysubstituted (penta-) cyclopropanes possessing two quaternary carbon stereocenters with excellent diastereoselectivities.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website. Experimental procedures, characterization data for all new compounds, along with copies of spectra.

Computational methods and data, geometries of computed intermediates and transition states (PDF).

Accession Codes

CCDC 2176014 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033

AUTHOR INFORMATION

Corresponding Authors

Ilan Marek. Email: chilanm@technion.ac.il

Dominique Cahard. Email: dominique.cahard@univ-rouen.fr

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

Funding Sources

This project received funding from the Israel Science Foundation administrated by the Israel Academy of Sciences and Humanities (Grant No.487/21), from the graduate school of research XI-Chem (ANR-18-EURE-0020 XL CHEM), and by Region Normandie.

REFERENCES

(1) For recent reviews, see: (a) O'Hagan, D. Understanding organofluorine chemistry. An introduction to the C–F bond. *Chem. Soc. Rev.* **2008**, *37*, 308–319. (b) Ojima, I. Exploration of fluorine chemistry at the multidisciplinary interface of chemistry and biology. *J. Org. Chem.* **2013**, *78*, 6358–6383. (c) Emerging Fluorinated Motifs. Properties, Synthesis and Applications, Volumes 1&2. Editors: Dominique Cahard & Jun-An Ma, **2020**. Wiley-VCH Verlag GmbH & Co. KGaA, Boschste. 12, 69469 Weinheim, Germany. ISBN 978–3–527–34681–3. (d) Cahard, D.; Bizet, V. The influence of fluorine in asymmetric catalysis. *Chem. Soc. Rev.* **2014**, *43*, 135–147.

(2) (a) Wang, J.; Sánchez-Roselló, M.; Aceña, J. L.; del Pozo, C.; Sorochinsky, A. E.; Fustero, S.; Soloshonok, V. A.; Liu, H. Fluorine in pharmaceutical industry: fluorine-containing drugs introduced to the market in the last decade (2001–2011). *Chem. Rev.* **2014**, *114*, 2432–2506. (b) Zhou, Y.; Wang, J.; Gu, Z.; Wang, S.; Zhu, W.; Aceña, J. L.; Soloshonok, V. A.; Izawa, K.; Liu, H. Next generation of fluorine-containing pharmaceuticals, compounds currently in phase II–III clinical trials of major pharmaceutical companies: new structural trends and therapeutic areas. *Chem. Rev.* **2016**, *116*, 422–518.

(3) (a) Wong, H. N. C.; Hon, M. Y.; Tse, C. W.; Yip, Y. C.; Tanko, J.; Hudlicky, T. Use of cyclopropanes and their derivatives in organic synthesis. *Chem. Rev.* **1989**, *89*, 165–198. (b) Talele, T. T. The “Cyclopropyl fragment” is a versatile player that frequently appears in preclinical/clinical drug molecules. *J. Med. Chem.* **2016**, *59*, 8712–8756.

(4) (a) Pons, A.; Delion, L.; Poisson, T.; Charette, A. B.; Jubault, P. Asymmetric synthesis of fluoro, fluoromethyl, difluoromethyl, and trifluoromethylcyclopropanes. *Acc. Chem. Res.* **2021**, *54*, 2969–2990. (b) Bos, M.; Poisson, T.; Pannecoucke, X.; Charette, A. B.; Jubault, P. Recent progress toward the synthesis of trifluoromethyl- and difluoromethyl-substituted cyclopropanes. *Chem. Eur. J.* **2017**, *23*, 4950–4961; (c) David, E.; Milanole, G.; Ivashkin, P.; Couve-Bonnaire, S.; Jubault, P.; Pannecoucke, X. Syntheses and applications of monofluorinated cyclopropanes. *Chem. Eur. J.* **2012**, *18*, 14904–14917; (d) Pons, A.; Poisson, T.; Pannecoucke, X.; Charette, A. B.; Jubault, P. Synthesis and applications of fluorocyclopropanes. *Synthese* **2016**, 4060–4071.

(5) Inoue, M.; Sumii, Y.; Shibata, N. Contribution of organofluorine compounds to pharmaceuticals. *ACS Omega.* **2020**, *5*, 10633–10640.

(6) Abutaleb, A.; Kottlil, S.; Wilson, E. Glecaprevir/pibrentasvir expands reach while reducing cost and duration of hepatitis C virus therapy. *Hepatology International* **2018**, *12*, 214–222.

(7) Sánchez, R. A.; Banner, D.; Ceccarelli, S. M.; Grether, U.; Haap, W.; Hartman, P.; Hartmann, G.; Hilpert, H.; Kuehne, H.; Mauser, H.; Plancher, J.-M. in *Novel Proline Derivatives*, US2010/0267722, 2010.

(8) Davies, J. C.; Moskowitz, S. M.; Brown, C.; Horsley, A.; Mall, M. A.; McKone, E. F.; Plant, B. J.; Prais, D.; Ramsey, B. W.; Taylor-Cousar, J. L.; Tullis, E.; Uluer, A.; McKee, C. M.; Robertson, S.; Shilling, R. A.; Simard, C.; Van Goor, F.; Waltz, D.; Xuan, F.; Young, T.; Rowe, S. M. CX-659-Tezacaftor-ivacaftor in patients with cystic fibrosis and one or two Phe508del alleles. *N. Engl. J. Med.* **2018**, *379*, 1599–1611.

(9) (a) Huang, W.-S.; Schlinquer, C.; Poisson, T.; Pannecoucke, X.; Charette, A. B.; Jubault, P. General catalytic enantioselective access to monohalomethyl and trifluoromethyl cyclopropanes. *Chem. Eur. J.* **2018**, *24*, 10339–10343. (b) Chen, S.; Zhang, J.; Yang, M.; Liu, F.; Xie, Z.; Liu, Y.-L.; Lin, W.; Wang, D.; Li, X.; Wang, J. Diastereoselective synthesis of cyclopropanes bearing trifluoromethyl-substituted all-carbon quaternary centers from 2-trifluoromethyl-1,3-enynes beyond fluorine elimination. *Chem. Commun.* **2019**, *55*, 3879. (c) Wu, W.; Lin, J.; Xiao, J.; Cao, Y.; Ma, Y. Recent advances in the synthesis of CF₃- or HCF₂- substituted cyclopropanes. *Asian J. Org. Chem.* **2021**, *10*, 485–495.

(10) For selected reviews, see: (a) Decaens, J.; Couve-Bonnaire, S.; Charette, A. B.; Poisson, T.; Jubault, P. Synthesis of fluoro-, monofluoromethyl-, difluoromethyl-, and trifluoromethyl-substituted three-membered rings. *Chem. Eur. J.* **2021**, *27*, 2935–2962. For recent examples: (b) Cyr, P.; Flynn-Robitaille, J.; Boissarie, P.; Marinier, A. mild and diazo-free synthesis of trifluoromethyl cyclopropanes using diafonium ylides. *Org. Lett.* **2019**, *21*, 2265–2268. (c) Chen, G.-S.; Yan, X.-X.; Chen, S.-J.; Mao, X.-Y.; Li, Z.-D.; Liu, Y.-L. Diastereoselective synthesis of 1,3-diyne-tethered trifluoromethylcyclopropanes through a sulfur ylide mediated cyclopropanation/DBU-mediated epimerization sequence. *J. Org. Chem.* **2020**, *85*, 6252–6260. (d) Chernykh, A. V.; Olifir, O. S.; Kuchkovska, Y. O.; Volochnyuk, D. M.; Yarmolchuk, V. S.; Grygorenko, O. O. Fluoroalkyl-substituted cyclopropane derivatives: synthesis and physicochemical properties. *J. Org. Chem.* **2020**, *85*, 12692–12702.

(11) (a) Müller, P.; Grass, S.; Shahi, S. P.; Bernardinelli, G. Rh(II)-Catalyzed asymmetric carbene transfer with ethyl 3,3,3-trifluoro-2-diazopropionate. *Tetrahedron* **2004**, *60*, 4755–4763. (b) Le Maux, P.; Juillard, S.; Simonneaux, G. R. Asymmetric Synthesis of Trifluoromethylphenyl Cyclopropanes Catalyzed by Chiral Metalloporphyrins. *Synthesis* **2006**, 1701–1704. (c) Denton, J. R.; Sukumaran, D.; Davies, H. M. L. Enantioselective synthesis of trifluoromethyl-substituted cyclopropanes. *Org. Lett.* **2007**, *9*, 2625–2628. (d) Duncton, M. A. J.; Singh, R. Synthesis of trans-2-(trifluoromethyl)cyclopropanes via Suzuki reactions with an N-methyliminodiacetic acid boronate. *Org. Lett.* **2013**, *15*, 4284–4287. (e) Morandi, B.; Mariampillai, B.; Carreira, E. M. Enantioselective Cobalt-Catalyzed Preparation of Trifluoromethyl-Substituted Cyclopropanes. *Angew. Chem. Int. Ed.* **2011**, *50*, 1101–1104. (f) Kotozaki, M.; Chanthamath, S.; Fujii, T.; Shibatomi, K.; Iwasa, S. Highly Enantioselective Synthesis of Trifluoromethyl Cyclopropanes by Using Ru(II)–Pheox Catalysts. *Chem. Commun.* **2018**, *54*, 5110–5111.

(12) (a) Tinoco, A.; Steck, V.; Tyagi, V.; Fasan, R. Highly diastereo and enantioselective synthesis of trifluoromethyl-substituted cyclopropanes via myoglobin-catalyzed transfer of trifluoromethylcarbene. *J. Am. Chem. Soc.* **2017**, *139*, 5293–5296. (b) Carminati, D. M.; Decaens, J.; Couve-Bonnaire, S.; Jubault, P.; Fasan, R. Biocatalytic strategy for the highly stereoselective synthesis of CHF₂-containing trisubstituted cyclopropanes. *Angew. Chem. Int. Ed.* **2021**, *60*, 7072–7076.

- (13) (a) Cohen, Y.; Marek, I. Directed regioselective carbometallation of 1,2-dialkyl-substituted cyclopropenes. *Angew. Chem. Int. Ed.* **2021**, *60*, 26368–26372. (b) Müller, D. S.; Marek, I. Asymmetric copper-catalyzed carbocation of cyclopropenes en route to the formation of diastereo- and enantiomerically enriched polysubstituted cyclopropanes. *J. Am. Chem. Soc.* **2015**, *137*, 15414–15417. (c) Dian, L.; Müller, D. S.; Marek, I. Asymmetric copper-catalyzed carbomagnesiation of cyclopropenes. *Angew. Chem. Int. Ed.* **2017**, *56*, 6783–6787. (d) Müller, D. S.; Werner, V.; Akyol, S.; Schmalz, H.-G.; Marek, I. Tandem hydroalumination/Cu-catalyzed asymmetric vinyl metalation as a new access to enantioenriched vinylcyclopropane derivatives. *Org. Lett.* **2017**, *19*, 3970–3973.
- (14) (a) Yamazaki, T.; Ando, M.; Kitazume, T.; Kubota, T.; Imura, M. Conformational fixation of enolates by intramolecular metal fluorine interaction. *Org. Lett.* **1999**, *1*, 905–908; (b) Sini, G.; Tessier, A.; Pytkowicz, J.; Brigaud, T. Fluorine and π -alkali metal interactions control in the stereoselective amide enolate alkylation with fluorinated oxazolidines (Fox) as a chiral auxiliary. An experimental and theoretical study. *Chem. Eur. J.* **2008**, *14*, 3363–3370; (c) Zell, D.; Müller, V.; Dhawa, U.; Bursch, M.; Presa, R. R.; Grimme, S.; Ackermann, L. Mild cobalt(III)-catalyzed allylative C-F/C-H functionalizations at room temperature. *Chem. Eur. J.* **2017**, *23*, 12145–12148; (d) Takemura, H.; Iwanaga, T.; Shinmyozu, T. C-F Rb⁺ interaction in a fluorinated cage complex. *Tetrahedron Lett.* **2006**, *47*, 8989–8991. (e) Plenio, H. The coordination chemistry of fluorine in fluorocarbons. *ChemBioChem.* **2004**, *5*, 650–655. (f) Takemura, H.; Nakashima, S.; Kon, N.; Yasutake, M.; Shinmyozu, T.; Inazu, T. A Study of C–F \cdots M⁺ Interaction: Metal Complexes of Fluorine-Containing Cage Compounds. *J. Am. Chem. Soc.* **2001**, *123*, 9293–9298.
- (15) (a) Morandi, B.; Carreira, E. M. Rhodium-catalyzed cyclopropanation of alkynes: synthesis of trifluoromethyl-substituted cyclopropenes. *Angew. Chem. Int. Ed.* **2010**, *49*, 4294–4296. (b) Tanbouza, N.; Carreras, V.; Ollevier, T. Photochemical cyclopropanation of alkynes with diazirines as carbene precursors in continuous flow. *Org. Lett.* **2021**, *23*, 5420–5424. (c) Barroso, R.; Jiménez, A.; Carmen Pérez-Aguilar, M.; Cabal, M.-P.; Valdés, C. Synthesis of 1,3-diaryl-3-trifluoromethylcyclopropenes by transition-metal-free reaction of 2,2,2-trifluoroacetophenone tosylhydrazones with alkynes: the effect of the trifluoromethyl group. *Chem. Commun.* **2016**, *52*, 3677–3680. (d) Briones, J. F.; Davies, H. M. L. Silver Triflate-Catalyzed Cyclopropanation of Internal Alkynes with Donor-/Acceptor-Substituted Diazo Compounds. *Org. Lett.* **2011**, *13*, 3984–3987.
- (16) (a) Tran, U. P. N.; Hommelsheim, R.; Yang, Z.; Empel, C.; Hock, K. J.; Nguyen, T. V.; Koenigs, R. M. Catalytic synthesis of trifluoromethyl cyclopropenes and oligo-cyclopropenes. *Chem. Eur. J.* **2020**, *26*, 1254–1257. (b) Zhang, X.; Tian, C.; Wang, Z.; Sivaguru, P.; Nolan, S. P.; Bi, X. Fluoroalkyl N-triftosylhydrazones as easily decomposable diazo surrogates for asymmetric [2 + 1] cycloaddition: synthesis of chiral fluoroalkyl cyclopropenes and cyclopropanes. *ACS Catalysis* **2021**, *11*, 8527–8537. (c) Uehara, M.; Suematsu, H.; Tasutomi, Y.; Katsuki, T. Enantioenriched synthesis of cyclopropenes with a quaternary stereocenter, versatile building blocks. *J. Am. Chem. Soc.* **2011**, *133*, 170–171.
- (17) Didier, D.; Delaye, D.; Simaan, M.; Island, B.; Eppe, G.; Eijsberg, H.; Kleiner, A.; Knochel, P.; Marek, I. Modulable and highly diastereoselective carbometalations of cyclopropenes. *Chem. Eur. J.* **2014**, *20*, 1038–1048.
- (18) (a) Dian, L.; Marek, I. Asymmetric preparation of polysubstituted cyclopropanes based on direct functionalization of achiral three-membered carbocycles. *Chem. Rev.* **2018**, *118*, 8415–8434. (b) Müller, D. S.; Marek, I. Copper mediated carbometalation reactions. *Chem. Soc. Rev.* **2016**, *45*, 4552–4566.
- (19) Müller, D. S.; Marek, I. Asymmetric copper-catalyzed carbocation of cyclopropenes en route to the formation of diastereo- and enantiomerically enriched polysubstituted cyclopropanes. *J. Am. Chem. Soc.* **2015**, *137*, 49, 15414–15417.
- (20) (a) Hutton, H. M.; Schaefer, T. Proton coupling constants in substituted cyclopropanes. *Can. J. Chem.* **1963**, *41*, 684–689. (b) Lin, H.; Shen, Q.; Lu, L. β -(Trifluoromethyl)vinyl sulfonium salts: preparation and reactions with active methylene and methenyl compounds. *J. Org. Chem.* **2011**, *76*, 7359–7369. (c) Risse, J.; Fernández-Zúmel, M.A.; Cudré, Y.; Severin, K. Synthesis of trifluoromethyl-substituted cyclopropanes via sequential Kharasch-dehalogenation reactions. *Org. Lett.* **2012**, *14*, 3060–3063.
- (21) Müller, P.; Grass, S.; Shahi, S. P.; Bernardinelli, G. Rh (II)-Catalyzed asymmetric carbene transfer with ethyl 3,3,3-trifluoro-2-diazopropionate. *Tetrahedron* **2004**, *60*, 4755–476.
-