



HAL
open science

Sur la gerbe de Kaletha, la courbe et l'ensemble de Kottwitz

Laurent Fargues

► **To cite this version:**

| Laurent Fargues. Sur la gerbe de Kaletha, la courbe et l'ensemble de Kottwitz. 2023. hal-04234891

HAL Id: hal-04234891

<https://hal.science/hal-04234891>

Preprint submitted on 10 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Public Domain

Let $E_n|E$ be the degree n unramified extension and $X_n := X \otimes_E E_n$. Let $X'_n := X \otimes_E E(\pi^{1/n})$. There are two morphisms

$$X_n \rightarrow \mathfrak{X} \text{ and } X'_n \rightarrow \mathfrak{X}.$$

I think the following is true.

Proposition 1. *The category of vector bundles on \mathfrak{X} is equivalent to the category of vector bundles \mathcal{E} on X_n and \mathcal{E}' on X'_n together with an isomorphism of their pullback to $X \otimes_E E_n(\pi^{1/n})$.*

The point would be the following. Consider

$$X_n \times_{\mathfrak{X}} X_n \longrightarrow X_n \times_X X_n = \coprod_{\tau} X_n$$

where $\tau \in \text{Gal}(E_n|E)$. Now one has, with respect to this decomposition,

$$X_n \times_{\mathfrak{X}} X_n = \coprod_{\tau} T_{\tau}$$

where $T_{\tau} \rightarrow X_n$ is the μ_n -torsor of isomorphisms between $\tau^* \mathcal{O}_{X_n}(1)$ and $\mathcal{O}_{X_n}(1)$ as n -roots of $\mathcal{O}_X(1)$. Now, the isomorphism class of this torsor lies in

$$H^1(X_n, \mu_n) = E_n^*/(E_n^*)^n$$

via the identification with $H^1(E_n, \mu_n)$ and Kummer theory.

Lemma 1. *If $\tau = \sigma^i$, the preceding torsor is identified with the torsor of n -roots of π^{-i} .*

Proof. The functor from μ_n -torsors over $\text{Spec}(E_n)$ toward the one over X_n is an equivalence. Via the functors $\varphi\text{-mod}_L \rightarrow \text{Bun}_X$ and $\varphi^n\text{-mod}_L \rightarrow \text{Bun}_{X_n}$, the base change functor $\text{Bun}_X \rightarrow \text{Bun}_{X_n}$ is identified with $(D, \varphi) \mapsto (D, \varphi^n)$. Moreover the identification $\tau^* \mathcal{E}(D, \varphi^n) \xrightarrow{\sim} \mathcal{E}(D, \varphi^n)$ is given by $\varphi^i \otimes 1 : (D, \varphi^n) \otimes_{E_n, \sigma^i} E_n \xrightarrow{\sim} (D, \varphi^n)$.

Now, $\mathcal{O}_{X_n}(1)$ corresponds to the isocrystal $(L, \pi^{-1}\varphi^n)$ together with the identification $(L, \pi^{-1}\varphi^n)^{\otimes n} = (L, (\pi^{-1}\varphi^n)^n)$. Moreover $\tau^* \mathcal{O}_{X_n}(1)$ corresponds to the same but with the preceding identification multiplied by π^{-i} . Thus, our torsor sends the étale E -algebra A to the set of automorphisms of $(L, \pi^{-1}\varphi^n) \otimes_E A$ that respect the preceding identifications. \square

Thus,

$$T_{\tau} = X \otimes_E E_n[T]/(T^n - \pi^{-i}),$$

and in particular, $T_{\sigma} = X \otimes_E E_n(\pi^{1/n})$.

Now, we have to look at

$$X_n \times_{\mathfrak{X}} X_n \times_{\mathfrak{X}} X_n \longrightarrow \coprod_{\tau_1, \tau_2} X_n$$

which, I think, is equal to

$$\coprod_{\tau_1, \tau_2} T_{\tau_1} \times T_{\tau_2}$$

together with morphisms from T_{τ_1, τ_2} toward T_{τ_1} , T_{τ_2} and $T_{\tau_1 \tau_2}$. Try to conclude using this ! Maybe I'm wrong, I don't know, the cocycle condition has to be written carefully.