



HAL
open science

Reclassification of plasmacytoid dendritic cells as innate lymphocytes is premature

Boris Reizis, Juliana Idoyaga, Marc Dalod, Franck Barrat, Shalin Naik,
Giorgio Trinchieri, Roxane Tussiwand, Marina Cella, Marco Colonna

► **To cite this version:**

Boris Reizis, Juliana Idoyaga, Marc Dalod, Franck Barrat, Shalin Naik, et al.. Reclassification of plasmacytoid dendritic cells as innate lymphocytes is premature. *Nature Reviews Immunology*, 2023, 23 (5), pp.336-337. 10.1038/s41577-023-00864-y . hal-04236067

HAL Id: hal-04236067

<https://hal.science/hal-04236067>

Submitted on 12 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted version of the following correspondence published in Nature Immunology

Reizis B, Idoyaga J, Dalod M, Barrat F, Naik S, Trinchieri G, Tussiwand R, Cella M, Colonna M. Reclassification of plasmacytoid dendritic cells as innate lymphocytes is premature. *Nat Rev Immunol.* 2023 May;23(5):336-337. doi: 10.1038/s41577-023-00864-y. PMID: 36959479.

The published version of the manuscript can be accessed on the Journal website:
<https://www.nature.com/articles/s41577-023-00864-y>

Reclassification of plasmacytoid dendritic cells as innate lymphocytes is premature

Boris Reizis^{1†}, Juliana Idoyaga², Marc Dalod³, Franck Barrat⁴, Shalin Naik⁵, Giorgio Trinchieri⁶, Roxane Tussiwand⁷, Marina Cella⁸, Marco Colonna⁸

¹Department of Pathology, New York University Grossman School of Medicine, New York, NY, USA

²Department of Microbiology & Immunology, Stanford University School of Medicine, Stanford, CA, USA

³Aix-Marseille University, CNRS, INSERM, CIML, Centre d'Immunologie de Marseille-Luminy, Turing Center for Living Systems, Marseille, France

⁴HSS Research Institute and David Z. Rosensweig Genomics Research Center, Hospital for Special Surgery, New York, NY, USA

⁵The Walter and Eliza Hall Institute of Medical Research, Melbourne, VIC, Australia

⁶Laboratory of Integrative Cancer Immunology, Center for Cancer Research, National Cancer Institute, NIH, Bethesda, MD, USA

⁷National Institute of Dental and Craniofacial Research, National Institutes of Health, Bethesda, MD, USA

⁸Department of Pathology and Immunology, Washington University School of Medicine, St. Louis, MO, USA

†e-mail: boris.reizis@nyulangone.org

In a recent Comment article, Ziegler-Heitbrock and colleagues¹ challenge the current definition of plasmacytoid dendritic cells (pDCs) as a sublineage of dendritic cells (DCs), and instead propose to reclassify them as innate lymphocytes. That proposal represents an important discussion regarding the complex nature and lineage affiliation of pDCs in view of the emerging new data in the field. At the same time, we believe that the proposed reclassification of pDCs is not warranted by the currently available information, as discussed here.

Ontogeny of pDCs

As correctly noted by Ziegler-Heitbrock et al.¹, the progenitors of pDCs encompass a variety of phenotypes and may indeed comprise cells with predominantly myeloid or lymphoid potentials. We also agree that all approaches to define these progenitors, from adoptive transfers to clonal barcoding, are subject to caveats whose discussion goes beyond the scope of this Comment. Recent studies cited by the authors delineated the development of mouse pDCs from progenitors that are distinct from the progenitors of conventional DCs (cDCs) and that express lymphoid markers such as IL-7 receptor (also known as CD127) and Ly6D. The expression of these markers and several other 'lymphoid' genes indeed distinguishes pDCs from cDCs and suggests that pDCs have similarities with immature lymphocytes, particularly B cells. However, these shared transcriptomic and phenotypic features of pDCs and immature lymphocytes do not necessarily imply their common origin. Instead, the shared features may result from the reliance of pDCs and early B cells on distinct but homologous E protein transcription factors — E2-2 (also known as TCF4) and E2A (also known as TCF3), respectively — which are known to share target genes. Indeed, global gene expression profiling showed a closer proximity of pDCs at steady state to cDCs than to natural killer (NK) cells or B cells, in both mice and humans².

Irrespective of their transcriptional regulation, several considerations regarding the development of pDCs are relevant. First, pDCs and cDCs — and only these cells — can develop from bone marrow progenitors cultured with the growth factor FLT3 ligand (FLT3L). By contrast, FLT3L is not sufficient for commitment to or development of any lymphoid cells. Conversely, the development of pDCs requires neither the common cytokine receptor γ -chain³ nor the transcription factor JAK3 (Ref.⁴), which are both required for the development of all lymphocytes. Second, both pDCs and cDCs are very short lived; thus, their developmental kinetics must, by definition, be different from those of long-lived lymphocytes. Third, pDCs develop normally even when lymphoid progenitors have been depleted, for example by oestrogen treatment⁵. Fourth, as cited by Ziegler-Heitbrock et al.¹, a recent study showed that mature peripheral pDCs and cDCs emerge with similar kinetics and efficiency from progenitors expressing the 'myeloid' marker CX3CR1. Although these studies do not rule out the development of some pDCs from

lymphoid progenitors, they point to a close developmental affiliation of the pDC and cDC lineages.

Functions of pDCs

Ziegler-Heitbrock et al.¹ are correct in outlining that the patterns of pDC migration and recirculation in the steady state are more similar to those of lymphocytes than of cDCs. They also correctly note that steady-state pDCs, unlike cDCs, have little capacity for antigen presentation and T cell priming in the absence of activation. However, activated pDCs undergo transcriptional remodelling towards cDC-like cells⁶. Accordingly, they acquire antigen-presentation capacity, even when focusing on the fate of pDCs that have produced interferon⁷. Importantly, this acquisition of antigen-presentation capacity is observed even after the potentially contaminating pDC-like DCs (also known as transitional DCs, Axl⁺ DCs or pre-DCs) have been removed^{8,9}. Activated pDCs also acquire capacity for cross-presentation¹⁰, which is typically restricted to only a few myeloid cell types such as cDC1s. Notwithstanding the capacity of activated pDCs for T cell priming and cross-priming in vitro, the relevance of these events in vivo remains to be established. Overall, pDCs do not fit the definition of a DC as a cell that is capable of T cell priming in the steady state; that is, indeed, a distinguishing feature of cDCs. Nevertheless, the acquisition of cDC-like features and of cDC-like capacity for antigen presentation and cross-presentation by activated pDCs further highlights their close affiliation with the DC lineage.

pDCs as cytokine-producing cells

We agree with Ziegler-Heitbrock et al.¹ that the capacity for rapid and large-scale production of all subtypes of type I interferon is the most salient, defining property of pDCs. However, one key difference between cytokine production by pDCs and lymphocytes is the nature of the stimulus. Lymphocytes are typically triggered for cytokine production by a surface receptor recognizing antigen (in the case of T cells and B cells), cell-associated molecules (in the case of NK cells) or other cytokines (in the case of innate lymphoid cells). By contrast, pDCs are triggered by innate pattern recognition receptors (PRRs), specifically by endosomal Toll-like receptors (TLRs) for pathogen-derived nucleic acids. This mode of direct pathogen recognition is characteristic of myeloid cells, including

cDCs. For example, cDC1s directly recognize protozoan profilin and viral double-stranded RNA through endosomal TLRs to trigger the production of IL-12 and type III interferon, respectively; these unique cytokine responses are a salient feature of this cell type. Moreover, interferon production by pDCs is important for optimal antiviral T cell responses, which links the direct recognition of a pathogen through TLRs to adaptive immune responses in a manner that is similar to cDCs. Thus, the primary role of pDCs as cytokine-producing cells does not necessarily define them as innate lymphoid cells and is equally compatible with affiliation to the DC lineage.

Conclusions

As highlighted by Ziegler-Heitbrock et al.¹, the lineage identity of pDCs is complex and eludes simple definitions based on either ontogeny or function. Without doubt, pDCs differ from cDCs in their inferior antigen-presentation capacity at steady state, and they resemble lymphocytes in many aspects of their transcriptome and homeostasis. By contrast, pDCs seem to be closely affiliated with the DC lineage in terms of their ontogeny, gene expression programme and ultimate role as a bridge between innate PRR-mediated pathogen recognition and adaptive immunity. In view of this complex and unique nature of pDCs, their radical reclassification as a lymphoid cell type seems to be premature. Future studies may add important new pieces to the puzzle of pDC identity and warrant its further discussion and possible nomenclature changes.

References

1. Ziegler-Heitbrock, L., Ohteki, T., Ginhoux, F., Shortman, K. & Spits, H. Reclassifying plasmacytoid dendritic cells as innate lymphocytes. *Nat. Rev. Immunol.* **23**, 1-2 (2023).
2. Robbins, S. H. et al. Novel insights into the relationships between dendritic cell subsets in human and mouse revealed by genome-wide expression profiling. *Genome Biol.* **9**, R17 (2008).
3. Blasius, A. L., Barchet, W., Cella, M. & Colonna, M. Development and function of murine B220⁺CD11c⁺NK1.1⁺ cells identify them as a subset of NK cells. *J. Exp. Med.* **204**, 2561-2568 (2007).

4. Yamaoka, K., Min, B., Zhou, Y. J., Paul, W. E. & O'Shea J, J. Jak3 negatively regulates dendritic-cell cytokine production and survival. *Blood* **106**, 3227-3233 (2005).
5. Harman, B. C., Miller, J. P., Nikbakht, N., Gerstein, R. & Allman, D. Mouse plasmacytoid dendritic cells derive exclusively from estrogen-resistant myeloid progenitors. *Blood* **108**, 878-885 (2006).
6. Yun, T. J. et al. Human plasmacytoid dendritic cells mount a distinct antiviral response to virus-infected cells. *Sci. Immunol.* **6**, eabc7302 (2021).
7. Abbas, A. et al. The activation trajectory of plasmacytoid dendritic cells in vivo during a viral infection. *Nat. Immunol.* **21**, 983-997 (2020).
8. Alcantara-Hernandez, M. et al. High-dimensional phenotypic mapping of human dendritic cells reveals interindividual variation and tissue specialization. *Immunity* **47**, 1037-1050.e6 (2017).
9. Alculumbre, S. G. et al. Diversification of human plasmacytoid predendritic cells in response to a single stimulus. *Nat. Immunol.* **19**, 63-75 (2018).
10. Oberkampff, M. et al. Mitochondrial reactive oxygen species regulate the induction of CD8⁺ T cells by plasmacytoid dendritic cells. *Nat. Commun.* **9**, 2241 (2018).

Competing interests

The authors declare no competing financial interests.

There is a reply to this letter by Ziegler-Heitbrock, L., Ohteki, T., Ginhoux, F., Shortman, K. & Spits, H. *Nat. Rev. Immunol.* <http://dx.doi.org/10.1038/xxx> (2023).