



The prion paradigm and structural information transfer

Pierre Sibille

► To cite this version:

| Pierre Sibille. The prion paradigm and structural information transfer. Master. UE d'épigénétique, Paris, France. 2020. hal-04236436

HAL Id: hal-04236436

<https://hal.science/hal-04236436v1>

Submitted on 10 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The prion paradigm and structural information transfer: Structural Epigenetic and prion pathologies

How to propagate a biological information without
any support of DNA or RNA??

From a cellular point of view...

There are two transmission types for biological information:

Classical Mendelian transmission:

- Gain of function
- Loss of function
- Long-range transmission

Non-Mendelian transmission:

- Dominant


- Loss/gain of function

Cytoplasmic inheritance

(Cytoduction)

Non-Chromosomal transmission!

Example of cytoduction transmission



Recommended lecture:

Curr Biol. 1994 Aug 1;4(8):744-8.

Cytoplasmic inheritance. Prion-like factors in yeast.

Cox B¹.

What is the support of the information
?


Propagation of information by structural modification: Concept of alternative folding

Simple mechanism mainly based on the modulation of aggregation propensity of a protein :


Aggregation state could generate :

- Gain of function
- Inactivation

The system should be reversible but robust!




Alternative folding pathways




The concept of misfolding and alternative folding


Unfolded
without any specific structure.



Native-folding
conformation of a protein associated to
a physiological function




Normal folding





Physiological function

Abnormal folding



Toxicity



Misfolding
conformation of a protein in which the
physiological function has been lost.
could be associated to a pathology.



Les assemblages de type amyloïde

- Assemblages périodiques, non-covalents de type fibrillaire non-anastomosé
- Structure secondaire très riche en feuillets- β
- Feuillets- β orientés perpendiculairement à l'axe des fibres
- Structures thermodynamiquement très stables

Axe de la fibre



Amyloïde A murin

Amyloïdes pathologiques

Localisation non-neuronale
diabète type II (amylin or IAPP)

Systémique
Amyloïdose AA (SAA)



Amyloïdose ApoAI (ApoAI)

Neurodégénérative
Les EST (PrP)

Alzheimer (A β)

Parkinson (α -synuclein)


...



Amyloïdes fonctionnels

Eléments épigénétiques

Sup35 (Levures)



Nanomatériaux

Fibre de la toile
(*Nephila edulis*)




Facilitation synaptique

ApCPEB (Aplysia)



Biofilm bactérien: Curli (E. Coli)



Relation between protein misfolding and protein aggregation



- Protein folding is dictated by the general laws of thermodynamic.

For $t \rightarrow \infty$ a protein reaches its maximum of stability (i.e. lowest ΔG) through a pathway called “folding pathway”.

The search for this minimum does not correspond to a random exploration of all possible configuration.

- All along the folding pathway there exist different folding intermediates with specific properties.


Amyloidogenic pathway is one of them.



Introduction of prion concept in protein misfolding disease

The nucleation elongation process can be roughly divided in two process:

- A primary nucleation step responsible for the ignition
- A secondary nucleation responsible for elongation



From a biochemical point of view, in the secondary nucleation pathway if $C_{i>1}$ is structurally different from C_1 then process become a Templated Assisted Conversion :

Prion-like propagation

↳ from a molecular point of view $C_{i>1}$ transfer its structural information to C_1



Existence of multiple misfolding pathways

Things are more complicated than they appear!

For aggregation process there are several pathways in competition.

How are these pathways entangled ?

Identification of “relevant” intermediates, etc...






Formation des fibres amyloïdes : mécanisme de type nucléation polymérisation

1 - phase de **nucléation** : phase initiale, création d'un nucleus

peut être court-circuitée par l'ajout d'amorce (oligomères ou protofibrilles)

2 - phase d'elongation (rapide) : déterminée par la concentration en substrat


3 - phase stationnaire déterminée par l'épuisement du substrat




The Prion theory

The prion theory implies the perpetuation of a structural information

1. The existence of a Prion object that constitutes the support of the structural information
2. A non-Prion state that constitutes the “receiver”
3. A segment of interaction between the Prion and the receiver: This segment constitutes a zone through which the structural information transits



The process of prion spreading/amplification by fragmentation



→ The spreading process which is a part of prion propagation theory implicitly needs a fragmentation process to take place


Propagation of information by structural modification: Concept of alternative folding

Simple mechanism mainly based on the modulation of aggregation propensity of a protein :

Aggregation state could generate :

- Gain of function
- Inactivation

The system should be reversible-but robust → Pathology






Prion pathology

- 1 – The concept of protein misfolding and aggregation
- 2 – PrP^C protein : the “good” prion conformer
- 3 – Prion pathology and the “bad” conformer
- 4 – Validation of Prion hypothesis
- 5 - Molecular basis of the Prion strains

Les pathologies liées au mauvais repliement des protéines

- De nombreuses pathologies sont dues au mauvais repliement et à l'agrégation des protéines





- Toute perturbation de l'état natif peut favoriser le processus de conversion

→ Une des conséquences du mauvais repliement des protéines est la formation d'assemblages de type amyloïde

How protein misfolding could induce physiological disorder?

To date what we are sure of is that physiological disorder is the consequence of misfolding
However the nature of misfolded species involved is uncertain!

In terms of physiology the consequence of misfolding could result from either:
Loss of function (neuroprotective role, antioxidanting function...)
Gain of function. Toxicity could be considered as a gain of function





The prion pathology

Proof of transmissibility of the scrapie TSE agent



intraocular inoculation of
spinal cord material



1 year incubation



1st experimental transmission in 1936



Cuillé



Chelle

TSE due to a
slow virus???

kuru: EST acquired human disease



Fore tribe in Papua with endocannibalism rites

- Kuru mainly affects children and women (65%)
- Spongiosis of the brain at necropsy but no inflammation
(excluding infectious disease ?)




Gajdusek suspects that the disease spreads through mortuary cannibal ceremonies. But concludes with a slow neurotropic virus (Medicine Nobel Prize in 1976)

Preuve de transmissibilité des EST humaines



Gadjusek



Il démontre:

- la transmissibilité du kuru et de sCJD
- Que ces 2 maladies provoquent les mêmes lésions cérébrales chez le chimpanzé
- Conclut à un virus lent

Ces travaux lui ont valu le Prix Nobel de Médecine en 1976

The protein only hypothesis

scrapie due to a slow virus???



BUT: In 1967 Alper demonstrated that the agent responsible of TSE is **resistant** to all techniques used in sterilisation



No RNA & DNA!

In 1967



J. S. Griffith

scrapie agent was a **proteinaceous
infectious particle**


=

Prion

A protein that could exist under two conformations: one physiologic and one pathogenic




In 1982



S. Prusiner

The **PrP protein** found in amyloid plaque

Le prion synthétique: dernière preuve irréfutable de l'hypothèse « protein-only »



Legname et al. 2004



Prion pathology

1 – Pathologies due to protein misfolding and aggregation

2 – PrP^C protein : the “good” prion conformer

3 – Prion pathology and the “bad” conformer

4 – Validation of Prion hypothesis

5 - Molecular basis of the Prion strains

The PrP^C structure

Signal peptide

1 MVKSHIGSWI LVLFVAMWSD VGLCKKRPKP GGGWNTGGSR YPGQGSPGGN

51 RYPPQGGGGW GQPHGGGWQ PHGGGWQPHH GGGWGQPHH GGWGQGGSHS


101 QWNKPSKPKT NMKHVAGAAA AGAVVGGLGG YMLGSAMSRP LIHFGNDYED

151 RYYRENMYRY PNQVYYRPVD QYSNQNNFVH DCVNITVKQH TVTTTKGEN


201 FTETDIKIME RVVEQMCITQ YQRESQAYYQ RGASVILFSS PPVILLISFL

251 IFLIVG


↑
GPI



recPrP 3D NMR structure



PrP^C is a membrane anchored glycoprotein




(Rudd, 2001)

The physiology of PrP protein

- PrP^C protein is **an ubiquitous** protein. Mainly expressed in the CNS.
- In mammalian PrP is **highly conserved**
- The PrP knock out mice present not any relevant physiopathology.
- PrP^C is proposed to be involved in metals homeostasis through the octarepeat domain. Transporter, SOD activity, antiapoptotic effect etc...



In one sentence: the physiological role of PrP^C is still unknown !!



Rôle physiologique de la PrP^C ?

- Homéostasie du Cu²⁺
- Activité SOD ?
- Balance Redox

- Développement embryonnaire
- Biologie des cellules souches
- Adhérence cellulaire



(Halliez et al., 2015)

- Activité neuronale
- Neuritogénèse & différentiation
- Transmission synaptique
- Neuroprotection


- Cancers (pronostic -)
- Récepteur du peptide A β



Prion pathology


- 1 – Pathologies due to protein misfolding and aggregation
- 2 – PrP^C protein : the “good” prion conformer
- 3 – Prion pathology and the “bad” conformer
- 4 – Validation of Prion hypothesis
- 5 - Molecular basis of the Prion strains

Les maladies à prion




Prion pathology

- Neurodegenerative disease
- Motor and sensitive dysfunction
- Neuron loss, spongiosis and astrocytosis
- **Accumulation of misfolded PrP**
- Usually a long asymptomatic incubation period
- Fatal issue
- without conventional immune and inflammatory reaction
- Absence of serological response



• **Transmissible/infectious**



The triple determinism of human Prion disease

Sporadic (90 %) : 1-2 case / year / million
Creutzfeldt-Jakob disease (*CJD*)

Infectious

- Kuru : by cannibalism
- variant of Creutzfeldt-Jakob disease (*vCJD*) : ~ 200 case
- iatrogenic (ex. hGH etc.)

Genetic (> 30 mutations in PrP)

- familial CJD
- Gerstmann-Straußler-Scheinker (GSS)
- fatal familial insomnia (IFF)




Maladies à prion : une triple étiologie

SPORADIQUE



— ?

- Mutation somatique ?
- Défaut conformationnel ?
- Facteur non identifié ?




SNC infectieux

> 65 ans

Maladies à prion : une triple étiologie

ACCIDENTELLE



- Kuru (rites anthropophages)
- forme variante de la MCJ
- Hormone de croissance



incubation : 5-50 ans

Maladies à prion : une triple étiologie


GENETIQUE



20-60 ans


- MCJ familiale
- Gerstmann-Straußler-Scheinker (GSS)
- Insomnie fatale familiale (IFF)

Pathogenic mutations




Molecular basis of prion disease

- Prion disease belongs to pathologies due to proteins misassemblies as Alzheimer, and Parkinson etc ...
- It results from autocatalytic misfolding of an ubiquitous protein : PrP^C
- The misfolded conformer of PrP^C is named: PrP^{Sc}
- The evolution of the prion pathology is always associated to the misfolding process



Biochemical characteristic of PrP^{Sc}

Characteristics	PrP ^c	PrP ^{Sc}
% helix α	42 %	30 %
% β sheet	3 %	42 %
PK resistance	-	++
Solubility	++	-
Digestion PIPLC	+++	-



Infected Brain


+ detergents

+ PK


+ Ultracentrifugation

↓


Scrapie associated fibrils (SAF)



(Prusiner et al., 1982)




(Sim et al., 2007)



PrP^{Sc} structural models


a



Modèle en hélice- β senestre

Govaerts et al 2004


b



Modèle β -spirale

DeMarco & Daggett , 2004

c





Modèle en feuillets- β parallèles

Cobb et al 2007



La cryo microscopie électronique au secours des structuralistes

- Agencement parallèle In register des protomères


– 4 protofilaments associés



- Implication du core de la protéine (110-140)
- D'autres structures, avec des différences...




Replication model: *seeding-polymerization* ?




Site(s) de conversion subcellulaires : surface / ERC / Lysosomes ?

Mechanisms of cell death ?




La PrP est essentielle à la réPLICATION des prions et à leur toxicité



(Bueler, Cell 1993, Brandner Nature 1996)



PrP expression is essential for the infectivity



(Bueler, Cell 1993, Brandner Nature 1996)


Des souris exprimant une PrP mutée tombent spontanément malades - D178N (IFF)

- Développement d'une maladie spontanée (400J)
- Symptomatologie évocatrice de l'IFF humaine
- Désordres transmissibles à des souris *wt*



(Jackson et al. Neuron 2009)

Des prions synthétiques authentiquement infectieux



(Legname et al., 2004)



Prion pathology

- 1 – Pathologies due to protein misfolding and aggregation
- 2 – PrP^C protein : the “good” prion conformer
- 3 – Prion pathology and the “bad” conformer
- 4 – Validation of Prion hypothesis
- 5 - Molecular basis of the Prion strains

Historical arguments for prion hypothesis

- Absence of nucleic acids (RNA, DNA) + UV- resistance
- PrP^{Sc} and infectivity co-purified
- Relation between PrP mutation and the genetic form
- Absence of pathology in knock-out PrP mice

Definitive proof: synthetic prion

Science. 2004 Jul 30;305(5684):673-6.

Synthetic mammalian prions.

Legname G¹, Baskakov IV, Nguyen HO, Riesner D, Cohen FE, DeArmond SJ, Prusiner SB.

Résistance des prions aux procédés de décontamination

- **Résistance à la chaleur**

Pas d'inactivation à 160°C / 24h ou 360°C / 1h (chaleur sèche)

3,5-7,5 log réduction à 134-138°C / 3bar / 20 min (chaleur humide)

- **Résistance aux ultrasons**

- **Résistance aux rayonnements ultraviolets**

dose inactivante 37% = 40 000 J/m²

(15 pour le VIH)

- **Résistance aux radiations ionisantes**


dose inactivante 37% = 10 000 grays

(100 pour l'herpès virus, 5000 pour le VIH)

- **Résistance aux procédés chimiques d'inactivation**


Inactivation totale avec la soude 2N, l'eau de Javel 20 000 ppm

Amplification acellulaire des prions par PMCA : *protein misfolding cyclic amplification*




(Saborio et al., 2001)

L'amplification in-vitro des prions par PMCA (protein misfolding cyclic amplification)




L'amplification in-vitro des prions par PMCA (proteic misfolding cyclic amplification)



Comment agit la sonication???

- Fragmentation des fibres?
- Dénaturation de la PrP qui favorise la réplication?






Prion pathology

- 1 – Pathologies due to protein misfolding and aggregation
- 2 – PrP^C protein : the “good” prion conformer
- 3 – Prion pathology and the “bad” conformer
- 4 – Validation of Prion hypothesis
- 5 - Molecular basis of the Prion strains


Prion strains: structural diversity of prion assemblies



In prion diseases, the **pathological information** or **phenotype** is encoded in the structure of PrP^{Sc} assemblies .

→ Existence of strains in prion suggests the existence of a PrP^{Sc} structural diversity

Prion strains and conformational diversity of PrP^{Sc}




The strain information is encoded in the structure of PrP^{Sc} assemblies



Prion replication = transfer of structural information and adaptation/evolution

Evolution of the structural information : Prion evolution and adaptation



	Host A	Host B	Host C
Yellow	●		
Green		●	
Red			■

→ Best replicator selection

→ Question : how does this structural diversity occur?

State of the Art (2000)

- Anatomic Site of Prion replication :
 - CNS, and throughout !!!!
- Cellular Site of Prion replication :
 - Not yet established
- Mechanisms of PrP conversion :
 - Griffiths (1967) and Lansbury (1993) Model
- Molecular basis of Prion Strain :
 - Hypothetic PrPSc structural polymorphisms.
- Mechanisms of Neurotoxicity :
 - An open question
- in vitro neosyntheses of infectivity :
 - Not yet!

State of the Art (2020)

-Anatomic Site of Prion replication :

→ CNS, and throughout !!!!

- Cellular Site of Prion replication :

→ Sometime in cell surface and sometime in the endocytosis root.

-Mechanisms of PrP conversion :

→ Griffiths (1967) and Lansbury (1993) Model

- Molecular basis of Prion Strain :

→ Hypothetic PrPSc structural polymorphisms.

- Mechanisms of Neurotoxicity :


→ open questions, but converged to the mechanisms of other amyloidopathy.

- in vitro neosyntheses of infectivity :

→ Yes.....

structural diversity of prion assemblies within an individual: Prion quasispecies


- Different Biochemical and biological outcome suggest the existence of structural diversity within a given strain
- Introduction of quasispecies concept in prion paradigm (*Li et al. 2010, Science*).



- ➔ Prion adaptation based on the best replicator selection within the quasispecies

Les mécanismes de propagations des prions restent à ce jour inconnus


Le modèle actuel de la propagation des prions de mammifères repose sur l'observation des prions fonctionnels de levures:



Aucune preuve de l'existence de la **fragmentation** chez les prions de **mammifères**


Les fibres de PrP sont constituées d'une sous-unité

AFM (atomic force microscopy)



A. Igel-Egalon et D. Martin

Répétition de motif




Cobb et al. 2009

HET-s amyloid fibril

Contrairement aux prions de levures, les assemblages de mammifères ne sont pas homogènes et présentent des **motifs répétés** le long de la fibre.


Les fibres de PrP sont constituées d'une sous-unité

AFM (atomic force microscopy)



Répétition de motif

Igel-Egalon et al. 2017




La dénaturation de PrP^{Sc} → formation d'un oligomère (trimère) PrP

L'existence de ce motif remet en cause la séquence d'évènement conduisant à l'elongation de la fibre


L'existence d'une sous unité change la vision des prions!

- The existence **multi-scale organization** imposes immediately the question of **sequence of events** during the template assisted **conversion**.
 - We propose two models of templating process:

Shift of equilibrium








Fit-adjustment of monomers



Les fibres de PrP sont constituées d'une sous-unité qui existe en équilibre avec la forme fibrillaire



AFM (atomic force microscopy)

Igel-Egalon et al. 2017






Une nouvelle vision du mode de replication des prions

Igel-Egalon et al. 2017



- Les assemblages de Prion sont des **entités dynamiques** constitués de **sous-unités** exitant en **équilibre** avec la forme fibrillaire.
- La **dépolymérisation**, au lieu de la fragmentation, serait à l'origine de la **réPLICATION** des prions.

Prion structural heterogeneity




Structural heterogeneity of prion assemblies contributes to its adaptation and evolution


Prion host-adaptation and evolution

Questions:

- As a pathogen how prion evolves and adapts to a new host ?



- What is the prion adaptation mechanism?





- The actual dogma is based on replicator dynamic:
 - Best mutator
 - Best response
 - Adaptive dynamic
 - Brown-von Neumann-Nash (BNN) dynamic
 - **Best replicator**

Open questions


- What is the origin of prion structural diversity?
Is it deterministic or stochastic ?

- How structurally distinct assemblies could coevolve within the same environment?
How this diversity is maintained ?

Unusual depolymerization behaviour of Human Prion fibrils: Existence of oscillatory phenomenon





- **Existence of oscillation :**
 - System far from the equilibrium
 - **Multi-compound system with a constitutional exchange**
 - Dissipative process
 - Existence on non-linear process as autocatalysis



Mezache et al. 2019.

Prion replication is a complex process

D.Martin and J.Torrent et al. in progress



→ Prion replication is a dissipative process involving a network of exchange between structural distinct set PrP^{Sc} assemblies