



Scleromatobacter humisilvae gen. nov., sp. nov., a novel bacterium isolated from oak forest soil

Sophie Mieszkin, Blandine Trouche, Julien Ancousture, Youssef Raouf, Stéphane Uroz, Karine Alain

► To cite this version:

Sophie Mieszkin, Blandine Trouche, Julien Ancousture, Youssef Raouf, Stéphane Uroz, et al.. Scleromatobacter humisilvae gen. nov., sp. nov., a novel bacterium isolated from oak forest soil. International Journal of Systematic and Evolutionary Microbiology, 2023, 73 (3), pp.005793. <10.1099/ijsem.0.005793>. <hal-04236566>

HAL Id: hal-04236566

<https://hal.science/hal-04236566v1>

Submitted on 11 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

***Scleromatobacter humisilvae* gen. nov., sp. nov., a novel bacterium isolated from an oak forest soil**

Sophie Mieszkin^{1,2*}, Blandine Trouche¹, Julien Ancousture¹, Youssef Raouf¹, Stéphane Uroz² and Karine Alain¹

Author affiliations :

¹Univ Brest, CNRS, Ifremer, UMR6197 Biologie et Ecologie des Ecosystèmes marins Profonds, F-29280 Plouzané, France

²Centre INRAE-Grand Est-Nancy, Université de Lorraine, INRAE, UMR IAM, 54280 Champenoux, F-54000 Nancy, France

*** Correspondence:** Sophie.Mieszkin@univ-brest.fr

Keywords: *Scleromatobacter humisilvae*; *Betaproteobacteria*; *Burkholderiales*; *Comamonadaceae*; oak forest soil

Repositories:

The 16S rRNA gene sequence and the assembled genome sequences of strain BS-T2-15^T have been deposited in GenBank under the accession numbers OM630150 and JAJLJH000000000, respectively.

Abbreviations:

AAI, Average Amino acids Identity; AF, Alignment Fraction; ANI, Average Nucleotide Identity; CAPSO, N-cyclohexyl-3-aminopropanesulfonic acid; COGs, cluster of

26 orthologous groups; dDDH, Digital DNA-DNA hybridization; DMSO, dimethyl
27 sulfoxide; DPG, diphosphatidylglycerol; DSMZ, Deutsche Sammlung von
28 Mikroorganismen und Zellkulturen; HEPES, 4-(2-hydroxyethyl)-1-
29 piperazineethanesulfonic acid; HOMOPIPES, Homopiperazine-1,4-bis(2-
30 ethanesulfonic acid); HPIC, Hight Pressure Ion Chromatography; ICSP, International
31 Committee on Systematics of Prokaryotes; KEGG, Kyoto Encyclopedia of Genes and
32 Genomes; MES, 2-(N-Morpholino) EthaneSulfonic acid; MMN, Mannitol-Mobility-
33 Nitrate agar medium; OGRI, overall genome related indices; PE,
34 Phosphatidylethanolamine; PG, Phosphatidylglycerol; PHA, Polyhydroxyalkanoates;
35 PIPES, Piperazine-N,N'-bis(2-ethanesulfonic acid); POCP, Percentage Of Conserved
36 Proteins; TEM, Transmission Electron Microscopy; TSA, Tryptic Soy Agar; TSB,
37 Tryptic Soy Broth; UBOCC, UBO Culture Collection.

Abstract

A novel bacterial strain, designated BS-T2-15^T, isolated from forest soil in close proximity to decaying oak wood, was characterized using a polyphasic taxonomic approach. Phylogenetic analyses based on 16S rRNA gene sequences as well as phylogenomic analyses based on coding sequences of 340 concatenated core proteins indicated that strain BS-T2-15^T forms a distinct and robust lineage in the *Rubrivivax-Roseateles-Leptothrix-Azohydromonas-Aquicola-Ideonella* branch of the order *Burkholderiales*. The amino acid identity (AAI) and the percentage of conserved proteins (POCP) between genome of strain BS-T2-15^T and genomes of closely related type strains ranged from 54.27 to 66.57% and from 43.17 to 49.27%, respectively, providing genomic evidence that strain BS-T2-15^T represents a new genus. Its cells stain Gram-negative, are aerobic, motile by a polar flagellum, rod-shaped and form incrustated white to ivory colonies. Optimal growth is observed at 20-22°C, pH 6 and 0% NaCl. The predominant fatty acids of strain BS-T2-15^T are C_{16:1} ω7c, C_{16:0} and C_{14:0} 2-OH. Its polar lipid profile consists of a mixture of phosphatidylethanolamine, diphosphatidylglycerol and phosphatidylglycerol and its main respiratory quinone is ubiquinone 8. The estimated size of its genome is 6.28 Mb with a DNA G+C content of 69.56 mol%. Therefore, on the basis of phenotypic and genotypic properties, the new strain BS-T2-15^T represents a new genus and a novel species for which the name *Scleromatobacter humisilvae* gen. nov., sp. nov., is proposed. The type strain is BS-T2-15^T (DSM 113115^T =UBOCC-M-3373^T).

59 In forest ecosystems, wood decay is an important process that participates to the soil
60 fertility through the action of a complex guild of decomposers [1, 2]. Among them,
61 fungi are considered as the main actors, while the role of bacteria is much less
62 documented [3]. To fill this gap, the taxonomic diversity and the metabolic and
63 functional potential of bacterial strains isolated from a soil-decaying-wood continuum
64 have been investigated [4]. Overall, this study demonstrated a community dominated
65 by representatives of the genera *Paraburkholderia* (*Betaproteobacteria*),
66 *Streptomyces*, *Kitasatospora*, *Arthrobacter* and *Streptacidiphilus* (*Actinobacteria*),
67 *Dyella* (*Gammaproteobacteria*) and by uncharacterized bacterial strains [4]. At the
68 functional level, this study also revealed that soil bacterial communities have the
69 potential to decompose organic matter and a stronger effectiveness than those
70 isolated from decaying oak wood. The taxonomic assignation and functional
71 screening allowed the identification of various strains of interest, among which strain
72 BS-T2-15^T, one of the uncharacterized strains, was selected to be further
73 characterized in the present study to clarify its taxonomic position within the
74 *Rubrivivax-Roseateles-Leptothrix-Azohydromonas-Aquicola-Ideonella* branch of the
75 order *Burkholderiales*, in the class *Betaproteobacteria*. To date, there is a taxonomic
76 uncertainty regarding the classification of these genera at the family rank level.
77 Recently, Liu *et al.* [5] proposed to refer to this branch as the closest-to-
78 *Comamonadaceae* (CTC) group, as it forms a phylogenetically coherent group close
79 to genera belonging to the *Comamonadaceae* family. They proposed to classify the
80 CTC group as a new family, 'Sphaerotilaceae' fam. nov., based on phylogenomic
81 analyses, genome relatedness indices and phenotypic data. However, this new
82 delineation is not yet recognized by the International Committee on Systematics of
83 Prokaryotes (ICSP). In addition, they also proposed several taxonomic revisions at

the genera and species levels. Among remarkable physiological features of the *Rubrivivax-Roseateles-Leptothrix-Azohydromonas-Aquicola-Ideonella* branch, some species are known to be involved in the biodegradation of petroleum compounds or in the reduction of chlorate to chloride under anaerobic conditions [6-8].

In the present study, we performed a polyphasic taxonomic characterization of strain BS-T2-15^T and provided phenotypic, phylogenomic, and genomic evidences that it meets the criteria for delineating a new species of a new genus within the *Rubrivivax-Roseateles-Leptothrix-Azohydromonas-Aquicola-Ideonella* branch. We propose to name this new species *Scleromatobacter humisilvae* gen. nov., sp. nov.

ISOLATION AND ECOLOGY

Strain BS-T2-15^T is from a collection of 308 bacterial strains established as part of a study on the functional abilities of bacterial isolates from decaying wood and their comparison with the properties of bacteria from underlying soil [4]. The experimental set up consisted in oak discs that have been placed during 9 months on the soil surface of the forest experimental site of Champenoux (north-eastern France; Lat: 48.718420° N; Long: 6.346580° E; Alt: 248 m; 0.5 ha of surface). Strain BS-T2-15^T was isolated from the bulk soil in direct contact with decaying oak disc by serial dilutions on 1/10 diluted Tryptic Soy Agar (TSA) medium (Difco's Tryptic Soy Broth (TSB) 3 g.L⁻¹ and agar 15 g.L⁻¹) containing cycloheximide (100 g.L⁻¹, final concentration), with a pH adjusted to 5. The strain was then purified by 3 successive spreads on 1/10 diluted TSA plates at pH 5 to obtain a pure culture. The strain BS-T2-15^T was then grown routinely on 1/10 TSA or 1/10 TSB media adjusted to pH 5 during 3 days at 20°C, under agitation (250 rpm). Its purity was routinely confirmed

by microscopic observations, and by sequencing of its 16S rRNA gene and genome. Stock cultures were stored at -80°C in 1/10 TSB medium supplemented with 5% (v/v) dimethylsulfoxide (DMSO). Main chemical characteristics of the bulk soil samples were $\text{pH } 4.65 \pm 0.10$, $48.38 \pm 7.93 \text{ g.kg}^{-1}$ total carbon, $3.26 \pm 0.49 \text{ g.kg}^{-1}$ total nitrogen, $0.17 \pm 0.01 \text{ g.kg}^{-1}$ phosphorus extracted according to the Duchaufour method and $83.80 \pm 13.92 \text{ g.kg}^{-1}$ organic matter [4].

Strain BS-T2-15^T (DSM 113115^T = UBOCC-M-3373^T) is available in the public culture collections Deutsche Sammlung von Mikroorganismen und Zellkulturen (DSMZ; <https://www.dsmz.de/collection>) and UBO Culture Collection (UBOCC; <https://www.univ-brest.fr/ubocc>).

MORPHOLOGY AND PHYSIOLOGY

Colony morphology of the isolate BS-T2-15^T was observed on 1/10 TSA adjusted to pH 6. Cell morphology and motility were determined by light microscopy (Olympus BX60 and CX40) and transmission electron microscopy (TEM; JEOL JEM 1400). Motility was also observed by using the Mannitol-Motility-Nitrate agar medium (MMN, composed (per liter) of 10 g tryptic hydrolysate of casein, 1 g potassium nitrate, 7.5 g mannitol, 40 mg phenol red and 3.5 g agar), which was also used to evidence mannitol fermentation and nitrate reductase activity. Gram-staining was determined using standard procedures and confirmed with a KOH (3%) test. Catalase and cytochrome oxidase activities were respectively evaluated using H_2O_2 and strips of N, N, N',N'-tetramethyl-*p*-phenylenediamine dihydrochloride (Bio-Rad). The oxidase test was repeated several times using independent cultures to confirm the result. Colonies of BS-T2-15^T are circular and rough with a color between white and ivory. A major phenotypic feature is that its colonies are incrustated in the agar medium. To our

knowledge, this was not yet reported in the literature for genera belonging to the *Rubrivivax-Roseateles-Leptothrix-Azohydromonas-Aquicola-Ideonella* branch. Cells are Gram-negative coccobacilli that divide by binary fission and occur mainly singly, but can also form aggregates (Fig. 1A, B). Cell size range from 0.59 to 0.93 μm wide (mean 0.78 μm ; $n=45$) and from 1.71 to 2.73 μm long (mean 2.15 μm ; $n=45$). Strain BS-T2-15^T is motile, as confirmed by growth observations on MMN agar medium and the observation of a single polar flagellum per cell by TEM (Fig. 1C, D). Refrinent intracellular granules, which could be polyhydroxyalkanoates (PHA) storage granules, were also observed (Fig. 1A). The strain is catalase positive and oxidase negative. The latter phenotypic feature clearly distinguishes strain BS-T2-15^T from other genera of the *Rubrivivax-Roseateles-Leptothrix-Azohydromonas-Aquicola-Ideonella* branch which are all oxidase positive. However, it is important to note that, cytochrome C oxidase complex was identified in its genome. This point is discussed latter in the study.

Physiological characterization of the novel strain BS-T2-15^T was carried out aerobically, in triplicate, on 1/10 TSA or TSB adjusted to pH 6 at 20°C and under agitation (250 rpm). Determination of the temperature range for growth and salt tolerance were respectively tested over the range 0-45°C, at 5°C intervals and 0-10% NaCl (w/v), at 0.5% intervals, both for 10 days on 1/10 TSA at pH 6. Growth was observed from 4 to 30°C with optimal growth between 20-22°C. Concerning salt tolerance, the strain grew only at 0.5% NaCl and exhibited an optimal growth without sodium chloride. The pH range for growth was tested from pH 2.0 to pH 12 (at 20°C), with increments of 1 unit on 1/10 TSB medium for 8 days. Cells were routinely enumerated by direct cell counting using a modified Thoma chamber (Preciss,

France; surface: 0.0025 mm², depth: 10 µm). The following buffers (each at 20 mM, Sigma-Aldrich, Darmstadt, Germany) were used to adjust the required pH: pH 4.0 and 5.0 with HOMOPIPES buffer, pH 6.0 with MES buffer, pH 7.0 with PIPES buffer, pH 8.0 with HEPES buffer, and pH 9.0 and 10.0 with CAPSO buffer. For pH 2.0, 3.0, 11.0 and 12.0, no buffer was used. Growth of strain BS-T2-15^T was observed from pH 2.0 to 12 with optimal growth at pH 6. The growth kinetics of BS-T2-15^T under optimal conditions was then studied in triplicates, on 1/10 TSB adjusted to pH 6 at 22°C with no NaCl salt, under agitation (250 rpm) for 4 days. Cell growth was monitored by direct cell counting, every 3 h using a modified Thoma chamber, in order to determine the growth rate and doubling time of the strain under optimal culture conditions. The growth rate and the generation time of strain BS-T2-15^T are respectively, 0.17 h⁻¹ and 3 h 58 min.

Utilization of carbon sources was investigated using the mineral basis of the 1/10 TSB medium containing (L⁻¹): 0.25 g KH₂PO₄ and adjusted at pH 6. Each substrate (lactose, *D*-mannose, *D*-ribose, *D*-maltose, *D*-glucose, *N*-acetylglucosamine, butyric acid, gluconate, citrate, benzoate, acetate, pyruvate and urea) was supplied at a final concentration of 20 mM. Utilization of nitrogen sources was investigated using 1/10 TSB medium adjusted at pH 6 with nitrate and ammonium supplied at a final concentration of 20 mM, while nitrite was supplied at a final concentration of 5 mM. Before inoculation, cells at the end of their exponential growth phase were harvested and washed three times with distilled water and then the bacterial suspension was adjusted to obtain a Mac Farland index value of 1. A medium composed of mineral basis of the 1/10 TSB without carbon source was used as negative control for each carbon utilization bioassay and a positive control was performed using 1/10 TSB

medium adjusted at pH 6. The hydrolysis of cellulose was determined using Carboxymethyl Cellulose (CM Cellulose) as described by Mieszkina *et al.* [4]. These tests were completed with liquid or solid cultures allowing to determine: (i) mannitol fermentation and presence of nitrate reductase (MMN agar medium), (ii) fermentative pathways (mixed acids or butane-2,3-diol pathways; Clark and Lubs liquid medium), (iii) glucose and lactose fermentation, gas and H₂S production (Kligler-Hajna agar medium), and, (iv) citrate utilization (Simmons citrate agar medium). For these assays, culture media, cell cultures and washes and preparation of bacterial suspensions were performed as described above and by Mieszkina *et al.* [9]. Strain BS-T2-15^T is chemoorganoheterotrophic and grows by aerobic respiration. It catabolizes *D*-glucose, lactose and pyruvate while a moderate growth was obtained with *D*-ribose, *N*-acetylglucosamine, urea and gluconate. A weak growth was also observed with all the nitrogen substrates tested (nitrate, nitrite and ammonium). In addition, BS-T2-15^T is capable of using nitrate or nitrite as terminal electron acceptors showing that the nitrate reductase is functional (Table 1).

Volatile fatty acid production of strain BS-T2-15^T after growth in duplicate in 1/10 TSB was assessed by high-pressure ion chromatography (Dionex ICS-6000 HPIC) (Growth conditions: pH 6, 20°C, under agitation (250 rpm), for 3 days). HPIC analyses showed that the strain consumed all the pyruvate present in the medium during its growth (approximately 930 µM), but did not produce acetate, lactate, pyruvate or oxalate under these culture conditions. This is consistent with previous results showing that it is capable to grow with pyruvate as sole carbon source.

Strain BS-T2-15^T was not capable to use chlorate as sole terminal electron acceptor (at 20 mM), when grown anaerobically on 5 g.L⁻¹ NaCl, with 1 g.L⁻¹ tryptone as carbon and energy source (with respect to negative and positive controls).

Sensitivity to antibiotics of strain BS-T2-15^T was tested by the disc diffusion method after spreading cell suspensions (1 McFarland) on 1/10 TSA adjusted to pH 6 at 20°C for 7 days. Discs were impregnated with antibiotic solutions to obtain a final quantity of antibiotic per disc of: kanamycin (40µg), rifampicin (30 µg), novobiocin (30 µg), gentamicin (15 µg), ampicillin (10 µg), penicillin (10 U), oxacillin (5 µg), oxytetracycline (30 µg), chloramphenicol (30 µg), and streptomycin (10 µg). The discs were then placed on the agar surface. The inhibition zones were read after 3 days of incubation at 20°C. Among the antibiotics tested, strain BS-T2-15^T was sensitive to kanamycin, novobiocin, gentamicin and streptomycin.

CHEMOTAXONOMY

In order to analyze respiratory quinones, polar lipids, and fatty acids, cells of strain BS-T2-15^T were cultured in 1/10 TSB pH 6 at 20°C under agitation (250 rpm) and then harvested by centrifugation (800 g; 10 min) at the end of their exponential phase of growth. These analyses were carried out by the Identification Service of the DSMZ (Braunschweig, Germany) as described by Tindall [10, 11] and Kuykendall *et al.* [12].

The major respiratory quinone identified for strain BS-T2-15^T is ubiquinone 8 (Q-8).

The polar lipid profile of the novel isolate consisted in phosphatidylethanolamine (PE), diphosphatidylglycerol (DPG), and phosphatidylglycerol (PG). A common feature of the new strain and species belonging to the related genera *Leptothrix*, *Rubrivivax*, *Ideonella* and *Aquicola* is the high proportion of PE and PG, which is consistent with the *Comamonadaceae* family [13, 14]. However, a major

characteristic of strain BS-T2-15^T is its high content of DPG, a polar lipid that is absent or only present in low proportions in the type strains of the closest genera (Table S1 et Fig. S1).

The predominant cellular fatty acids (>10% of the total fatty acids) of strain BS-T2-15^T are C_{16:1} ω7c (40.45%), C_{16:0} (25.18%) and C_{14:0} 2-OH (10.49%) (Table 2). Noticeably, the main saturated fatty acid detected (*i.e.*, C_{16:0}) appeared dominant and common among BS-T2-15^T and the four closely related genera (*Leptothrix*, *Rubrivivax*, *Ideonella* and *Aquicola*), while the hydroxyl fatty acid (*i.e.* C_{14:0} 2-OH) was only detected in strain BS-T2-15^T (Table 1). In addition, the presence of the unsaturated fatty acid C_{16:1} ω7c is another major feature of the new strain but could also be represented by summed feature 3 (C16: 1 ω7c o and/or C16: 1 ω6c) in *Rubrivivax gelatinosus* DSM 1709^T and *Ideonella dechloratans* CGUG 30977^T.

16S rRNA GENE PHYLOGENY

A full 16S rRNA gene sequence (1512 nt) of strain BS-T2-15^T was extracted from the genome using the bacterial ribosomal predictor tseemann/barnnap on Galaxy v1.2.1 [19]. Pairwise 16S rRNA gene sequence similarity was calculated using the global alignment algorithm implemented at the EzTaxon-e server (<http://eztaxon-e.ezbiocloud.net/>; [20]). Phylogenetic analyses were performed using the software Seaview version 4.7 for trees reconstruction by the neighbor-joining (NJ) and maximum parsimony (MP) methods [21-23]. The evolutionary distances were respectively calculated using the Kimura two-parameters and the Dnapars modes. The robustness of the inferred topology was assessed by bootstrap analyses based on 1,000 replications. A third tree was build using the maximum likelihood (PhyML) method with the subtree-pruning-regrafting algorithm

(http://phylogeny.lirmm.fr/phylo.cgi/simple_phylogeny.cgi) [24, 25]. The 16S rRNA gene sequence analysis of strain BS-T2-15^T showed that it belongs to the class *Betaproteobacteria*, the order *Burkholderiales* and the family *Comamonadaceae*. The NJ and MP phylogenetic trees based on 16S rRNA gene sequences revealed that strain BS-T2-15^T does not form a monophyletic clade with any representatives of related taxa (Fig. S2 and S3). However, most of the bootstrap percentages of the trees had a very low value (<70%). Concerning the PhyML tree, a clade was obtained with strain BS-T2-15^T and *Leptothrix mobilis* (Feox-1; X97071) but it was distantly related to most of the genera of the *Rubrivivax-Roseateles-Leptothrix-Azohydromonas-Aquicola-Ideonella* branch (Fig. S4).

Comparative analysis of the 16S rRNA gene sequence of the novel strain with relatives having validly published names showed that BS-T2-15^T is equidistant to species belonging to the genera *Ideonella* (96.55–95.78%), *Leptothrix* (96.47–95.50%), *Aquicola* (96.47–96.13%), *Rubrivivax* (96.40–95.85%) and *Aquabacterium* (96.13–95.53%). Its closest relative is *Ideonella azotifigens* 1a22^T (96.55%) followed by *Leptothrix mobilis* FeOx-1^T (96.47%), *Aquicola tertiaricarbonis* L10^T (96.47%), and *I. dechloratans* CCUG 30898^T (96.47%), *Rubrivivax gelatinosus* ATCC 17011^T (96.40%) and *R. benzoatilyticus* JA2^T (96.40%).

Overall, these results demonstrate that the 16S rRNA gene-based phylogeny lacks of resolution to evaluate the taxonomic position of strain BS-T2-15^T among the *Rubrivivax-Roseateles-Leptothrix-Azohydromonas-Aquicola-Ideonella* branch, as previously underlined by Liu *et al.* [25]. Genome based phylogeny is therefore mandatory to gain a better resolution to assign with certainty the taxonomic position of the new strain BS-T2-15^T.

GENOME FEATURES

Genomic DNA of strain BS-T2-15^T was extracted with a standard PCI (Phenol: Chloroform: Isoamyl Alcohol (25:24:1)) protocol [26]. Whole genome sequencing was performed by the Novogene company (Cambridge, United Kingdom), using the Illumina NovaSeq 6000 PE150 technology (2×150 bp paired-end reads). Reads quality was then evaluated with Fast QC (v0.11.9; <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/> [27]). The whole genome was assembled by using the Shovill pipeline with SPAdes and default parameters [28]. Genome completeness and potential contamination were estimated with CheckM on the MicroScope Microbial Genome Annotation and Analysis Platform (MaGe; <https://mage.genoscope.cns.fr>) [29]. The overall genome size was 6,276,266 bp (for 100% completion and 0% contamination) for 28 contigs and the GC content was 69.56%. The N50 and L50 values were respectively 579,168 bp and 3 contigs. A total of 5,712 coding DNA sequences (CDSs), 3 rRNA and 55 tRNA genes (corresponding to the 20 essential amino acids) were detected using the MaGe pipeline (Table 3 and Fig S5). Most of the CDSs (78.29%) could be assigned to at least one cluster of orthologous groups (COGs). COGs categories related to metabolic processes were dominant (38.30% of the CDSs) and the main processes involved (>5% of the CDSs) were: (i) amino acid transport and metabolism (E; 9.04%); (ii) carbohydrate transport and metabolism (G; 6.45%); inorganic ion transport and metabolism (P; 5.45%) and (iv) energy production and conversion (C; 5.38%). Then, 24.56% of CDSs were allocated to the COGs category related to cellular process and signaling and the following processes were dominant: signal transduction mechanisms (T; 7.64%) and cell wall, membrane and envelope

biogenesis (M; 5.26%). Finally, 14.91% of the CDSs were predicted to be involved in the processes dedicated to the information storage and processing (Table S2).

In addition to phylogenetic trees based on the 16S rRNA sequences that do not provide robust phylogenetic reconstructions with this dataset, a phylogenomic tree for was constructed based on 340 core gene clusters (gene clusters appearing once in each genome).

This tree encompassed 21 reference genomes representative of the *Rubrivivax-Roseateles-Leptothrix-Azohydromonas-Aquicola-Ideonella* branch and phylogenetically related to strain BS-T2-15^T (*Aquabacterium commune* DSM 11901^T, *Aquicola rivuli* KYPY4^T, *Aquicola tertiaricarbonis* DSM 18512^T, *Azohydromonas lata* NBRC 102462^T, *Ideonella azotifigens* DSM 21438^T, *Ideonella benzenivorans* B7, *Ideonella dechloratans* CGUG 30977^T, *Ideonella livida* TBM-1^T, *Ideonella paludis* KCTC 32238^T, *Ideonella sakaiensis* 201-F6^T, *Leptothrix cholodnii* SP-6^T, *Leptothrix mobilis* DSM 10617^T, *Roseateles aquatilis* CCUG 48205^T, *Roseateles depolymerans* KCTC 42856^T, *Rubrivivax albus* ICH-3^T, *Rubrivivax benzoatilyticus* JA2^T, *Rubrivivax gelatinosus* DSM 1709^T, *Schlegelella brevitalea* DSM 7029^T, *Sphaerotilus hippei* DSM 566^T, *Sphaerotilus natans* ATCC 13338^T and *Sphaerotilus natans* subsp. *sulfidivorans* D-507^T), the target genome BS-T2-15^T and two closely related Metagenome Assembled Genomes (MAGs; from the Genome Taxonomy Database (GTDB, <https://gtdb.ecogenomic.org/>); accession numbers GB_GCA_013289985.1 and GB_GCA_903845345.1) artificially classified as genus CAIMXF01. These closely related MAGs were respectively recovered from a forest soil [30] and from stratified freshwater lakes and ponds [31-33]. The phylogenomic tree was built as follows: (i) amino acid sequences were extracted, aligned and concatenated using

command line *anvi-get-sequences-for-gene-clusters* on Anvi'o v. 7.1 [34], then (ii) positions with over 0.5 gap frequency were masked, and finally (iii) a maximum likelihood tree was computed using IQTREE (v2.0.3, [35, 36]) under the model WAG with 1,000 ultrafast bootstrap replicates (See SI for more details). The tree was rooted using the genome of *Alcaligenes faecalis* ZD02 (GCA_000967305.2) as an outgroup. A second phylogenomic tree was built by placing BS-T2-15^T in the GTDB tree using GTDB-tk (v2.1, [37]), then subsetting the tree to family *Burkholderiaceae* and genus *Escherichia*, extracting the amino acid alignment, masking positions with over 0.5 gap and computing a maximum likelihood (ML) tree with IQTREE under the model WAG as described above. This second tree was built with 229 MAGs and 60 cultivated bacterial strains from the *Rubrivivax-Roseateles-Leptothrix-Azohydromonas-Aquicola-Ideonella* branch.

The phylogenomic tree built with the 21 reference genomes and the two MAGs closely related to strain BS-T2-15^T, was very robust with high ultrafast bootstrap values, and showed a distinctly deeper branching of strain BS-T2-15^T with its two closely related MAGs and a close proximity to the *Ideonella* clade (Fig. 2). A similar result was also obtained with the phylogenomic tree built with the 229 MAGs (Fig. S6). Therefore, these two phylogenomic trees indicate that strain BS-T2-15^T indeed represents a new genus of the *Comamonadaceae* family within the order *Burkholderiales*, and belongs to the *Rubrivivax-Roseateles-Leptothrix-Azohydromonas-Aquicola-Ideonella* branch.

In addition to the phylogenomic trees, Overall Genome Related Indices (OGRI) were obtained by pairwise calculations between genome of strain BS-T2-15^T, the two closely related MAGs of strain BS-T2-15^T and the 21 reference genomes. The

average nucleotide identity scores (ANI) were obtained using the FastANI calculator tool from the GTDB web server (<https://gtdb.ecogenomic.org/tools/fastani>) [39]. In addition, FastANI calculator provided also the alignment fraction (AF) values between genomes. Digital DNA-DNA hybridization (dDDH) scores were determined by the genome-to-genome distance calculator (GGDC 2.1), using formula 2 [40]. The average amino acid identity (AAI) calculations between genomes were determined as described by Kim *et al.* [41] using the EzAAI pipeline (<http://leb.snu.ac.kr/ezaai>). The percentage of conserved proteins (POCP) was estimated as described by Qin *et al.* [42].

Digital DNA-DNA hybridization and ANI values between the strain BS-T2-15^T genome and the genomes of type strains of the closest genera ranged from 20.00 to 20.50% and from 78.50 to 79.70%, respectively (Table 3). These values are both far below the dDDH threshold level of 70% and the ANI value of 95-96% that are generally accepted for species delineation [43, 44]. Concerning the AAI and POCP values between genome of strain BS-T2-15^T and genomes of the type strains of the closest genera, they ranged, respectively, from 64.27 to 66.57% and from 43.17 to 49.27% (Table 3). AAI values are close to the 65% cutoff value generally accepted for delineating a new genus, while POCP values are below the 50% threshold recognized for demarcating a new genus [42, 45]. Finally, the AF values ranged between 0.40 and 0.48 between genomes of strain BS-T2-15^T and type strains of the closest genera (Table 3), and were observed in conjunction with low ANI values [46], which is not incompatible with the description of a new genus. In this case, POCP and, to a lesser extent, ANI, are the most relevant indices to demonstrate that strain BS-T2-15^T represents a new genus. Similar results were obtained when considering the 21 reference genomes (Fig. S7 and Tab. S3). In addition, all OGRI values

obtained between the BS-T2-15^T strain genome and the genomes of its two closely related MAGs were higher compared to the values obtained against genomes of cultured type strains. These OGRI values relative to the closest MAGs were respectively in the range of 30.10-40.40% and 87.40-88.90% for dDDH and ANI, and in the range of 84.14-86.43%, 70.43-74.76% and 0.70-0.80 for AAI, POCP and AF indices (Tab S4-S8). These dDDH and ANI values are below the threshold accepted for species delineation while the AAI, POCP and AF values are above the threshold for a new genus delineation. Overall, these OGRI analyses clearly show that strain BS-T2-15^T and its two closely related MAGs (i) belong to the same genus, (ii) represent a new genus of the *Rubrivivax-Roseateles-Leptothrix-Azohydromonas-Aquicola-Ideonella* branch and, (iii) belong to three distinct genomic species. Furthermore, values of the different OGRI indices calculated between the species of the genus *Ideonella* and strain BS-T2-15^T and its two closely related MAGs are much lower than those obtained between strain BS-T2-15^T and its two closely related MAGs (Tab S4-S8). Thus, OGRI indices, as well as phylogenomic analyses, lead all to the conclusion that strain BS-T2-15^T represents a new species of a new genus of the *Rubrivivax-Roseateles-Leptothrix-Azohydromonas-Aquicola-Ideonella* branch.

For gene function prediction, genome annotations were performed with the MaGe platform using KEGG and BioCyc databases. The genome of the new strain BS-T2-15^T encodes metabolic pathways for organoheterotrophic growth. Notably, the genome encodes a complete pentose phosphate pathway, and a complete Entner-Doudoroff pathway. The tricarboxylic acid cycle (TCA) pathway for aerobic respiration is almost complete (one enzyme missing). The genome of the novel isolate encodes also the nitrate VIII reduction pathway (dissimilatory) for anaerobic

405 nitrate respiration, which has been shown to be functional experimentally.
406 Degradation pathways for organic compounds such as amino acids (alanine,
407 arginine, asparagine, serine, glutamine, glycine, histidine, cysteine, lysine, ornithine,
408 taurine, threonine and tryptophan), carbohydrates (acetoin, chitin, glucose, glucose-
409 1-phosphate, lactose, ribose and xylose), aromatic compounds (anthranilate,
410 gentisate, phenylacetate and protocatechuate), alcohols (ethanol and glycerol),
411 aldehyde (L-lactaldehyde), amines and polyamines (choline, 4-aminobutyrate,
412 ethanolamine and urea), and carboxylates (D-gluconate, glutaryl CoA, glycolate and
413 glyoxylate) are evidenced. The cytochrome C oxidase complex (EC number: 7.1.1.9;
414 locus tags: COMBST215_v1_20371; 20374; 20375; 20376; 80152; 80156; 80157)
415 was also identified despite that the oxidase activity not being demonstrated
416 experimentally. The cyclopropane fatty acid (CFA) biosynthesis *via* the cyclopropane-
417 fatty-acyl-phospholipid synthase (EC number: 2.1.1.79; locus tag:
418 COMBST215_v1_10589; 50189) and the arginine dependent acid resistance
419 (arginine decarboxylase; EC number: 4.1.1.19; locus tag: COMBST215_v1_10270)
420 pathways are present and may give the strain a competitive advantage in acidic
421 environments such as the forest soil ecosystem. The new strain BS-T2-15^T has also
422 the genetic potential to detoxify arsenate using glutaredoxin (gene: arsC; EC number:
423 1.20.4.1; locus tag: COMBST215_v1_40055) and to degrade superoxide radicals
424 (identification of superoxide dismutase (EC number: 1.15.1.1; locus tags:
425 COMBST215_v1_20566; 40196; 50105) and catalase enzymes (EC number:
426 1.11.1.21; locus tag: COMBST215_v1_11754 and EC: 1.11.1.6; locus tags:
427 COMBST215_v1_60157; 140122)). In comparison with other closely related strains,
428 the new strain BS-T2-15^T is the only one with the genetic potential to achieve the

429 biosynthesis of betaxanthin (a secondary metabolite) and to degrade alanine and
430 anthranilate (Table S9).

431 In conclusion, from the clear genotypic distance with the closest genera, the
432 physiological similarities and some phenotypical and chemotaxonomic differences,
433 we comply with the phylo-phenetic concept that currently prevails for the description
434 of a new genus. Thus, we assigned strain BS-T2-15^T to a novel species, of a novel
435 genus, for which the name *Scleromatobacter humisilvae* gen. nov., sp. nov. is
436 proposed.

437 438 **DESCRIPTION OF *SCLEROMATOBACTER* GEN. NOV.**

439 *Scleromatobacter* sp. (scl.ro'mae. N.L. gen. n. scleromae, of scleroma, a hardened
440 part; derived from the colonies that are incrustated in solid medium when the strain
441 grows in Petri dishes).

442 Cells are Gram-negative, mesophilic, aerobic, chemoorganotrophic, straight rod-
443 shaped (0.6 to 0.9 µm by 1.7 to 2.7 µm in size) and motile with presence of storage
444 granules. Oxidase-negative and catalase-positive. The predominant fatty acids are
445 C_{16:1} ω7c, C_{16:0} and C_{14:0} 2-OH. The polar lipid profile consists of a mixture of
446 phosphatidylethanolamine, diphosphatidylglycerol and phosphatidylglycerol and the
447 main ubiquinone is Q-8. The estimated size of the genome is 6.28 Mb with a DNA
448 G+C content of 69.56 mol%. Phylogenetically, on the base of whole genome
449 comparisons, the genus belongs to the family *Comamonadaceae*, order
450 *Burkholderiales*. Overall, the genus belongs to the *Rubrivivax-Roseateles-Leptothrix-*
451 *Azohydromonas-Aquicola-Ideonella* branch. The type species is *Scleromatobacter*
452 *humisilvae* (BS-T2-15^T).

DESCRIPTION OF *SCLEROMATOBACTER HUMISILVAE* GEN. NOV. SP. NOV.

Scleromatobacter humisilvae (hu.mi.sil'vae. L. fem. n. *humus*, soil; L. fem. n. *silva*, forest; N.L. gen. fem. n. *humisilvae*; from forest soil; from the forest soil where the strain has been isolated).

Displays the following properties in addition to those given in the genus description:

Colonies are circular, rough and incrustated in the agar medium with a color between white and ivory. Optimal growth occurs at 20-22°C, pH 6 and 0% NaCl. Positive for nitrate reduction. Does not respire chlorate. The following carbon sources are used: L-arabinose, D-cellobiose, D-glucose, D-ribose, D-arabitol, adonitol, butyrate, gluconate, pyruvate, L-leucine, L-proline, N-acetylglucosamine, p-hydroxy phenylacetic acid, bromo succinic acid, L-pyroglutamic acid, tween 40 and 80, uridine, putrescine and dextrine. On the contrary, unable to use: citrate, acetate, aspartate, benzoate, formate, methyl-pyruvate, mono-methyl-succinate, cis-aconitic acid, D-galactonic acid lactone, D-galacturonic acid, D-gluconic acid, D-glucosaminic acid, D-glucuronic acid, α -, β -, γ -hydroxy butyric acid, p-hydroxyphenylacetic acid, itaconic acid, α -keto butyric acid, α -keto glutaric acid, α -keto valeric acid, D,L-lactic acid, malonic acid, propionic acid, quinic acid, D-saccharic acid, sebacic acid, succinic acid, succinamic acid, L-aspartic acid, L-glutamic acid, glycyl-L-Aspartic acid, glycyl-L-glutamic acid, L-pyroglutamic acid, γ -amino butyric acid, urocanic acid, phosphate, glucose-1-phosphate, glucose-2-phosphate, D-fructose, L-fucose, D-galactose, D-galactose, gentiobiose, CM-cellulose, D-psicose, D-raffinose, L-rhamnose, sucrose, D-maltose, D-trehalose, turanose, D-mannose, D-melibiose, α -D-lactose, lactulose, maltose, β -methyl-D-glucoside, m-inositol, xylitol, D-mannitol, i-erythritol, D-sorbitol, glycerol, D,L- α -glycerol, 2,3-butanediol, N-acetyl-D-glucosamine, N-acetyl-D-galactosamine, 2-aminoethanol, phenylethylamine,

Glucuronamide, L-alaninamide, D-, L-alanine, L-phenylalanine, L-alanyl-glycine, L-asparagine, L-histidine, L-ornithine, hydroxy-L-proline, D-, L-serine, L-threonine, D,L-carnitine, thymidine, α -cyclodextrin, inosine, glycogen, xylan, lignin and chitin. Produces the α -glucosidase but is negative for urease.

The type strain BS-T2-15^T (DSM 113115^T =UBOCC-M-3373^T) was isolated from bulk soil in direct contact with decaying oak placed for 9 months on the floor of the Champenoux forest experimental site (France). The 16S rRNA gene sequence and the assembled genome sequences of strain BS-T2-15^T have been deposited in GenBank under the accession numbers OM630150 and JAJLJH000000000, respectively.

Acknowledgements

We are grateful to Claire Geslin (UBO, LM2E) and Philippe Eliès (UBO, Plateforme d'imagerie et de mesures en microscopie (PIMM)) for their technical assistance in TEM imaging. We acknowledge Nadège Bienvenu (UBO, LM2E) for her help concerning the deposit of the strain at the UBOCC. We thank Erwann Vince and Xavier Philippon for their technical help concerning ionic chromatography. The LABGeM (CEA/Genoscope & CNRS UMR8030), the France Génomique and French Bioinformatics Institute national infrastructures (funded as part of Investissement d'Avenir program managed by Agence Nationale pour la Recherche, contracts ANR-10-INBS-09 and ANR-11-INBS-0013) are acknowledged for support within the MicroScope annotation platform. The UMR1136 is supported by the ANR through the Labex Arbre (ANR-11-LABX-0002-01).

Funding information

This research was funded by the Sino-French IRP 1211 MicrobSea to K.A. and by the UMR6197.

Conflict of interest

The authors declare that there are no conflicts of interest.

References

1. **Ulyshen MD.** Wood decomposition as influenced by invertebrates. *Biol Rev* 2016; 91:70–85.
2. **Tláskal V, Brabcová V, Větrovský T, Jomura M, López-Mondéjar R, et al.** Complementary roles of wood-inhabiting fungi and bacteria facilitate deadwood decomposition. *Msystems* 2021; 6:e01078–20.
3. **Johnston SR., Boddy L, Weightman AJ.** Bacteria in decomposing wood and their interactions with wood-decay fungi. *FEMS Microbiol. Ecol.* 2016; 92(11).
4. **Mieszkin S, Richet P, Bach C, Lambrot C, Augusto L, et al.** Oak decaying wood harbors taxonomically and functionally different bacterial communities in sapwood and heartwood. *Soil Biol Biochem* 2021; 155:108160.
5. **Liu Y, Du J, Pei T, Du H, Feng GD, et al.** Genome-based taxonomic classification of the closest-to-*Comamonadaceae* group supports a new family *Sphaerotilaceae* fam. nov. and taxonomic revisions. *Syst Appl Microbiol* 2022; 45:126352.
6. **Bedics A, Táncsics A, Tóth E, Banerjee S, Harkai P, et al.** Microaerobic enrichment of benzene-degrading bacteria and description of *Ideonella benzenivorans* sp. nov., capable of degrading benzene, toluene and ethylbenzene under microaerobic conditions. *Antonie von Leeuwenhoek* 2022; 115: 1113-1128.
7. **Lechner U, Brodkorb D, Geyer R, Hause G, Härtig C et al.** *Aquicola tertiaricarbonis* gen. nov., sp. nov., a tertiary butyl moiety-degrading bacterium. *Int J Syst Evol Microbiol* 2007; 57 :1295–1303.
8. **Malmqvist Å, Welander T, Moore E, Ternström A, Molin G et al.** *Ideonella dechloratans* gen. nov., sp. nov., a new bacterium capable of growing anaerobically with chlorate as an electron acceptor. *Syst Appl Microbiol* 1994; 17:58–64.
9. **Mieszkin S, Pouder E, Uroz S, Simon-Colin C, Alain K.** *Acidisoma silvae* sp. nov. and *Acidisoma cellulosilytica* sp. nov., two acidophilic bacteria isolated from decaying wood, hydrolyzing cellulose and producing poly-3-hydroxybutyrate. *Microorganisms* 2021; 9:2053.
10. **Tindall BJ.** A comparative study of the lipid composition of *Halobacterium saccharovorum* from various sources. *Syst Appl Microbiol* 1990;13:128–130.
11. **Tindall BJ.** Lipid composition of *Halobacterium lacus profundus*. *FEMS Microbiol Lett* 1990; 66: 199–202.

12. **Kuykendall LD, Roy MA, O'Neill JJ, Devine TE.** Fatty acids, antibiotic resistance, and deoxyribonucleic acid homology groups of *Bradyrhizobium japonicum*. *Int J Syst Bact* 1988; 38:358–361.
13. **Jeon CO., Park W, Giorse WC, Madsen EL.** *Polaromonas naphthalenivorans* sp. nov., a naphthalene-degrading bacterium from naphthalene-contaminated sediment. *Int J Syst Evol Microbiol* 2004; 54:93–97.
14. **Blümel S, Busse H J, Stolz A, Kämpfer P.** *Xenophilus azovorans* gen. nov., sp. nov., a soil bacterium that is able to degrade azo dyes of the Orange II type. *Int J Syst Evol Microbiol* 2001; 51:1831–1837.
15. **Chen WM, Chen LC, Sheu DS, Tsai JM, Sheu SY.** *Ideonella livida* sp. nov., isolated from a freshwater lake. *Int J of Syst Evol Microbiol* 2020;70:4942–4950.
16. **Sheu SY, Hsieh TY, Chen WM.** *Aquicola rivuli* sp. nov., isolated from a freshwater stream. *Int J Syst Evol Microbiol* 2019; 69:2226–2232.
17. **Spring S, Kämpfer P, Ludwig W, Schleifer KH.** Polyphasic characterization of the genus *Leptothrix*: new descriptions of *Leptothrix mobilis* sp. nov. and *Leptothrix discophora* sp. nov. nom. rev. and emended description of *Leptothrix cholodnii* emend. *Syst Appl Microbiol* 1996; 19:634–643.
18. **Willems A, Gillis M, De Ley J.** Transfer of *Rhodocyclus gelatinosus* to *Rubrivivax gelatinosus* gen. nov., comb. nov., and phylogenetic relationships with *Leptothrix*, *Sphaerotilus natans*, *Pseudomonas saccharophila*, and *Alcaligenes latus*. *Int J Syst Evol Microbiol* 1991; 41:65–73.
19. **Seemann T, Booth T.** BARNAP: BASic Rapid Ribosomal RNA Predictor [Internet]. 2020; Berlin: GitHub; 2013. p.
20. **Yoon SH, Ha SM, Kwon S, Lim J, Kim Y.** Introducing EzBioCloud: a taxonomically united database of 16S rRNA gene sequences and whole-genome assemblies. *Int J Syst Evol Microbiol* 2017; 67:1613.
21. **Gouy M, Guindon S, Gascuel O.** SeaView version 4: A multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Mol Biol Evol* 2010; 27:221–224
22. **Saitou N, Nei M.** The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol Biol Evol* 1987; 4:406–425.
23. **Kimura M.** The Neutral Theory of Molecular Evolution. Cambridge: Cambridge University Press; 1983.
24. **Dereeper A, Guignon V, Blanc G, Audic S, Buffet S, et al.** Phylogeny. fr: robust phylogenetic analysis for the non-specialist. *Nucleic Acids Res* 2008; 36: W465–W469.
25. **Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, et al.** New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst Biol* 2010; 59:307–321.
26. **Charbonnier F, Forterre P, Erauso G, Prieur D.** Purification of plasmids from thermophilic and hyperthermophilic archaeobacteria. In *Archaea: A Laboratory Manual*; Robb, F.T., Place, A.R., DasSarma, S., Schreier, H.J., Fleischmann, E.M., Eds.; Cold Spring Harbor Laboratory Press: Woodbury, NY, USA, 1995; pp. 87–90.
27. **Andrews, S.** FastQC: A quality control tool for high throughput sequence data. 2010.
28. **Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M.** SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 2012; 19:455–477.

29. **Vallenet D, Calteau A, Dubois M, Amours P, Bazin, A. et al.** MicroScope: an integrated platform for the annotation and exploration of microbial gene functions through genomic, pangenomic and metabolic comparative analysis. *Nucleic Acids Res* 2020; 48(D1):D579-D589.
30. **Alteio LV, Schulz F, Seshadri R, Varghese N, Rodriguez-Reillo W, et al.** Complementary metagenomic approaches improve reconstruction of microbial diversity in a forest soil. *msystems* 2020; 5:e00768-19.
31. **Martin G, Rissanen AJ, Garcia SL, Mehrshad M, Buck M, et al.** Candidatus *Methylumidiphilus* drives peaks in methanotrophic relative abundance in stratified lakes and ponds across Northern landscapes. *Front Microbiol* 2021; 12:669937.
32. **Garcia SL, Mehrshad M, Buck M, Tsuji JM, Neufeld JD, et al.** Freshwater Chlorobia exhibit metabolic specialization among cosmopolitan and endemic populations. *msystems* 2021; 6:e01196-20.
33. **Rissanen AJ, Saarela T, Jäntti H, Buck M, Peura S, et al.** Vertical stratification patterns of methanotrophs and their genetic controllers in water columns of oxygen-stratified boreal lakes. *FEMS Microbiol Ecol* 2021; 97:fiaa252.
34. **Eren AM, Kiefl E, Shaiber A, Veseli I, Miller S et al.** Community-led, integrated, reproducible multi-omics with anvi'o. *Nat Microbiol* 2021; 6:3–6.
35. **Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS.** UFBoot2: improving the ultrafast bootstrap approximation. *Mol Biol Evol* 2018; 35:518–522.
36. **Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, et al.** IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic era. *Mol Biol Evol* 2020; 37:1530–1534.
37. **Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH.** GTDB-Tk: a toolkit to classify genomes with the Genome Taxonomy Database. *Bioinform* 2020; 6:1925-1927.
38. **Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH, et al.** GTDB-Tk v2: memory friendly classification with the Genome Taxonomy Database. *bioRxiv* 2022.
39. **Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S.** High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun* 2018; 9:1-8.
40. **Meier-Kolthoff JP, Auch AF, Klenk HP, Göker M.** Genome sequence-based species delimitation with confidence intervals and improved distance functions. *BMC Bioinform* 2013; 14:60.
41. **Kim D, Park S, Chun J.** Introducing EzAAI: a pipeline for high throughput calculations of prokaryotic average amino acid identity. *J Microbiol* 2021; 59:476-480.
42. **Qin QL, Xie BB, Zhang XY, Chen XL, Zhou BC, et al.** A proposed genus boundary for the prokaryotes based on genomic insights. *J Bacteriol* 2014; 196:2210-2215.
43. **Richter M, Rosselló-Móra R.** Shifting the genomic gold standard for the prokaryotic species definition. *Proc Natl Acad Sci USA* 2009; 106:19126–19131.
44. **Wayne L, Brenner DJ, Colwell RR, Grimont PAD, Kandler O et al.** Report of the Ad Hoc Committee on Reconciliation of Approaches to Bacterial Systematics. *Int J Syst Bacteriol* 1987; 37:463–464.
45. **Konstantinidis KT, Rosselló-Móra R, Amann R.** Uncultivated microbes in need of their own taxonomy. *ISME J* 2017; 11:2399-2406.

643 46. **Barco RA, Garrity GM, Scott JJ, Amend JP, Nealson KH et al.** A genus
644 definition for bacteria and archaea based on a standard genome relatedness
645 index. *mBio* 2020; 11:e02475–19.
646

List of figures:




Figure 1. Transmission electron micrographs of cells of strain BS-T2-15^T showing (A) intracellular granules (indicated by black arrows), (B) a dividing cell (by binary fission), (C) a cell with its polar flagellum, and (D) the insertion point of the flagellum.




Fig. 2. Phylogenomic tree (concatenated alignment of 340 shared gene clusters) showing the phylogenomic position of strain BS-T2-15^T, its two closely related MAGs and representatives of other related taxa. The tree was built using IQ-Tree and the maximum likelihood method with 1,000 replicates for the calculation of the ultrafast bootstrap. Bootstrap values >95% are indicated at branching point by a dot. The tree was rooted using *Alcaligenes faecalis* ZD02 (GCA_000967305.2) as the outgroup. Bar, 0.04 substitutions per position.

List of tables:

Table 1. Differential characteristics of strain BS-T2-15^T and type strains of closely related genera.

Taxa : 1. BS-T2-15^T, 2. *Leptothrix mobilis* DSM 10617^T, 3. *Rubrivivax gelatinosus* DSM 1709^T, 4. *Ideonella dechloratans* CGUG 30977^T, 5. *Aquicola tertiaricarbonis* DSM 18512^T.

Data are from Mieszkina *et al.* [4], Lechner *et al.* [7], Malmqvist *et al.* [8], Chen *et al.* [15], Sheu *et al.* [16], Spring *et al.* [17] and Willems *et al.* [18]. +, Positive; –, negative; NA, not available; w, weak growth; ^aSummed feature 3 comprises C16: 1 ω 7c and/or C16: 1 ω 6c. All strains are motile, positive for tween 40 hydrolysis, and negative for urease activity and citrate utilization. [§]As all strains have not been grown under exactly the same conditions nor in the same media, percentages of fatty acids cannot be compared from one strain to another.

Characteristics	1	2	3	4	5
Colony color	White-ivory	Dark-brown	Orange-brown	White	White
Cell size (μm)	0.6-0.9 x 1.7-2.7	0.6-0.8 x 1.5-12	0.4-0.7 x 1-3	0.7-1 x 2.5-5	0.8-1.1 x 1.2-2.3
Temperature range (optimum) (°C)	4-30 (20-22)	10-37	10-45 (37-40)	15-30 (30)	4-40 (30)
pH growth range (optimum)	2-12 (6)	6.5-8.5	5-9 (6-7)	6-8 (6-7)	5-9 (6-7)
NaCl tolerance (optimum) (% w/v)	0-0.5 (0)	NA	0-2 (1)	0-2	0-1 (0)
Carbon sources utilization					
Acetate	–	–	+	+	+
Butyrate	+	–	+	+	+
Gluconate	W	–	NA	–	+
Pyruvate	+	–	+	+	+
Arabinose	+	NA	–	–	NA
Cellobiose	+	NA	–	–	NA
CM-cellulose	–	NA	–	+	–
D-Glucose	+	–	–	+	+
D-Maltose	–	–	+	–	+
D-Mannose	–	–	–	+	+
D-Mannitol	–	+/-	–	–	+
N-Acetylglucosamine	W	–	–	–	+
Indole formation	–	–	+	–	–
Enzymatic activities					
Catalase	+	NA	–	+	+
Oxidase	–	+	+	+	+
Nitrate reductase	+	NA	+	+	–
α -Glucosidase	+	NA	–	–	+
Quinone type	Q-8	Q-8	Q-8, MK-8	Q-8	Q-8
Polar lipids					
Diphosphatidylglycerol (DPG)	+	–	+	+	+
Major fatty acids (%)[§]					
14: 0 2-OH	10.49	–	–	–	–
16: 1 <i>cis</i> -9	–	53.04	–	–	39
16: 1 ω 7c	40.45	–	–	–	–
Summed feature 3 ^a	–	–	42.8	40.2	–
DNA G+C content (mol%)	69.6	68	71.9	68.1	70.5
Isolation source	Forest	Freshwater	Acetate	Activated	Aquifer

	soil	sediment	enrichment	sludge
--	------	----------	------------	--------

Table 2. Comparison of whole-cell fatty acid profiles (% of the total) of strain BS-T2-15^T with type strains of closely related genera.

Taxa : 1. BS-T2-15^T, 2. *Leptothrix mobilis* DSM 10617^T, 3. *Rubrivivax gelatinosus* DSM 1709^T, 4. *Ideonella dechloratans* CGUG 30977^T, 5. *Aquicola tertiarycarbonis* DSM 18512^T. Values are percentages of the fatty acids that were assigned to fatty acids in the peak-naming table of the MIS database (MIDI, Microbial ID, Newark, DE 19711 U.S.A.). The nomenclature is as follows: the first number indicates the number of carbon atoms in the molecule; 'OH' and 'cyclo' indicate hydroxy or cyclic fatty acids; the second number following the colon indicates the number of double bonds present. The position of the double bond is indicated by the carbon atom position starting from the methyl (ω) end of the molecule. *c*, cis isomer. Data are from Lechner *et al.* [7], Malmqvist *et al.* [8], Sheu *et al.* [16] and Spring *et al.* [18]. Major fatty acids (>10% of the total fatty acids) are indicated in bold.

Fatty acids	1	2	3	4	5
10: 0 3-OH	5.99	4.54	4.8	2.4	2
12: 0	—	2.43	3.7	2.1	4
12: 0 2-OH	0.47	—	—	2.5	—
12: 0 3-OH	3.17	—	—	4.1	—
14: 0	3.14	0.77	5.4	1.5	2
14: 0 2-OH	10.49	—	—	—	—
15: 0	—	—	—	—	3
15: 1	—	—	—	—	2
15: 1 ω 6 <i>c</i>	—	—	1.7	—	—
16: 0	25.18	30.29	33.1	31.5	37
16: 1 <i>cis</i> -9	—	53.04	—	—	39
16: 1 ω 7 <i>c</i>	40.45	—	—	—	—
17: 0	—	—	1.1	—	2
17: 0 cyclo	—	—	—	—	2
17: 0 cyclo ω 7 <i>c</i>	1.44	—	—	—	—
18: 0	—	1.19	—	—	1
18: 1	—	—	—	—	6
18: 1 <i>cis</i> -9,11 ^a	—	6.99	—	—	—
18: 1 ω 7 <i>c</i>	9.68	—	4.5	12.7	—
19: 1 <i>cis</i> -9,10 ^b	—	0.84	—	—	—
Summed feature	—	—	42.8	40.2	—

^{3c}

—, Not detected

^aSum of both compounds: *cis*-9octadecenoic acid and *cis*-11-octadecenoic acid

^b*cis*-9,10-methyleneoctadecanoic acid

^cSummed feature 3 comprises C16: 1 ω 7*c* o and/or C16: 1 ω 6*c*

Table 3. Genome statistics and overall genome relatedness indices (OGRI) between strain BS-T2-15^T and type strains of closely related genera and closely related MAGs.

Taxa : 1. BS-T2-15^T, 2. *Leptothrix mobilis* DSM 10617^T, 3. *Rubrivivax gelatinosus* DSM 1709^T, 4. *Ideonella dechloratans* CGUG 30977^T, 5. *Aquicola tertiarycarbonis* DSM 18512^T. 6. *Azohydromonas lata* NBRC 102462^T, 7. *Roseateles depolymerans* KCTC 42856^T, 8. MAG GCA_013289985.1, 9. MAG_GCA_903845345.1.

	1	2	3	4	5	6	7	8	9
Number of contigs	28	17	36	158	98	342	1	210	631
Size (Mb)	6.27	4.65	5.08	4.51	6.32	7.18	5.68	6	5.4
GC content (%)	69.56	69.00	71.40	69.30	70.20	69.00	66.60	69.55	69.50
Number of CDS	5,712	4,019	4,691	4,172	5,711	6,349	4,894	5,441	5,362
rRNA	3	6	3	7	3	4	12	3	1
tRNA	55	49	46	59	43	46	58	19	17
dDDH (%)	100	20.50	20.40	20.10	20.30	20.30	20.00	30.10	33.40
ANI (%)	100	79.50	79.70	79.70	79.60	78.60	78.50	87.80	88.90
AAI (%)	100	65.34	66.03	66.57	65.48	65.32	64.27	84.14	86.43
POCP (%)	100	43.17	44.84	47.33	49.02	40.89	49.27	72.60	74.76
AF	1.0	0.42	0.48	0.43	0.40	0.32	0.29	0.71	0.80