

Altered ovarian transcriptome is linked to early mortality and abnormalities in zebrafish embryos after maternal exposure to gamma irradiation

Noëmie Guirandy, Olivier Armant, Sandrine Frelon, Fabien Pierron, Benjamin Geffroy, Guillemine Daffe, Camille Houdelet, Patrice Gonzalez, Olivier Simon

▶ To cite this version:

Noëmie Guirandy, Olivier Armant, Sandrine Frelon, Fabien Pierron, Benjamin Geffroy, et al.. Altered ovarian transcriptome is linked to early mortality and abnormalities in zebrafish embryos after maternal exposure to gamma irradiation. Aquatic Toxicology, 2023, 262, pp.106660. 10.1016/j.aquatox.2023.106660. hal-04237612

HAL Id: hal-04237612 https://hal.science/hal-04237612

Submitted on 11 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

1 Abstract

Recent laboratory studies focusing on multigenerational approach demonstrated 2 3 drastic phenotypic effects after chronic fish irradiation exposure. No irradiation effect at phenotypic scale was observed for F0 (reproductive performances) while early 4 mortality and malformations were observed in F1 offspring whether they were 5 irradiated or not. The objective was to study molecular mechanisms likely to be 6 involved in these phenotypic effects induced by parental irradiation. Thus, F0 adult 7 zebrafish were irradiated for ten days until reproduction and maternal involvement in 8 offspring development was assessed. Levels of maternal provided cortisol and 9 vitellogenin, needed for embryo development, were not impacted by irradiation. 10 11 However, maternal transcriptome highlighted irradiation effect on processes involved in oocyte development, as well as on essential maternal factors needed for offspring 12 development. Therefore, this study highlighted the importance of parental exposure on 13 offspring fate and of the importance of multigenerational exposure in risk assessment. 14

Keywords: multigenerational, maternal transcripts, cortisol, vitellogenin, zebrafish,irradiation

- 17
- 18
- 19
- 20
- 21
- 22
- 23

24

25

Altered ovarian transcriptome is linked to early mortality and abnormalities in zebrafish embryos after maternal exposure to gamma irradiation

29

	-			-		_			
30	Guirandy	Noëmie ^{a*} ,	Armant	Olivier ^a ,	Frelon	Sandrine ^a ,	Pierron	Fabien ^b ,	Geffroy

- ³¹ Benjamin^c, Daffe Guillemine^b, Houdelet Camille^c, Gonzalez Patrice^b, Simon Olivier^a
- ³² ^a IRSN/PSE-ENV/SRTE/LECO, Centre de Cadarache-B.P. 3 Bat 183 13115 St
- 33 Paul Lez Durance, France
- ^b Univ. Bordeaux, CNRS, EPOC, EPHE, UMR 5805, F-33600 Pessac, France
- ³⁶ ^cMARBEC, Univ Montpellier, CNRS, Ifremer, IRD, Palavas-Les-Flots, France
- ^{*}Corresponding author.
- 38 E-mail addresses: noemie.guirandy@hotmail.fr
- 39

35

40 **1. Introduction**

As aquatic ecosystems are particularly impacted by chemical pollutants, many data are available on their phenotypic effects and toxic mechanisms of action. Recently, multigenerational consequences of chemical exposure is a growing area of interest for environmental risk assessment (ERA). Although nuclear accidents such as Chernobyl and Fukushima represent an ecological threat for wildlife populations, much less is known regarding the effects of ionizing radiations (IR) over generations (Beresford et al., 2020; Car et al., 2022; Cunningham et al., 2021; Geras'kin et al., 2021).

We previously showed that laboratory exposure of adult zebrafish to high dose rate (50 mGy.h-1, 12 Gy) for 10 days did not alter their reproductive performances nor survival (Guirandy et al. 2019). But it is likely that effects would be more tenuous since

DNA damages and oxidative stress were already detected following IRs exposure 51 (Simon et al., 2011). Moreover, these subtle effects could be maternally transmitted, 52 with potential consequences on offspring fitness. We indeed reported that the 53 irradiated or non-irradiated (recovery condition) offspring from irradiated genitors at a 54 high dose rate showed a high mortality from 24 hours post fertilization (hpf) (Guirandy 55 et al., 2019). Similar results were observed by Hurem et al. 2017 after an exposition to 56 53 mGy.h-1, which induced non-viable offspring. Another study at a lower dose rate (5 57 mGy.h-1, 30d, 3.6 Gy) showed an impact of IR on the irradiated or non-irradiated 58 offspring leading to high mortality rates at early stages of development (8 days post 59 60 fertilization (dpf)) as well as male-biased sex ratio at adulthood (Guirandy et al., 2022). All these studies highlighted a high mortality of the offspring from irradiated parents. 61 The mechanisms of parental effect on offspring still remain unclear. 62

Strong maternal effects are likely to occur following exposure to IRs since mothers not 63 only provide genomic material but also many essential cytoplasmic components and 64 65 mRNA needed to achieve the proper embryonic development of the offspring. First, females contain the future eggs, on which IR may induce oxidative stress and DNA 66 damages (Guirandy et al., 2019; Hurem et al., 2018, 2017). Damaged eggs would then 67 suffer higher mortality, as observed by Chen et al. 2015 and Vuori et al. 2008. 68 Secondly, some essential macro-molecules are provided by female and are needed 69 for embryo development ("Maternal Effect Genes in Development, Volume 140 - 1st 70 Edition," n.d.). Two macro-molecules are of particular interest, namely vitellogenin 71 (VTG) which needs to be cleaved by cathepsin D in order to form the yolk egg (Finn, 72 73 2007; Yilmaz et al., 2018); and cortisol (the main stress hormone in fish (Sadoul and Geffroy, 2019)). VTG is the yolk egg precursor and the developing offspring of 74 oviparous animals are entirely dependent on this energy store. A defect in VTG 75

quantity or in its metabolism is known to be a cause of early mortality (Soares et al., 76 77 2009; Yilmaz et al., 2018). Altered cortisol concentration could also play a key role in the described effects since a proper maternally-provided quantity of cortisol is needed 78 for early embryo development (Faught and Vijayan, 2018; McCormick, 1999; Nesan 79 and Vijayan, 2016, 2012). Finally, maternal mRNAs support embryonic development 80 until activation of the zygote genome (Abrams and Mullins, 2009; "Maternal Effect 81 Genes in Development, Volume 140 - 1st Edition," n.d.). Indeed, good quality eggs 82 production begins with oocyte development in females. For instance, many studies 83 highlighted the primordial role of maternal mRNAs and more generally of the Balbiani 84 85 body (Bb) in embryo axis establishment and progeny survival (Abrams and Mullins, 2009; Flores et al., 2008; Langdon and Mullins, 2011). Particularly, formation of the Bb 86 in developing oocytes allows establishment of the first polarity (animal/vegetal) of 87 88 embryo (Jamieson-Lucy and Mullins, 2019; Marlow, 2010). Two maternal genes are particularly involved in Bb formation; (i) buc, which allows Bb assembly and (ii) macf1, 89 which allows Bb translocation and dispersal in oocyte. Maternal mRNA needed later 90 for embryo development are accumulated in the Bb before to be dispersed to their 91 functional localization. 92

Here, we assessed maternally transmitted effects following parental exposure to 93 gamma irradiation with no offspring irradiation (recovery) as well as irradiation of both 94 genitors and offspring (irradiated). Maternal transcriptome was then analysed to get 95 mechanistic insights into the impact of irradiation. This study represents a first step to 96 define parental effects after irradiation, by assessing mother involvement. Hence, after 97 98 a gamma irradiation performed at 50 mGy.h-1 for 10 days, we sampled ovaries and eggs to study irradiation-induced impacts (i) on maternally deposited material through 99 cortisol and VTG measurements, (ii) on oocyte development and (iii) on specific 100

101 maternal transcripts needed for offspring development by analysing ovary 102 transcriptome (RNA-Seq) in order to get insights into the mechanisms that could 103 explain the transmitted effects from female to offspring. 50 mGy.h-1 dose rate was 104 used to underspin effects observed in our previous study and to assess a mechanistic 105 study at high dose rate not observed in the environment.

106 2. Materials and methods

107 **2.1. Fish husbandry**

The project (APAFIS#15821) was authorized by the Institut de Radioprotection et de 108 Surêté Nucléaire ethics committee no. 81 (EU 0520) in an application under the 109 directive 2010/63/UE relating to animal care. The study was conducted on AB strain 110 zebrafish that were kept, reproduced, and irradiated in a zebrafish housing system 111 (Zebtec Tecniplast Stand Alone) with recirculating oxygenated freshwater. Adult fish 112 113 were acclimatized over 3 weeks to tap water + 20% demineralized water renewed daily (Aquadem; pH 7.4 \pm 0.2, conductivity = 398 \pm 2 μ S cm⁻¹, temperature = 28.4 \pm 0.3 °C), 114 with a 12:12-h light:dark cycle photoperiod. The fish were fed ab libitum three times a 115 day with GEMMA Wean (Skretting®). 116

117 **2.2. Adult and embryo exposure**

Nominal dose rate was 50 mGy.h⁻¹ for the irradiated condition. Gamma-rays were 118 emitted from a 137Cs source (444 GBq, 662 keV; Institut de Radioprotection et de 119 Surêté Nucléaire MICADO-Lab platform). Dose rate was simulated using MCNP5 120 thermoluminescent software and measured using dosimeters 121 (Chiyoada Technologies), and the values represented between 80 and 120% of the nominal 122 values. Control condition was kept in a separate room (60–80 nGy.h⁻¹). The population 123 density of adult fish was 0.7 L.g⁻¹. 124

Twenty-four adult fish per condition (12 females : 12 males) were irradiated over 10 d.
Daily controls and feeding were conducted as described in Guirandy et al., 2019.

F1 offspring were obtained from 12 F0 spawning couples (24 fish, 1 female:1 male) 127 128 (i.e., replicates) per condition. Mating and eggs viability were determined as described in Guirandy et al., 2019. The 3hpf-embryos from 6 spawns per condition were 129 separated into 2 groups of 30 eggs per clutch. The first one F1 embryos were then 130 positioned in an incubator and irradiated at 50 mGy.h⁻¹ (irradiated) (Figure 1). The 131 second group was placed in non-irradiated incubator (recovery). F1 embryos were 132 maintained in both conditions over 6days. The same culture medium was used for 133 embryos and adults and 10% was replenished daily. 134

F0 adults were then placed in control condition and reproduction was done as explained before, at 6 days (resilience_6d) and 36 days (resilience_36d). Each F1 obtained offspring were kept in control conditions for 4 days.

138 **2.3. Ecological endpoints and morphological analysis**

Reproductive performances (reproductive success, fecundity: number of eggs laid per
female) of adults were measured after 10 days of exposure.

The progeny survival rate was assessed from 24 to 96 hours post fertilization (hpf) in the control, irradiated, and non-irradiated (recovery) conditions (30 eggs from 6 spawns per condition).

Photographs of F1 larvae were recorded at 72hpf (6 to 30 larvae per condition) under a stereomicroscope (Nikon SMZ800) connected to a high-resolution camera (acA1300-60gm, Basler, Germany). Recording was performed using Media Recorder Software (v.4.0.542.1, Noldus Information Technology, Netherlands) and the analysis of photos was performed using the DanioScope Software (v.1.1.110, Noldus Information Technology, Netherlands) in order to assess morphology measurements
(surface of yolk reserve) and malformations (cardiac edema). Analyses were made for
F1 irradiated, F1 recovery, F1 resilience 6d and F1 resilience 36d.

152 **2.4. Sequencing library and transcriptomic analysis of ovaries**

Ovaries were dissected after reproduction and gonads were kept in RNALater® (Sigma-aldrich) at -80°C until analysis. RNA extractions were made from female gonads (8 per condition). Each sample was homogenized in 600 μ L of ice-cold RTL buffer (Qiagen) with 6 μ L of β -mercaptoethanol using a tissue homogenizer for 3 times 155 Following centrifugation, DNA and RNA were extracted from the homogenate using the AllPrep DNA/RNA kit (Qiagen) according to manufactures' guidelines, with DNAsel's treatment (Qiagen) according to the manufacturer's recommendations.

Libraries were generated from 1 µg of total RNA following the TruSeg mRNA stranded 160 161 protocol (Illumina). RNA integrity (RIN) assessed using RNA Nano Chips (Bioanalyzer 2100, Agilent) did not show any sign of degradation (RIN > 7). Librarie quality and 162 concentration were measured on DNA1000 Chips (Bioanalyzer 2100, Agilent) and 163 sequenced on a NovaSeg 6000 platform to produce 50 bases long paired-end reads 164 (Clinical Research Sequencing Platform, Broad Institute, MIT, Cambridge, USA). 165 Good-quality reads (Q > 30) were produced for each sample (Table S1). Quality check 166 167 of paired end reads and quantification of the RNAseq data were performed as described before (Murat El Houdigui et al., 2019). Briefly, read quality was assessed 168 with FastQC (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/), adapter 169 TrimGalore! v0.6.4 170 sequences were removed with (http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/) and mapped against 171 the GRCz11 zebrafish reference genome using RNA-STAR v020201 and the known 172 exon-exon junctions from Ensembl release 103. Normalisation and differential 173

expression analysis were performed with DESeq2 v1.26.0. Genes with adjusted pvalue < 0.05 (false discovery rate) were considered as differentially expressed in all analysis (DEG). Genes with $|\log 2$ FoldChcange| \geq 1 and adjusted p-value < 0.05 (false discovery rate) were considered as up or down regulated.

178

2.5. Enrichment analysis

ClusterProfiler, a R package, was used to perform pathway enrichment using Gene ontology terms (GO terms), Kyoto Encyclopedia of Genes and Genomes pathway (KEGG pathway) on the DEG obtained from DESeq2 (Table S2). Enrichments with pvalue from Fisher's exact test \leq 0.05 were considered significant. Dot plots were produced using the R package ggplot2.

184 **2.6. RT-qPCR analysis**

RNA extraction for qPCR was performed as for the RNA-seq. A total of 1 µg of RNA 185 was used to synthesize first-strand cDNA using the GoScript Reverse Transcription 186 System (Promega), according to the manufacturer's guidelines. cDNA was then diluted 187 40-fold. Real-time PCR reactions were then performed in a thermocycler (LC480, 188 Roche; 95 °C for 2 min, followed by 45 cycles of 95 °C for 15 s and 60 °C for 1 min). 189 190 Each 15 µL reaction contained 7.5 µL of GoTag qPCR master mix (Promega), 5 µL template and the specific primer pairs at a final concentration of 0.33 µM each. 191 Amplification efficiencies for all primer sets were calculated to allow direct comparison 192 of amplification plots according to the $\Delta\Delta$ Ct method (Livak and Schmittgen, 2001). 193 Relative quantification of gene expression was achieved by amplification of 194 endogenous control genes chosen from our RNAseq dataset, i.e. genes for which their 195 transcription level was not impacted by irradiation (aspa, pdia6). During our 196 experiment, total RNAs were quantified and the same quantity was used to be reverse-197

transcribed. During the subsequent qPCR amplifications, the output cycles of the endogenous controls were examined, no significant difference was observed on the output cycle of aspa and pdia6 among conditions. The mean Ct value of the two latter genes was used as reference. Primers used are available in Table S3 and are involved in oogenesis and sex determination. Log2FC from RNAseq and qPCR analysis were compared in order to validate RNAseq data (Figure S1); they varied similarly between the two analyses.

205 **2.7. Steroid extraction and cortisol measurement**

A half female's gonad per replicate was flash-frozen in liquid nitrogen and kept at -206 80°C until extraction. Gonads were then thawed and transferred in sodocalcic glass 207 208 tube. 1ml of ethyl acetate/cyclohexane (50:50) was added per sample before grinding for 2 min in a potter (Eurostardigital, IKA Labortechnik,) at 2000 Tr/min. Lysate was 209 210 then transferred to new sodocalcic glass tube. This step was carried out twice in order to recover a maximum of material as recommended in Sadoul and Geffroy, 2019. 211 Tubes were vortexed for 10 sec then centrifuged for 5 min at 5000G. Supernatant was 212 213 collected in a sodocalcic glass tube. This step was performed twice. Each sample was evaporated to dryness under nitrogen at 40 °C and resuspended in buffer from the 214 "Cortisol Express ELISA Kit" from Cayman Chemical. Cortisol assay were then 215 216 performed according to manufacturer's guidelines.

217

2.8. Protein extraction and VTG western blot analysis of 24hpf eggs

Twenty 24hpf-eggs per replicate were quick frozen in liquid nitrogen and stored at -80°C until extraction. For protein extraction, samples were placed in a modified RIPA lysis buffer (Sigma Aldrich; with anti-protease mixture), with 0.5-mm diameter zirconium beads and homogenized by three 5000-rpm cycles at 4°C in a Precellys

grinder system (Ozyme). After 30mn incubation on ice, crude extracts were centrifuged 222 for 10 minutes at 4°C at 5000g. Protein concentration was determined using the BCA 223 kit (Thermo Scientific) with BSA as a standard, according to the manufacturer's 224 instructions and then adjusted to a same concentration to ensure the loading of the 225 same amount of proteins for western blot analysis. After denaturation (15mns, 95°C 226 with Laemli buffer (Tris (0,31M), DTT (0,25M), BBP (0,02%), SDS (10%) and glycerol) 227 9µg of proteins from each sample were loaded in precast gels (4-20%? TGX Biorad) 228 and electrophoresis was run in TGS buffrer during 50min (80V for 15 min and 150V 229 35min). 5µL of load marker (RNP800E, ECL Plex Fluo rainbow) were added in one 230 well. After migration, gel was transferred on a polyvinylidene fluoride (PDVF, GE 231 Healthcare) membrane (1h, 4°C, transfer buffer (TGS 1X, ethanol (20%)), for 1 hours 232 at 100V. Membrane was then blocked with BSA 1% (TBST-BSA buffer (TBS1X, 233 234 pH=7.5 - Tween 0.1% - 1% filtered BSA) for 1hand incubated overnight at 4°C in primary mouse antibody JE-10D4 (Abcam ab36804, dilution 1:1000 in TBST-BSA) 235 followed by incubation in secondary anti-mouse antibody Cy5® (Goat Anti-Mouse IgG 236 H&L preadsorbed (ab6563)dilution 1:6000 in TBST-BSA) for 2h at 4°C. Membrane was 237 finally rinsed and results were visualized with Typhoon FLA 9500 (GE Healthcare) at 238 670nm, PMT (). Integrations of bands (volume without background) were done using 239 ImageQuantTL software (1D gel analysis module). Recent bibliography shows that the 240 use of housekeeping protein to normalization is difficult (Eaton et al., 2013; Moritz, 241 2017). These findings were proved in our case with traditional proteins (β -actin and α -242 tubulin) which were highly impacted by gamma irradiation (Figure S2). Reproducibility 243 of lane was proved by multiple gel migrations, as described by (Eaton et al., 2013; 244 Moritz, 2017; Richon et al., 2000), using total protein gel stained in parallel (Table S4) 245 since the use of housekeeping protein to normalization is difficult after exposure to a 246

stressor (Eaton et al., 2013; Moritz, 2017). Homogeneity of variances was tested using bartlett's test for each membrane (p-val>0.05) and variances of two gels were tested using Anova (p-value>0.05).

250 **2.9. Statistical analysis**

For F1 survival rate, F1 morphology, F1 maturation products of vitellogenin intensity 251 and F0 cortisol concentrations, data were presented as mean values ± SE, with 252 significance taken as p < 0.05. For F1 vitellogenin ratio, results were presented by the 253 ratio 30kDa band/ 26kDa band. Concerning data relating to the F1 generation (survival 254 rate and vitellogenin ratio), conditions were compared using a GLM (Generalized 255 Linear Model, glm function in R) with binomial distribution. Concerning data related to 256 morphology, vitellogenin density results and cortisol, conditions were compared using 257 Anova with BoxCox transformation when normality and homogeneity were not verified. 258 A TukeyHSD, post-hoc test, was then used. Concerning qPCR data, data were 259 presented as log2FoldChange where FoldChange (FC) = (mean $2^{-\Delta Ct}_{irradiated}$ - mean $2^{-\Delta Ct}$ 260 $\Delta Ct_{control}$)/ mean 2^{- $\Delta Ct_{control}$}. Conditions were compared using Anova with BoxCox 261 262 transformation when normality and homogeneity were not verified. A Tukey (HSD), post-hoc test, was then used. When normality was not verified even after data 263 transformation, a non-parametric test was used (kruskal wallis). A Dunnet test with 264 265 Bonferroni correction was then used as post hoc test.

The analyses were performed using R software (R Core Team, 2021) with the following packages : "tidyverse", "here", "knitr", "Ime4", "MASS","car". Figures were produced using the package "ggplot2".

269 **3. Results**

270 **3.1. F0 reproductive performances and F1 survival**

F0 adult fish (n=24 per condition) and F1 embryos (n=30 eggs X 6 spawns per condition) were irradiated to a cumulative dose of 12Gy and 4.8Gy, respectively. The direct exposure to IRs did not induce F0 adult mortality over the experiment. The reproductive success (RS) of F0 irradiated adults was lower (n=12 mating pairs, 75%), but not significantly different to control fish (n=12, 92%). No effect on fecundity was observed after exposure to gamma irradiation (n=9, 367±170), compared to the control group (n=11, 344±153).

At 24 hpf, the percentage of F1 mortality for control, irradiated and recovery groups 278 reached 54.9%±5.6, 71.7%±10.7 and 77.0%±7.0, respectively (Figure 2). Despite 279 enhanced mortality for both irradiated and recovery conditions, no significant difference 280 281 was observed compared to the control group. However, an important variability was observed for all groups. The same trend was observed at 48 hpf (control: 56.1%±5.9; 282 283 irradiated: 77.8% \pm 8.2; recovery: 83.0% \pm 5.7), but a significant difference (p < 0.05) was observed only for the recovery condition. A significant difference in mortality (p < 0.05) 284 was observed from 72 hpf for both irradiated (89.9%±4.9) and recovery (98.4%±0.7) 285 groups compared to controls (56.1% ± 5.9). No statistical difference was observed 286 between the irradiated and recovery conditions. Control's mortality remained similar 287 until the last day (6 dpf) of the experiment (57.2%±5.7), whereas it reached 100% for 288 the irradiated group and 99.5%±0.5 for the recovery group. 289

- 290
- 291

3.2. F1 abnormalities after parental exposure

Malformations (pericardium edema and yolk reserve edema) were observed in F1 72 hpf larvae after parental exposure to IRs (Figure 3A). A significant increase (p < 0.05) of yolk reserve area was observed in both conditions, irradiated (n=9, 0.319mm² ±

0.007) and recovery (n=12, 0.349mm² ± 0.019), compared to the control group (n=6, 295 0.244 mm² ± 0.014), with no difference between them (Figure 3B). After setting F0 296 irradiated adult fish in control condition for 6 days, their offspring (Resilience_6d) still 297 showed a significant increase (p < 0.05) in yolk reserve area (n=11, 0.400 mm² ± 0.020) 298 compared to the control group (n=11, 0.241mm² ± 0.009). On the other hand, after 36 299 days of setting the adults in control condition, no morphological defects were observed 300 in their offspring (Resilience_36d). For pericardial area, statistical difference (p < 0.05) 301 was observed only for resilience 6d compared to control group (Figure 3C). 302

303 **3.3. Transmitted macro-molecules from F0 to F1**

304

3.3.1. Glucocorticoid regulation in the gonad

Cortisol concentration in F0 female gonads after reproduction was not impacted by IR (Figure 4A). Indeed, its concentration reached 0.036ng.mg⁻¹ \pm 0.005 for irradiated group (n=6) and 0.041ng.mg⁻¹ \pm 0.006 for control group (n=6). However, at the transcriptional level (RNAseq), a down-regulation of *nr3c1*, the glucocorticoid receptor, was observed in F0 female gonads (n=8) after irradiation (Figure 4B). Moreover, many genes involved in cortisol biosynthesis were also down-regulated (*cyp11a1, cyp11a2, apoa1a, apoeb, npc2, fdx1, fdx1b*).

312 3.3.2. VTG analysis in eggs

Western-blot analysis, using an anti-VTG antibody on 24 hpf F1 embryos (n=7 per condition), revealed two bands at around 26 and 30 kDa (Figure S3), but not the band corresponding to the whole VTG (around 150 kDa). Whatever the band considered, i.e. total one, 30kDa band intensity or 26 kDa band intensity, or the ratio between intensities, no significant difference was observed between control and irradiated groups (Figure S4). However, the transcription level of the gene encoding for the enzyme responsible for VTG cleavage, cathepsin D (*ctsd*), was down-regulated
(Log2FoldChange= -0.5, FDR <0.01).

321 3.4. **Ovarian transcriptome**

Differential gene expression was assessed in pairwise comparison between irradiated and control ovaries (n=8 per condition). The total number of Differentially Expressed Genes (DEG) (FDR < 0.05), up and down regulated genes with a fold change superior to 2 ($|\log 2$ fold change| \geq 1 and FDR < 0.05), up regulated genes ($\log 2$ fold change \geq 1 and FDR < 0.05) and down regulated genes ($\log 2$ fold change \leq 1 and FDR < 0.05) reached 9182, 5348, 3234 and 2114, respectively.

Pathways enrichment analyses were then performed using GO repositories (with zebrafish orthologues) and the KEGG database (with human orthologous genes).

Considering all dysregulated genes (FDR < 0.05) or only up regulated or down 330 regulated genes ($|\log 2$ fold change | ≥ 1 and FDR < 0.05), 304, 394 and 83 GO terms 331 were enriched, respectively (Table S2). The most dysregulated pathways from 332 dysregulated genes were related to ribosomes (GO:0022613, ribonucleoprotein 333 complex biogenesis, p.adjust=1.74E-22) and DNA damage (GO:0006281, DNA repair, 334 p.adjust= 7.25E-11) (Figure 5A). GO terms linked to ribosomes synthesis were 335 principally represented by down regulated genes (GO:0042254, ribosome biogenesis, 336 p.adjust= 5.28E-09). We also found other GO terms associated with DNA damage and 337 genotoxicity that were highly dysregulated (GO:0051403, stress-activated MAPK 338 cascade, p.adjust= 0.02; GO:0097190, apoptotic signalling pathway, p.adjust= 0.02; 339 GO:0006302, double-strand break repair, p.adjust= 2.72E-04). For DNA damage 340 group, transcripts of the p53 pathway were all up-regulated (Figure 5B, tp53bp1, 341 tp53bp2a, tp53bp2b, deltaNp63) while other genes involved in DNA repairs were 342

down-regulated (*gadd45ab*, *rad51*, *xrcc6*, *xrcc5*, *ddb2*, *creb1a*, *creb1b*, *polb*).
Transcripts involved in the fight against oxidative stress were also down-regulated
(*gpx1a*, *gpx1b*, *gsta.1*, *sod1*, *sod2*).

GO terms linked to reproduction (GO:0022414, reproductive process, p.adjust= 6.21E-04; GO:0048477, oogenesis, p.adjust= 3.12E-04) and germ line production (GO:0007281, germ cell development, p.adjust=0.03) were also impacted by irradiation.

350 Dysregulated genes involved in oogenesis and ribosomes are presented in Figure 5B.

They were mostly down-regulated (*buc*, *rpl, rps*) while *macf1a* was up-regulated.

Among the upregulated genes, many were involved in morphogenesis (GO:0060560, 352 developmental growth involved in morphogenesis, p.adjust= 1.13E-04) and embryo 353 axis establishment (GO:0009952, anterior/posterior pattern specification, 354 p.adjust=0.01; GO:0009953, dorsal ventral pattern formation, p.adjust= 2.73E-03, 355 GO:0016055, Wnt signalling pathway, p.adjust= 9.96E-03). Transcripts related to 356 embryonic development, and in particular in axis development, were mainly up-357 regulated (Figure 5B, *ctnnb2*, *runx2b*, *smad* genes), but some important factors for axis 358 establishment were down-regulated (grip2a, sybu, pou5f3, ndr1, gdf3). Transcripts 359 related to the maternal to zygotic transition (MZT) stage were also up-regulated 360 (drosha, dicer1, ago2, ago3b, ago4, ythdf2). 361

Considering dysregulated genes (FDR <0.05), or only up regulated genes or down regulated genes 20, 15 and 5 KEGG pathways were enriched, respectively and are presented in Figure 6. KEGG enrichment analysis with only upregulated genes were consistent to GO terms enrichment since pathway linked to embryo development were impacted (dre04068, FoxO signalling pathway, p.adjust= 0.04; dre04330, Notch signalling pathway, p.adjust= 4.88E-03; dre04340, Hedgehog signalling pathway,
p.adjust= 3.53E-03; dre04310, Wnt signalling pathway, p.adjust= 3.90E-04; dre04012,
ErbB signalling pathway, p.adjust= 1.56E-05). As previously described, many
dysregulated genes or down regulated genes were involved in oogenesis (dre04914,
Progesterone mediated oocyte maturation, p.adjust= 4.23E-03; dre04114, Oocyte
meiosis, p.adjust= 4.23E-03) and ribosomes (dre03008, Ribosome biogenesis in
eukaryotes, p.ajust= 3.61E-06).

374 **4. Discussion**

375 In order to investigate the molecular cascade that induce drastic effects on offspring, a high dose rate of 50 mGy.h⁻¹ was purposely chosen. Here we detected significant 376 differences on offspring survival rates at 72 hpf (Figure 2). Moreover, malformations, 377 mainly yolk reserve edema, were observed for irradiated, recovery and resilience_6d 378 conditions after parental irradiation (Figure 3). Offspring effects were therefore 379 observed even when only genitors were irradiated. Moreover, Resilience_6d condition 380 show altered offspring, while genitors were in control condition. This supports that 381 deleterious effects in genitors induce non-viable offspring. Several maternal factors 382 383 being transmitted to embryo could explain these phenotypes. This is why this study initially focused on maternal involvement. Paternal involvement cannot be totally 384 excluded but seems less obvious to us in the present case where effects are very early, 385 386 probably before zygote genome activation.

387

4.1. IR altered maternal provided molecules

388 4.1.1. IR led to oxidative stress and DNA damages in ovary

The RNAseq analysis on female gonads showed an evidence of DNA damage, confirming previous results using COMET assay (Guirandy et al., 2019; Hurem et al., 2018). Enrichment analysis on GO terms (Figure 5A) revealed typical biological processes related to DNA damage and oxidative stress (« stress-activated MAPK cascade »). Enrichment analysis on KEGG pathways (Figure 6) showed IR-induced effects on « FoxO signalling pathway » and « ErbB signalling pathway », which are involved in oxidative stress and cancer development, respectively, and already known to be impacted by radiations (Dent et al., 2003; Luo et al., 2007).

397 Several genes involved in apoptosis via the P53 pathway (tp53bp1, tp53bp2, deltaNp63) (Elersek et al., 2022)) were up-regulated (Figure 5B). p53 genes, required 398 for radiation-induced apoptosis in mouse (Lowe et al., 1993), are involved in cell cycle 399 400 arrest but also in damage repair in zebrafish (Canedo and Rocha, 2021; Sandrini et al., 2009). However, several genes involved in DNA repair mechanisms, mainly in 401 double strand break reparation (Canedo and Rocha, 2021; Reinardy et al., 2013; 402 Sandrini et al., 2009) were also down regulated (Figure 5B, gadd45ab, rad51, xrcc6, 403 xrcc5, ddb2, creb1a, creb1b, polb). In addition, genes involved in the fight against 404 405 oxidative stress (Figure 5B, gpx1a, gpx1b, gsta.1, sod1, sod2) were down regulated, suggesting an alteration of the antioxidant response and ROS imbalance. The altered 406 oxidative balance in genitors did not seem to disturb the reproductive performances of 407 408 adults but could contribute to adverse effects on oocyte DNA, as explain before, or on offspring development, as proposed by W. Huang et al. 2020. Thus, a direct effect of 409 ROS or an induced effect on the integrity of the maternal DNA in oocytes could occur, 410 thereby producing a future non-viable offspring. However, fertilization and first stages 411 of embryo development (gastrulation) seemed to occur normally. 412

413 4.1.2

4.1.2. IR did not impact cortisol and VTG

414 Maternal cortisol is necessary for proper embryo development (Colson et al., 2015;
415 Faught and Vijayan, 2018). Altered maternal concentration of cortisol may yield cardiac

oedema as described by Nesan and Vijayan, 2012. In our study, even if cardiac 416 oedema were observed in offspring at 72 hpf for resilience-6d (Figure 3E), no effect 417 was observed in ovarian cortisol level (Figure 4A). However, maternal glucocorticoid 418 receptor transcripts (nr3c1 or gr), that participate in the maternal programming of 419 embryo development (Pikulkaew et al. 2011), were down regulated in female gonads 420 (Figure 4B). Furthermore, other transcripts involved in cortisol biosynthetic process 421 were also differentially regulated (Figure 4B, cyp11a1, cyp11a2, apoa1a, apoeb, npc2, 422 fdx1, fdx1b) (Facchinello et al., 2017; Nesan and Vijayan, 2012). Thus, even if these 423 molecular effects were not associated with changes at the metabolite level, an effect 424 of cortisol could not be totally discarded. 425

Moreover, yolk reserve oedemas (Figure 3) were observed in offspring at 72 hpf for 426 irradiated, recovery and resilience-6d, suggesting a nutrient reserve disruption 427 (observed for many pollutants in fish). The gene encoding for the enzyme responsible 428 429 for VTG cleavage, cathepsin D (ctsd), was down regulated, suggesting less VTG conversion into storage forms that are required for early development (Yilmaz et al., 430 2018). However, the total quantity of vitellogenin products that can bind to the antibody 431 was not impacted by irradiation, and yolk reserve areas were higher after irradiation 432 suggesting rather a non-consumption of reserves in larvae from irradiated parents. 433 Since maternally-provided *ctsd* transcripts allows proper treatment and consumption 434 of yolk proteins by embryos (Follo et al., 2011), down-regulation of this gene could 435 explain the non-consumption. Cortisol and VTG were non-impacted by IRs and cannot 436 explain mortality and malformation. Molecular alteration on cortisol biosynthesis and 437 VTG degradation encourage us to further investigations. 438

Another hypothesis to identify mechanisms of transmitted effects was that altered
maternal transcriptome, after irradiation exposure, could have early and delayed
consequences on embryo development.

442 **4.2. IR altered oocyte development**

Enrichment analysis showed GO terms and KEGG pathways linked to reproduction 443 and oogenesis, including « oocyte meiosis » and «oocyte development » (Figure 5A, 444 Figure 6). Embryo development success is dependent on mRNA, molecules, 445 germplasm and organelles (golgi and mitochondria) aggregated in the Balbiani body 446 (Bb) during maternal oogenesis (Jamieson-Lucy and Mullins, 2019; Marlow, 2010). 447 Three maternal genes are involved in Bb development (Marlow, 2010). Among them, 448 bucky ball (buc), that is necessary to establish the polarity of the egg (animal-vegetal 449 axis), was down regulated after irradiation (Figure 5B) (Marlow, 2010). Maternal buc 450 deficiency is known to induce polyspermy with normal cell division after fertilization and 451 ultimately embryo death (Jamieson-Lucy and Mullins, 2019). This could explain death 452 453 at around 72 hpf here and at around 24 hpf in the study of Guirandy et al., 2019. Microtubule actin crosslinking factor 1 (macf1) is critical for Bb disassembly ensuring 454 the dispersion of mRNA along the animal-vegetal axis (Jamieson-Lucy and Mullins, 455 456 2019). In our study, macf1a was up-regulated (Figure 5B), potentially leading to disturbed mRNA distribution and finally to embryo death. 457

Oogenesis also allows the accumulation of maternal ribosomal proteins (RP) in order to support early embryo development until zygotic genome activation (ZGA) (Mercer et al., 2021). Precise control of ribosome biogenesis is vital for cell survival (Danilova et al., 2008) and therefore for preventing abnormalities during zebrafish embryonic development. For example, a deficiency in *rpl11* induces an increase in yolk reserve size in mutants, reflecting a slow metabolism, and a decrease in larvae size (Palasin

et al., 2019). These two phenotypes were also observed in this study and by Guirandy 464 et al. 2019. Enrichment analysis (Figure 5A, Figure 6) highlighted different processes 465 linked to ribosome synthesis (« ribonucleoprotein complex biogenesis », « ribosome 466 biogenesis », « Ribosome biogenesis in eukaryotes »). Several RP transcripts that 467 were down-regulated (Figure 5B, rpl10a, rpl11, rpl23) are known to activate the p53 468 pathway (as observed in 4.1.1) (Chakraborty et al., 2009; Palasin et al., 2019). Down-469 regulation of rps19, rps8a, rps8b, rps11 and rps18 were associated with an up-470 regulation of deltaNp63, as previously observed (Danilova et al., 2008). RP deficiency 471 via its impact on cell division and on the p53 pathway, could represent a further 472 explanation for the observed early effects on offspring after maternal gamma 473 irradiation. 474

It is clear that irradiation altered proper oocyte development. Thus, it could produce
defective offspring. Next, we wondered if altered maternal mRNAs could induce later,
i.e; after spawning and fertilization, effects in addition to direct impacts on oocytes.

478 **4.3. IR altered maternal mRNAs needed for offspring development**

479 *4.3.1. Embryo axis establishment*

Other maternal mRNAs needed for embryo development were also impacted by irradiation. Among them, many are accumulated in Bb and are involved in embryo axis establishment (Marlow, 2010). Enrichment analysis revealed GO terms and KEGG pathways involved in embryo development (Figure 5A, Figure 6), in particular with genes that were up-regulated. Several biological processes concerning the morphogenesis were impacted ("cell projection morphogenesis", "embryonic organ morphogenesis", "dorsal/ventral pattern formation","Wnt signalling pathway"). Since 487 oocyte polarity and maternal mRNAs were impacted (4.2), it appears not aberrant that
488 maternal mRNAs governing embryo axis establishment were also disturbed.

Maternal β -catenin and grip2a are the first indication of dorso-ventral polarity (Langdon 489 490 and Mullins, 2011). β-catenin genes (*ctnnb1* and *ctnnb2*) were down-regulated and upregulated, respectively (Figure 5B). *ctnnb1* is known to act upstream at the molecular 491 level and ctnnb2 then acts on the physical establishment of the axis. Opposite 492 regulations for these two key stages could alter proper embryo development (Zhang et 493 al., 2012). For example, up-regulation of β -catenin leads to hyper-dorsalization (Kelly 494 et al., 1995; Nojima et al., 2004; Sumoy et al., 1999). grip2a that acts upstream or in 495 parallel to maternal β -catenin was down-regulated (Figure 5B). In the same way, 496 syntabulin (sybu) was down regulated (Figure 5B) and loss of maternal sybu was found 497 to induce ventralization and failure to activate the Wnt signalling pathway, which is 498 required to initiate dorsal cell fate specification (Langdon and Mullins, 2011). 499

In contrast, the Bone Morphogenetic Protein signalling pathway, involved in ventral 500 axis specification, was not impacted. However, an initiator gene of this pathway, pou5f3 501 (Abrams and Mullins, 2009; Flores et al., 2008; "Maternal Effect Genes in 502 Development, Volume 140 - 1st Edition," n.d.) was down-regulated (Figure 5B). 503 Moreover, several ventral genes were also dysregulated (Figure 5B, runx2b, smad1, 504 smad2, smad4a, smad5, smad6a) (Pelegri, 2003). As for dorsal axis specification, 505 genes involved in ventral axis specification seemed to be highly dysregulated that 506 could lead to the observed malformations and non-viable embryos. 507

508 Finally, the left/right axis, needed to organ asymmetry such as brain and heart, is 509 mainly governed by the nodal signalling pathway (Bisgrove et al., 2017; Miccoli et al., 510 2017), which was not impacted. However, a key gene of this pathway, *ndr1* was downregulated (Figure 5B). This was not surprising since its positive co-factor, *gdf*3 (Bisgrove et al., 2017), was also down-regulated. *Gdf*3 mutation leads to embryo death. Gamma irradiation thus impacted maternal transcripts necessary to asymmetry development and could explain observed mortality.

515 4.3.2. Maternal mRNA degradation

516 Failure of maternal mRNA degradation concomitantly to zygotic genome activation (ZGA) can lead to miscoordinated gene expression (Chang et al., 2018; Walser and 517 518 Lipshitz, 2011). The best-known degradation mechanism is deadenylation. The majority of maternal transcripts are therefore deadenylated and degraded between 4 519 and 6 hpf (Chang et al., 2018; Walser and Lipshitz, 2011) at the same time as ZGA. 520 521 Interestingly, enrichment analysis showed an impact on the biological process "positive regulation of nuclear-transcribed mRNA catabolic process, deadenylation-dependent 522 decay" (Figure 5), which could reflect a deregulation of maternal mRNA catabolic 523 process after irradiation. We also detected the dysregulation of many factors involved 524 in the maturation of miRNA (drosha, dicer1, ago2, ago3b, ago4) as mentioned after 525 parental irradiation in Martín et al., 2021. As MiR-430 facilitates the deadenylation of 526 maternal mRNAs (Giraldez et al., 2006; Laue et al., 2019; Liu et al., 2020; Yao et al., 527 2014), it is possible that deregulation of miRNA production contribute to the increased 528 529 in embryonic mortality. Another degradation way during maternal to zygotic transition stage is uridylation, done by *tut7* which was down regulated (Figure 5B) (Chang et al., 530 2018). Furthermore, *igf2bp3*, a gene encoding for a protein involved in maternal mRNA 531 532 stability (Ren et al., 2020), was up-regulated (Figure 5B). whereas ythdf2, involved in maternal transcript degradation, was down-regulated (Figure 5B) (Ren et al., 2020). 533 These results suggest a disturbed mRNA degradation process in female gonads. 534 Maternal mRNA could thus be rapidly degraded and could not support early 535

development or could alter the temporal pattern of gene expression during earlydevelopment.

538 **5. Conclusions**

This paper highlighted transmitted effect from genitors to offspring after 10 days of 539 gamma irradiation at 50 mGy.h-1. No phenotypic effect was observed in genitors, 540 notably on reproductive performances (reproductive success and fecundity). Early 541 mortality in offspring in link to parental transmitted effects was observed as no 542 difference was observed between irradiated and recovery conditions. Moreover, early 543 malformations were observed in offspring after parental irradiation. The main objective 544 545 was to get mechanistic insights into the impacts of irradiation at the cellular and molecular levels in order to identify mechanisms behind phenotypic effects on 546 offspring. 547

Cortisol level in F0 female gonads after reproduction was not impacted by irradiation. 548 However, it should be necessary to measure cortisol in embryos to refute its 549 550 involvement in F1 mortality. Despite a down-regulation of cathepsin D in maternal gonads, VTG analysis in embryos did not highlight any different amount nor distribution 551 of maturation products which suggests no nutrition defect for embryos. This non-effect 552 should be however confirmed by measuring the whole VTG protein. Oogenesis and 553 maternal factors (mRNA) required for production of viable offspring and their 554 development were significantly altered, supporting maternal transmission of IR-555 induced effects. 556

557 This first multigenerational approach to link phenotype to molecular events highlighted 558 an important effect of female irradiation on offspring viability with (i) an impact of 559 irradiation on oocyte development and (ii) an impact on maternal mRNAs needed later for embryonic development. However, paternal contribution should not be excluded and should be investigated in future experiments, since they were also exposed. Further experiments would be to perform a cross-exposure (only female exposed) to assess maternal role. Finally, offspring transcriptomic analysis before and after the maternal-to-zygotic transition should be interesting to define more precisely the role of maternal mRNAs. Our data could be used to improve the Adverse Outcome Pathways and challenge the environmental risk assessment method currently deployed.

567 Funding sources

568 Funding: This work was supported by the Institut de Radioprotection et de Sûreté

569 Nucléaire (IRSN).

570 **References**

571	Abrams, E.W., Mullins, M.C., 2009. Early zebrafish development: It's in the maternal genes. Curr.
572	Opin. Genet. Dev. 19, 396–403. https://doi.org/10.1016/j.gde.2009.06.002
573	Beresford, N.A., Horemans, N., Copplestone, D., Raines, K.E., Orizaola, G., Wood, M.D., Laanen, P.,
574	Whitehead, H.C., Burrows, J.E., Tinsley, M.C., Smith, J.T., Bonzom, JM., Gagnaire, B., Adam-
575	Guillermin, C., Gashchak, S., Jha, A.N., de Menezes, A., Willey, N., Spurgeon, D., 2020.
576	Towards solving a scientific controversy – The effects of ionising radiation on the
577	environment. J. Environ. Radioact. 211, 106033.
578	https://doi.org/10.1016/j.jenvrad.2019.106033
579	Bisgrove, B., Su, YC., Yost, H., 2017. Maternal Gdf3 is an obligatory cofactor in Nodal signaling for
580	embryonic axis formation in zebrafish. eLife. https://doi.org/10.7554/eLife.28534
581	Canedo, A., Rocha, T.L., 2021. Zebrafish (Danio rerio) using as model for genotoxicity and DNA repair
582	assessments: Historical review, current status and trends. Sci. Total Environ. 762, 144084.
583	https://doi.org/10.1016/j.scitotenv.2020.144084
584	Car, C., Gilles, A., Armant, O., Burraco, P., Beaugelin-Seiller, K., Gashchak, S., Camilleri, V., Cavalié, I.,
585	Laloi, P., Adam-Guillermin, C., Orizaola, G., Bonzom, JM., 2022. Unusual evolution of tree
586	frog populations in the Chernobyl exclusion zone. Evol. Appl. 15, 203–219.
587	https://doi.org/10.1111/eva.13282
588	Chakraborty, A., Uechi, T., Higa, S., Torihara, H., Kenmochi, N., 2009. Loss of Ribosomal Protein L11
589	Affects Zebrafish Embryonic Development through a p53-Dependent Apoptotic Response.
590	PLOS ONE 4, e4152. https://doi.org/10.1371/journal.pone.0004152
591	Chang, H., Yeo, J., Kim, JG., Kim, H., Lim, J., Lee, M., Kim, H.H., Ohk, J., Jeon, HY., Lee, H., Jung, H.,
592	Kim, KW., Kim, V.N., 2018. Terminal Uridylyltransferases Execute Programmed Clearance of
593	Maternal Transcriptome in Vertebrate Embryos. Mol. Cell 70, 72-82.e7.
594	https://doi.org/10.1016/j.molcel.2018.03.004

- Chen, M.-F., Huang, C.-Z., Pu, D.-Y., Zheng, C.-Y., Yuan, K.-M., Jin, X.-X., Zhang, Y.-G., Jin, L., 2015.
 Toxic effects of CdSe/ZnS QDs to zebrafish embryos. Huanjing KexueEnvironmental Sci. 36,
 719–726. https://doi.org/10.13227/j.hjkx.2015.02.046
- Colson, V., Valotaire, C., Geffroy, B., Kiilerich, P., 2015. Egg Cortisol Exposure Enhances Fearfulness in
 Larvae and Juvenile Rainbow Trout. Ethology 121, 1191–1201.
 https://doi.org/10.1111/eth.12437
- Cunningham, K., Hinton, T.G., Luxton, J.J., Bordman, A., Okuda, K., Taylor, L.E., Hayes, J., Gerke, H.C.,
 Chinn, S.M., Anderson, D., Laudenslager, M.L., Takase, T., Nemoto, Y., Ishiniwa, H., Beasley,
 J.C., Bailey, S.M., 2021. Evaluation of DNA damage and stress in wildlife chronically exposed
 to low-dose, low-dose rate radiation from the Fukushima Dai-ichi Nuclear Power Plant
 accident. Environ. Int. 155, 106675. https://doi.org/10.1016/j.envint.2021.106675
- Danilova, N., Sakamoto, K.M., Lin, S., 2008. Ribosomal protein S19 deficiency in zebrafish leads to
 developmental abnormalities and defective erythropoiesis through activation of p53 protein
 family. Blood 112, 5228–5237. https://doi.org/10.1182/blood-2008-01-132290
- Dent, P., Yacoub, A., Contessa, J., Caron, R., Amorino, G., Valerie, K., Hagan, M.P., Grant, S., SchmidtUllrich, R., 2003. Stress and Radiation-Induced Activation of Multiple Intracellular Signaling
 Pathways1. Radiat. Res. 159, 283–300. https://doi.org/10.1667/00337587(2003)159[0283:SARIAO]2.0.CO;2
- Eaton, S.L., Roche, S.L., Hurtado, M.L., Oldknow, K.J., Farquharson, C., Gillingwater, T.H., Wishart,
 T.M., 2013. Total Protein Analysis as a Reliable Loading Control for Quantitative Fluorescent
 Western Blotting. PLOS ONE 8, e72457. https://doi.org/10.1371/journal.pone.0072457
- Facchinello, N., Skobo, T., Meneghetti, G., Colletti, E., Dinarello, A., Tiso, N., Costa, R., Gioacchini, G.,
 Carnevali, O., Argenton, F., Colombo, L., Dalla Valle, L., 2017. Nr3c1 null mutant zebrafish are
 viable and reveal DNA-binding-independent activities of the glucocorticoid receptor. Sci. Rep.
 7. https://doi.org/10.1038/s41598-017-04535-6
- Faught, E., Vijayan, M., 2018. Maternal stress and fish reproduction: The role of cortisol revisited.
 Fish Fish. https://doi.org/10.1111/FAF.12309
- Finn, R.N., 2007. Vertebrate Yolk Complexes and the Functional Implications of Phosvitins and Other
 Subdomains in Vitellogenins1. Biol. Reprod. 76, 926–935.
 https://doi.org/10.1095/biolreprod.106.059766
- Flores, M.V.C., Lam, E.Y.N., Crosier, K.E., Crosier, P.S., 2008. Osteogenic transcription factor Runx2 is
 a maternal determinant of dorsoventral patterning in zebrafish. Nat. Cell Biol. 10, 346–352.
 https://doi.org/10.1038/ncb1697
- Follo, C., Ozzano, M., Mugoni, V., Castino, R., Santoro, M., Isidoro, C., 2011. Knock-Down of Cathepsin
 D Affects the Retinal Pigment Epithelium, Impairs Swim-Bladder Ontogenesis and Causes
 Premature Death in Zebrafish. PLOS ONE 6, e21908.
- 631 https://doi.org/10.1371/journal.pone.0021908
- Geras'kin, S.A., Fesenko, S.V., Volkova, P.Yu., Isamov, N.N., 2021. What Have We Learned about the
 Biological Effects of Radiation from the 35 Years of Analysis of the Consequences of the
 Chernobyl NPP Accident? Biol. Bull. 48, 2105–2126.
- 635 https://doi.org/10.1134/S1062359021120050
- Giraldez, A.J., Mishima, Y., Rihel, J., Grocock, R.J., Van Dongen, S., Inoue, K., Enright, A.J., Schier, A.F.,
 2006. Zebrafish MiR-430 promotes deadenylation and clearance of maternal mRNAs. Science
 312, 75–79. https://doi.org/10.1126/science.1122689
- Guirandy, N., Gagnaire, B., Camilleri, V., Cavalié, I., Pierron, F., Gonzalez, P., Simon, O., 2022.
 Multigenerational exposure to gamma radiation affects offspring differently over generations in zebrafish. Aquat. Toxicol. 244, 106101. https://doi.org/10.1016/j.aquatox.2022.106101
- Guirandy, N., Gagnaire, B., Frelon, S., Munch, T., Dubourg, N., Camilleri, V., Cavalié, I., Floriani, M.,
 Arcanjo, C., Houdigui, S.M.E., Armant, O., Adam-Guillermin, C., Gonzalez, P., Simon, O., 2019.
 Adverse effects induced by chronic gamma irradiation in progeny of adult fish not affecting
 parental reproductive performance. Environ. Toxicol. Chem. 38, 2556–2567.
- 646 https://doi.org/10.1002/etc.4562

647 Huang, W., Zheng, S., Wang, X., Cai, Z., Xiao, J., Liu, C., Wu, K., 2020. A transcriptomics-based analysis 648 of toxicity mechanisms of zebrafish embryos and larvae following parental Bisphenol A 649 exposure. Ecotoxicol. Environ. Saf. 205. https://doi.org/10.1016/j.ecoenv.2020.111165 650 Hurem, S., Gomes, T., Brede, D.A., Lindbo Hansen, E., Mutoloki, S., Fernandez, C., Mothersill, C., 651 Salbu, B., Kassaye, Y.A., Olsen, A.-K., Oughton, D., Aleström, P., Lyche, J.L., 2017. Parental 652 gamma irradiation induces reprotoxic effects accompanied by genomic instability in zebrafish 653 (Danio rerio) embryos. Environ. Res. 159, 564–578. 654 https://doi.org/10.1016/j.envres.2017.07.053 Hurem, S., Gomes, T., Brede, D.A., Mayer, I., Lobert, V.H., Mutoloki, S., Gutzkow, K.B., Teien, H.-C., 655 656 Oughton, D., Aleström, P., Lyche, J.L., 2018. Gamma irradiation during gametogenesis in 657 young adult zebrafish causes persistent genotoxicity and adverse reproductive effects. 658 Ecotoxicol. Environ. Saf. 154, 19–26. https://doi.org/10.1016/j.ecoenv.2018.02.031 659 Jamieson-Lucy, A., Mullins, M.C., 2019. Chapter One - The vertebrate Balbiani body, germ plasm, and 660 oocyte polarity, in: Lehmann, R. (Ed.), Current Topics in Developmental Biology, The 661 Immortal Germline. Academic Press, pp. 1–34. https://doi.org/10.1016/bs.ctdb.2019.04.003 Kelly, G.M., Erezyilmaz, D.F., Moon, R.T., 1995. Induction of a secondary embryonic axis in zebrafish 662 663 occurs following the overexpression of β -catenin. Mech. Dev. 53, 261–273. 664 https://doi.org/10.1016/0925-4773(95)00442-4 Langdon, Y.G., Mullins, M.C., 2011. Maternal and zygotic control of zebrafish dorsoventral axial 665 patterning. Annu. Rev. Genet. 45, 357–377. https://doi.org/10.1146/annurev-genet-110410-666 667 132517 668 Laue, K., Rajshekar, S., Courtney, A.J., Lewis, Z.A., Goll, M.G., 2019. The maternal to zygotic transition 669 regulates genome-wide heterochromatin establishment in the zebrafish embryo. Nat. 670 Commun. 10, 1551. https://doi.org/10.1038/s41467-019-09582-3 671 Liu, Y., Zhu, Z., Ho, I.H.T., Shi, Y., Li, J., Wang, X., Chan, M.T.V., Cheng, C.H.K., 2020. Genetic Deletion 672 of miR-430 Disrupts Maternal-Zygotic Transition and Embryonic Body Plan. Front. Genet. 11. 673 https://doi.org/10.3389/fgene.2020.00853 674 Lowe, S.W., Ruley, H.E., Jacks, T., Housman, D.E., 1993. p53-dependent apoptosis modulates the 675 cytotoxicity of anticancer agents. Cell 74, 957–967. https://doi.org/10.1016/0092-676 8674(93)90719-7 677 Luo, X., Puig, O., Hyun, J., Bohmann, D., Jasper, H., 2007. Foxo and Fos regulate the decision between 678 cell death and survival in response to UV irradiation. EMBO J. 26, 380–390. 679 https://doi.org/10.1038/sj.emboj.7601484 680 Marlow, F.L., 2010. Oocyte Polarity and the Embryonic Axes: The Balbiani Body, an Ancient Oocyte 681 Asymmetry, Maternal Control of Development in Vertebrates: My Mother Made Me Do It! 682 Morgan & Claypool Life Sciences. 683 Martín, L., Kamstra, J.H., Hurem, S., Lindeman, L.C., Brede, D.A., Aanes, H., Babiak, I., Arenal, A., 684 Oughton, D., Salbu, B., Lyche, J.L., Aleström, P., 2021. Altered non-coding RNA expression 685 profile in F1 progeny 1 year after parental irradiation is linked to adverse effects in zebrafish. 686 Sci. Rep. 11, 4142. https://doi.org/10.1038/s41598-021-83345-3 687 Maternal Effect Genes in Development, Volume 140 - 1st Edition [WWW Document], n.d. URL 688 https://www.elsevier.com/books/maternal-effect-genes-in-development/marlow/978-0-12-689 815220-1 (accessed 2.9.22). 690 McCormick, M.I., 1999. Experimental test of the effect of maternal hormones on larval quality of a 691 coral reef fish. Oecologia 118, 412–422. https://doi.org/10.1007/s004420050743 692 Mercer, M., Jang, S., Ni, C., Buszczak, M., 2021. The Dynamic Regulation of mRNA Translation and 693 Ribosome Biogenesis During Germ Cell Development and Reproductive Aging. Front. Cell 694 Dev. Biol. 9. https://doi.org/10.3389/fcell.2021.710186 695 Miccoli, A., Dalla Valle, L., Carnevali, O., 2017. The maternal control in the embryonic development of 696 zebrafish. Gen. Comp. Endocrinol. 245, 55–68. https://doi.org/10.1016/j.ygcen.2016.03.028 697 Moritz, C.P., 2017. Tubulin or Not Tubulin: Heading Toward Total Protein Staining as Loading Control 698 in Western Blots. PROTEOMICS 17, 1600189. https://doi.org/10.1002/pmic.201600189

- Murat El Houdigui, S., Adam-Guillermin, C., Loro, G., Arcanjo, C., Frelon, S., Floriani, M., Dubourg, N.,
 Baudelet, E., Audebert, S., Camoin, L., Armant, O., 2019. A systems biology approach reveals
 neuronal and muscle developmental defects after chronic exposure to ionising radiation in
 zebrafish. Sci. Rep. 9, 20241. https://doi.org/10.1038/s41598-019-56590-w
- Nesan, D., Vijayan, M.M., 2016. Maternal Cortisol Mediates Hypothalamus-Pituitary-Interrenal Axis
 Development in Zebrafish. Sci. Rep. 6, 22582. https://doi.org/10.1038/srep22582
- Nesan, D., Vijayan, M.M., 2012. Embryo exposure to elevated cortisol level leads to cardiac
 performance dysfunction in zebrafish. Mol. Cell. Endocrinol. 363, 85–91.
 https://doi.org/10.1016/j.mce.2012.07.010
- Nojima, H., Shimizu, T., Kim, C.-H., Yabe, T., Bae, Y.-K., Muraoka, O., Hirata, T., Chitnis, A., Hirano, T.,
 Hibi, M., 2004. Genetic evidence for involvement of maternally derived Wnt canonical
 signaling in dorsal determination in zebrafish. Mech. Dev. 121, 371–386.
 https://doi.org/10.1016/j.mod.2004.02.003
- Palasin, K., Uechi, T., Yoshihama, M., Srisowanna, N., Choijookhuu, N., Hishikawa, Y., Kenmochi, N.,
 Chotigeat, W., 2019. Abnormal development of zebrafish after knockout and knockdown of
 ribosomal protein L10a. Sci. Rep. 9, 18130. https://doi.org/10.1038/s41598-019-54544-w
- Pelegri, F., 2003. Maternal factors in zebrafish development. Dev. Dyn. Off. Publ. Am. Assoc. Anat.
 228, 535–554. https://doi.org/10.1002/dvdy.10390
- Pikulkaew, S., Benato, F., Celeghin, A., Zucal, C., Skobo, T., Colombo, L., Dalla Valle, L., 2011. The
 knockdown of maternal glucocorticoid receptor mRNA alters embryo development in
 zebrafish. Dev. Dyn. 240, 874–889. https://doi.org/10.1002/dvdy.22586
- R Core Team, R.C.T., 2021. R: A language and environment for statistical ## computing. R
 Foundation for Statistical Computing, Vienna, Austria. [WWW Document]. URL URL
 https://www.R-project.org/
- Reinardy, H.C., Syrett, J.R., Jeffree, R.A., Henry, T.B., Jha, A.N., 2013. Cobalt-induced genotoxicity in
 male zebrafish (Danio rerio), with implications for reproduction and expression of DNA repair
 genes. Aquat. Toxicol. 126, 224–230. https://doi.org/10.1016/j.aquatox.2012.11.007
- Ren, F., Lin, Q., Gong, G., Du, X., Dan, H., Qin, W., Miao, R., Xiong, Y., Xiao, R., Li, X., Gui, J.-F., Mei, J.,
 2020. Igf2bp3 maintains maternal RNA stability and ensures early embryo development in
 zebrafish. Commun. Biol. 3, 94. https://doi.org/10.1038/s42003-020-0827-2
- Richon, V.M., Sandhoff, T.W., Rifkind, R.A., Marks, P.A., 2000. Histone deacetylase inhibitor
 selectively induces p21WAF1 expression and gene-associated histone acetylation. Proc. Natl.
 Acad. Sci. 97, 10014–10019. https://doi.org/10.1073/pnas.180316197
- Sadoul, B., Geffroy, B., 2019. Measuring cortisol, the major stress hormone in fishes. J. Fish Biol. 94,
 540–555. https://doi.org/10.1111/jfb.13904
- Sandrini, J.Z., Trindade, G.S., Nery, L.E.M., Marins, L.F., 2009. Time-course Expression of DNA Repair related Genes in Hepatocytes of Zebrafish (Danio rerio) After UV-B Exposure. Photochem.
 Photobiol. 85, 220–226. https://doi.org/10.1111/j.1751-1097.2008.00422.x
- Simon, O., Massarin, S., Coppin, F., Hinton, T.G., Gilbin, R., 2011. Investigating the embryo/larval toxic
 and genotoxic effects of γ irradiation on zebrafish eggs. J. Environ. Radioact. 102, 1039–1044.
 https://doi.org/10.1016/j.jenvrad.2011.06.004
- Soares, J., Coimbra, A.M., Reis-Henriques, M.A., Monteiro, N.M., Vieira, M.N., Oliveira, J.M.A.,
 Guedes-Dias, P., Fontaínhas-Fernandes, A., Parra, S.S., Carvalho, A.P., Castro, L.F.C., Santos,
 M.M., 2009. Disruption of zebrafish (Danio rerio) embryonic development after full life-cycle
 parental exposure to low levels of ethinylestradiol. Aquat. Toxicol. 95, 330–338.
 https://doi.org/10.1016/j.aquatox.2009.07.021
- Sumoy, L., Kiefer, J., Kimelman, D., 1999. Conservation of intracellular Wnt signaling components in dorsal-ventral axis formation in zebrafish. Dev. Genes Evol. 209, 48–58.
 https://doi.org/10.1007/s004270050226
- Vuori, K.A., Kanerva, M., Ikonen, E., Nikinmaa, M., 2008. Oxidative stress during baltic salmon feeding
 migration may be associated with yolk-sac fry mortality. Environ. Sci. Technol. 42, 2668–
 2673. https://doi.org/10.1021/es702632c

- Walser, C.B., Lipshitz, H.D., 2011. Transcript clearance during the maternal-to-zygotic transition. Curr.
 Opin. Genet. Dev., Differentiation and gene regulation 21, 431–443.
 https://doi.org/10.1016/j.gde.2011.03.003
- Yao, Y., Ma, L., Jia, Q., Deng, W., Liu, Z., Zhang, Y., Ren, J., Xue, Y., Jia, H., Yang, Q., 2014. Systematic
 characterization of small RNAome during zebrafish early developmental stages. BMC
 Genomics 15, 117. https://doi.org/10.1186/1471-2164-15-117
- Yilmaz, O., Patinote, A., Nguyen, T., Bobe, J., 2018. Multiple vitellogenins in zebrafish (Danio rerio):
 quantitative inventory of genes, transcripts and proteins, and relation to egg quality. Fish
 Physiol. Biochem. 44, 1509–1525. https://doi.org/10.1007/s10695-018-0524-y
- Zhang, M., Zhang, J., Lin, S.-C., Meng, A., 2012. β-Catenin 1 and β-catenin 2 play similar and distinct
 roles in left-right asymmetric development of zebrafish embryos. Development 139, 2009–
 2019. https://doi.org/10.1242/dev.074435

763

764			
765			
766			
767			
768			
769			
770			
771			
772			
773			
774			
775			
776			
777			
778			
779			
780			
781			
782			
783			

Figures 784

785



786

787 Figure 1: Experimental design and exposure conditions (duration and dose rate (mGy.h⁻¹). F0 adults (24 per 788 conditions) were exposed over 10 days until reproduction. F1 progenies (30 eggs X 6 spawns) were then placed 789 in irradiated (irradiated condition) and non-irradiated (recovery condition) exposure conditions over 6 days. F0 adults 790 791 were then moved to non-irradiated condition for 6 and 36 days before reproduction. F1 progenies from 6 days F0

(resilience_6d condition) and from 36 days F0 (resilience_36d condition) were then placed in control condition.



Figure 2: Box-plot of cumulative mortality rates of F1 (%)) at 24, 48, 72, 96, 120, 144 hpf after control, dose rate exposure to 50 mGy.h⁻¹ (irradiated) and recovery conditions. The boxplot represents the 25th and the 75th percentile with the median indicated by the horizontal black line. For control, irradiated and recovery, n=9, n=5, n=6, respectively. * (p < 0.05), ** (p < 0.01), *** (p < 0.001).

797



800 Figure 3: (A) Photographs of F1 larvae at 72 hpf after Control, F0 exposure to 50 mGy.h⁻¹ (10 days) with F1 exposure to 50 mGy. h^{-1} (Irradiated), F0 exposure to 50 mGy. h^{-1} (10 days) follow by non-irradiation 6 days (Resilience_6d) or 36 days (Resilience_36d). (B and C) Bar plot of yolk reserve area (mm²) and pericardial area (mm²) for control 801 802 803 (n=6), control_resilience_6d (n=11), control_resilience_36d (n=45), irradiated (F0 + F1 exposed to 50 mGy.h⁻¹, 804 n=9), recovery (F0 exposed to 50 mGy.h⁻¹, F1 non-irradiated, n=12), resilience_6d (F0 exposed to 50 mGy.h⁻¹ follow by non- irradiation exposure 6 days, F1 non-irradiated, n=11), resilience_36d (F0 exposed to 50 mGy.h⁻¹ follow by 805 806 non-exposure 36 days, F1 non-exposed, n=32). Letters indicate statistical differences (pl<0.05) among conditions 807 (Anova test and Tukey post-hoc).

808





Figure 4: (A) Bar plot of F0 ovaries cortisol concentration after control and exposure to 50 mGy.h⁻¹ (irradiated), n=6 per condition. (B) Barplot of log2FoldChange (as compared to controls) for genes involved in cortisol metabolism; n=8 per condition. Data came from transcriptomic data on F0 ovaries. Red line represents cut off value chosen for interpretation of gene regulation. * (p < 0.05), ** (p < 0.01), *** (p < 0.001). 812 813 814



Figure 5: (A) Dot plot of selected GO (Gene Ontology) term enrichment from dysregulated, up and down genes in
exposed F0 ovaries. Colours indicate the p-values from Fisher's exact test, and dots size is the number of genes
constituting the given pathway. n=8 per condition. (B) Change in gene transcription levels (FC as compared to
controls) in female gonad F0 from transcriptomic data. For all genes, FDR<0.01. Red line represents cut off value
chosen for interpretation of gene regulation. n=8 per condition.



Figure 6: Dot plot of selected KEGG enrichment using human orthologues from misregulated, up and down genes
in irradiated ovaries. Colours indicate the enrichment p-values from exact Fisher's test, and dots size is the number
of genes constituting the given pathway. n=8 per condition.