



**HAL**  
open science

# Cyclin G: A potential mitotic bookmark and its role in developmental stability through transcriptional regulation

Léa Pautet, Delphine Dardalhon-Cuménal, Valérie Ribeiro, Marco da Costa,  
F. Peronnet

## ► To cite this version:

Léa Pautet, Delphine Dardalhon-Cuménal, Valérie Ribeiro, Marco da Costa, F. Peronnet. Cyclin G: A potential mitotic bookmark and its role in developmental stability through transcriptional regulation. VIP&DIF 2023, Jun 2023, Paris, France. hal-04240295

**HAL Id: hal-04240295**

**<https://hal.science/hal-04240295v1>**

Submitted on 13 Oct 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Léa Pautet<sup>✉</sup>, Delphine Dardalhon-Cuménal<sup>✉</sup>, Valérie Ribeiro, Marco Da Costa and Frédérique Peronnet.


Developmental Biology Laboratory (LBD), Biology Paris Seine Institute (IBPS), CNRS & Sorbonne Université, 75005 Paris, France.

✉ These authors contributed equally to this work.

In *Drosophila melanogaster*, the atypical cyclin, Cyclin G (CycG), is a transcriptional regulator that maintains cell growth homeostasis and developmental stability. The expression of a potentially more stable form of CycG, CycG<sup>ΔPEST</sup>, has been shown to inhibit the G1/S transition of the cell cycle, slow development<sup>1</sup>, and lead to a loss of correlation between cell size and total cell number in wings of flies<sup>2</sup>. Furthermore, flies expressing CycG<sup>ΔPEST</sup> have a high fluctuating asymmetry, notably of wings, indicating a high developmental noise<sup>2</sup>. CycG also interacts genetically and physically with several subunits of the Polycomb PRC1 and PR-DUB transcriptional repressor complexes, which regulates developmental processes<sup>3</sup>. Mutants for these complexes exhibit increased CycG-induced fluctuating asymmetry in wings, with extreme effects in mutants for the ubiquitinase Sce of PRC1 and the deubiquitinase Calypso of PR-DUB. CycG physically interacts with both enzymes and can be ubiquitinated, suggesting that it might be ubiquitinated/deubiquitinated by PRC1 and PR-DUB, respectively. The regulation of CycG by ubiquitination might be crucial for developmental stability. Furthermore, RNA-seq experiments on wing imaginal discs expressing CycG<sup>ΔPEST</sup> show that the expression of genes involved in metabolism and mitochondrial activity decreases whereas expression of genes involved in translation increases<sup>3</sup>. ChIP-seq experiments also reveal that CycG is enriched on the promoters of some of these genes. Strikingly, these promoters are bound by PRC1 in the same tissue. Interestingly, studies on HeLa cells suggest that promoters of homologous genes are bookmarked during mitosis by PRC1 ubiquitination of an unknown protein<sup>4</sup>. It is hypothesized that this ubiquitinated protein could be CycG that would serve as a mitotic bookmark on its transcriptional targets.


## I - Developmental Stability and Fluctuating Asymmetry

In bilaterians, developmental noise is assessed by Fluctuating Asymmetry in a population of isogenic flies raised in a standardized environment.




## II - Cyclin G genetically interacts and shares transcriptional targets with PRC1 and PR-DUB

Mutations in the Polycomb complexes PRC1 and PR-DUB increase CycG-induced Fluctuating Asymmetry




## Functional subnetwork identified in wing imaginal discs expressing CycG




## III - Cyclin G, a mitotic bookmark?

Hypothesis:



CycG, ubiquitinated by PRC1, would bind its transcriptional targets during mitosis and act as a bookmark that triggers activation of their transcription in early G1 phase.



## CycG physically interacts with PRC1 and PR-DUB



## Development of the CUT&RUN technique to study the presence of CycG on its transcriptional targets during the cell cycle



## References

- 1- Faradji, F et al. Cell Cycle 10, 805–818 (2011).
- 2- Debat, V. et al. PLoS Genet 7, e1002314 (2011).
- 3- Dardalhon-Cuménal, D. et al. PLoS Genet 14, e1007498 (2018).
- 4- Arora, M., et al. Nucleic Acids Res 44, 2136–2144 (2016).
- 5- Dupont CA et al. Epigenetics Chromatin. 2015 May 7;8:18. doi: 10.1186/s13072-015-0008-6.
- 6- Skene, P. J. et al. E life 6, e21856 (2017)

## Ongoing projects

- ✓ Analysis of CycG ubiquitination : Is CycG ubiquitinated/deubiquitinated during the cell cycle? and if so, are PRC1 and PR-DUB involved in this process?
- ✓ Binding of CycG to its transcriptional targets along the cell cycle by CUT&RUN on flow cytometry sorted cells : Is CycG involved in mitosis bookmarking of these genes?