



**HAL**  
open science

## Caution is advised with prevalence comparisons: the case of psychological distress with students and COVID in France, a network approach

Quentin Facon-Barillot, Lucia Romo, Camille Vansimaey, Astrid Chevance, Ariel Frajerman, Yannick Morvan

### ► To cite this version:

Quentin Facon-Barillot, Lucia Romo, Camille Vansimaey, Astrid Chevance, Ariel Frajerman, et al.. Caution is advised with prevalence comparisons: the case of psychological distress with students and COVID in France, a network approach. ICPS 2023: International Convention of Psychological Science, Mar 2023, Bruxelles, Belgium. OSF, 2023, 10.17605/OSF.IO/MT5YU . hal-04240892

**HAL Id: hal-04240892**

**<https://hal.science/hal-04240892v1>**

Submitted on 16 Oct 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Comparison of prevalences since the pandemic may not be relevant!

## Caution is advised with prevalence comparisons: The Case of Psychological Distress with Students and COVID in France, a Network Approach

Quentin Facon-Barillot<sup>1</sup> Lucia Romo<sup>1,2,3</sup> Camille Vansimaey<sup>4,5</sup> Astrid Chevance<sup>6,7</sup> Ariel Frajerman<sup>8,9,10</sup> Yannick Morvan<sup>1,10,11</sup>

<sup>1</sup> Université Paris Nanterre, UFR SPSE, UR4430 CLIPSYD, Nanterre, France

<sup>2</sup> Inserm U1018, CESP Soins primaires et prévention, France

<sup>3</sup> Hôpital Raymond-Poincaré AP-HP, France

<sup>4</sup> Université Paris Cité, Institut de psychologie, LPPS, Boulogne, France

<sup>5</sup> Nightline France

<sup>6</sup> Inserm U1153 CRESS, France

<sup>7</sup> Service d'épidémiologie clinique, APHP, France

<sup>8</sup> Inserm U1018, CESP, Equipe MOODS, France

<sup>9</sup> SHU de Psychiatrie, APHP, Kremlin-Bicêtre, France

<sup>10</sup> Institut de psychiatrie, CNRS GDR 3557, France

<sup>11</sup> Inserm U1018, CESP, Psychiatrie du développement et trajectoires, France

## Introduction

The Covid pandemic has motivated many studies to measure its psychological impact, especially among students.<sup>1,2,3</sup> However, it is important to ensure that the measured constructs are comparable (measurement invariance); otherwise, the comparison of total scores or prevalences is biased<sup>4,5</sup>. We propose to use the network approach<sup>6</sup> to model, assess and compare psychological distress in a way that circumvents the problem of measurement non-invariance while informing the potential psychological impact of the pandemic among students. Finally, based on post-pandemic networks, we used simulation techniques to identify the most relevant intervention and prevention targets<sup>7</sup>.




Figure 1: Changes in psychological distress between 2020 and 2021

## Methods

OVE surveys measured Psychological Distress in nationally representative French student samples, over the last 2 weeks, in 2020 (n=60,014) and 2021 (n=4,901) using the MHI-5<sup>8</sup>. Measurement Invariance (latent modeling with ordinal indicators)<sup>9</sup>, Network Modeling (GGM, corMethod=spearman)<sup>10,11</sup>, Network Comparison Tests (NCT with GGM)<sup>12</sup>, and simulations (NIRA method with Ising Model)<sup>7</sup> were performed with R software.

## Results

### Measurement invariance

There is a maximum difference of 12.7% in the prevalence of Psychological Distress between 2020 and 2021 based on sum score indicators (Figure 1). However, measurement invariance in a latent framework is only partial after correction (Figure 2).

Comparison	Weak <sup>1</sup>	Strong <sup>2</sup>	Strict <sup>3</sup>	Partial <sup>4</sup>
2020 - 2021	Partial	Partial	No	Loadings : D & T   Intercepts : T, D & H

<sup>1</sup> Weak -> Loadings equals  
<sup>2</sup> Strong -> Loading + Intercepts equals  
<sup>3</sup> Strict -> Loadings + Intercepts + Residuals equals  
<sup>4</sup> Equality constraints removed between groups  
N = Nervous ; S = Sad ; C = Calm ; H = Happy ; D = Discouraged

Figure 2: Measurement invariance between 2020 and 2021

### Network comparisons

For networks, if the results were not conclusive for global connectivity, however, changes were observed for the global network structure and for edge strength, specifically the relationship between nervousness - sadness / nervousness - discouragement / nervousness - happy / discouragement - calm (Figure 3).




Figure 3: Networks graphs from 2020 and 2021

## Simulations

Simulations (NIRA method) identified nervousness as the most effective intervention target for Psychological Distress, while discouragement is more relevant to prevent the system from transitioning to a “pathological” state (Figures 4).




Figure 4: Alleviating (left) and Aggravating (right) simulations for 2021 network

## Limits and conclusion

We must remain cautious with the results of surveys that compare prevalences when measurement invariance is not established<sup>13</sup>. Indeed, confusion remains between 1) an increase in prevalence linked to a worsening of the situation and/or 2) a different way of answering questions due to contextual effects. Using network modeling makes it possible to represent the system's evolution without resorting to prevalence comparisons, which would imply measurement invariance. However, and regardless of the chosen modeling technique, without qualitative data or cognitive interviews, it is difficult to gain information regarding the confusion mentioned above.

Simulation techniques make it possible to identify how the system would behave in a given context and with a given intervention, thus opening the way to more targeted interventions. However, the generalist approach used here captures only a global trend that is difficult to apply to each individual<sup>14</sup>. Finally, psychotherapeutic interventions cannot target a symptom in isolation<sup>15</sup>, but rather a set of processes in complex interactions, confirming the choice of complex systems-based modeling techniques<sup>16-18</sup>.

## References

- Prati, G. & Mancini, A. D. The psychological impact of COVID-19 pandemic lockdowns: A review and meta-analysis of longitudinal studies and natural experiments. *Psychol Med* 51, 201-211 (2021).
- Belghith, F., Couto, M.-P., Ferry, O., Morvan, Y. & Patros, T. Une année seuls ensemble. Enquête sur les effets de la crise sanitaire sur l'année universitaire 2020-2021. (2021).
- Jia, Q. et al. Mental health among medical students during COVID-19: A systematic review and meta-analysis. *Front Psychol* 13, 846789 (2022).
- McNeish, D. & Wolf, M. G. Thinking twice about sum scores. *Behav Res* 52, 2287-2305 (2020).
- Fried, E. I. et al. Measuring depression over time . . . Or not? Lack of unidimensionality and longitudinal measurement invariance in four common rating scales of depression. *Psychol Assess* 28, 1354-1367 (2016).
- Borsboom, D. A network theory of mental disorders. *World Psychiatry* 16, 5-13 (2017).
- Lunansky, G. et al. Intervening on psychopathology networks: Evaluating intervention targets through simulations. *Methods* (2021) doi:10.1016/j.jymeth.2021.11.006.
- Lepège, A., Ecosse, E., Verdier, A. & Perneger, T. V. The french SF-36 health survey: Translation, cultural adaptation and preliminary psychometric evaluation. *Journal of Clinical Epidemiology* 51, 1013-1023 (1998).
- Rossee, Y. et al. Lavaan: Latent variable analysis. (2022).
- Epskamp, S. & Fried, E. I. *Bootnet: Bootstrap methods for various network estimation routines.* (2021).
- Isvoranu, A.-M. & Epskamp, S. Which estimation method to choose in network psychometrics? Deriving guidelines for applied researchers. *Psychological Methods* (2021) doi:10.1037/met0000439.
- Borkulo, C. D. van et al. Comparing network structures on three aspects: A permutation test. *Psychological Methods* (2022) doi:10.1037/met0000476.
- Fried, E. I., Flake, J. K. & Robinaugh, D. J. Revisiting the theoretical and methodological foundations of depression measurement. *Nat Rev Psychol* 1-11 (2022) doi:10.1038/s44159-022-00050-2.
- Beltz, A. M., Wright, A. G. C., Sprague, B. N. & Molenaar, P. C. M. Bridging the nomothetic and idiographic approaches to the analysis of clinical data. *Assessment* 23, 447-458 (2016).
- Eronen, M. I. Causal discovery and the problem of psychological interventions. *New Ideas in Psychology* 59, 100785 (2020).
- Borsboom, D., Haslbeck, J. M. B. & Robinaugh, D. J. Systems-based approaches to mental disorders are the only game in town. *World Psychiatry* 21, 420-422 (2022).
- Fried, E. I. & Robinaugh, D. J. Systems all the way down: Embracing complexity in mental health research. *BMC Medicine* 18, 205 (2020).
- Lunansky, G., Borkulo, C. D. van, Blanken, T., Cramer, A. & Borsboom, D. Bouncing back from life's perturbations: Formalizing psychological resilience from a complex systems perspective. (2022) doi:10.31234/osf.io/8vaxj.

