



**HAL**  
open science

## Combination of DES and macrocyclic host molecules: Review and perspectives

Sophie Fourmentin, Miriana Kfoury, David Landy

► **To cite this version:**

Sophie Fourmentin, Miriana Kfoury, David Landy. Combination of DES and macrocyclic host molecules: Review and perspectives. *Current opinion in green and sustainable chemistry*, 2022, 36, pp.100630. 10.1016/j.cogsc.2022.100630 . hal-04242259

**HAL Id: hal-04242259**

**<https://hal.science/hal-04242259v1>**

Submitted on 22 Jul 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0  
International License



10 **Abstract**

11  
12 Solvation in Deep Eutectic Solvents (DESs) and encapsulation in macrocycles are efficient  
13 strategies for molecule's solubilization. Recently, new systems combining DES and  
14 macrocycles are being constructed using two approaches: *a*) solubilizing hosts in preformed  
15 DES or *b*) using hosts as a DES component. This strategy could evolve in new outcomes in  
16 DES properties, more precisely provide the solvents with supramolecular capacity, and  
17 translate into interesting performances. In the present review, we will give the reader an  
18 overview of the studies accomplished in this growing research area, with a focus on the  
19 experimental perspectives that we consider to be the most important in order to enlarge  
20 knowledge on the properties and behavior of these neoteric systems.

21  
22 **Keywords:** cucurbituril, crown ether, cyclodextrin, deep eutectic solvents, low melting  
23 mixtures.




24  
25  
26

## 27 **Introduction**

28  
29 Within the perspective of green chemistry, Deep Eutectic Solvents (DESs) are gaining a  
30 rapidly growing interest and popularity as sustainable substitutes to organic solvents and ionic  
31 liquids [1,2]. DESs are considered for their negligible vapor pressures, nonflammability,  
32 thermal stability, chemical inertness, recyclability and high solubilizing ability. Such a class  
33 of emerging solvents has many desirable applications in analytical chemistry, catalysis,  
34 extraction, biomass pretreatment and conversion and gas separation. DESs are low melting  
35 mixtures (LMMs) characterized by a large depression of the melting temperature of the  
36 mixture at the eutectic point compared to the melting temperature of the separated pure  
37 constituents and their ideal liquid mixture. DESs are classified according to the constituents  
38 used in their preparation. Type I and II are formed from quaternary ammonium salts and  
39 either metal chloride or metal chloride hydrate, respectively. Type III and IV are obtained by  
40 combining either quaternary ammonium salts or metal chloride hydrate, respectively, with  
41 hydrogen bond donors. More recently, type V eutectics have been proposed. They are  
42 obtained using non-ionic hydrogen bond acceptors (HBA) and hydrogen bond donors (HBD)  
43 [3].

44 The most commonly studied DES is a blend of choline chloride and urea (1:2 molar ratio)  
45 called reline [3]. The huge panel of cheap, safe, sustainable, renewable and biodegradable  
46 materials (*e.g.* carbohydrates, polyalcohols, amino acids) that are able to form eutectic  
47 mixtures makes the possibility of creating new DESs exceedingly large. However, changing  
48 the nature and ratio of the components can influence the physical properties and behavior but  
49 do not grant the solvents with additional functionalities. Hence, it is crucial to find a new base  
50 to reinforce DESs with new features.

51 Supramolecular chemistry is one of today's main growing research areas, with great  
52 application potential in various domains (*e.g.* food, medicine, environment) [4]. It is the  
53 chemistry beyond the molecule that deals with the association of two or more chemicals  
54 through non-covalent interactions [5]. This leads to organized entities of higher complexity  
55 [6]. Host/guest chemistry, a branch of supramolecular chemistry, allows the construction of  
56 inclusion complexes with improved physicochemical properties compared to those of the free  
57 guest. Host molecules are various [7]. Their efficient functioning was traditionally  
58 accomplished in conventional media *i.e.* water or organic solvents.



Macrocycle	Cyclodextrins	Cucurbiturils	Crown ethers
Origin	Natural	Synthetic	Synthetic
Structure	Hydroxyl groups rims and hydrophobic cavity	Hydrophobic cavity and two identical carbonyl-laced portals	Macrocyclic polyethers with hydrophobic external ring
Guests	Hydrophobic guests	Hydrophobic guests	Cations
Interactions	Size complementarity	Size complementarity	Coordination with oxygen atoms
Media	Aqueous media	Aqueous media	Non polar solvents


59

60 **Figure 1. General features of the macrocycles used in the literature for the construction**  
 61 **of ternary or binary systems.**

62


63 Transition to sustainable chemistry conducted researchers to attempt studying the solubility of  
 64 macrocyclic hosts in green solvents such as DESs or to prepare LMMs using macrocyclic  
 65 hosts where theoretically these molecules can act as both functional monomers and cage  
 66 molecules. Figure 1 shows the general features of host molecules studied in or used for the  
 67 design of eutectics. The two strategies for the combination of DESs and macrocycles are  
 68 illustrated in Figure 2. Ternary systems refer to HBA + HBD + host while binary systems  
 69 refer to HBA (host) + HBD . A literature overview of the field can be summarized in Figure  
 70 3, showing a climbing interest over the last five years (keywords used: eutectic and either  
 71 macrocycle or host or cyclodextrin or crown ether or cucurbituril or calixarene).

72



73  
74  
75

**Figure 2. The two existing combination strategies of DESs and macrocyclic hosts.**



76  
77  
78  
79

**Figure 3. Number of papers related to the combination of DESs and macrocyclic host molecules associated to the countries (source: Scopus, October 2021).**

80 However, two main questions arise from these forerunner studies; what are the main driving  
81 forces of the solubilizing mechanisms of these hosts in such non-conventional solvents? Do  
82 these macrocycles still preserve their peculiarity to form host/guest inclusion complexes? In  
83 this brief opinion, we will not simply summarize the published results related to the  
84 combination of DESs and macrocycles but also try to answer these questions. We will  
85 additionally attempt to gain a better understanding of what is required to get desired  
86 formulation characteristics and good performance and to point out the gaps which need to be  
87 filled by further investigations.

88  
89

### **Solubilization of macrocycles in DESs: Ternary systems**

90 In 2016, a patent described for the first time the efficiency of solubilizing CDs in reline in  
91 order to improve the ability of this DES to solubilize Volatile Organic Compounds (VOCs)

92 [8]. Then, experimental investigations conducted in 2017 focused on the solvation of native  
93 CDs and CB[n]s in reline. These two host families are insoluble in organic solvents and suffer  
94 from a moderate solubility in water limiting their application to a specific concentration range  
95 [9]. Interestingly, the solubility of the most widely used CD,  $\beta$ -CD, was increased 55 times  
96 compared to water [10]. The absence of  $\beta$ -CD aggregates or clusters in reline was later  
97 demonstrated, proving that  $\beta$ -CDs are maintained as isolated entities even at elevated  
98 concentrations [11].

99 DESs possess a highly ordered liquid structure comprising strong hydrogen-bonding  
100 interactions [12,13]. This network might have the ability to include and stabilize high amount  
101 of macrocycles *via* extra hydrogen bonding. Molecular dynamics results showed that all reline  
102 moieties (choline, chloride, and urea) favorably interact with  $\beta$ -CD, with urea developing the  
103 most favorable dispersive interactions by extenuating inter-CD hydrophobic interactions,  
104 while still maintaining the potency to establish strong hydrogen bonds with both the  
105 constituents of DES and  $\beta$ -CD itself [11]. In other terms, urea is able to accumulate inside the  
106 hydrophobic  $\beta$ -CD cavity as well as its exterior forming two solvation layers. Since urea  
107 exerts the best solubilizing potential why not only using urea aqueous solution to enhance  $\beta$ -  
108 CD concentration? It is true that urea has much more strongly solubilizing effect towards  $\beta$ -  
109 CD in comparison to choline chloride when tested alone. But, the solubility of urea in water is  
110 limited and the very high concentration of urea found in reline cannot be attained in aqueous  
111 solutions. Replacing water with choline chloride enhances  $\beta$ -CD's stabilization by urea,  
112 leading to overall much higher  $\beta$ -CD solubility in reline at low hydration [11]. The dissolution  
113 process of  $\beta$ -CD in reline was shown to be an exothermic process, meaning that interactions  
114 between the DES and CD are highly favorable and raising temperature does not facilitate the  
115 dissolution [14].

116 However, results obtained with a macrocycle in a defined DES could not be projected to  
117 another system. If  $\beta$ -CD was perfectly soluble in choline:urea (1:2), it was insoluble  
118 in choline chloride:ethylene glycol (1:2), choline chloride:glycerol (1:2) and  
119 tetrabutylammonium chloride:lactic acid (1:2) DESs [11,15]. Thus, each system should be  
120 considered separately. Additionally, a panel of urea-based DESs should be tested in order to  
121 strengthen the role of urea in the solvation process and evaluate the potential influence of the  
122 DES partner.

123 The ability of DESs to dissolve macrocycles at higher concentrations opens up additional  
124 interesting exploitations of such hosts. Nowadays, innovative extraction techniques and  
125 catalysis are substituting well-known organic solvents with green, environment-friendly DESs

126 [16]. Researchers took profit from the ability of DESs to solubilize CDs to dope extraction  
127 media with these macrocycles aiming to boost the extraction efficiency [17–21] or use them  
128 as supramolecular catalysts in the green reaction medium [22]. Others synthesized water-  
129 soluble  $\beta$ -CD-based nanosponges in natural DESs [23,24].

130 The usefulness of the use of a macrocycle resides in its peculiar encapsulation ability. In this  
131 context, the new hybrid systems combining DES and CD would be mutually armed with the  
132 beneficial properties of each component. Deeper queries were conducted trying to answer the  
133 question: Is the high added value from CDs flow from their complexing ability? A positive  
134 answer was given in 2019 by two different techniques (static-headspace gas chromatography  
135 and NMR) [14,25]. Another experimental evidence of the formation of inclusion complexes  
136 with native CDs and derivatives in reline was obtained later [26]. However, a drastic decrease  
137 of formation constants ( $K_f$ ) values of inclusion complexes was observed in DES compared to  
138 water [14,25]. No correlation was observed between  $K_f$  values obtained in reline neither with  
139  $K_f$  in water nor with the decrease magnitude. This indicates that several mechanisms, that  
140 should be revealed, affect the complexation in reline.

141 Adding water to DESs could modify their physicochemical properties (*e.g.* viscosity, polarity)  
142 [27]. While high amount of water jeopardizes the integrity of DES hydrogen bond network, it  
143 seems like adding some water to reline preserves a residual molecular network while offering  
144 CD an aqueous-like environment with an increase in molecular rotational motion allowing it  
145 to form inclusion complexes [28]. Since the integrity of DES is still preserved in the presence  
146 of moderate amount of water, this leads to the apparition of the appellation of hydrated reline.  
147 Interestingly, it seems like the integration of water in the system is full of benefits; *a)* at low  
148 amounts it does not threaten the DES self-organization, *b)* it gives CD enough mobility to  
149 form inclusion complexes and *c)* it decreases the high viscosity of DES, permitting  
150 consequently an easy use of traditional NMR for quantitative measure of binding strength.  
151 ROESY spectrum showed that  $\beta$ -CD cavity protons interact with protons of several guests in  
152 hydrated reline such as piroxicam [28], VOCs (aniline and toluene) [25] and drugs  
153 (amitriptyline and cyclobenzaprine) [29]. NMR results also proved that  $\beta$ -CD retained its  
154 enantio-recognition ability in hydrated DES [29].

155 Yes, strong experimental findings evidenced that CDs maintain their encapsulation ability in  
156 hydrated reline.  $K_f$  values for  $\beta$ -CD/toluene in water  $287 \text{ M}^{-1}$  [30], hydrated reline  $79 \text{ M}^{-1}$  [25]  
157 and pure reline  $11 \text{ M}^{-1}$  [14] seem to support a linear correlation between the  $K_f$  and the  
158 amount of water in the system. However, to undoubtedly elucidate this tendency more

159 investigations need to be undergone with stable inclusion complexes (with higher  $K_f$  values in  
160 water) in order to increase the accuracy.

161 Electron paramagnetic resonance and spin probe studies underlined that  $\beta$ -CD and  $\gamma$ -CD did  
162 not show the same tendency to encapsulate probe molecules in DESs [26]. This indicated that  
163 one could not generalize from findings of data collected for one CD in one DES.  
164 Additionally, caution should be taken when dealing with other host molecules, more precisely  
165 CB[7]. Results showed that a small amount of DES in water ( $H_2O:DES$  20:1, V:V) was  
166 sufficient to completely obstruct guest inclusion even at a relatively large amount of  
167 macrocycle [26].

168 From an application point of view, formulation with improved solubilizing ability should also  
169 ensure sustained and controlled release. Literature lacks any data about the release profiles of  
170 entrapped guest in DES/macrocycle systems. Table 1 summarizes the topical areas of ternary  
171 systems research fields.

172

### 173 **Formation of DESs with macrocycles as a partner: Binary systems**

174 Development of binary systems that are efficient requires less assay development, less time,  
175 and less consumption of reagents than optimizing ternary formulations. However, totally  
176 shifting to binary systems could present some inconvenience. This could mainly reduce the  
177 number of possible combinations that could be made to control the properties of the medium  
178 (*e.g.* the concentration of the host) as compared to ternary systems.

179 Hydrogen bonding is one of the most important requirements to create a DES, macrocycles  
180 possessing functional groups that can be involved in hydrogen bonding could be used as  
181 DESs components. Additionally, the host/guest recognition is a kind of non-covalent  
182 interaction that could be applied to form DESs [31]. Even if the first attempts have started as  
183 early as 2005 when  $\alpha$ -CD was considered among other sugars used to prepare LMMs [32], the  
184 use of macrocycles as host partners in LMMs have not blow up before 2014 [33,34]. In 2020,  
185 the era of SUPRADES (supramolecular DES) was officially launched [35,36]. Table 2  
186 summarizes all DESs or LMMs based on macrocycles.

187

188 **Table 1. The topical areas of ternary systems research fields.**

Research field	Macrocycle	DES	Reference
Absorption process of gaseous effluents by combining macrocycles and DESs	$\beta$ -CD, RAMEB	Choline chloride:urea	[8]
Solubilization of macrocycles in DESs	$\alpha$ -CD, $\beta$ -CD, $\gamma$ -CD CB[6], CB[7], CB[8]	Choline chloride:urea	[9]
Experimental and computational structural study of the dissolution mechanism of macrocycles in DESs	$\beta$ -CD	Choline chloride:urea	[10,11]
Investigation of the thermodynamic properties of macrocycles dissolution in DESs	$\beta$ -CD	Choline chloride:urea	[14]
Study of the complexation mechanism with or without water	$\alpha$ -CD, $\beta$ -CD, $\gamma$ -CD, HP- $\beta$ -CD, and CRYSMEB	Choline chloride:urea	[14,25,26,28,29]
	Methyl- $\beta$ -CD	Glycine:glycerol	[17]
	$\beta$ -CD	Ammonium acetate:L-lactic acid	[18]
	HP- $\beta$ -CD	Sodium acetate:glycerol	[19]
New extraction media combining macrocycles and DESs	$\beta$ -CD	Nicotinamide:L-lactic acid Choline chloride:L-lactic acid Ammonium acetate:L-lactic acid Sodium acetate: L-lactic acid L-alanine:L-lactic acid Glycine:L-lactic acid	[20]
New catalytic media combining macrocycles and DESs	$\alpha$ -CD, $\beta$ -CD, $\gamma$ -CD, HP- $\beta$ -CD, and MCD	17 NADES	[22]
Synthesis of macrocycles-based polymers in DESs	$\beta$ -CD	Choline chloride:citric acid	[23]

**Table 2. DESs or LMMs prepared using macrocycles.**

Macrocycle	% (wt)	Partner	% (wt)	Liquid at RT	Reference
$\alpha$ -CD	30	DMU	70	No	[32]
$\beta$ -CD/ CRYSMEB/ RAMEB/ TRIMEB/ HP- $\beta$ -CD/ HTMAP- $\beta$ -CD	30	DMU	70	No	[33]
$\alpha$ -CD/ $\beta$ -CD/ $\gamma$ -CD HP- $\alpha$ -CD/ HP- $\beta$ -CD/ HP- $\gamma$ -CD/ TRIMEB/ TRIAC- $\beta$ -CD/ CRYSMEB/ RAMEB	30		70		
$\beta$ -CD/ CRYSMEB/ RAMEB	20	DMU	80	No	[34]
$\beta$ -CD/ CRYSMEB/ RAMEB	50		50		
$\beta$ -CD/ CRYSMEB/ RAMEB	70		30		
CRYSMEB/ RAMEB	80		20		
HP- $\beta$ -CD/ CRYSMEB/ RAMEB/ Captisol®	30	Levulinic acid	70	Yes	[36]
$\beta$ -CD/ RAMEB/ TRIMEB / HP- $\beta$ -CD	20	L-lactic acid	80	Yes	[15]
	3	PEG 200	97		
15-crown-5:NaOH	1.5	PEG 400	98.5		
	1	PEG 600	99	Yes	[37]
	3.5	PEG 200	96.5		
18-crown-6:KOH	2	PEG 400	98		
	1	PEG 600	99		

191 DMU: *N,N'*-dimethylurea; PEG: Polyethyleneglycol; Captisol®: Sulfobutylether- $\beta$ -cyclodextrin; CRYSMEB: Low methylated- $\beta$ -cyclodextrin;  
 192 HP- $\alpha$ -CD: 2-Hydroxypropyl- $\alpha$ -cyclodextrin; HP- $\beta$ -CD: 2-Hydroxypropyl- $\beta$ -cyclodextrin; HP- $\gamma$ -CD: 2-Hydroxypropyl- $\gamma$ -cyclodextrin;  
 193 HTMAP- $\beta$ -CD: 2-Hydroxy-3-(trimethylammonio)propyl- $\beta$ -cyclodextrin; RAMEB: Randomly methylated- $\beta$ -cyclodextrin; TRIMEB: Trimethyl-  
 194  $\beta$ -cyclodextrin; TRIAC- $\beta$ -CD: Triacetyl- $\beta$ -cyclodextrin.

195 A series of LMMs based on native CDs ( $\alpha$ -CD,  $\beta$ -CD and  $\gamma$ -CD) as well as  $\beta$ -CD derivatives  
196 and *N,N'*-dimethylurea was successfully evaluated as inert catalytic solvents [33,34].  
197 Interestingly, the yield of 1-decene hydroformylation varied significantly according to the  
198 nature of CD incorporated in the LMMs [33]. This proved that CDs are not a dump partner in  
199 the green solvent. Surprisingly, the values of conversion did not follow the size of the cavity  
200 but were affected by the type and degree of CD substitution as well as solute solubility and  
201 solvent viscosity [34]. Still, the aim was not reached since all these new solvents have a  
202 melting point higher than 67 °C, thus they are not liquid at ambient temperature.

203 SUPRADES, DESs with supramolecular properties were then introduced [35]. Interestingly,  
204 the new RAMEB:levulinic acid system was liquid at room temperature. 2D ROESY (Rotating  
205 frame Overhauser Enhancement Spectroscopy) and DOSY (Diffusion ordered spectroscopy)  
206 experiments confirmed the interaction between the studied guest and the CD cavity protons  
207 while no interaction was detected with levulinic acid. This constituted the first experimental  
208 evidence that the solubilized guest is embedded within the host cavity. Then, in addition to  
209 this SUPRADES, three other LMMs, HP- $\beta$ -CD:levulinic acid, CRYSMEB:levulinic acid and  
210 Captisol®:levulinic acid, were successfully obtained [36]. They showed distinct storage and  
211 thermal stability as well as physicochemical properties (density, viscosity) depending on the  
212 type of CD. The viscosity of levulinic acid based SUPRADES were around 200, 400, 600 and  
213 1400 mPa s at 30 °C when respectively using RAMEB, CRYMEB, HP- $\beta$ -CD and Captisol®  
214 for the construction of the solvents. Interestingly, these viscosity values were shown to  
215 significantly decrease with temperature [34]. The effect of the host molecule on the viscosity  
216 of the solvents was also observed when using dimethylurea as a partner for LMM formation  
217 [31]. Authors showed that adding water efficiently decreases the viscosity of these solvents.

218 On the other hand, El Achkar et al. reported for the first time the ability of a CD-based LMMs  
219 to delay the release of a solute over time. Other SUPRADES, comprised of CDs and lactic  
220 acid were obtained and tested for their efficacy to extract aromatic organic pollutants from  
221 aqueous solutions [15].

222 Beside CDs, crown ethers took also their share in the preparation of DESs [31,37]. Crown  
223 ether based DESs showed desirable characteristics, more precisely low viscosity and melting  
224 point. They were evaluated for their ability to extract *N*-compounds from fuel oils. Active  
225 hydrogen-bonding interaction was shown to be formed between the macrocycle and solute  
226 while no evidence proved the encapsulation of extracts in crown ethers [37].

227 Generally, titration experiments, where the concentration of the host is varied, are applied to  
228 obtain  $K_f$  for host/guest inclusion complexes. When incorporated in DESs, hosts are available

229 in large amounts in the medium (>20%) but their concentration cannot be varied. Thus, new  
230 methods should be envisioned allowing the evaluation of the complexing ability of a  
231 macrocycle incorporated in a LMM. Additionally, a systematic and clear comparison of the  
232 performance of host-based DESs and host solubilized in DESs should be realized. This would  
233 clarify whether all the advantages deriving from the increased amount of hosts achieved in  
234 DESs could be or not counterbalanced by a decrease in the host properties of macrocycles. In  
235 host-based DES, macrocycles may be more toughly engaged in a strong hydrogen bond  
236 network that might further reduce their encapsulation performance. Also, the effect of water  
237 addition should be investigated.

238

### 239 **Conclusions and future perspective**

240

241 The combination of novel deep eutectic systems and historical supramolecular chemistry  
242 fields is currently an exciting scientific challenge. While pursuing the development and  
243 optimization of innovative ternary and binary systems with promising and various application  
244 fields, it will be important for future work to address deeper fundamental issues. Despite  
245 numerous efforts executed in order to solubilize and elucidate the solubilization mechanism of  
246 macrocycles in DESs or to prepare new host-based systems, the factors needed to elaborate  
247 the optimal formulation are still not fully discovered.

248 Authors are encouraged to reveal the requirements needed to develop any binary system  
249 because it becomes clear that hydrogen bonding is not the only must; a DES could not be  
250 obtained using all hydrogen bond acceptor and donor couples. Various characteristics of the  
251 building blocks should be considered (structure, mesomeric effect, hygroscopy, etc.).  
252 Additionally when using macrocycles, like CDs for example, for the construction of eutectics,  
253 different hosts do not form a DES with a same partner. Also, authors should give more  
254 insights on the interaction mechanisms between hosts and DESs components and different  
255 factors that influence the encapsulation properties (binding strength, isomers discrimination,  
256 release) of macrocycles either when they are dissolved in DESs or used as partners.

257 On the other hand, these systems are growing in order to achieve safety and sustainability.  
258 However, information on their storage stability, environmental degradation and toxicity is still  
259 missed. Even if researchers take critical care in the selection of the building blocks, the  
260 system safety statement should be proved. Completing fundamental and toxicological  
261 investigations would boost the scaling up of DES/macrocycle systems.

262

263

264 **Conflict of interest statement**

265 Nothing declared.

266

267 **References**

268

269 Papers of particular interest, published within the period of review, have been highlighted as:

270 \* of special interest

271 \*\* of outstanding interest

272


- 273 [1] F.M. Perna, P. Vitale, V. Capriati, Deep eutectic solvents and their applications as  
274 green solvents, *Curr. Opin. Green Sustain. Chem.* 21 (2020) 27–33.  
275 <https://doi.org/10.1016/j.cogsc.2019.09.004>.
- 276 [2] A. Paiva, A.A. Matias, A.R.C. Duarte, How do we drive deep eutectic systems towards  
277 an industrial reality?, *Curr. Opin. Green Sustain. Chem.* 11 (2018) 81–85.  
278 <https://doi.org/10.1016/j.cogsc.2018.05.010>.
- 279 [3] A.P. Abbott, G. Capper, D.L. Davies, R.K. Rasheed, V. Tambyrajah, Novel solvent  
280 properties of choline chloride/urea mixtures, *Chem. Commun.* (2003) 70–71.  
281 <https://doi.org/10.1039/b210714g>.
- 282 [4] N. Morin-Crini, S. Fourmentin, É. Fenyvesi, E. Lichtfouse, G. Torri, M. Fourmentin,  
283 G. Crini, 130 years of cyclodextrin discovery for health, food, agriculture, and the  
284 industry: a review, *Environ. Chem. Lett.* (2021). [https://doi.org/10.1007/s10311-020-](https://doi.org/10.1007/s10311-020-01156-w)  
285 [01156-w](https://doi.org/10.1007/s10311-020-01156-w).
- 286 [5] J.-M. Lehn, *Supramolecular Chemistry-Scope and Perspectives Molecules,*  
287 *Supermolecules, and Molecular Devices (Nobel Lecture)*, *Angew. Chemie Int. Ed.*  
288 *English.* 27 (1988) 89–112. <https://doi.org/10.1002/anie.198800891>.
- 289 [6] J.-M. Lehn, From supramolecular chemistry towards constitutional dynamic chemistry  
290 and adaptive chemistry, *Chem. Soc. Rev.* 36 (2007) 151–160.  
291 <https://doi.org/10.1039/B616752G>.
- 292 [7] J.-M. Lehn, *Supramolecular Chemistry: Concepts and perspectives*, Wiley-VCH,  
293 Weinheim, 1995.
- 294 [8]\*\* S. Fourmentin, D. Landy, L. Moura, S. Tilloy, H.H. Bricout, M. Ferreira, Procédé  
295 d'épuration d'un effluent gazeux, FR3058905B1. (2016).  
296 This patent considered for the first time the solubilization of CD in DES and more  
297 specifically the efficiency of this combination in the entrapment and retention of  
298 volatile organic compounds (VOCs).
- 299 [9] J.A. McCune, S. Kunz, M. Olesińska, O.A. Scherman, DESolution of CD and CB  
300 macrocycles, *Chem. - A Eur. J.* 23 (2017) 8601–8604.  
301 <https://doi.org/10.1002/chem.201701275>.
- 302 [10]\* A. Triolo, F. Lo Celso, O. Russina, Structural features of  $\beta$ -cyclodextrin solvation in  
303 the deep eutectic solvent, *reline*, *J. Phys. Chem. B.* 124 (2020) 2652–2660.  
304 <https://doi.org/10.1021/acs.jpcc.0c00876>.  
305 This article made certain the capacity of reline to solubilize  $\beta$ -CD at very high  
306 concentrations with no clustering or aggregations.  $\beta$ -CD was maintained as isolated  
307 entities as proved by the SAXS (Small angle X-ray scattering) technique.
- 308 [11]\*\* I. Shumilin, D. Harries, Cyclodextrin solubilization in hydrated reline: Resolving the  
309 unique stabilization mechanism in a deep eutectic solvent, *J. Chem. Phys.* 154 (2021)

- 310 224505. <https://doi.org/10.1063/5.0052537>.
- 311 This work showed the importance of the use of DES, more precisely, reline and not  
312 only simple urea-in-water solutions to solubilize  $\beta$ -CD. It indicated that both partners  
313 of DES are necessary as a) they synergistically enhance each other's stabilization effect  
314 on  $\beta$ -CD and b) the very high urea concentration found in reline cannot be attained in  
315 aqueous urea solutions as its solubility in water is limited.
- 316 [12] O.S. Hammond, D.T. Bowron, K.J. Edler, Liquid structure of the choline chloride-urea  
317 deep eutectic solvent (reline) from neutron diffraction and atomistic modelling, *Green*  
318 *Chem.* 18 (2016) 2736–2744. <https://doi.org/10.1039/C5GC02914G>.
- 319 [13] Y. Zhang, D. Poe, L. Heroux, H. Squire, B.W. Doherty, Z. Long, M. Dadmun, B.  
320 Gurkan, M.E. Tuckerman, E.J. Maginn, Liquid Structure and Transport Properties of  
321 the Deep Eutectic Solvent Ethaline, *J. Phys. Chem. B.* 124 (2020) 5251–5264.  
322 <https://doi.org/10.1021/acs.jpcc.0c04058>.
- 323 [14]\*\*T. Moufawad, L. Moura, M. Ferreira, H. Bricout, S. Tilloy, E. Monflier, M. Costa  
324 Gomes, D. Landy, S. Fourmentin, First evidence of cyclodextrin inclusion complexes  
325 in a deep eutectic solvent, *ACS Sustain. Chem. Eng.* 7 (2019) 6345–6351.  
326 <https://doi.org/10.1021/acssuschemeng.9b00044>.
- 327 This article was the first quantitative evidence of the formation of CD inclusion  
328 complexes in DES. Formation constants values for several complexes were  
329 successfully determined using UV–Visible spectroscopy and static headspace gas  
330 chromatography.
- 331 [15] M.Q. Farooq, V.R. Zeger, J.L. Anderson, Comparing the Extraction Performance of  
332 Cyclodextrin-containing Supramolecular Deep Eutectic Solvents Versus Conventional  
333 Deep Eutectic Solvents by Headspace Single Drop Microextraction, *J. Chromatogr. A.*  
334 1658 (2021) 462588. <https://doi.org/10.1016/j.chroma.2021.462588>.
- 335 [16] L. Nakhle, M. Kfoury, I. Mallard, D. Landy, H. Greige-Gerges, Microextraction of  
336 bioactive compounds using deep eutectic solvents: a review, *Environ. Chem. Lett.* 19  
337 (2021) 3747–3759. <https://doi.org/10.1007/s10311-021-01255-2>.
- 338 [17] V. Athanasiadis, S. Grigorakis, S. Lalas, D.P. Makris, Methyl  $\beta$ -cyclodextrin as a  
339 booster for the extraction for *Olea europaea* leaf polyphenols with a bio-based deep  
340 eutectic solvent, *Biomass Convers. Biorefinery.* 8 (2018) 345–355.  
341 <https://doi.org/10.1007/s13399-017-0283-5>.
- 342 [18] D. Chakroun, S. Grigorakis, S. Loupassaki, D.P. Makris, Enhanced-performance  
343 extraction of olive (*Olea europaea*) leaf polyphenols using L-lactic acid/ammonium  
344 acetate deep eutectic solvent combined with  $\beta$ -cyclodextrin: screening, optimisation,  
345 temperature effects and stability, *Biomass Convers. Biorefinery.* (2019).  
346 <https://doi.org/10.1007/s13399-019-00521-2>.
- 347 [19] I. Karageorgou, S. Grigorakis, S. Lalas, I. Mourtzinos, D.P. Makris, Incorporation of 2-  
348 hydroxypropyl  $\beta$ -cyclodextrin in a biomolecule-based low-transition temperature  
349 mixture (LTTM) boosts efficiency of polyphenol extraction from *Moringa oleifera*  
350 *Lam* leaves, *J. Appl. Res. Med. Aromat. Plants.* 9 (2018) 62–69.  
351 <https://doi.org/10.1016/j.jarmap.2018.02.005>.
- 352 [20] C. Georgantzi, A.-E. Lioliou, N. Paterakis, D.P. Makris, Combination of lactic acid-  
353 based deep eutectic solvents (DES) with  $\beta$ -cyclodextrin: performance screening using  
354 ultrasound-assisted extraction of polyphenols from selected native greek medicinal  
355 plants, *Agronomy.* 7 (2017) 54. <https://doi.org/10.3390/agronomy7030054>.
- 356 [21] V. Athanasiadis, S. Grigorakis, S. Lalas, D.P. Makris, Stability effects of methyl  $\beta$ -  
357 cyclodextrin on *Olea europaea* leaf extracts in a natural deep eutectic solvent, *Eur.*  
358 *Food Res. Technol.* 244 (2018) 1783–1792. [https://doi.org/10.1007/s00217-018-3090-](https://doi.org/10.1007/s00217-018-3090-8)  
359 8.

- 360 [22] L. Xiong, X. Kong, H. Liu, P. Wang, Efficient biosynthesis of (S)-1-[2-  
361 (trifluoromethyl)phenyl]ethanol by a novel isolate *Geotrichum silvicola* ZJPH1811 in  
362 deep eutectic solvent/cyclodextrin-containing system, *Bioresour. Technol.* 329 (2021)  
363 124832. <https://doi.org/10.1016/j.biortech.2021.124832>.
- 364 [23] C. Cecone, G. Hoti, I. Krabicová, S.L. Appleton, F. Caldera, P. Bracco, M. Zanetti, F.  
365 Trotta, Sustainable synthesis of cyclodextrin-based polymers by exploiting natural  
366 deep eutectic solvents, *Green Chem.* 22 (2020) 5806–5814.  
367 <https://doi.org/10.1039/d0gc02247k>.
- 368 [24] A. Rubin Pedrazzo, F. Trotta, G. Hoti, F. Cesano, M. Zanetti, Sustainable  
369 mechanochemical synthesis of  $\beta$ -cyclodextrin polymers by twin screw extrusion,  
370 *Environ. Sci. Pollut. Res.* (2021). <https://doi.org/10.1007/s11356-021-15187-5>.
- 371 [25]\*\*M.E. Di Pietro, G. Colombo Dugoni, M. Ferro, A. Mannu, F. Castiglione, M. Costa  
372 Gomes, S. Fourmentin, A. Mele, Do cyclodextrins encapsulate volatiles in deep  
373 eutectic systems?, *ACS Sustain. Chem. Eng.* 7 (2019) 17397–17405.  
374 <https://doi.org/10.1021/acssuschemeng.9b04526>.  
375 This article showed how different NMR techniques can contribute to study structural  
376 and supramolecular features of DES/ $\beta$ -CD systems. It proved that  $\beta$ -CD preserves its  
377 supramolecular peculiarities in both hydrated and non hydrated reline with a higher  
378 encapsulation efficiency in the latter.
- 379 [26] A. Cantelli, P. Franchi, E. Mezzina, M. Lucarini, Determination of binding strengths of  
380 host-guest complexes in deep eutectic solvents using spin probe methodology,  
381 *ChemPhysChem.* 22 (2021) 517–521. <https://doi.org/10.1002/cphc.202001007>.
- 382 [27] T. El Achkar, H. Greige-Gerges, S. Fourmentin, Basics and properties of deep eutectic  
383 solvents: a review, *Environ. Chem. Lett.* (2021). [https://doi.org/10.1007/s10311-021-](https://doi.org/10.1007/s10311-021-01225-8)  
384 01225-8.
- 385 [28]\*\*G.C. Dugoni, M.E. Di Pietro, M. Ferro, F. Castiglione, S. Ruellan, T. Moufawad, L.  
386 Moura, M. Costa Gomes, S. Fourmentin, A. Mele, Effect of water on deep eutectic  
387 solvent/ $\beta$ -cyclodextrin systems, *ACS Sustain. Chem. Eng.* 7 (2019) 7277–7285.  
388 <https://doi.org/10.1021/acssuschemeng.9b00315>.  
389 This article evaluated the impact of water addition on DES/ $\beta$ -CD systems.  
390 Interestingly, it proved that water strengthens the individual features of each  
391 component of the system DES/ $\beta$ -CD/H<sub>2</sub>O: it provides  $\beta$ -CD with enough mobility to  
392 form inclusion complexes with solutes while maintaining a residual DES structure.
- 393 [29] M.E. Di Pietro, M. Ferro, A. Mele, Drug encapsulation and chiral recognition in deep  
394 eutectic solvents/ $\beta$ -cyclodextrin mixtures, *J. Mol. Liq.* 311 (2020) 113279.  
395 <https://doi.org/10.1016/j.molliq.2020.113279>.
- 396 [30] M. Kfoury, D. Landy, S. Fourmentin, Characterization of cyclodextrin/volatile  
397 inclusion complexes: A review, *Molecules.* 23 (2018) 1204.  
398 <https://doi.org/10.3390/molecules23051204>.
- 399 [31] D. Yu, H. Mou, H. Fu, X. Lan, Y. Wang, T. Mu, “Inverted” deep eutectic solvents  
400 based on host-guest interactions, *Chem. – An Asian J.* 14 (2019) 4183–4188.  
401 <https://doi.org/10.1002/asia.201901365>.
- 402 [32] G. Imperato, E. Eibler, J. Niedermaier, B. König, Low-melting sugar-urea-salt  
403 mixtures as solvents for Diels–Alder reactions, *Chem. Commun.* (2005) 1170–1172.  
404 <https://doi.org/10.1039/B414515A>.
- 405 [33] F. Jérôme, M. Ferreira, H. Bricout, S. Menuel, E. Monflier, S. Tilloy, Low melting  
406 mixtures based on  $\beta$ -cyclodextrin derivatives and N,N'-dimethylurea as solvents for  
407 sustainable catalytic processes, *Green Chem.* 16 (2014) 3876–3880.  
408 <https://doi.org/10.1039/c4gc00591k>.

- 409 [34] M. Ferreira, F. Jérôme, H. Bricout, S. Menuel, D. Landy, S. Fourmentin, S. Tilloy, E.  
410 Monflier, Rhodium catalyzed hydroformylation of 1-decene in low melting mixtures  
411 based on various cyclodextrins and N,N'-dimethylurea, *Catal. Commun.* 63 (2015) 62–  
412 65. <https://doi.org/10.1016/j.catcom.2014.11.001>.
- 413 [35]\*\*T. El Achkar, T. Moufawad, S. Ruellan, D. Landy, H. Greige-Gerges, S. Fourmentin,  
414 Cyclodextrins: From solute to solvent, *Chem. Commun.* 56 (2020) 3385–3388.  
415 <https://doi.org/10.1039/d0cc00460j>.  
416 This study opened up new orisons as it announced a new generation of SUPRADES  
417 (DES based on CDs): liquid at room temperature and reinforced with supramolecular  
418 properties. CD incorporated in the solvent maintained its encapsulation properties as  
419 undoubtedly evidenced by NMR results.
- 420 [36] T. El Achkar, L. Moura, T. Moufawad, S. Ruellan, S. Panda, S. Longuemart, F.X.  
421 Legrand, M. Costa Gomes, D. Landy, H. Greige-Gerges, S. Fourmentin, New  
422 generation of supramolecular mixtures: Characterization and solubilization studies, *Int.*  
423 *J. Pharm.* 584 (2020) 119443. <https://doi.org/10.1016/j.ijpharm.2020.119443>.
- 424 [37] M.C. Ali, R. Liu, J. Chen, T. Cai, H. Zhang, Z. Li, H. Zhai, H. Qiu, New deep eutectic  
425 solvents composed of crown ether, hydroxide and polyethylene glycol for extraction of  
426 non-basic N-compounds, *Chinese Chem. Lett.* 30 (2019) 871–874.  
427 <https://doi.org/10.1016/j.ccllet.2019.02.025>.  
428

429 **Graphical abstract**  
430  
431



432