



HAL
open science

The phosphatidylserine receptor TIM1 promotes infection of enveloped hepatitis E virus

Laura Corneillie, Irma Lemmens, Claire Montpellier, Martin Ferrié, Karin Weening, Freya van Houtte, Xavier Hanouille, Laurence Cocquerel, Ali Amara, Jan Tavernier, et al.

► To cite this version:

Laura Corneillie, Irma Lemmens, Claire Montpellier, Martin Ferrié, Karin Weening, et al.. The phosphatidylserine receptor TIM1 promotes infection of enveloped hepatitis E virus. *Cellular and Molecular Life Sciences*, 2023, 80 (11), pp.326. 10.1007/s00018-023-04977-4 . hal-04245784

HAL Id: hal-04245784

<https://hal.science/hal-04245784>

Submitted on 19 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

The phosphatidylserine receptor TIM1 promotes infection of enveloped hepatitis E virus

Laura Corneillie^{1,*}, Irma Lemmens², Claire Montpellier³, Martin Ferrié³, Karin Weening¹, Freya Van Houtte¹, Xavier Hanouille⁴, Laurence Cocquerel³, Ali Amara⁵, Jan Tavernier², Philip Meuleman^{1,*}

Affiliations

¹Laboratory of Liver Infectious Diseases, Department of Diagnostic Sciences, Faculty of Medicine and Health Sciences, Ghent University, C. Heymanslaan 10, Ghent, Belgium

²VIB-UGent Center for Medical Biotechnology, Department of Biomolecular medicine, Faculty of Medicine and Health Sciences, Ghent University, Zwijnaarde 75, Ghent, Belgium

³Univ. Lille, CNRS, INSERM, CHU Lille, Institut Pasteur de Lille, U1019 - UMR 8204 - CIIL- Center for Infection and Immunity of Lille, 1 Rue du Professeur Calmette, Lille, France

⁴Univ. Lille, Inserm, CHU Lille, Institut Pasteur Lille, U1167 - RID-AGE - Facteurs de risque et déterminants moléculaires des maladies liées au vieillissement, F-59000, Lille, France.; CNRS, EMR9002 - BSI - Integrative Structural Biology, 59000 Lille, France

⁵Université de Paris Cité, INSERM U944, CNRS UMR 7212, Institut de Recherche Saint-Louis, Hôpital Saint-Louis, 75010 Paris, France

*Corresponding authors

Laura Corneillie

Prof. Dr. Philip Meuleman

Laboratory of Liver Infectious Diseases (LLID),

Department of Diagnostic Sciences, Faculty of Medicine and Health Sciences,

Ghent University.

Building MRBII, Corneel Heymanslaan 10, B-9000 Ghent, Belgium.

E-mail: laura.corneillie@ugent.be; philip.meuleman@ugent.be;

Telephone: +32 9 332 02 05

30 <https://orcid.org/0000-0003-3937-7000>; <https://orcid.org/0000-0001-6821-234X>; [https://orcid.org/0000-0001-](https://orcid.org/0000-0001-6821-234X)
31 [6821-234X](https://orcid.org/0000-0001-6821-234X)

32

33

34 **Acknowledgements**

35 We would like to thank Suzanne U. Emerson (NIH, United States) for providing the p6 clone, Takaji Wakita for
36 the 83-2-27 clone; Jane McKeating for HEK293T cells and Charles M. Rice (The Rockefeller University, USA)
37 for Huh-7.5 cells. We acknowledge the CORE Flow Cytometry from the Faculty of Medicine and Health Sciences
38 at Ghent University for use and support of the flow cytometry and sorting devices.

39

Abstract

41 The hepatitis E virus (HEV) is an underestimated RNA virus of which the viral life cycle and pathogenicity remain
42 partially understood and for which specific antivirals are lacking. The virus exists in two forms: nonenveloped
43 HEV that is shed in feces and transmits between hosts; and membrane-associated, quasi-enveloped HEV that
44 circulates in the blood. It is suggested that both forms employ different mechanisms for cellular entry and
45 internalization but little is known about the exact mechanisms. Interestingly, the membrane of enveloped HEV is
46 enriched with phosphatidylserine, a natural ligand for the T-cell immunoglobulin and mucin domain-containing
47 protein 1 (TIM1) during apoptosis and involved in ‘apoptotic’ mimicry, a process hijacked by viruses to promote
48 infection. We here investigated the role of TIM1 in the entry process of HEV. We determined that HEV infection
49 with particles derived from culture supernatant, which are cloaked by host-derived membranes (eHEV), was
50 significantly impaired after knockout of TIM1, whereas infection with intracellular HEV particles (iHEV) was
51 unaffected. eHEV infection was restored upon TIM1 expression; and enhanced after ectopic TIM1 expression.
52 The significance of TIM1 during entry was further confirmed by viral binding assay, and point mutations of the
53 PS-binding pocket diminished eHEV infection. Additionally, Annexin V, a PS-binding molecule also significantly
54 reduced infection. Taken together, our findings support a role for TIM1 in eHEV-mediated cell entry, facilitated
55 by the PS present on the viral membrane, a strategy HEV may use to promote viral spread throughout the infected
56 body.

57

Keywords

59 Hepatitis E virus; viral hepatitis; T-Cell Immunoglobulin and Mucin Domain-Containing Protein 1; HAVCR1;
60 virus-host interaction; viral entry; cell surface receptor; apoptotic mimicry

61

62 1. Introduction

63 The hepatitis E virus (HEV) is annually responsible for infection of at least 20 million people and is increasingly
64 recognized as a medical problem in the Western world [1-3]. Most of the human cases are attributed to 4 viral
65 genotypes (gts) that differ in geographical distribution, routes of transmission, disease course and pathogenicity.
66 Gt-1 and gt-2 transmit fecal-orally and are sources of outbreaks in developing countries with high mortality seen
67 in gt-1 infected pregnant women [4]. Gt-3 and gt-4 are zoonotic and predominant in industrialized countries and
68 chronicity has been observed in gt-3 infected immunocompromised patients [5]. No HEV-specific therapy exists,
69 necessitating the identification and development of novel and more effective therapies [1].

70 HEV is a positive-sense, single-stranded RNA virus with a 7.2 kb genome. Depending on viral gt, the viral genome
71 comprises three or four open reading frames (ORF1-4). ORF1 encodes a nonstructural polyprotein, comprised of
72 several domains essential for viral replication [6]. ORF2 and ORF3 are encoded from a bicistronic subgenomic
73 RNA. ORF2 encodes the viral capsid protein, binds to the host cells and elicits an antibody response [7]. ORF3
74 encodes a small phosphoprotein important for viral release. The more recent discovered ORF4 is only expressed
75 in gt-1 HEV and influences viral replication by interacting with both host and viral proteins [8].

76 HEV has long been neglected as it seemed a virus confined to developing countries, and the lack of appropriate
77 cell culture systems as well as animal models made it difficult to study this virus in detail. Recent advances in this
78 field have been made but parts of the viral life cycle remain poorly understood, such as viral entry. The virus is
79 considered quasi-enveloped, but in fact different forms of the viral particle exist. Non-enveloped virus particles
80 are retrieved from feces and are the major form present in (intra)cellular lysate (designated iHEV), whereas HEV
81 particles circulating in the blood stream and cell culture supernatant are cloaked with membrane lipids, therefore
82 considered enveloped (eHEV) [9]. Naked viral particles mediate HEV spread in the population, whereas enveloped
83 viral particles are presumably more important for viral spread within the body. Experimental evidence suggests
84 that both forms have distinct ways of entering a host cell and understanding both of these processes is necessary
85 for insight into viral pathogenesis and developing highly needed antivirals [10]. Although several host factors have
86 been shown to be involved in cell attachment and/or entry of iHEV, not one protein is generally accepted as the
87 true HEV receptor. The cell entry of enveloped particles is even less studied, and it seems a more inefficient
88 process compared to naked virus [9, 11]. It was previously described that eHEV particles contain
89 phosphatidylserine (PS) [11]. This negatively charged phospholipid is a marker for apoptosis and normally present
90 at the cytosolic side of the membrane. Upon apoptosis, PS is exposed and binds to PS-receptors, thereby
91 modulating phagocytosis. This process is hijacked by a variety of viruses for their entry and is termed 'apoptotic

92 mimicry' [12]. Not only enveloped viruses such as Ebola virus, Dengue virus or Marburg virus use this process
93 but also hepatitis A virus (HAV) has been shown to bind the PS-receptor TIM1 (T cell immunoglobulin mucin
94 domain 1) [13-16]. HEV is similar to HAV in that they both circulate as quasi-enveloped viruses [17]. The presence
95 of PS in the eHEV particle might bind PS-receptors such as TIM1 and thereby mediate eHEV entry, but this has
96 not previously been experimentally determined. In this study, we explored the role of TIM1 by using CRISPR-
97 Cas9 technology, as well as TIM1 rescue and ectopic expression. We provide evidence that TIM1 mediates the
98 infection of enveloped HEV but not intracellular HEV and facilitates eHEV internalization. Moreover, we show
99 that PS present in viral membrane is crucial in this process. Our results suggest that TIM1 acts as an entry cofactor
100 for eHEV.

101

102

103 **2. Materials & Methods** – (A complete overview of the materials and methods can be found in the online
104 supplement)

105

106 2.1 Virus production and infection

107 The gt-3 HEV Kernow-C1 Passage 6 (p6) plasmid (GenBank accession number JQ679013) was linearized with
108 *MluI* (New England Biolabs), the gt-3 HEV 83-2-27 plasmid (GenBank accession number AB740232) was
109 linearized with *HindIII*. Capped RNA was produced either using T7 RiboMAX Express Large Scale RNA
110 production (Promega), followed by ScriptCap m7G capping system (Cellsript) or using T7 mMACHINE
111 mMACHINE kit (Ambion).

112 Capped RNAs were electroporated into PLC3 using a Gene Pulser (Bio-Rad).
113 Extracellular viral particles (eHEV) were harvested from centrifuged and filtered (0.22 µm) supernatant twice a
114 week, and concentrated using Amicon Ultra Centrifugal Filters (Merck). Lysis of transfected cells was performed
115 to release intracellular viral particles (iHEV) by 3 freeze-thaw cycles in distilled water, followed by addition of
116 10X phosphate-buffered saline (PBS) and centrifugation at 10.000 x g for 30 minutes to separate cell debris. A
117 detergent/protease-treated viral preparation (tHEV) was made by treating culture supernatant with 0.1% trypsin
118 and 0.1% sodium deoxycholic acid at 37°C for 1 hour.

119 To determine the HEV infectivity titer, Huh-7.5 cells were seeded in Poly-L-Lysine-coated black 96 well plates
120 and exposed to serial dilutions of respectively eHEV, iHEV and tHEV preparations. Three (p6) or five (83-2-27)
121 days after infection, indirect immunofluorescence staining was used to check for HEV ORF2 positive cells.

122

123 2.2 Neutralization assays

124 Huh-7.5 cells were pre-incubated for 30 minutes with a polyclonal anti-TIM1 (AF1750, R&D systems) or isotype
125 control (R&D systems) antibody at concentrations of 1 µg/ml, 3.3 µg/mL and 10 µg/mL after which eHEV or
126 iHEV viral particles were added to the cell culture. Culture media was refreshed and antibodies were replenished.
127 After three days, infection was quantified as described. In order to monitor cell viability under antibody
128 neutralization, a MTT assay (Roche) was performed according to manufacturer's instructions. Cells were
129 incubated with the indicated amounts of anti-TIM1 or isotype control antibody or left untreated. After incubation
130 time, cells were incubated with 10 µL MTT solution, solubilization solution was added after 4 hours and incubated

131 overnight. Absorbance was measured by ELISA and cell viability of anti-TIM1 or isotype control were normalized
132 to the untreated wells.

133

134 *2.3 TIM1 rescue and ectopic expression*

135 The TIM1 expressing plasmid or empty vector control (cloning information in supplemental information) were
136 transfected into Huh-7.5^{TIM1-KO} or HEK293T cells, seeded in 96 well-plates, using Lipofectamine 2000 (Thermo
137 Fisher Scientific). Cells were exposed to eHEV or iHEV 24 hours post transfection. Three days after infection
138 TIM1 and ORF2 staining was performed.

139

140 *2.4 Generation of HEK293T cells stably expressing TIM1 WT or TIM1 ND mutant*

141 pTRIP lentiviral vectors containing wild-type or mutated TIM1 were described previously [13]. In the mutated
142 variant further referred to as TIM1 ND, both the Asparagine and Aspartate residues are substituted by Alanine
143 (N114A and D115A) within the metal ion-dependent ligand-binding site (MILIBS) of TIM1. The pTRIP vectors
144 were transfected together with pCMV-VSV-G envelope and pCMV-gag-pol packaging vectors (Sigma), in
145 HEK293T using JetPEI (Polyplus) according to manufacturer's instructions. Viral supernatant was harvested 48
146 and 72h post transfection. HEK293T cells expressing these variants were generated by lentiviral transduction,
147 followed by cell sorting of the populations expressing TIM1 by TIM1 antibody staining (P365D) and a FACS Aria
148 II (BD Biosciences). After expansion, cells were exposed to eHEV or iHEV followed by determination of infection
149 levels.

150

151 *2.5 Annexin V blocking assay*

152 Equal infectious units of eHEV or iHEV preparations were pre-incubated for 30 minutes at 4°C with 1 or 10 µL
153 of Annexin V (ANX5) (A35108 – Invitrogen, concentration proprietary information) in the presence of Ca²⁺. After
154 incubation, eHEV or iHEV virus was added to plated Huh-7.5 WT cells and infection was determined three days
155 later by ORF2 immunostaining.

156

157 *2.6 Virus attachment and internalization assays*

158 For virus attachment assays, 10^6 cells were incubated with eHEV or iHEV for 60 minutes at 4°C. Cells were
159 washed thrice to remove unbound virus and lysed, followed by total RNA extraction using the RNeasy kit
160 (Qiagen). For internalization assays, virus was allowed to adhere for 1 hour at 4°C and temperature was shifted to
161 37°C to allow endocytosis for 2 hours, then treated by trypsin to eliminate non-internalized virus and total RNA
162 was extracted as stated above. Bound and internalized virus was then quantified by qPCR as stated in the online
163 supplement.

164

165 *2.7 Statistical Analyses*

166 Statistical analysis was performed using GraphPad Prism 8.0.1 software. Statistical significance of differences
167 between groups was tested with a t-test or two-way ANOVA. A probability value of $p < 0.05$ was considered
168 significant.

169

3. Results

3.1 *TIM1 is involved in the infectious cycle of eHEV*




To get a general idea of the potential role(s) of TIM1 in the HEV life cycle, we used Huh-7.5 cells that are permissive for HEV and in which TIM1 expression was knocked-out by CRISPR/Cas9, as verified by TIM1 flow cytometry and western blot (**Fig. 1a**, left panels). Cells were exposed to virus preparations of different type (**Fig. 1a**). The first viral preparation consisted of extracellular viral particles (eHEV) isolated from supernatant of gt-3 HEV p6 electroporated cells, of which the majority are quasi-enveloped virions (**Supplemental Figure 1a**) [9]. The second viral preparation consisted of intracellular particles (iHEV) isolated from lysate of HEV p6-electroporated cells; most of the iHEV particles are considered non-enveloped [9].

Three days after virus inoculation, infection levels were determined by immunofluorescence staining against intracellular HEV ORF2 protein. Compared to wild-type (WT) Huh-7.5 cells, we observed a 97% reduction of eHEV infection in Huh-7.5^{TIM1-KO} cells ($p < 0.0001$) (**Fig. 1a**). Interestingly, infection rates were similar in both cells types infected with iHEV ($p = 0.5126$) (**Fig. 1a**).

Given this intrinsic difference between eHEV and iHEV, we performed a treatment of eHEV (tHEV) with a solution comprising sodium deoxycholate and trypsin, thereby mimicking the process viral particles undergo in the bile duct and duodenum; i.e. the removal of lipid membranes and incorporated proteins. Upon treatment, a 50% reduction of infection in Huh-7.5^{TIM1-KO} was seen compared to Huh-7.5^{WT} ($p = 0.0006$), and is consequently positioned between the eHEV or iHEV infection levels, relating to presumably only a partial loss of lipid membranes (**Fig. 1a**).

We corroborated our findings in A549^{WT} and A549^{TIM1-KO}, where TIM1 expression was verified by flow cytometry and western blot (**Supplemental Fig. 2**). In accordance with eHEV infections in Huh-7.5 cells, infection was inhibited by more than 95% in A549^{TIM1-KO} cells compared to A549^{WT} ($p < 0.009$) (**Supplemental Fig. 2**). iHEV or tHEV preparations failed to establish robust infection levels (**Supplemental Fig. 2**).

These findings were also further confirmed with the gt-3 viral isolate 83-2-27. General infection levels were much lower compared to p6 but upon eHEV inoculation, we could again demonstrate a decrease in infection of 99% in the Huh-7.5^{TIM1-KO} cells, compared to Huh-7.5^{WT} ($p = 0.004$) (**Supplemental Fig. 3**). iHEV infection was the same in the two cell lines ($p = 0.9271$).



199

200

201 **Fig. 1**

202 Knockout or blocking of TIM1 decreases eHEV infection in Huh-7.5 cells.

203 (A) Upper left panel shows an overlay of TIM1 cell surface expression in Huh-7.5^{WT} (green) and Huh-7.5^{TIM1-KO} (purple)
 204 cells. Expression evaluated by flow cytometry using a TIM1 recombinant rabbit monoclonal antibody. Middle left panel
 205 shows confirmation of TIM1 KO by western blot, 1 = Huh-7.5^{WT} 2 = Huh-7.5^{TIM1-KO}. Lower left panel shows infection in
 206 Huh-7.5^{WT} and Huh-7.5^{TIM1-KO} exposed to different preparations of HEV inoculum: extracellular HEV (eHEV),
 207 intracellular HEV (iHEV) and extracellular HEV treated with deoxycholate/trypsin (tHEV). Infection was determined 3
 208 days post infection by indirect immunostaining against the HEV ORF2 protein. Representative confocal images of the
 209 different HEV preparations are on the right with DAPI staining in blue and HEV ORF2 in green.

210 (B) Quantification of HEV RNA by qRT-PCR in supernatants of Huh-7.5^{WT} and Huh-7.5^{TIM1-KO} at two different time-points
 211 after challenge with eHEV on the left or iHEV on the right.

212 (C) Huh-7.5^{WT} were pre-incubated with different concentrations (1 µg/mL, 3 µg/mL or 10 µg/mL) of a polyclonal anti-TIM1
 213 antibody or normal IgG as control for 30 minutes. Pre-incubated cells were challenged with eHEV (left panel) or iHEV
 214 (middle panel) and infection was determined 3 days later by HEV ORF2 immunostaining. Right panel shows cell viability
 215 as measured by MTT assay and normalized to untreated control.

216 Data are represented as mean ± SEM of at least 2 independent experiments. Each experiment consisted of at least 3 technical
 217 replicates. Significance was calculated using a two-tailed t-test (A&B) or two-way ANOVA (C). ****p<0.0001, ***p<0.001,
 218 **p<0.01, *p<0.05 ns non-significant

219
 220 Next we made different ratios of co-cultures of Huh-7.5^{WT} and Huh-7.5^{TIM1-KO} cells and exposed them to either
 221 eHEV or iHEV to check the distribution of infection with regard to the TIM1 status. The ratio of WT cells ranged
 222 from 52% to 17% in the total cell population (**Supplemental Fig. 4a and 4c**). Interestingly, eHEV preferentially
 223 infects the TIM1 positive population with roughly 90% of the infected cells being TIM1-positive, irrespectively
 224 of the percentage of TIM1-cells within the total population (**Supplemental Fig. 4b**). On the other hand, iHEV
 225 infection does not preferentially infect the TIM1-positive cells as the percentage of infected TIM1 cells correlated
 226 more-or-less with the percentage of TIM1-positive cells within the total population (**Supplemental Fig. 4d**). For
 227 example, when a population consisting of 15,4-17,7% TIM1-expressing cells was exposed to eHEV, 87% of the
 228 infected cells were of the TIM1-positive type, whereas if exposed to iHEV, only 25% of the infected cells belonged
 229 to the TIM1-positive subpopulation (**Supplemental Fig. 4a-d, fourth column**).

230 Additionally, we quantified the level of secreted HEV into culture supernatant of eHEV and iHEV infected cultures
 231 at 2- and 7-days post-infection. For eHEV infection, extracellular RNA titers were on average 1-log₁₀ lower in
 232 supernatant of Huh-7.5^{TIM1-KO} cell cultures compared to Huh-7.5^{WT} cultures at both time points (p=0.0098 and
 233 p=0.0378 for day 2 and day 7 respectively) (**Fig. 1b, left panel**). For iHEV infection, viral load on day 2 and day
 234 7 were comparable in both Huh-7.5^{TIM1-KO} and Huh-7.5^{WT} cell cultures (p=0.0608 and p=0.4705 respectively)(**Fig.**
 235 **1b, right panel**).

236 The involvement of TIM1 in HEV infection was further explored by targeting the protein with increasing amounts
 237 of a neutralizing anti-TIM1 polyclonal antibody, followed by eHEV or iHEV inoculation. We found a
 238 concentration-dependent decrease in eHEV infection. The lowest concentration of antibody tested (1 µg/mL)
 239 already showed a 60% reduction in infection compared to control IgG (p=0.0097), while a reduction of more than

240 95% was observed at the highest antibody concentration ($p=0.0029$) (**Fig. 1c, left panel**) (**Supplemental Fig. 5**).
241 TIM1 neutralization followed by iHEV infection did not diminish the infection in any of the treated conditions (p
242 values 0.3709, 0.5251, >0.9999 from high to low concentration) (**Fig. 1c, middle panel**) (**Supplemental Fig. 5**).
243 Cell viability under both anti-TIM1 and IgG antibody treatment was determined by an MTT assay. Viability only
244 slightly reduced under treatment, but the TIM-1 antibody was not considered more cytotoxic than the IgG control
245 in all tested conditions (**Fig. 1c, right panel**).

246

247 *3.2 eHEV infection is rescued by TIM1 expression in KO-cells.*

248 To prove that the absence of TIM1 is responsible for decreased eHEV infection in TIM1-KO cells, these cells
249 were transiently transfected with the pcDNA3.1-TIM1 expression vector. Successful TIM1 transfection was
250 verified via western blot (**Fig. 2a, upper panel**). To investigate the effect of HEV infection in transiently
251 transfected Huh-7.5^{TIM1-KO} cells, both HEV ORF2 and TIM1 were visualized by immunofluorescence staining in
252 Huh-7.5^{TIM1-KO} MOCK transfected or Huh-7.5^{TIM1-KO} pcDNA-TIM1 transfected cells. At the time-point of HEV
253 analysis on day three, an average of 6% of the cells showed TIM1 expression (**Fig. 2, blue bars**).

254 In accordance with our previous infection experiments in TIM1-KO cells, an average eHEV infection rate of only
255 4 foci/well could be observed in Huh-7.5^{TIM1-KO} cells that were MOCK transfected. However, upon TIM1 rescue,
256 we observed a nearly 150-fold increase in eHEV infection ($p=0.0135$) (**Fig. 2a, green bars**). Moreover, it is clear
257 that the HEV infection is observed in the cells expressing TIM1, as illustrated by the immunostaining images
258 where HEV and TIM1 signals are predominantly observed in the same cells (**Fig. 2a, confocal images**). On the
259 contrary, rescue of TIM1 had no effect on iHEV infection levels, as infection was similar in MOCK or pcDNA-
260 TIM1 transfected cells ($p=0.9849$) (**Fig. 2b, green bars**) and infected cells are generally not the TIM1-positive
261 cells (**Fig. 2b, confocal images**) This further confirms the role of TIM1 in eHEV infection.

262



263 *3.1 Ectopic expression of TIM1 facilitates eHEV infection in HEK293T cells*

264 We further evaluated the ability of ectopic TIM1 expression to mediate eHEV infection in HEK293T cells, which
265 do not express TIM1. We first confirmed that HEK293T could support HEV genome replication (**Fig. 3a**). Next,
266 we transfected HEK293T cells transiently with the pcDNA3.1-TIM1 expression vector and confirmed TIM1
267 expression first via western blot (**Fig. 3b**). To investigate the effect of HEV infection in transiently transfected
268 HEK293T cells, both HEV ORF2 and TIM1 were visualized by immunofluorescence staining in HEK293T



269 MOCK transfected or HEK293T pcDNA-TIM1 transfected cells. At the time-point of HEV analysis on day three,
270 almost 20% of the cells showed TIM1 expression (**Fig 3c and 3d, blue bars**). Parental MOCK transfected
271 HEK293T were nearly completely resistant to eHEV infection, whereas TIM1 expression enhanced eHEV
272 infection level to on average 15% ($p=0.0063$) (**Fig. 3c**). iHEV infection in parental MOCK transfected cells was
273 generally low, but infected cell foci could be observed and transient TIM1 expression did not alter iHEV infection
274 ($p=0.7889$) (**Fig. 3d**).

275

A



B



276

277

278 **Fig. 2**


279 Rescue of TIM1 enhances HEV infection.

- 280 (A) Huh-7.5^{TIM1-KO} cells were transfected with an empty plasmid (MOCK) or pcDNA3.1-TIM1 expression vector.
281 Expression was verified via western blot, 1= Huh-7.5^{TIM1-KO}/MOCK and 2= Huh-7.5^{TIM1-KO}/pcDNA-TIM1.
282 eHEV infection was performed 24h after transfection and analyzed three days later. eHEV infection levels
283 determined by HEV ORF2 indirect immunofluorescence staining (right axis, green). TIM1 expression on the
284 same day is depicted on the left y-axis (blue) and was measured by TIM1 immunofluorescence staining. On
285 the right side, representative confocal microscopy images of MOCK and TIM1 transfected Huh-7.5^{TIM1-KO}
286 cells are depicted. Cells were fixed and permeabilized, followed by immunofluorescence staining against
287 TIM1 (red) and HEV ORF2 (green), DAPI nuclei in blue. Lowest panel shows a merged image where a region
288 of interest was acquired by 40x objective lens instead of 20x objective (white square) to demonstrate overlap
289 between the cells expressing TIM1 and being infected with HEV.
- 290 (B) Huh-7.5^{TIM1-KO} cells were transfected with an empty plasmid (MOCK) or pcDNA3.1-TIM1 expression vector
291 and infected with iHEV. Experiment and graph same as for the upper panel. On the right side, representative
292 confocal microscopy images are depicted, the same as described for the upper panel.

293 Data are represented as means \pm SEM from two independent experiments. Each experiment included at least 3
294 technical replicates. Significance was calculated using a two-tailed t-test. *p<0.05 ns non-significant

295

296



297

298 **Fig. 3**

299 Ectopic expression of TIM1 facilitates eHEV infection in HEK293T cells.

300 (A) HEK293T cells support HEV replication after transfection with HEV RNA, comparison made with electroporated Huh-
 301 7.5 cells. Percentages indicate expression of HEV, as determined by flow cytometry after intracellular staining using an
 302 HEV ORF2 monoclonal antibody, followed by labeling with an Alexa Fluor 488-conjugated secondary antibody.

303 (B) HEK293T were transiently transfected with an empty plasmid (MOCK) or pcDNA3.1-TIM1 expression vector and
 304 expression was verified via western blot. 1= HEK293T/MOCK 2=HEK293T/pcDNA-TIM1

305 (C) HEK293T were transfected with an empty plasmid (MOCK) or pcDNA3.1-TIM1 expression vector one day before eHEV
 306 infection. left panel shows results with eHEV infection percentage on the right y-axis 3 days after, as measured by HEV
 307 ORF2 immunofluorescence staining. TIM1 expression on the same day is depicted on the left y-axis and was measured
 308 by TIM1 immunofluorescence staining. Right panel are representative confocal images of eHEV infected
 309 HEK293T/MOCK (parental) compared to the HEK293T/pcDNA-TIM1, DAPI staining in blue, HEV ORF2 in green and
 310 TIM1 in red.

311 (D) HEK293T cells transfected the same way as described and infected with iHEV. iHEV percentage on the right y-axis on
 312 day 3 and TIM1 expression on the same day depicted on the left y-axis. Right panel shows representative confocal images
 313 of iHEV infected HEK293T/MOCK compared to HEK293T/pcDNA-TIM1.

314


315 Data represented as means \pm SEM from two independent experiments. Each experiment consisted of at least 3 technical
 316 replicates. Significance was calculated using a t-test. ** $p < 0.01$ ns non-significant

317

318 3.1 *TIM1 facilitates eHEV internalization*

319 The above-mentioned data clearly indicates the role of TIM1 during infection of eHEV. To examine the exact
 320 function of TIM1 during viral entry in more detail, we performed viral binding assays. eHEV or iHEV were
 321 allowed to adhere to Huh-7.5^{WT} and Huh-7.5^{TIM1-KO} cells at 4°C. Perhaps unexpectedly, we could not observe a
 322 difference in eHEV binding between Huh-7.5^{WT} and Huh-7.5^{TIM1-KO} ($p = 0.4697$) (**Fig. 4a, left panel**). As a control,
 323 iHEV attachment assays were performed in parallel and also no difference in viral binding was observed
 324 ($p = 0.3728$) (**Fig. 4a, left panel**). In a next step, we carried out an internalization assay, where the virus was first
 325 attached, followed by a shift to 37°C to permit virus internalization. Cell-surface bound virions were removed by
 326 trypsinization and after RNA extraction, qPCR showed a marked decrease of almost 1-log in internalized eHEV
 327 in the Huh-7.5^{TIM1-KO}, compared to Huh-7.5^{WT} ($p = 0.0046$), while internalized iHEV was similar in the two cell
 328 types ($p = 0.4462$) (**Fig. 4a, right panel**). These findings were confirmed in 293T stably transduced with TIM1 and
 329 compared to parental 293T. For both eHEV and iHEV, no difference in viral binding could be observed ($p = 0.1254$
 330 and $p = 0.1864$ respectively), but a decrease in internalized eHEV was observed ($p = 0.0463$), while iHEV
 331 internalization was the same ($p = 0.0911$) (**Fig. 4b**). Similarly, we verified our observations with another gt-3
 332 isolate, 83-2-27. Again no differences in attachment between Huh-7.5^{WT} and Huh-7.5^{TIM1-KO} could be observed
 333 ($p = 0.9684$ for eHEV and $p = 0.1230$ for iHEV) but, similar as with the p6 clone, less enveloped virus internalizes
 334 in TIM1 KO cells ($p = 0.0139$) (**Supplementary fig. 6**).

335



336

337 **Fig. 4**

338 TIM1 facilitates eHEV entry

- 339 (A) Adherence of eHEV and iHEV to Huh-7.5^{WT} and Huh-7.5^{TIM1-KO} at 4°C for 1 hour. After washing, RNA was
340 extracted and determined by qRT-PCR (left panel). Right panel shows viral binding of eHEV and iHEV to Huh-
341 7.5^{WT} and Huh-7.5^{TIM1-KO}, followed by a shift to 37°C for 2 hours. Cells were washed, trypsinized and total RNA
342 was extracted and measured by qRT-PCR.
- 343 (B) Adherence of eHEV and iHEV to HEK293T^{parental} and HEK293T^{TIM1-WT} at 4°C for 1 hour. After washing, RNA was
344 extracted and determined by qRT-PCR (left panel). Right panel shows viral binding of eHEV and iHEV to Huh-
345 7.5^{WT} and Huh-7.5^{TIM1-KO}, followed by a shift to 37°C for 2 hours. Cells were washed, trypsinized and total RNA
346 was extracted and measured by qRT-PCR.

347 Data represented as means ± SEM from two independent experiments. Each experiment consisted of at least 2 technical
348 replicates. Significance was calculated using a t-test. **p<0.01 *p<0.05 ns non-significant

349

350 3.2 The PS-binding cavity of TIM1 is important for eHEV viral infection

351 TIM proteins bind phosphatidylserine (PS) present on the surface of apoptotic cells or viruses [12]. eHEV as well
352 has been described to be associated with PS [11]. To further explore the potential role of eHEV-associated PS and
353 the effect on viral infection, we pre-incubated eHEV or iHEV with Annexin V (ANX5), a PS-binding protein that
354 has often been used to explore the role of TIM1 in viral entry and infection. Using the highest amount of ANX5,
355 a reduction in eHEV infection of more than 50 % could be observed (p=0.0101), while a 10-fold less amount of
356 ANX5 had an intermediate effect (p=0.1502) (**Fig. 5a**), indicating dose-dependent inhibition. In contrast, control
357 infections with iHEV showed similar infection levels in all tested conditions (p=0.3321 for 10 µL and p=0.6465
358 for 1µL)

368 Data represented as means \pm SEM from two independent experiments. Each experiment consisted of at least 3 technical
369 replicates. Significance was calculated using a two-way ANOVA. * $p < 0.05$ ns non-significant


370

371 To study the role of PS in greater detail in the context of eHEV, we further focused on the metal ion-dependent
372 ligand-binding site (MILIBS) situated in a conserved cavity that is part of the extracellular IgG-V-like domain of
373 TIM receptors, which has been well characterized as the binding site for PS, including in the context of viral entry
374 and infection [12]. A HEK293T cell line was generated that expresses a previously characterized mutated TIM1
375 receptor [13], harboring two mutations (N114A and D115A; termed TIM1-ND) in the conserved cavity, rendering
376 it unable to bind PS. TIM1 was correctly expressed at the cell surface, as confirmed via flow cytometry and protein
377 expression was also verified via western blot (**Fig. 6a**). We determined the level of HEV infection compared to a
378 HEK293T cell line stably expressing TIM1-WT and included parental HEK293T cells as a control. eHEV infection
379 in the parental cells was almost non-existent compared to the HEK293T^{TIM1-WT} ($p < 0.0001$) (**Fig. 6b, left panels**),
380 but infection decreased almost 95% in HEK293T^{TIM1-ND} compared to HEK293T^{TIM1-WT} ($p < 0.0001$) (**Fig. 6b**). We
381 compared our results to iHEV infection. As with transient expression in HEK293T, infection in parental cells was
382 low, though remained the same in HEK293T^{TIM1-WT} or HEK293T^{TIM1-ND} expressing cells (**Fig 6b, right panels**).

383 To confirm these findings, Huh-7.5^{TIM1-KO} cells were also transduced with the TIM1-ND mutant and infection
384 levels were compared to Huh-7.5^{TIM1-KO} stably transduced with TIM1-WT and control Huh-7.5^{TIM1-KO} (Fig. 7a).
385 We observed a decrease of 99% in Huh-7.5^{TIM1-ND} compared to Huh7.5^{TIM1-WT} upon eHEV infection (**Fig. 7b, left**
386 **panels**). iHEV infection levels were the same in all three conditions (**Fig. 7b, right panels**), indicating that the
387 PS-binding pocket is required only for the TIM1-mediated eHEV infection.

388

389



390

391 **Fig. 6**

392 The PS-binding cavity is important for viral binding to TIM1.

393 (A) TIM1 cell surface expression of parental HEK293T (purple) compared to TIM1-WT transduced HEK293T (green) and
 394 TIM1-ND mutant transduced HEK293T (grey). Expression was evaluated by flow cytometry using a TIM1 recombinant
 395 monoclonal antibody (left panel). Right panel shows expression verified by western blot. 1=HEK293T^{parental}
 396 2=HEK293T^{TIM1-WT} 3=HEK293T^{TIM1-ND}

397 (B) Stable transduced HEK293T^{TIM1-WT} and HEK293T^{TIM1-ND} mutant as well as parental HEK293T cells were challenged with
 398 eHEV (left panels) or iHEV (right panels) and analyzed by HEV ORF2 immunofluorescence staining three days later.
 399 Lower panels are representative confocal images of the different conditions, DAPI staining in blue, HEV ORF2 in green
 400 and TIM1 in red.


401 Data represented as means \pm SEM from two independent experiments. Each experiment consisted of at least 3 technical
 402 replicates. Significance was calculated using a two-way ANOVA. **** $p < 0.0001$ ns non-significant

403

404

405

406



407

408 **Fig. 7**

409

410 The PS-binding cavity is important for viral binding to TIM1.

411 (A) TIM1 cell surface expression of Huh-7.5^{TIM1-KO} (purple) compared to Huh-7.5^{TIM1-WT} (green) and Huh-7.5^{TIM1-ND} (grey).

412 Expression was evaluated by flow cytometry using a TIM1 recombinant monoclonal antibody (left panel). Right panel

413 shows expression verified by western blot. 1=Huh-7.5^{TIM1-WT} 2= Huh-7.5^{TIM1-ND} 3= Huh-7.5^{TIM1-KO}

414

415 (B) Huh-7.5^{TIM1-KO}, Huh-7.5^{TIM1-WT}, and Huh-7.5^{TIM1-ND} were challenged with eHEV (left panels) or iHEV (right panels) and

416 analyzed by HEV ORF2 immunofluorescence staining three days later. Lower panels are representative confocal images

417 of the different conditions, DAPI staining in blue, HEV ORF2 in green and TIM1 in red.

418 Data represented as means ± SEM from two independent experiments. Each experiment consisted of at least 3 technical

419 replicates. Significance was calculated using a two-way ANOVA. ****p<0.0001 ns non-significant

420

421

4. Discussion

422
423 In this study, we provide the first experimental evidence that TIM1 plays a role in the infectivity of enveloped
424 HEV particles. We found that absence of TIM1 significantly decreased eHEV infection in Huh-7.5^{TIM1-KO} cells,
425 but not iHEV infection (**Fig. 1a**), indicating that the inhibition of infection is probably limited to viral particles
426 surrounded by a host membrane-derived envelope. This finding was further corroborated by the fact that infection
427 by deoxycholate/trypsin-treated eHEV (tHEV) was less dependent on TIM1 (only 50% inhibition) compared with
428 non-treated eHEV (more than 90% inhibition) (**Fig. 1a**). Of note, tHEV infection levels were intermediate between
429 those of eHEV, consisting merely of enveloped particles, and those of iHEV, comprising mainly non-enveloped
430 virus (**Supplemental Fig. 1**). The fact that we observed a significant difference in infection between WT and
431 TIM1-KO cells using tHEV may be explained by a partial loss of the viral membrane upon deoxycholate/trypsin
432 treatment. This has been illustrated by tHEV density profile analysis, showing a mixture of both enveloped and
433 non-enveloped particles [18]. We as well observed a peak shift for the density profile (**Supplemental Fig. 1B**).

434 We confirmed eHEV infection inhibition in A549^{TIM1-KO} cells (**Supplemental Fig. 2**), but unfortunately, iHEV or
435 tHEV preparations failed to establish robust infection levels in these cells. Therefore, we focused the rest of our
436 infection studies mainly on Huh-7.5 cells and subsequently on HEK293T cells. We also confirmed eHEV infection
437 inhibition for the gt-3 83-2-27 isolate in Huh-7.5^{TIM1-KO} cells (**Supplemental Fig. 3**).

438 A reduced virus secretion from Huh-7.5^{TIM1-KO} cells when they were infected with eHEV could be detected (**Fig.**
439 **1b**).

440 We also showed that an antibody targeting TIM1 blocks eHEV infection but not iHEV infection in cells naturally
441 expressing the receptor (**Fig. 1c**). Furthermore, eHEV infection could be restored in Huh-7.5^{TIM1-KO} cells upon
442 transfection with a TIM1-expressing plasmid (**Fig. 2a**); and ectopic TIM1 expression in HEK293T cells enhances
443 eHEV infection (**Fig. 3c**). Taken together, it is clear that TIM1 is involved in the infectious cycle of enveloped
444 HEV particles.

445 In context of viral infections, TIM1 has been shown to enhance entry by a process called viral apoptotic mimicry
446 where PS exposed on viral envelope facilitate viral engulfment, either by directly interacting with TIM1 or via
447 bridging molecules [12]. The MILIBS of TIM1 is well characterized as the essential part of PS recognition [19,
448 20]. It was previously shown that HEV particles exit cells as exosome-like vesicles and contain PS, possibly
449 acquired from membranes derived from multivesicular bodies, key players of the endosome fraction that play a
450 central role during enveloped particle biogenesis and viral egress [11, 21]. Using Annexin V, which binds PS with

451 high affinity, we showed that eHEV infection decreases (**Fig. 5a**). Mutation of the critical residues in the PS-
452 binding pocket inhibited eHEV infection by more than 90% (**Fig. 6b** and **7b**). Both experiments indicate the
453 importance of this cavity. Based on current knowledge about the interaction between PS and the binding pocket
454 of TIM1 in context of viral infection, one could assume that PS present on enveloped HEV viral particles would
455 act as such. In general, it is not always clear whether TIM1 only enhances virus attachment or serves as an authentic
456 receptor. For the latter a direct interaction between a viral protein and TIM1 is needed, as has been described for
457 Dengue virus and Ebola virus [22, 23]. Our data suggests the role of PS in entry (**Fig. 5, Fig. 6** and **Fig. 7**), which
458 would favor a role of TIM1 as attachment factor. However, our attachment assays in both Huh-7.5 and HEK293T
459 cells show no difference in viral binding in absence of TIM1. A difference between WT and KO only became
460 apparent upon analysis of viral internalization (**Fig. 4** and **Supplementary Fig. 6**). A similar finding was observed
461 for hepatitis A virus (HAV) where an analogous distinction of both enveloped and non-enveloped viral particles
462 is made [17]. TIM1-KO reduces entry of enveloped HAV, but not non-enveloped HAV, and the role of TIM1 in
463 enveloped HAV entry is considered to involve interaction with PS present in the viral envelope [14, 24]. A
464 comparison between the two viruses might therefore seem conceivable with an analogous role for PS present in
465 the viral envelope of HEV. Interestingly, in the context of HAV, no difference in attachment to Huh-7.5^{TIM1-KO}
466 cells compared to WT was detected either. Only when a kidney-derived cell line was used, a difference in
467 attachment was observed, which was attributed to the fact that expression of TIM1 is higher in the kidney compared
468 to the liver [14]. We used the kidney-derived HEK293T cell line but were unable to observe a difference in
469 attachment upon TIM1 transduction. However, parental HEK293T cells do not express TIM1 as verified with
470 flow cytometric staining and Western blotting (**Fig. 6a**) and reports from other groups [13], precluding a similar
471 observation as with HAV. The fact that we do observe that TIM1 facilitates viral internalization, indicates that it
472 is not merely acting as attachment factor but is involved in the process of viral entry of enveloped HEV. Our
473 current understanding of this protein is incomplete to address all knowledge gaps. For example, the exact
474 mechanism of virus internalization and following uncoating in context of TIM1 should be further explored. It was
475 previously described that both eHEV and iHEV internalize via clathrin- and dynamin-dependent pathways but it
476 seems that the uncoating process differs. A previous study showed that iHEV seems to uncoat in early endosomes,
477 whereas eHEV entry is probably dependent on endosomal trafficking to early and late endosomes [25]. Further
478 uncoating may be achieved by either fusion of the viral membrane with the cellular membrane or by removal of
479 the envelope by lysosomal degradation. Previous work illustrated that lysosomal proteins Niemann-Pick disease
480 type C1 (NPC1) and lysosomal acid lipase (LAL) are involved in eHEV infection, and therefore the second

481 hypothesis may be more likely [9]. It is believed that the virus must encode specific targeting signals, and that the
482 interaction of a viral protein is possibly involved in trafficking [26]. Further study should explore a potential
483 additional role of TIM1 in viral trafficking. Interestingly, it was previously shown that TIM1 can traffic between
484 early endosomes as well as late endosome and lysosomal compartments, as they carry their cargo from the cell
485 surface to the lysosome [27]. This role of TIM1 has briefly been explored during Dengue virus infection, where
486 TIM1 seems to co-internalize with viral particles during endocytosis. Moreover, an interaction with STAM1 was
487 observed [22], a component of ESCRT-0 necessary for endosomal sorting of ubiquitinated membrane proteins
488 [28].

489 Although infection decreased by more than 90% after TIM1-KO or antibody neutralization, a limited number of
490 infected cells could still be observed. This may suggest that TIM1 is an important but not the only factor mediating
491 entry for enveloped particles. On the other hand, we cannot exclude that some of the eHEV preparations present
492 with a very low amount of naked HEV particles, which seem not to be dependent on TIM1 for entry. We were
493 unable to use density-gradient purified virus, merely because the infectivity of those purified preps was too low to
494 reveal potential significant differences in inhibition (data not shown). The *in vivo* significance of TIM1 during
495 HEV infection remains to be further explored. If endogenous TIM1 molecules are important for HEV infection,
496 one would expect a certain level of expression on hepatocytes. According to the Human Protein Atlas, hepatic
497 TIM1 expression is rather low and no TIM1 expression is detected on hepatocytes. We corroborated these findings
498 by cell surface staining of TIM1 on isolated primary human hepatocytes (PHH) and did not succeed to visualize
499 cell surface TIM1 on PHH (data not shown). However, also according to Human Protein Atlas, there should be
500 RNA expression in these cells in the liver. We sought out if this was the case by total RNA extraction from PHH,
501 followed by TIM1 amplification and sequencing. Our data confirmed that TIM1 mRNA is present in these cells
502 (data not shown). It is possible that TIM1 surface expression is indeed absent, as we observed, but natural surface
503 expression could also be below the detection limit of our detection method, and/or visualization of TIM1 cell
504 surface expression on isolated PHH is hampered by the proteinase digestion to isolate PHH from liver tissue.
505 Interestingly, it has been described that renal TIM1 expression level increases upon kidney injury. This
506 subsequently promotes transformation of the tubular epithelial cells into phagocytic cells that are able to recognize
507 PS-associated apoptotic cells and execute apoptotic clearance necessary for resolution of inflammation [29]. To
508 our knowledge, such effect has not yet been described in the context of virus infection, but it is an interesting
509 speculation that a virus would upregulate TIM1 surface expression on target cells to enhance its entry and
510 subsequent spread.

511 Even if hepatic TIM1 expression is low, this would not diminish the importance of the results we show as HEV
512 infection is also associated with a variety of extrahepatic manifestations, leading to e.g. neurological or renal
513 complications. In these tissues, TIM1 expression is notably high (Human Protein Atlas). During natural HEV
514 infection, infection is likely mediated by fecal-oral transmission between individuals with an exposure to non-
515 enveloped viral particles. However, after initial infection, enveloped particles are released from the infected liver
516 into the blood circulation. It is possible that viral spread within the body mainly occurs through enveloped HEV
517 particles that interact with a non-specific receptor such as TIM1, via a connection with PS incorporated in the viral
518 envelope. The enveloped nature of HEV particles in the blood is of importance since it makes them less
519 susceptible to ORF2-vaccine induced antibody mediated neutralization. Previously, PS-specific antibody therapy
520 was able to cure guinea pigs lethally infected with Pichinde virus, illustrating the potential of targeting PS in the
521 viral envelope [30]. On the other hand, given the importance of TIM1 in the entry of enveloped HEV, TIM1 might
522 also constitute a target for antiviral therapy.

523

524 **References**

- 525 1. Lhomme, S., et al., *Clinical Manifestations, Pathogenesis and Treatment of Hepatitis E Virus Infections*.
526 *Journal of clinical medicine*, 2020. **9**(2).DOI: 10.3390/jcm9020331.
- 527
- 528 2. Webb, G.W. and H.R. Dalton, *Hepatitis E: an expanding epidemic with a range of complications*. *Clinical*
529 *microbiology and infection : the official publication of the European Society of Clinical Microbiology and*
530 *Infectious Diseases*, 2020. **26**(7): p. 828-832.DOI: 10.1016/j.cmi.2020.03.039.
- 531
- 532 3. Sayed, I.M., et al., *Is hepatitis E virus an emerging problem in industrialized countries?* *Hepatology*
533 (Baltimore, Md.), 2015. **62**(6): p. 1883-92.DOI: 10.1002/hep.27990.
- 534
- 535 4. Navaneethan, U., M. Al Mohajer, and M.T. Shata, *Hepatitis E and pregnancy: understanding the*
536 *pathogenesis*. *Liver international : official journal of the International Association for the Study of the Liver*, 2008.
537 **28**(9): p. 1190-9.DOI: 10.1111/j.1478-3231.2008.01840.x.
- 538
- 539 5. Kamar, N., et al., *Hepatitis E virus infection*. *Nature reviews. Disease primers*, 2017. **3**: p. 17086.DOI:
540 10.1038/nrdp.2017.86.
- 541
- 542 6. Kenney, S.P. and X.J. Meng, *Hepatitis E Virus Genome Structure and Replication Strategy*. *Cold Spring*
543 *Harbor perspectives in medicine*, 2019. **9**(1).DOI: 10.1101/cshperspect.a031724.
- 544
- 545 7. Montpellier, C., et al., *Hepatitis E Virus Lifecycle and Identification of 3 Forms of the ORF2 Capsid*
546 *Protein*. *Gastroenterology*, 2018. **154**(1): p. 211-223 e8.DOI: 10.1053/j.gastro.2017.09.020.
- 547
- 548 8. Nair, V.P., et al., *Endoplasmic Reticulum Stress Induced Synthesis of a Novel Viral Factor Mediates*
549 *Efficient Replication of Genotype-1 Hepatitis E Virus*. *PLoS pathogens*, 2016. **12**(4): p. e1005521.DOI:
550 10.1371/journal.ppat.1005521.
- 551
- 552 9. Yin, X., et al., *Distinct Entry Mechanisms for Nonenveloped and Quasi-Enveloped Hepatitis E Viruses*.
553 *Journal of virology*, 2016. **90**(8): p. 4232-4242.DOI: 10.1128/JVI.02804-15.
- 554
- 555 10. Yin, X. and Z. Feng, *Hepatitis E Virus Entry*. *Viruses*, 2019. **11**(10).DOI: 10.3390/v11100883.
- 556
- 557 11. Nagashima, S., et al., *Characterization of the Quasi-Enveloped Hepatitis E Virus Particles Released by*
558 *the Cellular Exosomal Pathway*. *Journal of virology*, 2017. **91**(22).DOI: 10.1128/JVI.00822-17.
- 559
- 560 12. Amara, A. and J. Mercer, *Viral apoptotic mimicry*. *Nature reviews. Microbiology*, 2015. **13**(8): p. 461-
561 9.DOI: 10.1038/nrmicro3469.
- 562
- 563 13. Meertens, L., et al., *The TIM and TAM families of phosphatidylserine receptors mediate dengue virus*
564 *entry*. *Cell host & microbe*, 2012. **12**(4): p. 544-57.DOI: 10.1016/j.chom.2012.08.009.
- 565

- 566 14. Das, A., et al., *TIMI (HAVCR1) Is Not Essential for Cellular Entry of Either Quasi-enveloped or Naked*
567 *Hepatitis A Virions*. mBio, 2017. **8**(5).DOI: 10.1128/mBio.00969-17.
- 568
- 569 15. Costafreda, M.I. and G. Kaplan, *HAVCR1 (CD365) and Its Mouse Ortholog Are Functional Hepatitis A*
570 *Virus (HAV) Cellular Receptors That Mediate HAV Infection*. Journal of virology, 2018. **92**(9).DOI:
571 10.1128/JVI.02065-17.
- 572
- 573 16. Kondratowicz, A.S., et al., *T-cell immunoglobulin and mucin domain 1 (TIM-1) is a receptor for Zaire*
574 *Ebolavirus and Lake Victoria Marburgvirus*. Proceedings of the National Academy of Sciences of the United
575 States of America, 2011. **108**(20): p. 8426-31.DOI: 10.1073/pnas.1019030108.
- 576
- 577 17. Feng, Z., et al., *Naked Viruses That Aren't Always Naked: Quasi-Enveloped Agents of Acute Hepatitis*.
578 Annual review of virology, 2014. **1**(1): p. 539-60.DOI: 10.1146/annurev-virology-031413-085359.
- 579
- 580 18. Sayed, I.M., et al., *Study of hepatitis E virus infection of genotype 1 and 3 in mice with humanised liver*.
581 *Gut*, 2017. **66**(5): p. 920-929.DOI: 10.1136/gutjnl-2015-311109.
- 582
- 583 19. Kobayashi, N., et al., *TIM-1 and TIM-4 glycoproteins bind phosphatidylserine and mediate uptake of*
584 *apoptotic cells*. Immunity, 2007. **27**(6): p. 927-40.DOI: 10.1016/j.immuni.2007.11.011.
- 585
- 586 20. Santiago, C., et al., *Structures of T Cell immunoglobulin mucin receptors 1 and 2 reveal mechanisms for*
587 *regulation of immune responses by the TIM receptor family*. Immunity, 2007. **26**(3): p. 299-310.DOI:
588 10.1016/j.immuni.2007.01.014.
- 589
- 590 21. Nagashima, S., et al., *Tumour susceptibility gene 101 and the vacuolar protein sorting pathway are*
591 *required for the release of hepatitis E virions*. The Journal of general virology, 2011. **92**(Pt 12): p. 2838-2848.DOI:
592 10.1099/vir.0.035378-0.
- 593
- 594 22. Dejarnac, O., et al., *TIM-1 Ubiquitination Mediates Dengue Virus Entry*. Cell reports, 2018. **23**(6): p.
595 1779-1793.DOI: 10.1016/j.celrep.2018.04.013.
- 596
- 597 23. Zhang, Q., et al., *Stepwise Enzymatic-Dependent Mechanism of Ebola Virus Binding to Cell Surface*
598 *Receptors Monitored by AFM*. Nano letters, 2022. **22**(4): p. 1641-1648.DOI: 10.1021/acs.nanolett.1c04677.
- 599
- 600 24. Das, A., W. Maury, and S.M. Lemon, *TIMI (HAVCR1): an Essential "Receptor" or an "Accessory*
601 *Attachment Factor" for Hepatitis A Virus?* Journal of virology, 2019. **93**(11).DOI: 10.1128/JVI.01793-18.
- 602
- 603 25. Holla, P., et al., *Hepatitis E virus enters liver cells through a dynamin-2, clathrin and membrane*
604 *cholesterol-dependent pathway*. Traffic (Copenhagen, Denmark), 2015. **16**(4): p. 398-416.DOI:
605 10.1111/tra.12260.
- 606
- 607 26. Rivera-Serrano, E.E., et al., *Cellular entry and uncoating of naked and quasi-enveloped human*
608 *hepatoviruses*. eLife, 2019. **8**.DOI: 10.7554/eLife.43983.

609

610 27. Balasubramanian, S., et al., *TIM family proteins promote the lysosomal degradation of the nuclear*
611 *receptor NUR77*. *Science signaling*, 2012. **5**(254): p. ra90.DOI: 10.1126/scisignal.2003200.

612

613 28. Schmidt, O. and D. Teis, *The ESCRT machinery*. *Current biology : CB*, 2012. **22**(4): p. R116-20.DOI:
614 10.1016/j.cub.2012.01.028.

615

616 29. Ichimura, T., et al., *Kidney injury molecule-1 is a phosphatidylserine receptor that confers a phagocytic*
617 *phenotype on epithelial cells*. *The Journal of clinical investigation*, 2008. **118**(5): p. 1657-68.DOI:
618 10.1172/JCI34487.

619

620 30. Soares, M.M., S.W. King, and P.E. Thorpe, *Targeting inside-out phosphatidylserine as a therapeutic*
621 *strategy for viral diseases*. *Nature medicine*, 2008. **14**(12): p. 1357-62.DOI: 10.1038/nm.1885.

622

623

624

625 **Statements and Declarations**

626 *Funding*

627 PM was supported by the Special Research Fund of Ghent University (BOFEXP2017001002), the Research
628 Foundation-Flanders (FWO-Vlaanderen; Excellence of Science (EOS) projects VirEOS and VirEOS2.0).

629

630 *Competing interests*

631 The authors declare no conflict of interest.

632

633 *Author Contributions*

634 The study was conceived by PM. All authors contributed to the experimental design. LCor, IL, CM, XH, MF and
635 FVH performed experiments. LCor, IL, CM, MF, KW, XH, LCoc and PM analyzed and interpreted the
636 experimental data. LCor and PM wrote the manuscript. All authors critically reviewed and contributed to the
637 manuscript.

638

639 *Data Availability*

640 The datasets generated and analyzed during the current study are available from the corresponding author on
641 reasonable request.