



**HAL**  
open science

## Partial-body cryostimulation does not impact peripheral microvascular responsiveness but reduces muscular metabolic O<sub>2</sub> consumption ( $m\dot{V} \cdot O_2$ ) at rest

Dimitri Theurot, Olivier Dupuy, Julien Louis, Wafa Douzi, Renée Morin, Coralie Arc-Chagnaud, Benoit Dugué

### ► To cite this version:

Dimitri Theurot, Olivier Dupuy, Julien Louis, Wafa Douzi, Renée Morin, et al.. Partial-body cryostimulation does not impact peripheral microvascular responsiveness but reduces muscular metabolic O<sub>2</sub> consumption ( $m\dot{V} \cdot O_2$ ) at rest. *Cryobiology*, 2023, 112, pp.104561. 10.1016/j.cryobiol.2023.104561 . hal-04246147

**HAL Id: hal-04246147**

**<https://hal.science/hal-04246147v1>**

Submitted on 21 Mar 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Partial-body cryostimulation does not impact peripheral microvascular responsiveness**  
2 **but reduces muscular metabolic O<sub>2</sub> consumption (mVO<sub>2</sub>) at rest.**

3 **Authors:** *Dimitri Theurot*<sup>1</sup>, *Olivier Dupuy*<sup>1, 2</sup>, *Julien Louis*<sup>3</sup>, *Wafa Douzi*<sup>1</sup>, *Renée Morin*<sup>4</sup>,  
4 *Coralie Arc-Chagnaud*<sup>1</sup>, *Benoit Dugué*<sup>1</sup>

5 **Affiliations:**

6 <sup>1</sup> *University of Poitiers, laboratory MOVE (UR 20296), Faculty of Sport Sciences, Poitiers.*  
7 *France.*

8 <sup>2</sup> *Ecole de Kinésiologie et des Sciences de l'Activité Physique (EKSAP). Faculté de*  
9 *Medicine. Université de Montreal. Canada.*

10 <sup>3</sup> *Research Institute for Sport and Exercise Sciences (RISES), Liverpool John Moores*  
11 *University, Byrom Street, Liverpool L3 3AF, UK*

12 <sup>4</sup> *School of Human Kinetics, Faculty of Health Sciences, University of Ottawa, Ottawa, ON,*  
13 *Canada*

14

15 **Corresponding author:**

16 Dimitri Theurot Ph.D.

17 Laboratory MOVE (UR 20296)

18 Faculty of Sport Sciences

19 University of Poitiers.

20 8 allée Jean Monnet 86000 Poitiers. France

21 Email: [dimitri.theurot@univ-poitiers.fr](mailto:dimitri.theurot@univ-poitiers.fr)

22 **Abstract.**

23 This study aimed to investigate the effect of partial-body cryostimulation (PBC) on  
24 microvascular responsiveness and muscular metabolic O<sub>2</sub> consumption rate (mVO<sub>2</sub>). Twenty  
25 healthy young adults (ten males and ten females) underwent post-occlusive reactive  
26 hyperaemia (PORH) test at the flexor digitorum superficialis area before and after a 3-min  
27 PBC session and a 3-min control session. Using near-infrared spectroscopy, occlusion and  
28 reperfusion slopes were calculated: oxyhaemoglobin ([HbO<sub>2</sub>]) decrease rate ([HbO<sub>2</sub>] slope 1),  
29 deoxyhaemoglobin ([HHb]) increase rate ([HHb] slope 1), [HbO<sub>2</sub>] increase rate ([HbO<sub>2</sub>] slope  
30 2), and [HHb] increase rate ([HHb] slope 2). Using HbO<sub>2</sub> kinetics during the occlusion, mVO<sub>2</sub>  
31 was also calculated to characterize myocytes' metabolic O<sub>2</sub> consumption. HbO<sub>2</sub> slope 1 value  
32 was lower after PBC than before PBC ( $-0.15 \pm 0.08$  vs  $-0.24 \pm 0.11$  s<sup>-1</sup>; respectively; P < 0.05)  
33 in male participants only. A lower [HHb] slope 1 was also observed after PBC compared to  
34 before PBC ( $0.18 \pm 0.10$  vs  $0.24 \pm 0.16$  s<sup>-1</sup>; P < 0.05) with no interaction for sex categories.  
35 mVO<sub>2</sub> was significantly lower after PBC than before (pre values  $14.75 \pm 3.94$  vs  $18.47 \pm 5.73$   
36  $\mu\text{MO}_2\text{Hb}\cdot\text{s}^{-1}$ ; respectively; P < 0.01) with no interaction between sex categories. No changes  
37 in the calculated slope 2 were observed. These findings suggest that a single session of PBC  
38 reduces the muscular metabolic O<sub>2</sub> needs at rest; however, it does not alter the vascular ability  
39 to provide O<sub>2</sub> to the myocytes.

40

41 **Keywords:** Cryotherapy, Cryostimulation, Vascular responsiveness, Muscular oxygenation,

42

## 43 **Introduction**

44 Over the last decades, cryostimulation has gained popularity in sports medicine as a recovery  
45 technique for athletes or an adjunct treatment to relieve symptoms from various chronic  
46 diseases [2]. A growing number of studies have shown the beneficial effects of whole-body  
47 cryotherapy/cryostimulation (WBC) and partial-body cryotherapy/cryostimulation (PBC) on  
48 physiological mechanisms such as oxidative stress [8,35], inflammation [1,9,36], and cardiac  
49 autonomic control [15,20,21,39,41]. Such physiological responses to cryotherapy result from  
50 thermoregulatory processes triggered by skin temperature reduction [29]. Feedforward  
51 mechanisms, initiated by the brain's thermoregulatory center, lead to sympathetic stimulation  
52 of peripheral blood vessels and vasoconstriction [6]. Consequently, blood volume is  
53 redirected towards the core, increasing central blood pressure and decreasing heart rate  
54 through autonomic modulations [32].

55 Another proposed hypothesis concerning the beneficial physiological effects of  
56 cryostimulation is that changes in muscle vascularization, such as a reduction of muscle blood  
57 flow, might participate in the decrease of inflammatory responses [30,40]. Muscular blood  
58 flow following a WBC session has previously been studied, especially in lower limb muscles.  
59 It has been reported that a WBC session reduces superficial femoral artery blood flow, as well  
60 as vastus lateralis and gastrocnemius oxygenation, immediately after the cold exposure [24].  
61 However, it also demonstrated superior muscle oxygenation in rugby players up to 20 minutes  
62 after the WBC session compared to the baseline pre-exposure levels [31]. This increased  
63 oxygenation following cold exposure is still not fully understood but might be linked to  
64 vascular responsiveness mechanisms such as nitric oxide (NO) dependent vasodilation  
65 regulated by endothelial function. Although the literature is scarce about the effect of WBC  
66 on endothelium functions, previous research reported a beneficial impact of WBC on  
67 endothelium inflammatory parameters in various populations [19,20]. NO production capacity  
68 was also increased in a healthy older population after a 24 WBC exposure[21]. This greater  
69 muscle oxygenation after exposure to cold may also be due to a greater vasodilatory effect  
70 than before exposure, also called the 'Hunting effect' [7,19]. To date, the extent of these  
71 results seems contradictory, and the effects of cryostimulation on muscular microcirculatory  
72 hemodynamic responses are poorly understood. Although the FMD technique is the gold  
73 standard measurement of vascular responsiveness, it cannot assess the downstream  
74 microvascular function. Therefore, the microvascular function is classically studied through  
75 near-infrared spectroscopy (NIRS) by monitoring the reperfusion slope after a vascular

76 occlusion [24–26]. Previous studies have already shown a correlation between FMD and the  
77 reperfusion slope assessed by vascular occlusion combined with NIRS [24,27] providing  
78 evidence for the reliability of this technique to assess endothelial function at the  
79 microcirculatory level.

80 .

81 Several studies have already evaluated that microvascular perfusion and muscle metabolic  
82 activity were reduced following cold-water immersion (CWI) protocols [17,18,42]. However,  
83 physiological responses to cold water immersion and WBC are not comparable and there still  
84 are significant research gaps concerning the effect of PBC on said variables, as well as a need  
85 to investigate whether sex differences impact microvascular and metabolic responses  
86 following PBC interventions.

87 Therefore, this study aimed to investigate the effect of a 3-min session of PBC at -150 °C on  
88 the microvascular responsiveness of the flexor digitorum superficialis muscle, blood pressure,  
89 skin temperature, and perceived sensation to cold in young, healthy male and female adults.

90

## 91 **Materials and Methods**

### 92 **Participants**

93 Twenty healthy male and female volunteers participated in this study (see Table 1 for  
94 participants' characteristics). All participants were young (mean age:  $22.3 \pm 3.3$  years),  
95 physically active, and unaccustomed to PBC or cold exposure. Height (m), Body mass (kg),  
96 and BMI ( $\text{kg}/\text{m}^2$ ) were significantly higher ( $p < 0.01$ ) in male participants compared to female  
97 participants. As recommended by the working group of cryotherapy/cryostimulation from the  
98 International Institute Refrigeration [3], volunteers presenting contraindications to cold  
99 exposure such as hypertension, coronary disease, arrhythmia, circulatory disorders, Raynaud's  
100 phenomenon (decreased blood flow to the extremities), cold allergies, pulmonary disease, and  
101 obstruction of the bronchus caused by cold exposure were excluded. Only female participants  
102 under monophasic contraception were included in the study to control the effects of  
103 oestrogens and progesterone in women. Before each experimental session, volunteers were  
104 expected to follow pre-experimental criteria such as restraining from physical exercise,  
105 caffeine, and alcohol 24h before the session. All volunteers were given an in-depth  
106 explanation of the protocol and experimental sessions (including risks and benefits). During

107 the first visit, participants provided written consent and anthropometrics characteristics were  
108 measured. The study protocol was approved by local Ethics committee (CERSTAPS, number:  
109 2019-26-05-34).

### 110 *Experimental design*

111 The experimental design is presented in Figure 1. This study was a randomized controlled  
112 crossover experiment consisting of two experimental sessions separated by one week and  
113 conducted at the same time of the day. Volunteers were randomly assigned to a control  
114 session (CONT, where the volunteers stood 3-min in a cryo-cabin at ambient temperature) or  
115 to a treatment session (PBC session, where the volunteers stood 3-min at -150°C) for the first  
116 session. They switched conditions for the second experimental session. Both experimental  
117 sessions were organized in Figure 1 in a counterbalanced fashion for the control and  
118 cryotherapy sessions. Upon arrival at the laboratory, male participants were asked to remain  
119 in their undergarments while female participants wore their two-piece swimsuits. Every  
120 participant was given a bathrobe to be worn between exposures.

### 121 **Experimental measurements**

#### 122 *Partial body cryotherapy*

123 The PBC session consisted of a 3-min cold exposure in a cryo-cabin (Cryojet, ICE MINI,  
124 Paris, France) at -150°C. Following the pre-cooling phase of the cryo-cabin set at -70 °C,  
125 participants were asked to enter the cryo-cabin wearing only a swimsuit (women) or  
126 underwear (men), gloves, socks, and slippers. Participants were also asked to remove all  
127 metallic jewellery to avoid cold-related injuries. The temperature of the cryo-cabin was set at  
128 -150°C for the experimental session and remained stable throughout the entire duration of the  
129 3-min exposure. During the control condition, participants were asked to follow the exact  
130 instructions. However, the cryostimulation cabin was set at room temperature.

#### 131 *Vascular responsiveness*

132 Microvascular responsiveness was assessed through the NIRS technique during a post-  
133 occlusive reactive hyperemia test. Oxyhemoglobin ([HbO<sub>2</sub>]) and deoxyhemoglobin ([HHb])  
134 values were recorded continuously during the post-occlusive reactive hyperemia test. The  
135 NIRS probe was placed on the muscle belly of the flexor digitorum superficialis. It was  
136 placed 5 cm away from the medial epicondyle toward the distal insertion. A tightened elastic  
137 strap secured the probe to reduce movement and external light intrusion. NIRS data were

138 continuously recorded during the test at a 10Hz frequency. A manual pneumatic cuff was  
139 placed around the arm, approximately 5 cm above the elbow, to perform the occlusion. The  
140 post-occlusive reactive hyperemia test lasted 9 minutes. It consisted of 1-minute baseline  
141 measurements, a 5-min occlusion with cuff pressure set at 250mmHg, and a 3-min  
142 reperfusion. The variation of [HbO<sub>2</sub>] and [HHb] values obtained during the occlusion and the  
143 beginning of the reperfusion phase represent a linear response allowing slope calculations and  
144 characterization of microvascular responsiveness (Figure 2). As previously described, the rate  
145 of decrease in [HbO<sub>2</sub>] Slope 1 is an indication of the O<sub>2</sub> consumption rate, whereas the rate of  
146 increase in [HbO<sub>2</sub>] slope 2 characterizes the microvascular responsiveness [13]. O<sub>2</sub>  
147 consumption can also be estimated through [HHb] variation since venous and arterial outflow  
148 are blocked during the occlusion. [HHb] variations can therefore be explained by changes in  
149 O<sub>2</sub> consumption, allowing the use of [HHb] slope 1 to characterize the O<sub>2</sub> consumption rate.  
150 [HHb] Slope 2 is also used as an indicator of microvascular responsiveness as the decrease  
151 rate of [HHb] is impacted by the vascular motricity of capillaries.

152 Muscular metabolic O<sub>2</sub> consumption during arterial occlusion can also be calculated from the  
153 [HbO<sub>2</sub>] slope decrease. [HbO<sub>2</sub>] concentration variations are expressed in μM.s<sup>-1</sup> and  
154 converted to mlO<sub>2</sub>.min<sup>-1</sup>.100g<sup>-1</sup> using the mVO<sub>2</sub> equation:

$$(mVO_2 = Abs(((\Delta O_2 Hb * 60) / (10 * 1.04)) * 4) * 22.4 / 1000)$$

155 Where 1.04 kg.L<sup>-1</sup> represents muscle density, 22.4 L is the molar volume of gas, and 1:4 is the  
156 molecular ratio between Hb and O<sub>2</sub>.

### 157 *Skin temperatures*

158 Using a thermal camera Fluke Tis65 (FLUKE Corp, Washington), skin temperatures of the  
159 head, left and right arms and trunk were recorded immediately before and after the PBC  
160 exposure or control condition. The thermal camera emissivity was set to 0.98 which is the  
161 recommended level for the assessment of skin temperature by thermal imaging. The Fluke  
162 connect software (FLUKE Corp, Washington) calculated average temperatures by area.  
163 Participants were asked to remain standing in the anatomical position approximately 3 meters  
164 away from the thermal camera during the capture of the thermogram.

### 165 *Perceptual measures*

166 Subjective measures for thermal perception were assessed through a visual analogue scale  
167 (VAS). Immediately before and after the PBC session or control session, participants' thermal

168 perception was assessed and recorded using an 11 points VAS from 0 to 10 (0 = neutral, 2 =  
169 slightly cool, 4 = cool, 6 = cold, 8 = very cold, 10 = unbearably cold) (Lundgren et al. 2014).

## 170 **Statistical analysis and data distribution tests**

171 Data were presented as mean  $\pm$  standard deviation (SD). Data normality was evaluated by the  
172 Shapiro-Wilk test. A 3-way analysis of variance (Condition [CONT VS PBC] x sex [Male VS  
173 Female] x Time [Pre VS Post]) for repeated measures was performed to analyze the effect of  
174 cryotherapy, contingent upon sex, on vascular responsiveness, muscular O<sub>2</sub> consumption rate,  
175 HRV parameters, blood pressure, skin temperature, and cold perception. When a significant  
176 interaction between Time, Condition, or Sex was found, post-hoc analysis was performed  
177 using the Tukey HSD test. For all statistical analyses, alpha was set at  $p < 0.05$  to establish  
178 significance.

## 179 **Results**

### 180 *Skin temperatures*

181 Skin temperatures of the right and left arms, head, and trunk are displayed in Figure 3. Before  
182 the PBC exposure, all skin temperatures were similar between the control and PBC group.  
183 The ANOVA revealed a significant interaction between time and condition for the right and  
184 left arm and the head ( $P < 0.01$ ). Skin temperatures significantly decreased Post-PBC for the  
185 right arm (Pre-PBC:  $33.32 \pm 1.74$  vs Post-PBC:  $23.67 \pm 3.33$  °C,  $P < 0.01$ ), left arm (Pre-PBC:  
186  $32.89 \pm 2.06$  vs Post-PBC:  $24.07 \pm 3.07$  °C,  $P < 0.01$ ), and head (Pre-PBC:  $33.10 \pm 2.02$  vs  
187 Post-PBC:  $29.87 \pm 3.45$  °C,  $P < 0.01$ ) resulting in significantly lower skin temperatures for the  
188 right and left arm and for the head following *post-PBC* compared to Post-control. A  
189 significant interaction between time, condition, and sex were also found for trunk skin  
190 temperatures ( $P < 0.05$ ). Trunk skin temperature significantly decreased *post-PBC* both in  
191 males (*pre-PBC*:  $34.75 \pm 0.91$  vs *post-PBC*:  $25.70 \pm 1.18$  °C,  $P < 0.01$ ) and females (*pre-PBC*:  
192  $32.48 \pm 1.49$  vs Post-PBC:  $21.31 \pm 3.94$  °C,  $P < 0.01$ ) and significant lower skin  
193 temperature values were observed *post-PBC* compared to Post-control. After the PBC  
194 session, trunk skin temperature was also lower in the female group compared to the male  
195 group (*post-PBC* female:  $21.31 \pm 3.94$  vs *post-PBC* male:  $25.70 \pm 1.18$  °C,  $P < 0.01$ ).

### 196 *mVO<sub>2</sub>*

197 Data for mVO<sub>2</sub> are presented in Figure 4. Baseline values for mVO<sub>2</sub> were identical between  
198 the control and PBC groups. The ANOVA revealed a significant interaction between time and



199 condition ( $P < 0.01$ ). Post-hoc analyses indicated a significantly lower value of  $mVO_2$  *post-*  
200 *PBC* compared to *pre-PBC* value (*pre-PBC*:  $18.47 \pm 5.73$  vs. *post-PBC*:  $14.75 \pm 3.94$   
201  $\mu\text{MO}_2\text{Hb}\cdot\text{s}^{-1}$ ,  $P < 0.01$ )

#### 202 *Slopes [HbO<sub>2</sub>]*

203 Data of [HbO<sub>2</sub>] slopes 1 and 2 are presented in Figure 5.

204 The ANOVA revealed a significant interaction between time, condition, and sex. Post-hoc  
205 analyses revealed a significant reduction of [HbO<sub>2</sub>] slope 1 after the PBC session only in male  
206 subjects (*pre-PBC*:  $-0.24 \pm 0.11$  vs *post-PBC*:  $-0.15 \pm 0.08$   $\text{s}^{-1}$ ,  $P < 0.05$ ). Slope 2 was not  
207 significantly different between condition and time of measurements.

#### 208 *Slope [HHb]*

209 Data of [HHb] slopes 1 and 2 are presented in Figure 5.

210 A significant interaction between time and condition ( $P < 0.01$ ) was found, and post-hoc  
211 analyses revealed statistical differences between groups. A significant reduction of the slope 1  
212 in *post-PBC* compared to *pre-PBC* (*pre-PBC*:  $0.24 \pm 0.16$  vs *post-PBC*:  $0.18 \pm 0.10$   $\text{s}^{-1}$ ,  $P$   
213  $< 0.05$ ) was found with no interaction between sex categories. Slope 2 was not significantly  
214 different between condition and time of measurements.

#### 215 *Thermal sensation*

216 Thermal sensation data are presented in Figure 6. Baseline values were not statistically  
217 different between the control and PBC groups, and no effect of sex was found in cold thermal  
218 sensation ratings. The ANOVA revealed a significant interaction between time and condition  
219 ( $P < 0.01$ ), and post-hoc analyses showed a significantly higher perception of cold *post-PBC*  
220 when compared to *pre-PBC* and *post-control* (*post-PBC*:  $8.25 \pm 0.64$  vs *pre-PBC*:  $0.80 \pm 1.00$   
221 vs *post-control*:  $1.25 \pm 1.07$ ;  $P < 0.01$  for each comparison).

222

## 223 **Discussion**

224 This study aimed to investigate the effects of partial-body cryostimulation (PBC) on  
225 microvascular responsiveness and muscular metabolic O<sub>2</sub> consumption rate ( $mVO_2$ ). Using a  
226 randomized controlled crossover design, the effects of a 3-min PBC session at  $-150$  °C were  
227 tested on microvascular responsiveness of a superficial muscle, muscular metabolic O<sub>2</sub>

228 consumption, skin temperature, and perceived cold sensation in young, healthy male and  
229 female subjects. The main results of this study are a reduction of [HHb] slope 1 and  $mVO_2$   
230 after a PBC session, both in male and female participants. Moreover, a PBC session only  
231 reduced [HbO<sub>2</sub>] slope 1 in male participants.

### 232 *Microvascular responsiveness and metabolic consumption*

233 In our male participants, *post-PBC* slope 1 of [HbO<sub>2</sub>] was lower than *pre-PBC* slope 1 of  
234 [HbO<sub>2</sub>], demonstrating that cryostimulation induced a lower oxygen consumption in these  
235 subjects. However, this phenomenon was not observed in our female participants.  
236 Nonetheless, while [HbO<sub>2</sub>] slope 2 was similar under control and PBC conditions for both  
237 sexes, a lower value was observed in females compared to men in both conditions. This result  
238 highlights that men exhibit a higher reperfusion capacity than women, with or without  
239 previous cryotherapy exposure. While gender differences in reactive hyperaemia have already  
240 been highlighted, with females showing reduced endothelial reactivity compared to men [22],  
241 our study, to our knowledge, is the first to highlight a between-sex difference in vascular  
242 responsiveness in healthy young adults.

243 Moreover, our study revealed that cryostimulation affects other parameters of microvascular  
244 function independently of the participant's gender. PBC impacted [HHb] response during the  
245 occlusion phase, as demonstrated by a reduced [HHb] slope 1 following PBC. In addition,  
246  $mVO_2$  was reduced *post-PBC* compared to *pre-PBC*, indicating a lower O<sub>2</sub> metabolic  
247 consumption induced by cryotherapy. These results are consistent with previous studies that  
248 have established that cold exposure leads to immediate peripheral vasoconstriction and blood  
249 flow restriction [12]. Yanagisawa et al. (2007) reported that tissue oxygenation decreased  
250 during cooling. The total Hb drop was mainly due to [HbO<sub>2</sub>] decline [42]. Also, Stanley et al.  
251 (2014) observed that cold water immersion slowed VO<sub>2</sub> on-kinetics, leading to increased  
252 anaerobic contribution during high-intensity intermittent exercise compared to the control  
253 condition [37].

### 254 *Skin temperature and cold perception*

255 Skin temperature was significantly lower in male and female participants following the PBC  
256 session. Although both males and females displayed significant reductions in trunk skin  
257 temperature following PBC, female participants had significantly lower trunk skin  
258 temperature than male participants. This particular result highlights a difference in response to  
259 cryotherapy between sexes, a phenomenon that has already been reported. Hammond et al.

260 (2014) observed that skin temperature decrease was more significant in females than males  
261 after a 2-min WBC session [14]. They also found a negative association between the  
262 percentage of skin temperature variation after cryotherapy and body fat %. Indeed, apart from  
263 the effect of *sex per se*, differences in body composition could partially explain differences  
264 observed in skin temperature after cold exposure. Although we do not have data on body fat  
265 % in our participants, we assume that females had a higher fat composition than male subjects  
266 (as women generally usually present 14% more body fat than men [5]). Therefore, our  
267 findings support the hypothesis that individuals with higher adiposity significantly reduce  
268 skin temperature after cooling [4,23,38]*Specific responses in males compared to women*

269 Our results confirm that the physiological responses to cold differ between men and women.  
270 Indeed, as mentioned above, women seem to have a greater drop in skin temperature than men  
271 at an equivalent single dose. This decrease in temperature, probably due to the difference in  
272 body composition, will not have the same impact on physiological responses to cold. Authors  
273 have reported that hormonal responses differ between men and women [34]. The question  
274 then arises whether the duration of 3 minutes is the optimal duration for men and women.  
275 According to some authors [28], the duration for women should be shorter than for men. Our  
276 results demonstrated that an equivalent dose of cryotherapy could have different effects  
277 depending on the participant's anthropometric characteristics. These confirm that  
278 cryostimulation should not be a one-size-fits-all approach and that more efficient results will  
279 be yielded from a more individualized approach.

280 *What is the implication of the responses obtained in the recovery process?*

281 Following the physical activity, a lower vascularization induced by cryostimulation at the  
282 skeletal muscle level may be involved in the recovery process. Several studies demonstrated a  
283 faster and more effective recovery from exercise-induced muscle damage following a WBC  
284 session. Improvements in recovery have been characterized by an attenuation of DOMS,  
285 muscle swelling, and pain [10,11,16]. Such improvements may be linked to a localized  
286 reduced blood flow following PBC. Moreover, since reduced muscle vascularization leads to  
287 reduced O<sub>2</sub> consumption by muscles, it is plausible to assume that there is an associated  
288 reduction in the production of reactive oxygen species. Consequently, oxidative stress would  
289 be reduced, representing one of the mechanisms involved in the improvements observed for  
290 muscle recovery following cryostimulation [26,33]. With that said, further research is needed  
291 to determine the potential effect of cryotherapy on reducing the production and release of free  
292 radicals by the muscles. Future studies would be interesting to understand the possible

293 implication of the vascular mechanisms observed in this study and the effects of cryotherapy  
294 on post-exercise recovery of power qualities, marked recently by the meta-analysis of Moore  
295 and al. (2022) [27].The results of this study must be interpreted in the light of certain  
296 experimental considerations. The NIRS technique used in the protocol to assess microvascular  
297 reperfusion after the occlusion should be coupled with FMD techniques for a better  
298 understanding of the endothelial function. However, FMD is a highly operator-dependent  
299 measure, and the time to set up the measurement may vary between sessions and be  
300 incompatible with the experimental time points.

### 301 **Conclusion**

302 This study provides novel data concerning the effect of PBC on peripheral microvascular  
303 physiology. Our main finding is that a PBC session does not alter the microvascular  
304 vasodilation ability both in male and female subjects. These results suggest that a PBC session  
305 does not change the ability of the capillaries to provide O<sub>2</sub> to the muscles. The second  
306 noteworthy finding of this study is that a PBC session reduces the O<sub>2</sub> consumption of  
307 peripheral muscles at rest, especially in male participants. This could be explained by a  
308 reduction of muscle's metabolic rate at rest. The results provided in this study, in their  
309 globality, provide original information concerning the effect of PBC on peripheral vascular  
310 physiology.

311

### 312 **Disclosures**

313 No conflicts of interest, financial or otherwise, are declared by the authors

### 314 **Author contributions**

315 DT, BD, OD conceived and designed the research protocol; DT, BD, OD performed  
316 experiments; analyzed data; interpreted results of the investigation; DT, OD prepared the  
317 figures; DT, BD, WD, JL, OD, RM,CAC drafted the manuscript; DT, BD, WD, JL, OD,  
318 RM,CAC edited and revised the manuscript; DT, BD, WD, JL, OD, RM,CAC approved the  
319 final version of manuscript

320

321 **References**

- 322 [1] G. Banfi, G. Melegati, A. Barassi, G. Dogliotti, G. Melzi d'Eril, B. Dugué, M.M. Corsi,  
323 Effects of whole-body cryotherapy on serum mediators of inflammation and serum  
324 muscle enzymes in athletes, *Journal of Thermal Biology*. 34 (2009) 55–59.  
325 <https://doi.org/10.1016/j.jtherbio.2008.10.003>.
- 326 [2] R. Bouzigon, F. Grappe, G. Ravier, B. Dugué, Whole- and partial-body  
327 cryostimulation/cryotherapy: Current technologies and practical applications, *Journal of*  
328 *Thermal Biology*. 61 (2016) 67–81. <https://doi.org/10.1016/j.jtherbio.2016.08.009>.
- 329 [3] R. Bouzigon, O. Dupuy, I. Tiemessen, M. Nardi, J.P. Bernard, M. Thibaud, D. Theurot,  
330 E. Miller, G. Lombardi, B. Dugué, Cryostimulation for Post-exercise Recovery in  
331 Athletes: A Consensus and Position Paper, *Frontiers in Sports and Active Living*.  
332 3:688828 (2021). <https://doi.org/10.3389/fspor.2021.688828>.
- 333 [4] R. Bouzigon, G. Ravier, B. Dugué, F. Grappe, Thermal Sensations During a Partial-  
334 Body Cryostimulation Exposure in Elite Basketball Players, *Journal of Human Kinetics*.  
335 62 (2018) 55–63. <https://doi.org/10.1515/hukin-2017-0158>.
- 336 [5] R.L. Burse, Sex Differences in Human Thermoregulatory Response to Heat and Cold  
337 Stress, *Hum Factors*. 21 (1979) 687–699. <https://doi.org/10.1177/001872087912210606>.
- 338 [6] N. Charkoudian, Skin Blood Flow in Adult Human Thermoregulation: How It Works,  
339 When It Does Not, and Why, *Mayo Clinic Proceedings*. 78 (2003) 603–612.  
340 <https://doi.org/10.4065/78.5.603>.
- 341 [7] A.S. Dana Jr., I.H. Rex Jr., M.H. Samitz, The Hunting Reaction, *Archives of*  
342 *Dermatology*. 99 (1969) 441–450.  
343 <https://doi.org/10.1001/archderm.1969.01610220069012>.
- 344 [8] B. Dugué, J. Smolander, T. Westerlund, J. Oksa, R. Nieminen, E. Moilanen, M.  
345 Mikkelsen, Acute and long- term effects of winter swimming and whole- body  
346 cryotherapy on plasma antioxidative capacity in healthy women, *Scandinavian Journal*  
347 *of Clinical and Laboratory Investigation*. 65 (2005) 395–402.  
348 <https://doi.org/10.1080/00365510510025728>.
- 349 [9] O. Dupuy, W. Douzi, D. Theurot, L. Bosquet, B. Dugué, An Evidence-Based Approach  
350 for Choosing Post-exercise Recovery Techniques to Reduce Markers of Muscle  
351 Damage, Soreness, Fatigue, and Inflammation: A Systematic Review With Meta-  
352 Analysis, *Front. Physiol*. 9 (2018) 403. <https://doi.org/10.3389/fphys.2018.00403>.

- 353 [10] J.B. Ferreira-Junior, M. Bottaro, J.P. Loenneke, A. Vieira, C.A. Vieira, M.G. Bembem,  
354 Could whole-body cryotherapy (below  $-100^{\circ}\text{C}$ ) improve muscle recovery from muscle  
355 damage?, *Front Physiol.* 5 (2014) 247. <https://doi.org/10.3389/fphys.2014.00247>.
- 356 [11] B. Fonda, N. Sarabon, Effects of whole-body cryotherapy on recovery after hamstring  
357 damaging exercise: A crossover study: Whole-body cryotherapy after EIMD,  
358 *Scandinavian Journal of Medicine & Science in Sports.* (2013) n/a-n/a.  
359 <https://doi.org/10.1111/sms.12074>.
- 360 [12] B. Freire, J. Geremia, B.M. Baroni, M.A. Vaz, Effects of cryotherapy methods on  
361 circulatory, metabolic, inflammatory and neural properties: a systematic review, *Fisioter.*  
362 *Mov.* 29 (2016) 389–398. <https://doi.org/10.1590/0103-5150.029.002.AO18>.
- 363 [13] V. Gerovasili, S. Dimopoulos, G. Tzanis, M. Anastasiou-Nana, S. Nanas, Utilizing the  
364 vascular occlusion technique with NIRS technology, *International Journal of Industrial*  
365 *Ergonomics.* 40 (2010) 218–222. <https://doi.org/10.1016/j.ergon.2009.02.004>.
- 366 [14] L.E. Hammond, S. Cuttell, P. Nunley, J. Meyler, Anthropometric Characteristics and  
367 Sex Influence Magnitude of Skin Cooling following Exposure to Whole Body  
368 Cryotherapy, *BioMed Research International.* 2014 (2014) 1–7.  
369 <https://doi.org/10.1155/2014/628724>.
- 370 [15] C. Hausswirth, K. Schaal, Y. Le Meur, F. Bieuzen, J.-R. Filliard, M. Volondat, J. Louis,  
371 Parasympathetic Activity and Blood Catecholamine Responses Following a Single  
372 Partial-Body Cryostimulation and a Whole-Body Cryostimulation, *PLoS ONE.* 8 (2013)  
373 e72658. <https://doi.org/10.1371/journal.pone.0072658>.
- 374 [16] C. Hausswirth, J. Louis, F. Bieuzen, H. Pournot, J. Fournier, J.-R. Filliard, J.  
375 Brisswalter, Effects of Whole-Body Cryotherapy vs. Far-Infrared vs. Passive Modalities  
376 on Recovery from Exercise-Induced Muscle Damage in Highly-Trained Runners, *PLoS*  
377 *ONE.* 6 (2011) e27749. <https://doi.org/10.1371/journal.pone.0027749>.
- 378 [17] E. Hohenauer, T. Deliens, P. Clarys, R. Clijsen, Perfusion of the skin's microcirculation  
379 after cold- water immersion ( $10^{\circ}\text{C}$ ) and partial- body cryotherapy ( $-135^{\circ}\text{C}$ ), *Skin Res*  
380 *Technol.* 25 (2019) 677–682. <https://doi.org/10.1111/srt.12703>.
- 381 [18] M. Ihsan, G. Watson, M. Lipski, C.R. Abbiss, Influence of Postexercise Cooling on  
382 Muscle Oxygenation and Blood Volume Changes:, *Medicine & Science in Sports &*  
383 *Exercise.* 45 (2013) 876–882. <https://doi.org/10.1249/MSS.0b013e31827e13a2>.
- 384 [19] J.M. Johnson, C.T. Minson, D.L. Kellogg, Cutaneous Vasodilator and Vasoconstrictor  
385 Mechanisms in Temperature Regulation, in: R. Terjung (Ed.), *Comprehensive*  
386 *Physiology*, 1st ed., Wiley, 2014: pp. 33–89. <https://doi.org/10.1002/cphy.c130015>.

- 387 [20] J. Louis, K. Schaal, F. Bieuzen, Y. Le Meur, J.-R. Filliard, M. Volondat, J. Brisswalter,  
388 C. Hausswirth, Head Exposure to Cold during Whole-Body Cryostimulation: Influence  
389 on Thermal Response and Autonomic Modulation, PLOS ONE. 10 (2015) e0124776.  
390 <https://doi.org/10.1371/journal.pone.0124776>.
- 391 [21] J. Louis, D. Theurot, J.-R. Filliard, M. Volondat, B. Dugué, O. Dupuy, The use of  
392 whole-body cryotherapy: time- and dose-response investigation on circulating blood  
393 catecholamines and heart rate variability, Eur J Appl Physiol. 120 (2020) 1733–1743.  
394 <https://doi.org/10.1007/s00421-020-04406-5>.
- 395 [22] B.-J. Martin, T.J. Anderson, Risk prediction in cardiovascular disease: The prognostic  
396 significance of endothelial dysfunction, Canadian Journal of Cardiology. 25 (2009) 15A-  
397 20A. [https://doi.org/10.1016/S0828-282X\(09\)71049-9](https://doi.org/10.1016/S0828-282X(09)71049-9).
- 398 [23] T. Matsumoto, T. Miyawaki, H. Ue, T. Kanda, C. Zenji, T. Moritani, Autonomic  
399 responsiveness to acute cold exposure in obese and non-obese young women, Int J Obes.  
400 23 (1999) 793–800. <https://doi.org/10.1038/sj.ijo.0800928>.
- 401 [24] C. Mawhinney, D.A. Low, H. Jones, D.J. Green, J.T. Costello, W. Gregson, Cold Water  
402 Mediates Greater Reductions in Limb Blood Flow than Whole Body Cryotherapy.,  
403 Medicine & Science in Sports & Exercise. 49 (2017) 1252–1260.  
404 <https://doi.org/10.1249/MSS.0000000000001223>.
- 405 [25] K.M. McLay, F.Y. Fontana, J.P. Nederveen, F.F. Guida, D.H. Paterson, S. Pogliaghi,  
406 J.M. Murias, Vascular responsiveness determined by near-infrared spectroscopy  
407 measures of oxygen saturation: Assessment of vascular responsiveness by near-infrared  
408 spectroscopy, Exp Physiol. 101 (2016) 34–40. <https://doi.org/10.1113/EP085406>.
- 409 [26] E. Miller, Ł. Markiewicz, J. Saluk, I. Majsterek, Effect of short-term cryostimulation on  
410 antioxidative status and its clinical applications in humans, Eur J Appl Physiol. 112  
411 (2012) 1645–1652. <https://doi.org/10.1007/s00421-011-2122-x>.
- 412 [27] E. Moore, J.T. Fuller, J.D. Buckley, S. Saunders, S.L. Halson, J.R. Broatch, C.R.  
413 Bellenger, Impact of Cold-Water Immersion Compared with Passive Recovery  
414 Following a Single Bout of Strenuous Exercise on Athletic Performance in Physically  
415 Active Participants: A Systematic Review with Meta-analysis and Meta-regression,  
416 Sports Med. 52 (2022) 1667–1688. <https://doi.org/10.1007/s40279-022-01644-9>.
- 417 [28] G. Polidori, R. Elfahem, B. Abbes, F. Bogard, F. Legrand, B. Bouchet, F. Beaumont,  
418 Preliminary study on the effect of sex on skin cooling response during whole body  
419 cryostimulation (−110 °C): Modeling and prediction of exposure durations,  
420 Cryobiology. 97 (2020) 12–19. <https://doi.org/10.1016/j.cryobiol.2020.10.014>.

- 421 [29] A.A. Romanovsky, Skin temperature: its role in thermoregulation, *Acta Physiol (Oxf)*.  
422 210 (2014) 498–507. <https://doi.org/10.1111/apha.12231>.
- 423 [30] K.-D. Schaser, A.C. Disch, J.F. Stover, A. Lauffer, H.J. Bail, T. Mittlmeier, Prolonged  
424 superficial local cryotherapy attenuates microcirculatory impairment, regional  
425 inflammation, and muscle necrosis after closed soft tissue injury in rats, *Am J Sports*  
426 *Med.* 35 (2007) 93–102. <https://doi.org/10.1177/0363546506294569>.
- 427 [31] J. Selfe, J. Alexander, J.T. Costello, K. May, N. Garratt, S. Atkins, S. Dillon, H. Hurst,  
428 M. Davison, D. Przybyla, A. Coley, M. Bitcon, G. Littler, J. Richards, The Effect of  
429 Three Different (-135°C) Whole Body Cryotherapy Exposure Durations on Elite Rugby  
430 League Players, *PLoS ONE*. 9 (2014) e86420.  
431 <https://doi.org/10.1371/journal.pone.0086420>.
- 432 [32] N. Shibahara, H. Matsuda, K. Umeno, Y. Shimada, T. Itoh, K. Terasawa, The responses  
433 of skin blood flow, mean arterial pressure and R-R interval induced by cold stimulation  
434 with cold wind and ice water, *J Auton Nerv Syst.* 61 (1996) 109–115.  
435 [https://doi.org/10.1016/s0165-1838\(96\)00065-3](https://doi.org/10.1016/s0165-1838(96)00065-3).
- 436 [33] A.F. Siqueira, A. Vieira, G.V. Ramos, R. de C. Marqueti, T. de F. Salvini, G.O. Puntel,  
437 J.L.Q. Durigan, Multiple cryotherapy applications attenuate oxidative stress following  
438 skeletal muscle injury, *Redox Rep.* 22 (2017) 323–329.  
439 <https://doi.org/10.1080/13510002.2016.1239880>.
- 440 [34] R. Solianik, A. Skurvydas, A. Vitkauskienė, M. Brazaitis, Gender-specific cold  
441 responses induce a similar body-cooling rate but different neuroendocrine and immune  
442 responses, *Cryobiology.* 69 (2014) 26–33.  
443 <https://doi.org/10.1016/j.cryobiol.2014.04.015>.
- 444 [35] A. Stanek, A. Cholewka, T. Wielkoszyński, E. Romuk, A. Sieroń, Decreased Oxidative  
445 Stress in Male Patients with Active Phase Ankylosing Spondylitis Who Underwent  
446 Whole-Body Cryotherapy in Closed Cryochamber, *Oxidative Medicine and Cellular*  
447 *Longevity.* 2018 (2018) 1–9. <https://doi.org/10.1155/2018/7365490>.
- 448 [36] A. Stanek, A. Cholewka, T. Wielkoszyński, E. Romuk, A. Sieroń, Whole-Body  
449 Cryotherapy Decreases the Levels of Inflammatory, Oxidative Stress, and  
450 Atherosclerosis Plaque Markers in Male Patients with Active-Phase Ankylosing  
451 Spondylitis in the Absence of Classical Cardiovascular Risk Factors, *Mediators of*  
452 *Inflammation.* 2018 (2018) 1–11. <https://doi.org/10.1155/2018/8592532>.



- 453 [37] J. Stanley, J.M. Peake, J.S. Coombes, M. Buchheit, Central and peripheral adjustments  
454 during high-intensity exercise following cold water immersion, *Eur J Appl Physiol.* 114  
455 (2014) 147–163. <https://doi.org/10.1007/s00421-013-2755-z>.
- 456 [38] J.M. Stocks, N.A.S. Taylor, M.J. Tipton, J.E. Greenleaf, Human physiological responses  
457 to cold exposure, *Aviat Space Environ Med.* 75 (2004) 444–457.
- 458 [39] D. Theurot, B. Dugué, D. Wafa, P. Guitet, J. Louis, O. Dupuy, Impact of acute partial-  
459 body cryostimulation on cognitive performance, cerebral oxygenation, and cardiac  
460 autonomic activity, *Scientific Reports.* 11 (2021) 1–11. [https://doi.org/10.1038/s41598-  
461 021-87089-y](https://doi.org/10.1038/s41598-021-87089-y).
- 462 [40] G. Vieira Ramos, C.M. Pinheiro, S.P. Messa, G.B. Delfino, R. de C. Marqueti, T. de F.  
463 Salvini, J.L.Q. Durigan, Cryotherapy Reduces Inflammatory Response Without Altering  
464 Muscle Regeneration Process and Extracellular Matrix Remodeling of Rat Muscle, *Sci  
465 Rep.* 6 (2016) 18525. <https://doi.org/10.1038/srep18525>.
- 466 [41] T. Westerlund, A. Uusitalo, J. Smolander, M. Mikkelsen, Heart rate variability in  
467 women exposed to very cold air (−110°C) during whole-body cryotherapy, *Journal of  
468 Thermal Biology.* 31 (2006) 342–346. <https://doi.org/10.1016/j.jtherbio.2006.01.004>.
- 469 [42] O. Yanagisawa, T. Homma, T. Okuwaki, D. Shima, H. Takahashi, Effects of cooling  
470 on human skin and skeletal muscle, *European Journal of Applied Physiology.* 100  
471 (2007) 737–745. <https://doi.org/10.1007/s00421-007-0470-3>.
- 472
- 473
- 474

475 **Figures**


476 **Figure 1**

477

478

479

480



481

482

483 **Figure 2**

484


485

486

487

488

489



490


491

492 **Figure 3**

493

494

495



496

497

498


499

500 **Figure 4**

501

502

503



504 **Figure 5**


505

506

507

508

509



510 **Figure 6**

511


512

513

514

515

516



517

518 **Captions**

519 **Figure 1:** Experimental protocol

520 **Figure 2:** Presentation of slope 1 and slope 2 during a post-occlusive reactive hyperemia test.

521 **Figure 3:** Right and left arm (panel A and B), trunk (panel C), and head (panel D)  
522 temperature before and after a PBC session for male and female participants. \* Significant  
523 ( $P<0.05$ )

524 **Figure 4:**  $m\dot{V}O_2$  before and after experimental conditions. \* Significant ( $P<0.05$ )

525 **Figure 5:**  $[HbO_2]$  slope 1 (panel A) and slope 2 (panel B),  $[HHb]$  slope 1 (panel C) and slope  
526 2 (panel D) before and after PBC session for male and female participants. \* Significant  
527 ( $P<0.05$ )

528 **Figure 6:** Thermal sensation rating before and after PBC session in male and female  
529 participants. \* Significant ( $P<0.05$ )

530



531 **Tables**

532

Table 1 : Participant characteristics

	Age (Years)	Height (m)	Weight (Kg)	BMI (Kg/m <sup>2</sup> )
Males	23.10 ± 4.51	1.78 ± 0.09 *	81.00 ± 12.88 *	25.55 ± 2.36 *
Females	21.50 ± 1.27	1.65 ± 0.60	58.26 ± 8.49	21.40 ± 2.41

Data are presented mean ±SD; \* significant difference between male and female p<0.05

533

