



HAL
open science

Nd isotopes as a tracer of dust trapped by peat bogs: A reminder of basics. Comment on: “ Neodymium isotopes in peat reveal past local environmental disturbances ” by Marcisz et al. (2023)

Gaël Le Roux, Nathalie Fagel, Jack Longman, Nadine Mattielli, Robert Mcculloch, François de Vleeschouwer

► To cite this version:

Gaël Le Roux, Nathalie Fagel, Jack Longman, Nadine Mattielli, Robert Mcculloch, et al.. Nd isotopes as a tracer of dust trapped by peat bogs: A reminder of basics. Comment on: “ Neodymium isotopes in peat reveal past local environmental disturbances ” by Marcisz et al. (2023). *Science of the Total Environment*, 2023, 882, pp.163379. 10.1016/j.scitotenv.2023.163379 . hal-04246661

HAL Id: hal-04246661

<https://hal.science/hal-04246661>

Submitted on 17 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Public Domain

THIS IS A NON PEER-REVIEWED PREPRINT.

Nd isotopes as a tracer of dust trapped by peat bogs: A reminder of basics. Comment on: « Neodymium isotopes in peat reveal past local environmental disturbances » by Marcisz et al. (2023)

Gaël Le Roux¹, Nathalie Fagel², Jack Longman³, Nadine Mattielli⁴, Robert D. McCulloch⁵, François De Vleeschouwer^{6*}

1. Laboratoire Ecologie Fonctionnelle et Environnement (UMR5245 CNRS/UPS/INPT), Campus Ensat, Avenue de l'Agrobiopole, BP 32607, Auzeville tolosane, 31326 Castanet-Tolosan, France
2. Laboratoire Argiles, Géochimie et Environnements sédimentaires (AGEs). Département de Géologie, Université de Liège, Allée du 6 août 14, Liège 4000, Belgique
3. Department of Geography and Environmental Sciences, Northumbria University, Newcastle-upon-Tyne, NE1 8ST, UK.
4. Laboratoire G-Time (*Geochemistry & Geophysics: Tephra, Isotopes, Minerals and Earthquakes*), DGES, Université Libre de Bruxelles (ULB), Av. FD Roosevelt, 1050 Brussels, Belgium
5. Centro de Investigación en Ecosistemas de la Patagonia (CIEP), José de Moraleda, Coyhaique, Aysén, Chile.
6. Instituto Franco-Argentino para el Estudio del Clima y sus Impactos (IRL IFAECI/CNRS-CONICET-UBA-IRD), Dpto. de Ciencias de la Atmosfera y los Océanos, FCEN, Universidad de Buenos Aires, Intendente Guiraldes 2160 - Ciudad Universitaria, Pabellon II - 2do. Pis, (C1428EGA) Ciudad Autónoma de Buenos Aires, Argentina

*Corresponding author: fdevleeschouwer@cima.fcen.uba.ar

Neodymium (Nd) isotopes are one of a suite of radiogenic isotopes whose ratios in any geomaterial constitute a sensitive and faithful fingerprint signature of the material's original reservoir. A plethora of studies have demonstrated the relevance of Nd isotopes as powerful geochemical tracers of rock and sediment provenance at various geological time scales (e.g. O'Nions et al., 1983; Goldstein et al., 1984; Grousset et al., 1988; Dia et al. 1992). More recently, they have been applied to a range of palaeo-source studies, including palaeoceanography (e.g. Bazhenova et al., 2017; Fagel et al., 2004; Frank, 2002; Grousset et al., 1992; Lacan and Jeandel, 2004) and palaeo-dust studies (e.g. Aarons et al., 2016; Beaudon et al., 2022; Grousset et al., 1992; Longman et al., 2022; Saitoh et al., 2011; Sharifi et al., 2018; Simonsen et al., 2019; Stewart et al., 2023; Struve et al., 2022). However, since the first Nd isotope study by Le Roux et al. (2012), only a few studies have been published in peat bogs (e.g. Allan et al., 2013; Fagel et al., 2014; Fialkewicz-Koziel 2016, Pratte et al., 2017, Sharifi et al., 2018; Vanneste et al., 2015, 2016). There are several reasons for this: firstly, Nd isotopes must be measured using multicollector mass spectrometer such as MC-ICP-MS (Multicollector Inductively Coupled Plasma Mass Spectrometer) or TIMS (Thermal Ionization mass Spectrometer), which are not readily available in every laboratory. Secondly, Nd has to be separated by a chemical process to ensure that there are no Rare Earth Elements (REE) interferences. These procedures are not available to everyone, expensive and time consuming. Another simpler reason is that the dust trapped by peat bogs can be traced by other, more easily measured proxies, and that in fact the use of Nd isotopes is only the final step in a long process of geochemical characterization of the past atmospheric fallout in a peat bog. These other tracers are: lithogenic and REE concentrations and ratios, acid insoluble ash content or mineral characterization by XRD, SEM or grain size analyses. Presently, artificial intelligence has also been employed (see our recent dialog with Chat GPT in S.I.).

Prior to any inorganic geochemical investigation in peat aiming at reconstructing atmospheric dust fluxes (mineral dust or pollutants), one should apply rigorous site selection criteria. This is because atmospheric deposition reconstruction using peat bogs is based on 2 paradigms:

1. The selected site is ombrotrophic (i.e. it is a peat bog, not a minerotrophic fen): it is exclusively fed by atmospheric inputs, in very acidic (pH generally below 4) and anoxic conditions;
2. The dust that is deposited in a peat bog does not - most of the time - represent local dust, but instead originate from various distal sources.

The first paradigm has been verified many times by independent geochemical comparisons, in particular the comparison between airborne pollutants measured in precipitation and herbaria vs. records in peat bogs or mosses (e.g. Bindler, 2011; Dunlap et al., 1999; Farmer et al., 2002; Kempter et al., 2010, 2017; Markert, 1991; Novak et al., 2008; Le Roux et al., 2005; Pech et al., 2022; Rausch et al., 2005; Shotyk et al., 2015; Steinnes, 1995; Weiss et al., 1999, 2002; Zupančič and Bozau, 2022). It should be noted that simply the presence of *Sphagnum* moss occurrence in a peatland does not systematically indicate the site is ombrotrophic. Important differences can occur between the botanical and geochemical trophic status of a peatland because proxies are forming/accumulating in different ways; this was recognized several decades ago (e.g. Shotyk et al., 1998, 2002). Therefore, in any atmospheric reconstruction study using a peat bog, the geochemical trophic status of a site must always be assessed beforehand using an array of geochemical indicators (e.g. major and trace elements abundances and ratios, mineralogy, pH etc.) and a topographic assessment to exclude the possibility of local mineral inputs prior to embarking on a reconstruction of atmospheric sources of dust (e.g. De Vleeschouwer et al., 2007, Fialkewicz-Koziel, 2018, 2020, 2022; Li et al., 2020; Longman et al., 2017; Pratte et al., 2016, 2017a,b, 2020). All other types of peatland records (i.e. not ombrotrophic) should be considered as 'classical' sediment archives. As such, any atmospheric input will be mixed or possibly overprinted by lateral mineral inputs from the catchment, elemental diffusion from the bedrock, etc. 'Classical' sediment archives make it extremely challenging (virtually impossible) to separate an atmospheric signal from local inputs. Records from minerotrophic peatlands should be considered with caution and supported by additional geochemical proxies such as loss-on-ignition (i.e. ash weight-%), lithogenic element (Al, Ti, Sc, REE) concentrations, mineral identification and/or Principal Component Analysis (PCA), to support an atmospheric proxy reconstruction (e.g. De Vleeschouwer et al., 2014; Vanneste et al., 2015; Fagel et al., 2014; Fialkewicz-Koziel et al., 2022; Sharifi et al., 2018). However, Marcisz et al. (2023) have ignored these key points, and based their study on a kettle hole peatland, which could be geochemically minerotrophic (e.g. Lindsay et al., 2015) as well as having top-down peat accumulation from vegetation rafts (Gaudig et al., 2006). Their site is furthermore disturbed by several fires and floods (that can provide lateral mineral inputs disturbing a possible atmospheric signal) as mentioned by the authors. Marcisz et al. (2023) then make a rather speculative conclusion from a single atmospheric dust proxy without providing any geochemical evidence of ombrotrophy. The only proxy presented by Marcisz et al. (2023) that would help the reader assess the geochemical trophic status of the peatland, although not sufficient because not geochemical, is the ash content, but the authors incorrectly report the data in grams instead of in weight-%. This makes it impossible to evaluate the organic content (and therefore the mineral content) of the core. Loss of organic content reported in grams is strongly affected by variations in the peat accumulation rate. Ignoring this basic concept and reporting an ash content in grams removes any interpretative value from the data. One can however estimate the LOI percentage in Marcisz et al. (2023) peat core from the ash weight and ash free density reported in figure 3 (and knowing that 2cm³ were taken for these measurements). This would lead some levels to have an ash/mineral content of 75% which question if those levels are peat (see SI for detailed calculation).

The second paradigm, first expressed by Mattson and Koutler-Anderson (1954), is that peat bogs are located in humid climate zones, where a cover of native vegetation stabilizes the soil and, therefore, minimizing wind erosion and surface runoff. For example, Aubert et al. (2006) have shown that despite the pronounced geological differences between two peatlands located in the Black Forest (granite bedrock) and the Jura (limestone bedrock), similar REE signatures and deposition rates were found in both peatlands. Given that both sites confirm an Upper Continental Crust signature, the data suggests both sites are influenced by regional and not local, soil-derived lithogenic aerosols. In the case of the Etang de la Gruère study (Le Roux et al., 2012), the term local corresponds to the catchment area (10 km), regional to 100 km and global to thousands of km (as defined by Lawrence and Neff, 2009). The larger the peatland and the more it is in a protected area, the less risk there is for recent inputs of local dust. For example in his seminal studies, Gorham foresaw this, based on a transect of a bog to a road and a transect of collected *Sphagnum* mosses in bogs in Minnesota, where high concentrations of lithogenic elements were associated with the presence or absence of cultivated land (Gorham and Tilton, 1978; Santelmann and Gorham, 1988). It should also be noted that local sources have sometimes

been identified in dust records from peat bogs (e.g. De Vleeschouwer et al., 2009; Vanneste et al. 2015) but those local sources had been identified beforehand by studying the geographical and geomorphological settings, sometimes supported by geochemical analyses of the local source itself to make sure to decipher it from potential distal dust sources (e.g. Vanneste et al., 2015, 2016).

By measuring Nd isotopes alone (i.e. with no other geochemical proxies as mentioned above) in a peatland without an in-depth geochemical study of the trophic status (i.e. possibly geochemically minerotrophic), Marcisz et al. (2023) ignore two widely accepted paradigms and do not use the standard practice supported by a large body of literature and approved by the community of peat geochemists. Without a proper geochemical characterization of their record (e.g. lithogenic elements, insoluble ash content, mineralogy, grain size, PCA to determine the driver of each element's behavior) as well as the geomorphological and geographical setting of their site (i.e. lateral/local mineral inputs), it is impossible to assess the suitability of their record for atmospheric reconstruction. In addition, any mixing process of dust sources could be identified if the Nd isotopic data were supported by elemental composition/ratios and by other isotopic systems (for example Pb isotopes). But Marcisz et al. (2023) have employed Nd as a stand-alone proxy, therefore, any interpretation not supported by any in-depth geochemical characterization of the records is speculative.

Given that, we are therefore rather surprised by the unsupported criticisms of Marcisz et al. (2023) towards our previous studies in paragraphs 4.2 and 4.3, for which we provide a point-by-point rebuttal in the supplementary information. We would however like to highlight two facts that are valid for our studies:

- Long-term peat records do indeed provide local signatures by having access to the deeper sediments (whether it is minerotrophic peat or lake sediments), thus providing a geochemical and isotopic signature of the catchment. This is the case for example for Le Roux et al., (2012) and Stewart et al (2023) where the basal sediment has a clear distinct signature from upper peat layers.
- Our studies always provide complete peat records, i.e. it includes the upper part of the profile characterized by living or poorly decomposed moss layers

In addition, we would like to point out that Nd isotopes have been successfully used as cryptotephra identifiers in various archives and time scales (e.g. D'Antonio et al., 2016; Hart et al., 1993). Moreover, the distinction of young volcanism from old shields is somewhat basic for the use of Nd isotopes as provenance tracers. The Nd isotopic signature of a rock is indeed primarily controlled by its age, with fresh mantle-derived rocks having a much more positive ϵ_{Nd} than older cratonic rock (Bizimis & Scher, 2016). Therefore, the isotopic compositions of Nd (as well as Sr and Pb) radiogenic isotopes are significantly different in mantle- vs. in crust-derived rocks, permitting the distinction between the two extreme origins and of lithologies in between: young volcanic areas and old continental shields (Grousset and Biscaye, 2005). Marcisz's et al. (2023) seem however to somehow ignore these critical basics of radiogenic isotopic geochemistry. Moreover, Nd isotopes have also been used elsewhere to distinguish two distal dust sources with different origin (volcanic Iceland dust vs Saharan dust) in peat deposits (Stewart et al., 2023). In the case of the Vasset-Killian tephra, it has moreover been previously identified in Swiss lakes and French peatlands (e.g. Hajdas et al., 1993, Hajdas and Michczynski, 2010, Juvigné, 1991). The unambiguous volcanic ϵ_{Nd} and REE signature reported by Le Roux et al. (2012) is supported by an array of evidence: literature data showing that eruptive rocks from the "Chaîne des Puys" (where the Vasset-Killian complex belongs) display positive Nd signatures (e.g. Hamelin et al., 2009), Nd isotopes, peak in ash content and density, and ultimately radiocarbon dating.

In conclusion, Nd continues to be a very useful and sensitive geochemical tracer in peat bogs, but it is dependent on choosing a suitable site where one can have a high degree of confidence that any mineral material separated from the peat should be from dust sources transported atmospherically. This should be assessed by a range of geochemical and mineralogical proxies to characterize the studied site to avoid any erroneous interpretations. Finally, reference material from potential local and distal dust sources must be isotopically characterized and their pathways to the site either excluded (as for local sources) or atmospheric routes understood (as for distal sources).

References

- Aarons S.M., Aciego S.M., Gabrielli P., Delmonte B., Koornneef J.M., Wegner A. and Blakowski M.A. (2016). The impact of glacier retreat from the Ross Sea on local climate: Characterization of mineral dust in the Taylor Dome ice core, East Antarctica. *Earth and Planetary Science Letters* 444: 34-44.
- Allan M., Le Roux G., Piotrowska N., Beghin J., Javaux E., Court-Picon M., Mattielli N., Verheyden S., and Fagel N. (2013). Mid- and late Holocene dust deposition in western Europe: the Misten peat bog (Hautes Fagnes – Belgium), *Clim. Past*, 9, 2285–2298, <https://doi.org/10.5194/cp-9-2285-2013>
- Aubert D. Le Roux G., Krachler M., Cheburkin A., Kober B. Shotyk W. and Stille P. (2006). Origin and fluxes of atmospheric REE entering an ombrotrophic peat bog in Black Forest (SW Germany): Evidence from snow, lichens and mosses. *Geochimica et Cosmochimica Acta* 70: 2815-2826.
- Bacardit M. and Camarero L. (2009). Fluxes of Al, Fe, Ti, Mn, Pb, Cd, Zn, Ni, Cu, and As in monthly bulk deposition over the Pyrenees (SW Europe): The influence of meteorology on the atmospheric component of trace element cycles and its implications for high mountain lakes. *Journal of Geophysical Research: Biogeosciences* 114. <https://doi.org/10.1029/2008JG000732>
- Beaudon E., Sheets J.M., Martin E., Sierra-Hernández M.R., Mosley-Thompson E. and Thompson L.G. (2022). Aeolian Dust Preserved in the Guliya Ice Cap (Northwestern Tibet): A Promising Paleo-Environmental Messenger. *Geosciences* 12: 366. <https://doi.org/10.3390/geosciences12100366>
- Bazhenova, E., Fagel, N. and Stein, R. (2017). North American origin of “pink–white” layers at the Mendeleev Ridge (Arctic Ocean): New insights from lead and neodymium isotope composition of detrital sediment component *Marine Geology* 386, 44–55.
- Bindler, R. (2011). Contaminated lead environments of man: reviewing the lead isotopic evidence in sediments, peat, and soils for the temporal and spatial patterns of atmospheric lead pollution in Sweden. *Environ Geochem Health* 33, 311–329. <https://doi.org/10.1007/s10653-011-9381-7>
- Bizimis M. and Scher H.D. (2016). Neodymium Isotopes. In: White, W. (eds) *Encyclopedia of Geochemistry. Encyclopedia of Earth Sciences Series.* Springer, Cham. https://doi.org/10.1007/978-3-319-39193-9_124-1
- D'Antonio M., Mariconte R., Arienzo I., Mazzeo F.C., Carandente A., Perugini D., Petrelli M., Corselli C., Orsi G., Principato M.S. and Civetta L. (2016). Combined Sr-Nd isotopic and geochemical fingerprinting as a tool for identifying tephra layers: Application to deep-sea cores from Eastern Mediterranean Sea. *Chemical Geology* 443: 121-136.
- De Angelis M. and Gaudichet A. (1991). Saharan dust deposition over Mont Blanc (French Alps) during the last 30 years. *Tellus B: Chemical and Physical Meteorology* 43, 61–75. <https://doi.org/10.3402/tellusb.v43i1.15246>
- De Vleeschouwer F., Vanneste H., Mauquoy D., Piotrowska N., Torrejon F., Roland T., Stein A. and Le Roux G. (2014). Emissions from pre-Hispanic metallurgy in the South American atmosphere. *PLOS ONE* <http://dx.doi.org/10.1371/journal.pone.0111315>
- De Vleeschouwer F., Piotrowska N., Sikorvsky J., Pawlyta J., Cheburkin A.K., Le Roux G., Lamentowicz, M., Fagel, N. and Mauquoy D. (2009). Multiproxy evidence of ‘Little Ice Age’ palaeoenvironmental changes in a peat bog from northern Poland. *The Holocene*, 19-4: 625-637.
- De Vleeschouwer F., Gérard L., Goormaghtigh C., Mattielli N., Le Roux G. and Fagel N. (2007). Atmospheric lead and heavy metal pollution records from a Belgian peat bog spanning the last

- two millennia: Human impact on a regional to global scale. *The Science of the Total Environment* 377 2-3: 297-310.
- Dia, A., Dupré B. and Allegre, C.J. (1992). Nd isotopes in Indian Ocean sediments used as a tracer of supply to the ocean and circulation paths. *Mar. Geol.* 103, 349-359.
- Dunlap C.E., Steinnes E. and Flegal A.R. (1999). A synthesis of lead isotopes in two millennia of European air. *Earth Planet. Sci. Lett.* 167: 81–88.
- Fagel, N., C. Hillaire-Marcel, M. Humblet, R. Brasseur, D. Weis, and R. Stevenson (2004), Nd and Pb isotope signatures of the clay-size fraction of Labrador Sea sediments during the Holocene: Implications for the inception of the modern deep circulation pattern, *Paleoceanography*, 19, PA3002, doi:10.1029/2003PA000993
- Fagel N., Allan M., Le Roux G., Mattielli N., Piotrowska N. and Sikorski J. (2014). Deciphering human-climate interactions in an ombrotrophic peat record: REE, Nd and Pb isotope signatures of dust supplies over the last 2500 years (Misten bog, Belgium). *Geochim. Cosmochim. Acta* 135, 288–306.
- Farmer J.G., Eades, L.J., Atkins, H., Chamberlain, D.F. (2002). Historical trends in the lead isotopic composition of archival Sphagnum mosses from Scotland (1838–2000). *Environmental Science and Technology* 36: 152–157.
- Fiałkiewicz-Kozieł B., Bao K., Smieja-Król B. (2022). Geographical drivers of geochemical and mineralogical evolution of Motianling peatland (Northeast China) exposed to different sources of rare earth elements and Pb, Nd, and Sr isotopes. *Sci. Total Environ.* 807 : 150481.
- Fiałkiewicz – Kozieł B., Łokas E., Gałka M., Kołaczek P., De Vleeschouwer F., Le Roux G., Smieja-Król B.(2020). Influence of transboundary transport of trace elements on mountain peat geochemistry (Sudetes, Central Europe). *Quaternary Science Reviews* 230, 106162.
- Fiałkiewicz-Kozieł B., De Vleeschouwer F., Mattielli N., Fagel N, Palowski B., Pazdur A. Smieja-Król, B., (2018). Record of Anthropocene pollution sources of lead in disturbed peatlands from Southern Poland. *Atmospheric Environment* 179:61-68.
- Fiałkiewicz-Kozieł B., Smieja-Król B., Frontasyeva M., Słowiński M., Marcisz K., Lapshina E., Gilbert D., Buttler A., Jassey V.E.J., Kaliszan K., Laggoun-Defarge F., Kołaczek P., Lamentowicz M. (2016). Anthropogenic and natural sources of dust in peatland during the Anthropocene. *Nature Scientific Reports* 6: 38731.
- Frank, M. (2002), Radiogenic isotopes: Tracers of past ocean circulation and erosional input, *Rev. Geophys.*, 40(1), 1001, doi:10.1029/2000RG000094.
- Gaudig G., Couwenberg J. and Joosten H. (2006). Peat accumulation in kettle holes: bottom up or top down? *Mires and Peat* 1: art. 6.
- Goldstein, S.L., O’Nions, R.K., Hamilton, P.J., 1984. A Sm/Nd isotopic study of atmospheric dust particulates from major river systems. *Earth Planet. Sci. Lett.* 70, 221-236.
- Gorham E. and Tilton D.L., (1978). The mineral content of Sphagnum fuscum as affected by human settlement. *Can. J. Bot.* 56, 2755–2759. <https://doi.org/10.1139/b78-328>
- Grousset F.E., Biscaye P.E., Revel M., Petit J.-R., Pye K., Joussaume S. and Jouzel J. (1992). Antarctic (Dome C) ice-core dust at 18 k.y. B.P.: Isotopic constraints on origins. *Earth and Planetary Science Letters* 111: 175-182,
- Grousset, F.E., Biscaye, P.E., Zindler, A., Prospero, J. and Chester, R. (1988). Neodymium isotopes as tracers in marine sediments and aerosols. *Earth Planet. Sci. Lett.* 87, 367-378.

- Grousset F.E. and Biscaye P.E. (2005). Tracing dust sources and transport patterns using Sr, Nd and Pb isotopes: Chemical Geology 222: 149–167, doi:10.1016/j.chemgeo.2005.05.006.
- Hajdas, I., Ivy, S.D., Beer, J. *et al.* AMS radiocarbon dating and varve chronology of Lake Soppensee: 6000 to 12000 ¹⁴C years BP. *Climate Dynamics* 9, 107–116 (1993). <https://doi.org/10.1007/BF00209748>
- Hajdas I. and Michczynski, A. (2010) Age-depth model of Lake Soppensee (Switzerland) based on ¹⁴C chronology compared with varve chronology. *Radiocarbon* 52: 1027–1040.
- Hamelin C., Seitz H.-M., Barrat J.-A., Dosso L., Maury R.C. and Chaussidon M.C. (2009). A low $\delta^{7}\text{Li}$ lower crustal component: Evidence from an alkalic intraplate volcanic series (Chaîne des Puys, French Massif Central). *Chemical Geology* 266: 205–217.
- Hart W.K., Walter R.C., Wolde Gabriel G. (1992). Tephra sources and correlations in Ethiopia: Application of elemental and neodymium isotope data. *Quaternary International* 13–14: 77–86.
- Juvigné E. (1991). Distribution de vastes retombées volcaniques originaires de l'Eifel et du Massif Central aux temps post-glaciaires dans le NE de la France et les régions voisines. Paris, Académie des Sciences Comptes Rendus, ser. II, v. 312, p. 415–420.
- Kempton H., Krachler M., Shotyk W. and Zaccone C. (2017). Major and trace elements in Sphagnum moss from four southern German bogs, and comparison with available moss monitoring data. *Ecological Indicators* 78: 19–25.
- Kempton H., Krachler M. and Shotyk W. (2010). Atmospheric Pb and Ti Accumulation Rates from *Sphagnum* Moss: Dependence upon Plant Productivity. *Environmental Science and Technology* 44: 5509–5515.
- Lacan F. and Jeandel C. (2004). Denmark Strait Water circulation traced by heterogeneity in neodymium isotopic compositions, *Deep Sea Res.*, 51, 71–82.
- Lawrence, C.R. and Neff J.C. (2009). The contemporary physical and chemical flux of aeolian dust: A synthesis of direct measurements of dust deposition. *Chemical Geology, Combined Ecological and Geologic Perspectives in Ecosystem Studies* 267, 46–63. <https://doi.org/10.1016/j.chemgeo.2009.02.005>
- Le Roux G., Fagel N., De Vleeschouwer F., Krachler M., Debaille V., Stille P., Mattiell, N., Van Der Knaap W.O., Van Leeuwen J.F.N., Shotyk W. (2012). Volcano- and climate-driven changes in atmospheric dust sources and fluxes since the Late Glacial in Central Europe. *Geology* 40, 335–338. <https://doi.org/10.1130/G32586.1>
- Le Roux G., Aubert D., Stille P., Krachler M., Kober B., Cheburkin A., Bonani G. and Shotyk W. (2005). Recent atmospheric Pb deposition at a rural site in southern Germany assessed using a peat core and snowpack, and comparison with other archives. *Atmos. Environ.*: 39, 6790–6801.
- Li C., Sonke J., Le Roux R., van der Putten N., Piotrowska N., Jeandel C., Mattielli N., Benoit M., Wiggs G.F.S. and De Vleeschouwer F. (2020). Holocene dynamics of the southern westerly winds over the Indian Ocean inferred from a peat dust deposition record. *Quaternary Science Reviews* 231: 106169.
- Lindsay R. (2016). Peatland (Mire Types): Based on Origin and Behavior of Water, Peat Genesis, Landscape Position, and Climate. In C.M. Finlayson *et al.* (eds.), *The Wetland Book*, Springer Science and Media Business, Dordrecht. DOI 10.1007/978-94-007-6173-5_279-1
- Longman J., Struve T. and Pahnke K. (2022). Spatial and temporal trends in mineral dust provenance in the South Pacific—Evidence from mixing models. *Paleoceanography Paleoclimatology*, (e2021PA004356).

- Longman J., Veres D., Ersek V., Salzmann U., Hubay K., Bormann M., Wennrich V. and Schäbitz F. (2017). Periodic input of dust over the Eastern Carpathians during the Holocene linked with Saharan desertification and human impact. *Climate of the Past*, 13(7), pp.897-917.
- Marcisz K., Belka Z., Dopieralska J., Jakubowicz M., Karpińska-Kołaczek M., Kołaczek P., Mauquoy D., Słowiński M., Zieliński M. and Lamentowicz M. (2023). Neodymium isotopes in peat reveal past local environmental disturbances. *Science of the Total Environment* 871: 161859.
- Markert, B. Inorganic chemical investigations in the Forest Biosphere Reserve near Kalinin, USSR. *Vegetatio* 95, 127–135 (1991). <https://doi.org/10.1007/BF00045211>
- Mattson S. and Koutler-Anderson E. (1954) Geochemistry of a raised bog. *Ann. Royal Agr. Coll. Sweden*, 21, 321-366.
- Novak M., Erel Y., Zemanova L., Bottrell S.H. and Adamova M. (2008). A comparison of lead pollution record in Sphagnum peat with known historical Pb emission rates in the British Isles and the Czech Republic. *Atmospheric Environment* 42: 8997-9006.
- O’Nions R.K., Hamilton P.J. and Hooker A. (1983). A Nd isotope investigation of sediments related to crustal development in the British Isles. *Earth Planet. Sci. Lett.* 63, 229-240.
- Pech, P., Wojtuń, B., Samecka-Cymerman, A. *et al.* Metals in Plant Functional Types of Ombrotrophic Peatlands in the Sudetes (SW Poland). *Arch Environ Contam Toxicol* 82, 506–519 (2022). <https://doi.org/10.1007/s00244-022-00928-5>
- Pratte S., Bao K., Sapkota A., Zhang W., Shen J., Le Roux G. and De Vleeschouwer F. (2020) 14 kyr of atmospheric mineral dust deposition in north-eastern China: A record of palaeoclimatic and palaeoenvironmental changes in the Chinese dust source regions. *The Holocene* 30 (guest Eds. F. De Vleeschouwer, J.-B. Stuut, F. Lambert): 492-506.
- Pratte S., De Vleeschouwer F. and Garneau M. (2017a). Geochemical characterization (REE, Nd and Pb isotopes) of atmospheric mineral dust deposited in two maritime peat bogs from the St. Lawrence North Shore (eastern Canada). *Journal of Quaternary Sciences* 15: 617-627.
- Pratte S., Garneau M. and De Vleeschouwer F. (2017b). Increased atmospheric dust deposition during the Neoglacial in a boreal peat bog from North-Eastern Canada. *Palaeogeography, Palaeoclimatology, Palaeoecology* 469: 34-46.
- Pratte S., Garneau M. and De Vleeschouwer F. (2016). Late-Holocene atmospheric dust deposition in eastern Canada (St. Lawrence North Shore). *The Holocene* 27 (1): 12-25.
- Rausch N., Nieminen T., Ukonmaanaho L., Le Roux G., Krachler, M., Cheburkin A.K., Bonani, G. and Shotyk W. (2005). Comparison of atmospheric deposition of copper, nickel, cobalt, zinc and cadmium recorded by Finnish peat cores with monitoring data and emission records. *Environ. Sci. Technol.* 39 5989–5998.
- Saitoh Y., Tamura T., Kodama Y. and Nakano T. (2011). Strontium and neodymium isotopic signatures indicate the provenance and depositional process of loams intercalated in coastal dune sand, western Japan. *Sedimentary Geology* 236: 272-278.
- Santelmann M.V. and Gorham, E. (1988). The Influence of Airborne Road Dust on the Chemistry of Sphagnum Mosses. *Journal of Ecology* 76, 1219–1231. <https://doi.org/10.2307/2260644>
- Simonsen, M.F., Baccolo, G., Blunier, T. *et al.* East Greenland ice core dust record reveals timing of Greenland ice sheet advance and retreat. *Nat Commun* 10, 4494 (2019). <https://doi.org/10.1038/s41467-019-12546-2>

- Sharifi A., Murphy L.N., Pourmand A., Clement A.C., Canuel E.A., Beni A.N., Lahijani H.A.K., Delanghe D. and Ahmady-Birgani H. (2018). Early-Holocene greening of the Afro-Asian dust belt changed sources of mineral dust in West Asia. *Earth and Planetary Science Letters* 481: 30-40.
- Shotyk W., Kempter H., Krachler M. and Zaccone C. (2015). Stable (^{206}Pb , ^{207}Pb , ^{208}Pb) and radioactive (^{210}Pb) lead isotopes in 1 year of growth of Sphagnum moss from four ombrotrophic bogs in southern Germany: Geochemical significance and environmental implications. *Geochim. Cosmochim. Acta* 163: 101–125.
- Shotyk W., Krachler M., Martinez-Cortizas A., Cheburkin A., Emons H. (2002). A peat bog record of natural, pre-anthropogenic enrichments of trace elements in atmospheric aerosols since 12 370 ^{14}C yr BP, and their variation with Holocene climate change. *Earth and Planetary Science Letters* 199: 21-37,
- Shotyk W., Weiss D., Appleby P.G., Cheburkin A.K., Frei R., Gloor M., Kramers J.D., Reese S. and Van Der Knaap W.O. (1998). History of atmospheric lead deposition since 12,370 ^{14}C yr BP recorded in a peat bog profile, Jura Mountains, Switzerland. *Science*: 281 1635–1640.
- Steinnes E. (1995). A critical evaluation of the use of naturally growing moss to monitor the deposition of atmospheric metals. *Science of The Total Environment* 160–161: 243-249.
- Stewart, H., Bradwell, T., Bullard, J., McCulloch, R. D., & Millar, I. (2023). Increased North Atlantic dust deposition linked to Holocene Icelandic glacier fluctuations. *The Holocene*, 33(2), 231–237. <https://doi.org/10.1177/09596836221131697>
- Struve T., Longman J., Zander M., Lamy F., Winckler G. and Pahnke K. (2022). Systematic changes in circumpolar dust transport to the Subantarctic Pacific Ocean over the last two glacial cycles. *Proceedings of the National Academy of Sciences*, 119(47), p.e2206085119.
- Vanneste H., De Vleeschouwer F., Bertrand S., Martinez-Cortizas A., Vanderstraeten A., Mattielli N., Coronato A., Piotrowska N., Jeandel C. and Le Roux G. (2016). Elevated dust deposition in Tierra del Fuego (Chile) resulting from Neoglacial Darwin Cordillera glacier fluctuations. *Journal of Quaternary Science* 31 (7): 713-722.
- Vanneste H., De Vleeschouwer F., Martinez-Cortizas A., von Scheffer C., Piotrowska N., Coronato A. and Le Roux G. (2015). Late-glacial elevated dust deposition linked to westerly wind shifts in southern South America. *Nature Scientific Reports* 5, 11670, doi:10.1038/srep11670
- Weiss D., Shotyk W., Boyle A.E., Kramers D.J., Appleby P.G. and Cheburkin, A.K. (2002). Comparative study of the temporal evolution of atmospheric lead deposition in Scotland and eastern Canada using blanket peat bog. *Science of the Total Environment* 292: 7–18.
- Weiss D., Shotyk W., Kramers J.D. and Gloor M. (1999). Sphagnum mosses as archives of recent and past atmospheric lead deposition in Switzerland. *Atmos. Environ.* 33 : 3751–3763.
- Zupančič N. and Bozau E. (2022). Effect of the coronavirus pandemic lockdown to elemental composition of peat mosses. *Environ Sci Pollut Res* 29, 25473–25485. <https://doi.org/10.1007/s11356-021-17564-6>

Supplementary information

S.I.1 : Chat GPT 05/02/2023

How to investigate dust in peat paleoenvironmental records?

There are several methods to determine if dust in peat paleoenvironmental records is of local origin or from a remote source:

Elemental and isotopic analysis: Comparing the elemental and isotopic composition of the dust to local soils and sediment, as well as known remote sources, can help identify its origin. For example, if the dust contains elements that are not typically found in the local area, but are common in known remote sources, it is likely that the dust was transported from a remote location.

Statistical analysis: Using statistical methods such as principal component analysis (PCA) or clustering algorithms, can help identify patterns and trends in the elemental and isotopic composition of the dust, which can help distinguish between local and remote sources.

Physical analysis: Examining the size and shape of the dust particles can help determine if they were transported over long distances. Dust particles that are larger and more angular are typically indicative of local sources, while smaller and more rounded particles are often associated with remote sources.

Comparison with other paleoclimate proxies: Comparing the results of the dust analysis with other paleoclimate proxies, such as ice cores, tree rings, and sediment records, can help confirm the origin of the dust. For example, if the dust is found in association with other proxies that are known to be from a remote source, it is likely that the dust is also from that source.

Modeling: Using atmospheric circulation models can help simulate the transport of dust from remote sources and predict the chemical and isotopic composition of the dust that would be expected to arrive at a particular location. This information can be compared to the actual elemental and isotopic composition of the dust in the peat record to help determine its origin.

Can we use Nd isotopes in peat to evaluate dust sources?

Yes, Nd (neodymium) isotopes can be used to evaluate dust sources in peat paleoenvironmental records. Nd isotopes are a valuable tool for tracing the sources of sediment in the environment because different geological regions have unique Nd isotopic signatures. By analyzing the Nd isotopes in the dust, it is possible to determine the source of the dust and whether it is from a local or remote source.

In peat records, Nd isotopes can be measured using techniques such as inductively coupled plasma mass spectrometry (ICP-MS). The Nd isotopic composition of the dust is then compared to the Nd isotopic composition of local soils and sediment, as well as known remote sources. A match with a known remote source would suggest that the dust was transported from that location, while a match with local soils and sediment would suggest a local origin.

It is important to note that while Nd isotopes can be a useful tool for evaluating dust sources, they should be used in conjunction with other methods, such as elemental analysis and physical analysis, to provide a more complete picture of the origin and transport of the dust.

S.I.2: attempt at calculating an ash content (in weight %)

In figure 3 of their paper, Marcisz et al. (2023) report various parameters that are not the common practice:

- “ash free density” instead of bulk density. Our understanding is that this density was derived from the sample after loss on ignition was performed. In other words that the mineral content of the sample was “removed” to calculate a density that only represent the organic matter content.
- Ash content is reported in grammes instead of weight %, which renders impossible to estimate the real proportion of mineral in the peat.

Marcisz et al. (2023) however report that 2cm³ were taken to perform LOI measurements. Reading from figure 3, there are levels containing 0.9g of ash with an ash free density of 0.15g cm⁻³. That means a 2cm³ sample is composed of 0.3g of organic matter and 0.9g of ash (i.e. ash/mineral content is 75%), which is not peat. It is possible we misunderstand the proxies reported by Marcisz et al. (2023), but this simple calculation is not made to prove that the studied core is not only composed of peat, rather than to show that once standard practices are not used, the reader is forced to navigate to try calculate a very basic proxy, while it would have just been very simple to report bulk density and ash content in weight-% to allow the reader to have a proxy of the geochemical trophic status of the peatland.

S.I.3: Point by point answer to paragraphs 4.2 and paragraphs 4.3 of Marcisz et al. (2023):

We agree that there seems to be a misquotation on the Allan et al. (2013) paper but as the main author is no longer in academic research, we do not wish to comment on that paper without his consent. We will therefore focus on the Etang de la Gruère record (Le Roux et al., 2012).

« However, these studies did not include reference samples collected in close proximity to the peatlands, and they refer solely to ϵ Nd data from the literature, measured at locations distant from the investigated sites (Allan et al., 2013; Fagel et al., 2014; Le Roux et al., 2012). »

As stated in the main text, In the 3 cited papers, we were able to collect the deepest sediments representing the geochemical signature thus the « local » signature, or that is to say, the average signature of the close proximity of the peatland.

« Moreover, the authors did not define what they mean by a “local source”, and how far from the site this local source may be located. For example, Le Roux et al. (2012) attributed the observed variability in the ϵ Nd values to shifts between a local source and Saharan dust, but, in the absence of additional data, provided no unequivocal evidence for the contribution of the latter. »

We deplore the cherry-picking of this paragraph by Marcisz et al. (2023), where the authors ignore decades of publications on Etang de la Gruère where lithogenic elements and Pb isotopes also show dust provenance from Sahara (see for example: Shotyk et al., 1998; 2002). In addition, it is well known that Sahara dust is a major dust source to European mountains (Bacardit and Camarero, 2009; De Angelis and Gaudichet, 1991) including the Alps, where Etang de la Gruyère is located.

« A comparison of these records shows that it is practically impossible to identify Saharan dust supply to a peatland based only on Nd isotopes alone; other, supporting evidence is needed to unequivocally detect Saharan derived materials. »

We fully agree that Nd alone is not enough (we note that it is however precisely what Marcisz et al. (2023) do: they only measure Nd isotopes, and therefore their argument is circular). That is why all our studies are based on a range of proxies before measuring Nd isotopes, such as: inorganic elemental geochemistry including metals and REE, grain size/mineralogy, acid insoluble ash content, Pb isotopes, etc.

« The apparent coincidence of a change in the ϵNd values and a reconstructed age of a specific peat layer is not direct evidence, as age-depth models are often not precise enough to allow such direct comparisons »

Once again Marcisz et al. (2023) were cherry-picking, and did not take the time to our paper in detail, especially the SI which details the work on age-depth modelling, including the comparison of the two Etang de la Gruère cores. In particular, the age-depth model developed for the master core shows sufficient resolution to detect centennial-scale events.

« Le Roux et al. (2012) attributed their Nd isotope record (ϵNd values of -9.7 and -9.6) to a supply of the Icelandic Vedde Ash, but this interpretation was supported only by the age of the peat layer in which the change in the ϵNd values occurred. »

This is again a misreading of our paper by Marcisz et al. (2023). We wrote: « The second peak (ca. 12 kyr B.P.) is a positive shift that **may** correspond to a volcanic signature (Vedde Ash) ». Moreover, not only do we have the Nd isotopic evidence, but this layer also has a density, ash content, and dust flux peak as well as distinct REE signatures suggesting a volcanic origin. This layer has already been attributed to a volcanic eruption by Shotyk et al (1998, 2002).

« Similarly, the highest measured ϵNd value of -1.0 was assigned to two eruptions from the Massif Central, because of the similar age of this peat layer, rather than the similarity of the ϵNd signatures (Le Roux et al., 2012). »

Here Marcisz et al. (2013) questions a tephra layer which is well established in the literature and obviously did not take the time to indeed check the literature we provide in our paper. This tephra was for example identified in peat bogs from Massif Central (Juvigné, 1991) and in sediments from Lake Soppensee (e.g. Hajdas et al., 1993, Hajdas and Michczynski, 2010). The Nd isotopic signature is without doubt volcanic (e.g. Grousset and Biscaye, 2005). Moreover, the Killian-Vasset Complex is part of the “Chaine des Puys” which eruptive rocks display positive ϵNd signatures up to $+3.8$ (eg. Hamelin et al., 2009) and there is no volcanic terrain in the peatland area since the watershed’s bedrock is mainly composed of carbonates (Shotyk et al., 1998).

References

- Allan M., Le Roux G., Piotrowska N., Beghin J., Javaux E., Court-Picon M., Mattielli N., Verheyden S., and Fagel N. (2013). Mid- and late Holocene dust deposition in western Europe: the Misten peat bog (Hautes Fagnes – Belgium), *Clim. Past*, 9, 2285–2298, <https://doi.org/10.5194/cp-9-2285-2013>
- Bacardit M. and Camarero L. (2009). Fluxes of Al, Fe, Ti, Mn, Pb, Cd, Zn, Ni, Cu, and As in monthly bulk deposition over the Pyrenees (SW Europe): The influence of meteorology on the atmospheric component of trace element cycles and its implications for high mountain lakes. *Journal of Geophysical Research: Biogeosciences* 114. <https://doi.org/10.1029/2008JG000732>
- De Angelis M. and Gaudichet A. (1991). Saharan dust deposition over Mont Blanc (French Alps) during the last 30 years. *Tellus B: Chemical and Physical Meteorology* 43, 61–75. <https://doi.org/10.3402/tellusb.v43i1.15246>
- Fagel N., Allan M., Le Roux G., Mattielli N., Piotrowska N. and Sikorski J. (2014). Deciphering human-climate interactions in an ombrotrophic peat record: REE, Nd and Pb isotope signatures of dust supplies over the last 2500 years (Misten bog, Belgium). *Geochim. Cosmochim. Acta* 135, 288–306.
- Grousset F.E. and Biscaye P.E. (2005). Tracing dust sources and transport patterns using Sr, Nd and Pb isotopes: *Chemical Geology* 222: 149–167, doi:10.1016/j.chemgeo.2005.05.006.
- Hajdas I., Ivy S.D., Beer J. *et al.* (1993). AMS radiocarbon dating and varve chronology of Lake Soppensee: 6000 to 12000 ¹⁴C years BP. *Climate Dynamics* 9: 107–116. <https://doi.org/10.1007/BF00209748>
- Hajdas I. and Michczynski, A. (2010) Age-depth model of Lake Soppensee (Switzerland) based on ¹⁴C chronology compared with varve chronology. *Radiocarbon* 52: 1027–1040.
- Hamelin C., Seitz H.-M., Barrat J.-A., Dosso L., Maury R.C. and Chaussidon M.C. (2009). A low $\delta^{71}\text{Li}$ lower crustal component: Evidence from an alkalic intraplate volcanic series (Chaîne des Puys, French Massif Central). *Chemical Geology* 266: 205-217,
- Juvigné E. (1991). Distribution de vastes retombées volcaniques originaires de l'Eifel et du Massif Central aux temps post-glaciaires dans le NE de la France et les régions voisines. Paris, Académie des Sciences Comptes Rendus, ser. II, v. 312, p. 415–420.
- Le Roux G., Fagel N., De Vleeschouwer F., Krachler M., Debaille V., Stille P., Mattiell, N., Van Der Knaap W.O., Van Leeuwen J.F.N., Shotyk W. (2012). Volcano- and climate-driven changes in atmospheric dust sources and fluxes since the Late Glacial in Central Europe. *Geology* 40, 335–338. <https://doi.org/10.1130/G32586.1>
- Marcisz K., Belka Z., Dopieralska J., Jakubowicz M., Karpińska-Kołaczek M., Kołaczek P., Mauquoy D., Słowiński M., Zieliński M. and Lamentowicz M. (2023). Neodymium isotopes in peat reveal past local environmental disturbances. *Science of the Total Environment* 871: 161859.
- Shotyk W., I Krachler M., Martinez-Cortizas A., Cheburkin A., Emons H. (2002). A peat bog record of natural, pre-anthropogenic enrichments of trace elements in atmospheric aerosols since 12 370 ¹⁴C yr BP, and their variation with Holocene climate change. *Earth and Planetary Science Letters* 199: 21-37,
- Shotyk W., Weiss D., Appleby P.G., Cheburkin A.K., Frei R., Gloor M., Kramers J.D., Reese S. and Van Der Knaap W.O. (1998). History of atmospheric lead deposition since 12,370 ¹⁴C yr BP recorded in a peat bog profile, Jura Mountains, Switzerland. *Science*: 281 1635–1640.