



HAL
open science

Best-of-both-Worlds Analysis of Online Search

Spyros Angelopoulos, Christoph Dürr, Shendan Jin

► **To cite this version:**

Spyros Angelopoulos, Christoph Dürr, Shendan Jin. Best-of-both-Worlds Analysis of Online Search. *Algorithmica*, 2023, 10.1007/s00453-023-01165-5 . hal-04248729

HAL Id: hal-04248729

<https://hal.science/hal-04248729>

Submitted on 18 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Best-of-both-Worlds Analysis of Online Search

Spyros Angelopoulos^{1*}, Christoph Dürr¹ and Shendan Jin²

^{1*}LIP6, Sorbonne Université, CNRS, 4 place Jussieu, Paris, 75252, State, France.

²Murex Inc., 25A Bd de l'Amiral Bruix, Paris, 75016, France.

*Corresponding author(s). E-mail(s): spyros.angelopoulos@lip6.fr;
Contributing authors: christoph.durr@lip6.fr; sjin@murex.com;

Abstract

In search problems, a mobile searcher seeks to locate a target that hides in some unknown position of the environment. Such problems are typically considered to be of an on-line nature, in that the target's position is unknown to the searcher, and the performance of a search strategy is usually analyzed by means of the standard framework of the competitive ratio, which compares the cost incurred by the searcher to an optimal strategy that knows the location of the target. However, one can argue that even for simple search problems, competitive analysis fails to distinguish between strategies which, intuitively, should have different performance in practice. Motivated by the above observation, in this work we introduce and study measures *supplementary* to competitive analysis in the context of search problems. In particular, we focus on the well-known problem of *linear search*, informally known as the *cow-path problem*, for which there is an infinite number of strategies that achieve an optimal competitive ratio equal to 9. We propose a measure that reflects the rate at which the line is being explored by the searcher, and which can be seen as an extension of the *bijective ratio* over an uncountable set of requests. Using this measure we show that a natural strategy that explores the line aggressively is optimal among all 9-competitive strategies. This provides, in particular, a strict separation from the competitively optimal doubling strategy, which is much more conservative in terms of exploration. We also provide evidence that this aggressiveness is requisite for optimality, by showing that any optimal strategy must mimic the aggressive strategy in its first few explorations.

Keywords: Online computation, search problems, analysis beyond the worst case, linear search, performance measures

1 Introduction

Searching for a hidden target is an important paradigm in computer science and operations research, with numerous applications. A typical search problem involves a search *environment*, a mobile *searcher* (who may, or may not, have knowledge of the environment) and a *hider* (sometimes also called *target*) who hides at some position within the environment that is unknown to the searcher. The objective is to define a *search strategy*, i.e., a traversal of the environment, that optimizes a certain efficiency criterion. A standard approach to the latter is by means of *competitive analysis*, in which we seek to minimize the worst-case cost for locating the target, divided by some concept of “optimal” solution; e.g., the minimum cost to locate the target once its position is known. Even prior to the advent of online computation and competitive analysis, search games had already been studied under such *normalized* measures within Operations Research [12]. Explicit studies of the competitive ratio and the closely related *search ratio* were given in [10] and [32], respectively, and led to the development of *online searching* [14, 27] as a subfield of online computation. See also [1] for an in-depth treatment of search games, including the role of payoff functions that capture the competitive ratio.

In this work we revisit one of the simplest, yet fundamental search problems, namely the *linear search* problem, informally also known as the *cow-path* problem. The setting involves an infinite (i.e., unbounded) line, with a point O designated as its origin, a searcher which is initially placed at the origin, and an immobile target which is at some position on the line that is unknown to the searcher. More specifically, the searcher does not know whether the hider is at the left branch or at the right branch of the line. The searcher’s *strategy* S defines its exploration of the line, whereas the hider’s strategy H is determined by its placement on the line. Given strategies S, H , the *cost* of locating the hider, denoted by $c(S, H)$ is the total distance traversed by the searcher at the first time it passes over H . Let H denote the distance of the hider from the origin. The competitive ratio of S , denoted by $\text{cr}(S)$, is the worst-case normalized cost of S , among all possible hider strategies. Formally,

$$\text{cr}(S) = \sup_H \frac{c(S, H)}{H}. \quad (1)$$

It has long been known [11, 23] that the competitive ratio of linear search is 9, and is achieved by a simple *doubling* strategy: in iteration i , the searcher starts from O , explores branch $i \bmod 2$ at a length equal to 2^i , and then returns to O . However, this strategy is not uniquely optimal; in fact, there is an infinite number of competitively optimal strategies for linear search (see Lemma 2

in Section 3). In particular, consider an “aggressive” strategy, which in each iteration searches a branch to the maximum possible extent, while maintaining a competitive ratio equal to 9. This can be achieved by searching, in iteration i , branch $i \bmod 2$ to a length equal to $(i + 2)2^{i+1}$ (see Corollary 3).

While both **doubling** and **aggressive** are optimal in terms of the competitive ratio, there exist realistic situations in which the latter may be preferable to the former. Consider, for example, a search-and-rescue mission for a missing backpacker who has disappeared in one of two (very long) concurrent, hiking paths. Assuming that we select our search strategy from the space of 9-competitive strategies, it makes sense to choose one that is tuned to discovering new territory, rather than a conservative strategy that tends to often revisit already explored lengths of the area. Another illustrative example is demining. Suppose that a demining team must locate and defuse a mine that is hidden along a very long line. Being competitive in this case should not be the only efficiency criterion, but rather one would like also to demine as large a length as possible, within a certain amount of time.

With the above observation in mind, we first need to quantify what constitutes efficiency in exploration. To this end, given a strategy S and $\ell \in \mathbb{R}^+$, we define $D(S, \ell)$ as the cost incurred by S the first time the searcher has explored an aggregate length equal to ℓ , combined in both branches. Formally, given a time t , and assuming a unit-speed searcher, let $L(S, t)$ and $R(S, t)$ denote the lengths that have been explored by the searcher at time t , to the left and to the right of the origin, respectively. Then $D(S, \ell)$ is the earliest time t for which $L(S, t) + R(S, t) = \ell$. An efficient strategy should be such that $D(S, \ell)$ is small, for all ℓ . Unfortunately, this criterion by itself is insufficient: Consider a strategy that first searches one branch to a length equal to L , where L is very large. Then $D(S, \ell)$ is as small as possible for all $\ell < L$; however, this is hardly a good strategy, since it all but ignores one of the branches (and thus its competitive ratio becomes unbounded as $L \rightarrow \infty$).

To remedy this situation, we will instead use the above definition in a way that will allow us a pairwise comparison of strategies, which also considers all possible explored lengths. More formally, we define the following measure.

Definition 1 Let S_1, S_2 denote two search strategies, we define the *discovery ratio* of S_1 against S_2 , denoted by $\text{dr}(S_1, S_2)$, as

$$\text{dr}(S_1, S_2) = \sup_{\ell \in \mathbb{R}^+} \frac{D(S_1, \ell)}{D(S_2, \ell)}.$$

Moreover, given a class \mathcal{A} of search strategies, the *discovery ratio* of S against the class \mathcal{A} is defined as

$$\text{dr}(S, \mathcal{A}) = \sup_{S' \in \mathcal{A}} \text{dr}(S, S').$$

In the case \mathcal{A} is the set of all possible strategies, we simply call $\text{dr}(S, \mathcal{A})$ the *discovery ratio* of S , and we denote it by $\text{dr}(A)$.

Intuitively, the discovery ratio preserves the worst-case nature of competitive analysis, and at the same time bypasses the need for an “offline optimum”

solution. Note that if a strategy S has competitive ratio c then it also has discovery ratio c ; this follows easily from the fact that for every hider position H , $c(S, H) \geq D(S, H)$. However, the opposite is not necessarily true.

It is worth pointing out that one could have defined the discovery ratio over a discrete, countable space (i.e., the target hides at some integer distance from the origin), which turns out to be identical to the *bijective ratio*. This performance measure was introduced in [8] as an extension of (exact) bijective analysis of online algorithms [7], and which in turn is based on the pairwise comparison of the costs induced by two online algorithms over all request sequences of a certain size. Bijective analysis has been applied in fundamental online problems (with a discrete, finite set of requests) such as paging and list update [9], k -server [8, 17], and online search¹ [18].

In what concerns linear search, in this work we choose to present the analysis over a “continuous” space of requests for two reasons. First, we demonstrate that this is indeed possible, which can be useful for other online problems which are defined over a continuous setting of requests (e.g., k -server problems defined over a metric space rather than over a finite graph). Second, the discretization introduces certain unnecessary and undesirable technical issues (e.g., in the choice of the “right” value t for strategy R_t , see Lemma 4). While the analysis is still tractable for our problem, for more complex search domains such as star search, the discrete analysis may be too complicated to yield results. We further discuss the connections between the discovery and the bijective ratios in Section 4.

The above observation implies that the discovery ratio inherits the appealing properties of bijective analysis, which further motivates its choice. In particular, note that bijective analysis has helped to identify theoretically efficient algorithms which also tend to perform well in practice (such as Least-Recently-Used for paging [9], and greedy-like k -server policies for certain types of metrics [8]). Furthermore, if an algorithm has bijective ratio c , then its average cost, assuming a uniform distribution over all request sequences of the same length, is within a factor c of the average cost of any other algorithm. Thus, bijective analysis can be used to establish “best of both worlds” types of performance comparisons. In fact, assuming again uniform distributions, much stronger conclusions can be obtained, in that bijective analysis implies a *stochastic dominance* relation between the costs of the two algorithms [8]. However, since the search domain is infinite, one must be careful in defining a uniform distribution of requests. More specifically, one could fix $L \geq 1$ and consider the uniform density function on the space $[-L, -1] \cup [1, L]$ (where the origin is assumed to be at 0). Thus, the probability that a request is at distance at most x from the origin is $(x - 1)/(L - 1)$. Our results then correspond to the setting in which L is unknown to the algorithm, and thus can be arbitrarily large. For known, and thus bounded L , the situation is much more complicated, since the optimal competitive ratio now depends on L and

¹In [18], online search refers to the problem of selling a specific item at the highest possible price, and is not related to the problem of searching for a target.

does not have a closed formula [16]. Our overall techniques still apply but the results unavoidably will be much more technical, and probably not tight.

It should be noted that the central question we study in this work is related to a phenomenon that is not unusual in online computation. Namely, for certain online problems, competitive analysis results in a very coarse performance classification of algorithms. This is due to the pessimistic, worst-case nature of the competitive ratio. The definitive example of an online problem in which this undesired situation occurs is the (standard) paging problem in a virtual memory system, which motivated the introduction of several analysis techniques alternative to the competitive ratio (see the surveys [30] and [22]). In our work we demonstrate that a similar situation arises in online search (which, perhaps surprisingly, has eluded attention so far) and we propose a remedy by means of the discovery ratio. We emphasize, however, that in our main results, we apply the discovery ratio as supplementary to the competitive ratio, instead of using it antagonistically as a measure that replaces the competitive ratio altogether. Thus, our analysis gives establishes *best-of-both-worlds* performance guarantees.

1.1 Contribution

We begin, in Section 2, by identifying the optimal tradeoff between the competitive ratio of a strategy and its discovery ratio (against all possible strategies). The result implies that there are strategies of discovery ratio $2 + \epsilon$, for arbitrarily small $\epsilon > 0$, which is tight. As corollary, we obtain that strategy **doubling** has discovery ratio equal to 3. These results allow us to set up the framework and provide some intuition for our main results, but also demonstrate that the discovery ratio, on itself, does not lead to a useful classification of strategies, when one considers the entire space of strategies.

Our main technical results are obtained in Section 3. Here, we apply synthetically both the competitive and the discovery ratios. More precisely, we restrict our interest to the set of competitively optimal strategies, which we further analyze using the discovery ratio as a supplementary measure. We prove that the strategy **aggressive**, which explores the branches to the furthest possible extent while satisfying the competitiveness constraint, has discovery ratio $\frac{8}{5} = 1.6$; moreover, we show that this is the optimal discovery ratio in this setting. In contrast, we show that the strategy **doubling** has discovery ratio $\frac{7}{3} = 2.\bar{3}$. In addition, we provide evidence that such “aggressiveness” is requisite. More precisely, we show that any competitively optimal strategy that is also optimal with respect to the discovery ratio must have the exact same behavior as the aggressive strategy in the first five iterations. Last, we show that the optimal strategy according to the discovery ratio is not unique.

In terms of techniques, the main technical difficulty in establishing the discovery ratios stems from answering the following question: given a length $\ell \in \mathbb{R}^+$, what is the strategy S that minimizes $D(S, \ell)$, and how can one express this minimum discovery cost? This is a type of a “dual” problem that can be of independent interest in the context of search problems, in the spirit

of a concept such as the *reach* of a strategy [26], also called *extent* in [27] (and which is useful in the competitive analysis of search strategies). We model this problem as a linear program for whose objective value we first give a lower bound; then we show this bound is tight by providing an explicit 9-competitive strategy which minimizes $D(S, \ell)$.

1.2 Related work

The linear search problem was first introduced and studied in works by Bellman [13] and Beck [11]. The generalization of linear search to m concurrent, semi-infinite branches is known as *star search* or *ray search*; thus linear search is equivalent to star search for $m = 2$. Optimal strategies for linear search under the (deterministic) competitive ratio were first given by [12]. Moreover [24] gave optimal strategies for the generalized problem of star search, a result that was rediscovered later [10]. Some of the related work includes the study of randomization [29]; multi-searcher strategies [34]; multi-target searching [31, 35]; searching with turn cost [2, 5, 21]; searching with an upper bound on the target distance [16, 26]; fault-tolerant search [20, 33] and searching with unreliable hints [4]; and the variant in which some probabilistic information on target placement is known [27, 28]. This list is not exclusive; see also Chapter 8 in the book [1].

Linear search and its generalization can model settings in which we seek an intelligent allocation of resources to tasks under uncertainty. For this reason, the problem and its solution often arises in the context of diverse fields such as AI (e.g., in the design of *interruptible algorithms* [3, 15]) and databases (e.g., *pipeline filter ordering* [19]).

Strategy **aggressive** has been studied in [26, 27] in the special case of maximizing the *reach* of a strategy (which informally is the maximum possible extent to which the branches can be searched without violating competitiveness) when we do not know the distance of the target from the origin. Although this gives some intuition that **aggressive** is indeed a good strategy, to the best of our knowledge, our work is the first that quantifies this intuition, in terms of comparing to other competitively optimal strategies using a well-defined, alternative performance measure.

1.3 Preliminaries

In the context of linear search, the searcher's strategy can be described as an (infinite) sequence of lengths at which the two branches (numbered 0,1, respectively) are searched. Formally, a search strategy is determined by a sequence of *search segments* (x_0, x_1, \dots) such that $x_i \in \mathbb{R}^+$ for all $i \in \mathbb{N}$, in the sense that in iteration i , the searcher starts from the origin, searches branch $i \bmod 2$ to distance x_i from the origin, and then returns back to O ; see Figure 1 for an illustration. We require that the search segments induce a complete exploration of both branches of the line, in that for every $d \in \mathbb{R}^+$, there exist $i, j \in \mathbb{N}$ such

that $x_{2i} \geq d$, and $x_{2j+1} \geq d$. Without loss of generality, we restrict ourselves to strategies which satisfy $x_{i+2} > x_i$ for every $i \in \mathbb{N}$.




Fig. 1 An illustration of a search strategy determined by the sequence $\{x_0, x_1, \dots\}$.

For convenience, we will define x_i to be equal to 0, for all $i < 0$ and introduce the following notation.

Definition 2 (prefix-sum $T_n(X)$) Given a strategy X , we define $T_n(X)$ (or simply T_n , when X is clear from context) to be equal to $\sum_{i=0}^n x_i$. For $n < 0$, we define $T_n := 0$.

We say that the searcher *turns* in iteration i at the moment it switches directions during iteration i , namely when it completes the exploration of length x_i and returns back to the origin. Moreover, assuming a searcher of unit speed, we can define at any point in time t the number of turns incurred by the searcher accordingly.

We will denote by \mathcal{S} the set of all search strategies, and by \mathcal{S}^* the set of competitively optimal strategies, namely strategies of competitive ratio equal to 9. When evaluating the competitive ratio, we will make the standard assumption that the target must be at distance at least 1 from O , since no strategy can have bounded competitive ratio if this distance can be arbitrarily small.

2 Strategies of optimal discovery ratio in \mathcal{S}

We begin by establishing the optimal tradeoff between the competitive ratio and the discovery ratio against all possible strategies. This will allow us to obtain strategies of optimal discovery ratio, and also setup some properties of the measure that will be useful in the context of competitively optimal strategies (Section 3).

Let X, Y denote two strategies in \mathcal{S} , with $X = (x_0, x_1, \dots)$. From the definition of the discovery ratio we have that

$$\text{dr}(X, Y) = \sup_{i \in \mathbb{N}} \sup_{\delta \in (0, x_i - x_{i-2}]} \frac{D(X, x_{i-1} + x_{i-2} + \delta)}{D(Y, x_{i-1} + x_{i-2} + \delta)}.$$

8 *Best-of-both-Worlds Analysis of Online Search*

Note that for $i = 0$, we have

$$\frac{D(X, x_{i-1} + x_{i-2} + \delta)}{D(Y, x_{i-1} + x_{i-2} + \delta)} = \frac{D(X, \delta)}{D(Y, \delta)} \leq \frac{\delta}{\delta} = 1.$$

This is because for all $\delta \leq x_0$, $D(X, \delta) = \delta$, and for all $\delta > 0$, $D(Y, \delta) \geq \delta$. Therefore,

$$\text{dr}(X, Y) = \sup_{i \in \mathbb{N}^+} \sup_{\delta \in (0, x_i - x_{i-2}]} \frac{D(X, x_{i-1} + x_{i-2} + \delta)}{D(Y, x_{i-1} + x_{i-2} + \delta)}. \quad (2)$$

The following theorem provides an expression of the discovery ratio in terms of the search segments of the strategy.

Theorem 1 (discovery ratio) *Let $X = (x_0, x_1, \dots)$. Then*

$$\text{dr}(X, \mathcal{S}) = \sup_{i \in \mathbb{N}^+} \frac{2 \sum_{j=0}^{i-1} x_j + x_{i-2}}{x_{i-1} + x_{i-2}}.$$

Proof Fix $Y \in \mathcal{S}$. From the definition of search segments in X , we have that

$$D(X, x_{i-1} + x_{i-2} + \delta) = 2 \sum_{j=0}^{i-1} x_j + x_{i-2} + \delta, \quad \text{for } \delta \in (0, x_i - x_{i-2}]. \quad (3)$$

Moreover, for every Y , we have

$$D(Y, x_{i-1} + x_{i-2} + \delta) \geq x_{i-1} + x_{i-2} + \delta. \quad (4)$$

Substituting (3) and (4) in (2) we obtain

$$\text{dr}(X, Y) \leq \sup_{i \in \mathbb{N}^+} \sup_{\delta \in (0, x_i - x_{i-2}]} \frac{2 \sum_{j=0}^{i-1} x_j + x_{i-2} + \delta}{x_{i-1} + x_{i-2} + \delta} \leq \sup_{i \in \mathbb{N}^+} \frac{2 \sum_{j=0}^{i-1} x_j + x_{i-2}}{x_{i-1} + x_{i-2}}. \quad (5)$$

For the lower bound, consider a strategy $Y_i = (y_0^i, y_1^i, \dots)$, for which $y_0^i = x_{i-1} + x_{i-2} + \delta$ (the values of y_j^i for $j \neq 0$ are not significant, as long as Y_i is a valid strategy). Clearly, $D(Y_i, x_{i-1} + x_{i-2} + \delta) = x_{i-1} + x_{i-2} + \delta$. Therefore, (2) implies

$$\text{dr}(X, Y_i) \geq \sup_{\delta \in (0, x_i - x_{i-2}]} \frac{2 \sum_{j=0}^{i-1} x_j + x_{i-2} + \delta}{x_{i-1} + x_{i-2} + \delta} = \frac{2 \sum_{j=0}^{i-1} x_j + x_{i-2}}{x_{i-1} + x_{i-2}}. \quad (6)$$

The lower bound on $\text{dr}(X, \mathcal{S})$ follows from $\text{dr}(X, \mathcal{S}) \geq \sup_{i \in \mathbb{N}^+} \text{dr}(X, Y_i)$. \square

In particular, note that for $i = 2$, Theorem 1 shows that for any strategy X ,

$$\text{dr}(X, \mathcal{S}) \geq \frac{3x_0 + 2x_1}{x_0 + x_1} \geq 2.$$

We will show that there exist strategies with discovery ratio arbitrarily close to 2, thus being optimal for \mathcal{S} . To this end, we will consider the geometric search strategy defined as $G_\alpha = (1, \alpha, \alpha^2, \dots)$, with $\alpha > 1$.

Lemma 1 (discovery ratio of geometric sequences) For G_α defined as above, we have $\text{dr}(G_\alpha, \mathcal{S}) = \frac{2\alpha^2 + \alpha - 1}{\alpha^2 - 1}$.

Proof From Theorem 1 we have

$$\begin{aligned} \text{dr}(G_\alpha, \mathcal{S}) &= \sup_{i \in \mathbb{N}^+} \frac{2 \sum_{j=0}^{i-1} \alpha^j + \alpha^{i-2}}{\alpha^{i-1} + \alpha^{i-2}} \\ &= \sup_{i \in \mathbb{N}^+} \frac{2 \left(\frac{\alpha^i - 1}{\alpha - 1} \right) + \alpha^{i-2}}{\alpha^{i-1} + \alpha^{i-2}} \\ &= \sup_{i \in \mathbb{N}^+} \frac{2(\alpha^i - 1) + \alpha^{i-1} - \alpha^{i-2}}{\alpha^i - \alpha^{i-2}}. \end{aligned}$$

The derivative of the function $f(i) := \frac{2(\alpha^i - 1) + \alpha^{i-1} - \alpha^{i-2}}{\alpha^i - \alpha^{i-2}}$ in i is

$$f'(i) = \frac{2\alpha^{2-i} \log \alpha}{\alpha^2 - 1},$$

which is positive. Thus, $\sup_{i \in \mathbb{N}^+} f(i) = \lim_{i \rightarrow \infty} f(i)$, which gives

$$\text{dr}(G_\alpha, \mathcal{S}) = \lim_{i \rightarrow +\infty} f(i) = \lim_{i \rightarrow +\infty} \frac{2(\alpha^i - 1) + \alpha^{i-1} - \alpha^{i-2}}{\alpha^i - \alpha^{i-2}} = \frac{2\alpha^2 + \alpha - 1}{\alpha^2 - 1}.$$

□

Lemma 1 shows that the discovery ratio of G_α tends to 2, as $\alpha \rightarrow \infty$, hence G_α has an asymptotically optimal discovery ratio. However, we can show a stronger result, namely that G_α achieves the optimal trade-off between the discovery ratio and the competitive ratio. This is established in the following theorem. Note that the competitive ratio of G_α is easily verified to be $1 + 2 \frac{\alpha^2}{\alpha - 1}$ (and is minimized for $\alpha = 2$).

Theorem 2 (dominance of geometric sequences) *For every strategy $X \in \mathcal{S}$, there exists $\alpha > 1$ such that $\text{dr}(X, \mathcal{S}) \geq \text{dr}(G_\alpha, \mathcal{S})$ and $\text{cr}(X) \geq \text{cr}(G_\alpha)$.*

In order to prove Theorem 2, we will use of a result by Gal [25] and Schuierer [36] which, informally, lower-bounds the supremum of an infinite sequence of functionals by the supremum of simple functionals of a certain geometric sequence, and which we state here in a simplified form. Given an infinite sequence $X = (x_0, x_1, \dots)$, define $X^{+i} = (x_i, x_{i+1}, \dots)$ as the suffix of the sequence X starting at x_i .

Theorem 3 ([25, 36]) *Let $X = (x_0, x_1, \dots)$ be a sequence of positive numbers, r an integer, and $\alpha = \limsup_{n \rightarrow \infty} (x_n)^{1/n}$, for $\alpha \in \mathbb{R} \cup \{+\infty\}$. Let F_i , $i \geq 0$ be a sequence of functionals which satisfy the following properties:*

1. $F_i(X)$ only depends on x_0, x_1, \dots, x_{i+r} ,
2. $F_i(X)$ is continuous for all $x_k > 0$, with $0 \leq k \leq i + r$,

3. $F_i(\lambda X) = F_i(X)$, for all $\lambda > 0$,
4. $F_i(X + Y) \leq \max(F_i(X), F_i(Y))$, and
5. $F_{i+k}(X) \geq F_i(X^{+k})$, for all $k \geq 1$,

then

$$\sup_{0 \leq i < \infty} F_i(X) \geq \sup_{0 \leq i < \infty} F_i(G_\alpha).$$

We do not reproduce the proof of the above theorem, but show how we can deduce Theorem 2 from it.

Proof of Theorem 2 Let $X = (x_0, x_1, \dots)$ denote a strategy in \mathcal{S} . From (6) we know that

$$\text{dr}(X, \mathcal{S}) \geq \sup_i F_i(X),$$

where $F_i(X)$ is defined as the functional $\frac{2 \sum_{j=0}^{i-1} x_j + x_{i-2}}{x_{i-1} + x_{i-2}}$. Moreover, the competitive ratio of X can be lower-bounded by

$$\text{cr}(X) \geq \sup_i F'_i(X), \quad \text{where } F'_i(X) = 1 + 2 \frac{\sum_{j=0}^{i+1} x_j}{x_i}.$$

This follows easily by considering a hider placed at distance $x_i + \epsilon$, with $\epsilon \rightarrow 0$, at the branch that is searched by X in iteration i .

It is easy to see that both $F_i(X)$ and $F'_i(X)$ satisfy the conditions of Theorem 3 (this also follows from Example 7.3 in [1]). Thus, there exists α defined as in the statement of Theorem 3 such that

$$\text{dr}(X, \mathcal{S}) \geq \sup_i F_i(G_\alpha) = \frac{2 \sum_{j=0}^{i-1} \alpha^j + \alpha^{i-2}}{\alpha^{i-1} + \alpha^{i-2}}, \quad \text{and} \quad (7)$$

$$\text{cr}(X, \mathcal{S}) \geq \sup_i F'_i(G_\alpha) = 1 + 2 \frac{\sum_{j=0}^{i+1} \alpha^j}{\alpha^i}. \quad (8)$$

It is easy to verify that if $\alpha = 1$, then $\text{dr}(X, \mathcal{S}), \text{cr}(X, \mathcal{S}) = \infty$. We can thus assume that $\alpha > 1$, and thus obtain from (7), (8), after some manipulations, that

$$\begin{aligned} \text{dr}(X, \mathcal{S}) &\geq \sup_i \frac{2(\alpha^2 - \frac{1}{\alpha^{i-2}}) + \alpha - 1}{\alpha^2 - 1} = \frac{2\alpha^2 + \alpha - 1}{\alpha^2 - 1}, \quad \text{and} \\ \text{cr}(X, \mathcal{S}) &\geq 1 + \sup_i \frac{2 \sum_{j=0}^{i+1} \alpha^j}{\alpha^i} = \sup_i 1 + 2 \frac{\alpha^2 - \frac{1}{\alpha^i}}{\alpha - 1} = 1 + 2 \frac{\alpha^2}{\alpha - 1}, \end{aligned}$$

which concludes the proof. \square

Figure 2 depicts the optimal tradeoff between the competitive and discovery ratios as attained by the search strategy G_α .

Note that although G_α with $\alpha \rightarrow \infty$ has optimal discovery ratio, its competitive ratio is unbounded. Furthermore, strategy `doubling` $\equiv G_2$ has optimal competitive ratio equal to 9, whereas its discovery ratio is equal to 3. This motivates the topic of the next section.




Fig. 2 Every strategy X corresponds to a point with coordinates $\text{dr}(X)$ and $\text{cr}(X)$, which belongs to the region depicted in gray. Its Pareto boundary is constituted by the geometric strategies G_α for $\alpha > 1$.

3 The discovery ratio of competitively optimal strategies

In this section we focus on competitively optimal strategies, namely strategies in \mathcal{S}^* . It is known that there is a characterization of such strategies in terms of an infinite set of linear inequalities that relate its search segments, as shown in the following lemma.

Lemma 2 (characterization of competitive optimal strategies [26]) The strategy $X = (x_0, x_1, x_2, \dots)$ is in \mathcal{S}^* if and only if its segments satisfy the following inequalities

$$1 \leq x_0 \leq 4 \quad \text{and} \quad x_n \leq 3x_{n-1} - \sum_{i=0}^{n-2} x_i, \quad \text{for all } n \geq 1.$$

There is an interesting corollary of this characterization, shown in [27]. Namely, the strategy **aggressive**, which maximizes the search segments in each iteration, has the property that its segment lengths satisfy all linear inequalities of Lemma 2 with equality, which provides a closed-form expression of its lengths:

Lemma 3 (closed expression for **aggressive** [27]) Strategy **aggressive** satisfies for every $n \geq 0$

$$\bar{x}_n = (4 + 2n)2^n \\ \text{and } T_n = (n + 1)2^{n+2}.$$

For the purpose of our analysis, we will define a class of strategies in \mathcal{S}^* that generalize the aggressive strategy. For given $t \in [1, 4]$, let R_t denote the

strategy whose search segments are determined by the linear recurrence

$$x_0 = t, \quad \text{and} \quad x_n = 3x_{n-1} - \sum_{i=0}^{n-2} x_i, \quad \text{for all } n \geq 1.$$

This class generalizes **aggressive** in the sense that R_4 equals **aggressive**, and the values x_n in R_t and R_4 differ exactly by the factor $t/4$. As a consequence, previously shown properties of **aggressive** hold for R_t as well, after applying the scaling factor $t/4$.

Corollary 1 (closed expression for R_t) The strategy R_t satisfies for every $n \geq 0$

$$x_n = t(1 + n/2)2^n \\ \text{and } T_n = t(n + 1)2^n.$$

In addition, we define the minimum number of turns that a strategy requires to discover a certain length.

Definition 3 (minimal number of turns m^*) Given strategy X and length $\ell \in \mathbb{R}^+$, define $m(X, \ell)$ as the number of turns that X has performed by the time it discovers a total length equal to ℓ . Also define

$$m^*(\ell) = \inf_{X \in \mathcal{S}^*} m(X, \ell),$$

that is, $m^*(\ell)$ is the minimum number of turns that a competitively optimal strategy is required to perform in order to discover length equal to ℓ .

From the constraint $x_0 \leq 4$, it follows that clearly $m^*(\ell) = 0$, for $\ell \leq 4$. The following corollary gives an expression for $m^*(\ell)$, for general values of ℓ .

Corollary 2 (turns of **aggressive**) For given $\ell > 4$,

$$m^*(\ell) = m(\mathbf{aggressive}, \ell) = \min\{n \in \mathbb{N}^+ : (3n + 5)2^n \geq \ell\}.$$

Proof The total length discovered by any $X \in \mathcal{S}^*$ at the turning point of the n -th iteration cannot exceed $\bar{x}_n + \bar{x}_{n-1}$ for $n \geq 1$, where x_i is the i -th step of **aggressive**. This implies that $m^*(\ell) = n$, if $\ell \in (\bar{x}_{n-1} + \bar{x}_{n-2}, \bar{x}_n + \bar{x}_{n-1}]$ for $n \geq 1$. In other words,

$$m^*(\ell) = \min\{n \in \mathbb{N}^+ : \bar{x}_n + \bar{x}_{n-1} \geq \ell\}.$$

From Lemma 3, we have $\bar{x}_n = (n + 2)2^{n+1}$, for $n \geq 0$. Hence,

$$m^*(\ell) = \min\{n \in \mathbb{N}^+ : (3n + 5)2^n \geq \ell\}.$$

□

We also define the minimum cost at which a competitively optimal strategy can discover a length equal to ℓ , namely:

Definition 4 (optimal cost for discovering length ℓ in \mathcal{S}^*) For a given $\ell \in \mathbb{R}^+$, define

$$d^*(\ell) = \inf_{X \in \mathcal{S}^*} D(X, \ell),$$

and note that, if $\ell \leq 4$, then trivially $d^*(\ell) = \ell$.

The following lemma is a central technical result that is instrumental in establishing the bounds on the discovery ratio. Specifically, Lemma 4 gives an expression of $d^*(\ell)$ for $\ell > 4$ in terms of $m^*(\ell)$; it also shows that there exists $t \in (1, 4]$ such that the strategy R_t attains this minimum cost.

We first give some motivation behind the purpose of the lemma. When considering general strategies in \mathcal{S} , we used a lower bound on the cost for discovering a length ℓ as given by (4), and which corresponds to a strategy that never turns. However, this lower bound is very weak when one considers strategies in \mathcal{S}^* . This is because a competitive strategy needs to turn sufficiently often, which affects considerably the discovery costs.

We also give some intuition about the proof. We show how to model the question by means of a linear program. Using the constraints of the LP, we first obtain a lower bound on its objective in terms of the parameters ℓ and $m^*(\ell)$. In this process, we also obtain a lower bound on the first segment of the strategy (x_0); this is denoted by t in the proof. In the next step, we show that the strategy R_t has a discovery cost that matches the lower bound on the objective, which suffices to prove the result.

Lemma 4 (discovery cost is optimized by some R_t) For any given $\ell > 4$, it holds

$$d^*(\ell) = D(R_t, \ell) = \ell \cdot \frac{6m^*(\ell) + 4}{3m^*(\ell) + 5}, \quad \text{where } t = \ell \cdot \frac{2^{2-m^*(\ell)}}{3m^*(\ell) + 5} \in (1, 4].$$

Proof Let $X = (x_0, x_1, \dots) \in \mathcal{S}^*$ denote the strategy which minimizes the quantity $D(X, \ell)$. Then there must exist a smallest $n \geq m^*(\ell)$ such that the searcher discovers a total length ℓ during the n -th iteration. More precisely, suppose that this happens when the searcher is at branch $n \bmod 2$, and at some position p (i.e., distance from O), with $p \in (x_{n-2}, x_n]$. Then we have $x_{n-1} + p = \ell$, and

$$d^*(\ell) = D(X, \ell) = 2 \sum_{i=0}^{n-1} x_i + p = 2 \sum_{i=0}^{n-1} x_i + (\ell - x_{n-1}) = 2 \sum_{i=0}^{n-2} x_i + x_{n-1} + \ell.$$

Therefore, $d^*(\ell)$ is the objective of the following linear program.

$$\begin{aligned} \min \quad & 2 \sum_{i=0}^{n-2} x_i + x_{n-1} + \ell \\ \text{subject to} \quad & x_n + x_{n-1} \geq \ell, \\ & 1 \leq x_0 \leq 4, \\ & x_{i-2} \leq x_i, \quad i \in [2, n] \end{aligned}$$

$$1 \leq x_i \leq 3x_{i-1} - \sum_{j=0}^{i-2} x_j, \quad i \in [1, n].$$

Recall that $n \geq m^*(\ell)$ is a fixed integer. Let Obj denote the objective value of the linear program. We claim that, for $1 \leq i \leq n$,

$$x_{n-i} \geq \frac{2^{2-i}}{3i+5}\ell + \frac{3i-1}{3i+5}T_{n-i-1} \quad \text{and} \quad \text{Obj} \geq \frac{6i+4}{3i+5}\ell + \frac{9 \cdot 2^i}{3i+5}T_{n-i-1}.$$

The claim provides a lower bound of the objective, since for $i = n$ it implies that

$$x_0 \geq \frac{2^{2-n}}{3n+5}\ell \quad \text{and} \quad \text{Obj} \geq \frac{6n+4}{3n+5}\ell \geq \frac{6m^*(\ell)+4}{3m^*(\ell)+5}\ell,$$

where the last inequality follows from the fact $n \geq m^*(\ell)$, and by monotonicity on n . We will argue later that this lower bound is tight.

First, we prove the claim, by induction on i , for all $i \leq n$. We first show the base case, namely $i = 1$. Since $x_n \leq 3x_{n-1} - T_{n-2}$ and $x_n + x_{n-1} \geq \ell$, it follows that

$$x_{n-1} \geq \ell - x_n \geq \ell - (3x_{n-1} - T_{n-2}) \Rightarrow x_{n-1} \geq \frac{\ell}{4} + \frac{T_{n-2}}{4}, \quad \text{hence}$$

$$\text{Obj} = \ell + 2T_{n-2} + x_{n-1} \geq \ell + 2T_{n-2} + \frac{\ell}{4} + \frac{T_{n-2}}{4} = \frac{5}{4}\ell + \frac{9}{4}T_{n-2},$$

thus the base case holds. For the induction step, suppose that

$$x_{n-i} \geq \frac{2^{2-i}}{3i+5}\ell + \frac{3i-1}{3i+5}T_{n-i-1} \quad \text{and} \quad \text{Obj} \geq \frac{6i+4}{3i+5}\ell + \frac{9 \cdot 2^i}{3i+5}T_{n-i-1}.$$

Then,

$$\begin{aligned} 3x_{n-i-1} - T_{n-i-2} &\geq x_{n-i} && \text{(by LP constraint)} \\ &\geq \frac{2^{2-i}}{3i+5}\ell + \frac{3i-1}{3i+5}T_{n-i-1} && \text{(ind. hyp.)} \\ &= \frac{2^{2-i}}{3i+5}\ell + \frac{3i-1}{3i+5}(T_{n-i-2} + x_{n-i-1}) && \text{(def. } T_{n-i-1}) \end{aligned}$$

By rearranging terms in the above inequality we obtain

$$\begin{aligned} \left(3 - \frac{3i-1}{3i+5}\right)x_{n-i-1} &\geq \frac{2^{2-i}}{3i+5}\ell + \left(1 + \frac{3i-1}{3i+5}\right)T_{n-i-2} \Rightarrow \\ \frac{6i+16}{3i+5}x_{n-i-1} &\geq \frac{2^{2-i}}{3i+5}\ell + \frac{6i+4}{3i+5}T_{n-i-2} \Rightarrow \\ x_{n-i-1} &\geq \frac{2^{1-i}}{3i+8}\ell + \frac{3i+2}{3i+8}T_{n-i-2}, \end{aligned}$$

and

$$\begin{aligned} \text{Obj} &\geq \frac{6i+4}{3i+5}\ell + \frac{9 \cdot 2^i}{3i+5}T_{n-i-1} && \text{(ind. hyp.)} \\ &= \frac{6i+4}{3i+5}\ell + \frac{9 \cdot 2^i}{3i+5}(T_{n-i-2} + x_{n-i-1}) && \text{(def. } T_{n-i-1}) \\ &\geq \frac{6i+4}{3i+5}\ell + \frac{9 \cdot 2^i}{3i+5}T_{n-i-2} + \frac{9 \cdot 2^i}{3i+5} \left(\frac{2^{1-i}}{3i+8}\ell + \frac{3i+2}{3i+8}T_{n-i-2} \right) && \text{(ind. hyp.)} \\ &= \frac{6i+10}{3i+8}\ell + \frac{9 \cdot 2^{i+1}}{3i+8}T_{n-i-2}. \end{aligned}$$

This concludes the proof of the claim, which settles the lower bound on $d^*(\ell)$. It remains to show that this bound is tight. Consider the strategy R_t , with $t = \frac{2^{2-m^*(\ell)}}{3m^*(\ell)+5}\ell$. In what follows we will show that R_t is a feasible solution of the LP, and that $D(R_t, \ell) = \frac{6m^*(\ell)+4}{3m^*(\ell)+5}\ell$.

First, we show that $t \in (1, 4]$. For the upper bound, from Corollary 2, we have $(3m^*(\ell) + 5)2^{m^*(\ell)} \geq \ell$, which implies that

$$1 \geq \ell \cdot \frac{2^{-m^*(\ell)}}{3m^*(\ell) + 5} \Rightarrow 4 \geq \ell \cdot \frac{2^{2-m^*(\ell)}}{3m^*(\ell) + 5} \Rightarrow 4 \geq t.$$

In order to show that $t > 1$, consider first the case $\ell \in (4, 5]$. Then $m^*(\ell) = 1$, which implies that

$$t = \frac{2^{2-m^*(\ell)}}{3m^*(\ell) + 5}\ell = \frac{\ell}{4} \geq 1.$$

Moreover, if $\ell > 5$, by Corollary 2, $m^*(\ell)$ is the smallest integer solution of the inequality $(3n + 5)2^n \geq \ell$, then $(3m^*(\ell) + 2)2^{m^*(\ell)-1} < \ell$, hence

$$\begin{aligned} t &= \frac{2^{2-m^*(\ell)}}{3m^*(\ell) + 5}\ell = \frac{4\ell}{(3m^*(\ell) + 5)2^{m^*(\ell)}} = \frac{2\ell}{(3m^*(\ell) + 2)2^{m^*(\ell)-1} \cdot \frac{3m^*(\ell)+5}{3m^*(\ell)+2}} \\ &> \frac{2\ell}{\ell \cdot \frac{3m^*(\ell)+5}{3m^*(\ell)+2}} = \frac{6m^*(\ell) + 4}{3m^*(\ell) + 5} > 1. \end{aligned}$$

The last inequality holds since we have $m^*(\ell) \geq 1$, for $\ell > 5$. This concludes that $t \in (1, 4]$, and R_t is a feasible solution of the LP since R_t satisfies all other constraints by its definition.

It remains thus to show that $D(R_t, \ell) = \frac{6m^*(\ell)+4}{3m^*(\ell)+5}\ell$. By Lemma 1, we have

$$\begin{aligned} x_{m^*(\ell)} + x_{m^*(\ell)-1} &= t \left(1 + \frac{m^*(\ell)}{2} \right) 2^{m^*(\ell)} + t \left(1 + \frac{m^*(\ell)-1}{2} \right) 2^{m^*(\ell)-1} \\ &= t \cdot 2^{m^*(\ell)} \cdot \frac{3m^*(\ell) + 5}{4} = \frac{2^{2-m^*(\ell)}}{3m^*(\ell) + 5}\ell \cdot 2^{m^*(\ell)} \cdot \frac{3m^*(\ell) + 5}{4} = \ell. \end{aligned}$$

Then R_t has exactly discovered a total length ℓ right before the $m^*(\ell)$ -th turn. Hence,

$$\begin{aligned} D(R_t, \ell) &= 2T_{m^*(\ell)-2} + x_{m^*(\ell)-1} + \ell \\ &= t \cdot (m^*(\ell) - 1) 2^{m^*(\ell)-1} + t \cdot \left(1 + \frac{m^*(\ell)-1}{2} \right) 2^{m^*(\ell)-1} + \ell \\ &\hspace{15em} \text{(by Lemma 1)} \\ &= t \cdot \frac{(3m^*(\ell) - 1)2^{m^*(\ell)}}{4} + \ell \hspace{10em} \text{(arranging terms)} \\ &= \frac{2^{2-m^*(\ell)}}{3m^*(\ell) + 5}\ell \cdot \frac{(3m^*(\ell) - 1)2^{m^*(\ell)}}{4} + \ell \hspace{10em} \text{(substituting } t) \\ &= \left(\frac{3m^*(\ell) - 1}{3m^*(\ell) + 5} + 1 \right) \cdot \ell = \frac{6m^*(\ell) + 4}{3m^*(\ell) + 5} \cdot \ell. \hspace{10em} \text{(arranging terms)} \end{aligned}$$

This concludes the proof of the lemma. \square

We are now ready to prove the main results of this section. Recall that for any two strategies X, Y , $\text{dr}(X, Y)$ is given by (2). Combining with (3), as

well as with the fact that for $Y \in \mathcal{S}^*$, we have that $D(Y, \ell) \geq d^*(\ell)$, (from the definition of d^*), we obtain that

$$\text{dr}(X, \mathcal{S}^*) = \sup_{i \in \mathbb{N}^+} \sup_{\delta \in (0, x_i - x_{i-2}]} F_i(X, \delta), \quad \text{where } F_i(X, \delta) = \frac{2 \sum_{j=0}^{i-1} x_j + x_{i-2} + \delta}{d^*(x_{i-1} + x_{i-2} + \delta)}. \quad (9)$$

3.1 Discovery ratio of strategies in \mathcal{S}^*

We first show that the discovery ratio of **aggressive** is equal to $8/5$.

Theorem 4 (discovery ratio of **aggressive**) *For the strategy **aggressive** it holds that*

$$\text{dr}(\text{aggressive}, \mathcal{S}^*) = 8/5.$$

Proof We will express the discovery ratio using (9). For $i = 1$, and $\delta \in (0, \bar{x}_1]$, we have that

$$F_1(\text{aggressive}, \delta) = \frac{2\bar{x}_0 + \delta}{d^*(\bar{x}_0 + \delta)} = \frac{8 + \delta}{d^*(4 + \delta)}.$$

From Lemma 4, $d^*(4 + \delta) = (4 + \delta) \cdot \frac{6 \cdot 1 + 4}{3 \cdot 1 + 5} = \frac{5(4 + \delta)}{4}$; this is because $1 \leq m^*(4 + \delta) \leq m^*(16) = 1$. Then,

$$F_1(\text{aggressive}, \delta) = \frac{8 + \delta}{\frac{5(4 + \delta)}{4}} = \frac{32 + 4\delta}{20 + 5\delta}, \quad \text{hence } \sup_{\delta \in (0, \bar{x}_1]} F_1(\text{aggressive}, \delta) = \frac{8}{5}. \quad (10)$$

For given $i \geq 2$, and $\delta \in (0, \bar{x}_i - \bar{x}_{i-2}]$, we have

$$F_i(\text{aggressive}, \delta) = \frac{2T_{i-1} + \bar{x}_{i-2} + \delta}{d^*(\bar{x}_{i-1} + \bar{x}_{i-2} + \delta)},$$

where T_{i-1} is given by Lemma 3. Moreover, from Lemma 4 we have that

$$\begin{aligned} d^*(\bar{x}_{i-1} + \bar{x}_{i-2} + \delta) &= (\bar{x}_{i-1} + \bar{x}_{i-2} + \delta) \cdot \frac{6m^*(\bar{x}_{i-1} + \bar{x}_{i-2} + \delta) + 4}{3m^*(\bar{x}_{i-1} + \bar{x}_{i-2} + \delta) + 5} \\ &= (\bar{x}_{i-1} + \bar{x}_{i-2} + \delta) \cdot \frac{6i + 4}{3i + 5}, \end{aligned}$$

where the last equality follows from the fact that $m^*(\bar{x}_{i-1} + \bar{x}_{i-2} + \delta) = i$. This is because

$$i \leq m^*(\bar{x}_{i-1} + \bar{x}_{i-2} + \delta) \leq m^*(\bar{x}_{i-1} + \bar{x}_{i-2} + \bar{x}_i - \bar{x}_{i-2}) = m^*(\bar{x}_i + \bar{x}_{i-1}) = i.$$

Substituting with the values of the search segments as well as T_{i-1} , we obtain that

$$F_i(\text{aggressive}, \delta) = \frac{i \cdot 2^{i+2} + i \cdot 2^{i-1} + \delta}{((i+1)2^i + i \cdot 2^{i-1} + \delta) \cdot \frac{6i+4}{3i+5}} = \frac{9i \cdot 2^{i-1} + \delta}{((3i+2)2^{i-1} + \delta) \cdot \frac{6i+4}{3i+5}}.$$

Since

$$\frac{\partial F_i(\text{aggressive}, \delta)}{\partial \delta} = -\frac{2^{i+1}(3i-1)(3i+5)}{(3i+2)(2^n(3i+2) + 2\delta)^2} \leq 0,$$

then $F_i(\mathbf{aggressive}, \delta)$ is monotone decreasing in δ . Thus

$$\sup_{\delta \in (0, \bar{x}_i - \bar{x}_{i-2}]} F_i(\mathbf{aggressive}, \delta) = \frac{9i \cdot 2^{i-1}}{((3i+2)2^{i-1}) \cdot \frac{6i+4}{3i+5}} = \frac{9i(3i+5)}{(3i+2)(6i+4)},$$

and

$$\sup_{i \geq 2} \sup_{\delta \in (0, \bar{x}_i - \bar{x}_{i-2}]} F_i(\mathbf{aggressive}, \delta) = \frac{(9 \cdot 2)(3 \cdot 2 + 5)}{(3 \cdot 2 + 2)(6 \cdot 2 + 4)} = \frac{99}{64} < \frac{8}{5}. \quad (11)$$

Combining (9), (10) and (11) yields the proof of the theorem. \square

The following theorem shows that **aggressive** has optimal discovery ratio among all competitively optimal strategies.

Theorem 5 (lower bound on discovery ratio) *For every strategy $X \in \mathcal{S}^*$, we have $dr(X, \mathcal{S}^*) \geq \frac{8}{5}$.*

Proof Let $X = (x_0, \dots)$. We will consider two cases, depending on whether $x_0 < 4$ or $x_0 = 4$. Suppose, first, that $x_0 < 4$. In this case, for sufficiently small ϵ , we have $m^*(x_0 + \epsilon) = 0$, which implies that $d^*(x_0 + \epsilon) = x_0 + \epsilon$, and therefore.

$$F_1(X, \epsilon) = \frac{2x_0 + \epsilon}{d^*(x_0 + \epsilon)} = \frac{2x_0 + \epsilon}{x_0 + \epsilon},$$

from which we obtain that

$$\sup_{\delta \in (0, x_1]} F_1(X, \delta) \geq F_1(X, \epsilon) \geq \frac{2x_0 + \epsilon}{x_0 + \epsilon} \rightarrow 2, \text{ as } \epsilon \rightarrow 0^+.$$

Next, suppose that $x_0 = 4$. In this case, for $\delta \in (0, x_1]$, it readily follows that $F_1(X, \delta) = F_1(\mathbf{aggressive}, \delta)$. Therefore, from (10), we have that

$$\sup_{\delta \in (0, x_1]} F_1(X, \delta) = \sup_{\delta \in (0, x_1]} \frac{32 + 4\delta}{20 + 5\delta} = \frac{8}{5}.$$

The lower bound follows directly from (9). \square

Recall that **doubling** $\equiv G_2 = (2^0, 2^1, 2^2, \dots)$. The following theorem shows that within \mathcal{S}^* , **doubling** has a worse discovery ratio than **aggressive**. The proof follows along the lines of the proof of Theorem 4, where instead of using the search segments \bar{x}_i of **aggressive**, we use the search segment $x_i = 2^i$ of **doubling**.

Theorem 6 (discovery ratio of **doubling**) *We have $dr(\mathbf{doubling}, \mathcal{S}^*) = \frac{7}{3}$.*

Proof We will express the discovery ratio using (9). For $i = 1$, and $\delta \in (0, x_1]$, we have that

$$F_1(G_2, \delta) = \frac{2x_0 + \delta}{d^*(x_0 + \delta)} = \frac{2 + \delta}{d^*(1 + \delta)}.$$

From the definition, $d^*(1 + \delta) = 1 + \delta$; this is because $0 \leq m^*(1 + \delta) \leq m^*(3) = 0$. Then,

$$F_1(G_2, \delta) = \frac{2 + \delta}{1 + \delta}, \text{ hence } \sup_{\delta \in (0, \bar{x}_1]} F_1(G_2, \delta) = 2. \quad (12)$$

For $i = 2$ and $\delta \in (0, x_2 - x_0]$, we have that

$$F_2(G_2, \delta) = \frac{3x_0 + 2x_1 + \delta}{d^*(x_0 + x_1 + \delta)} = \frac{7 + \delta}{d^*(3 + \delta)}.$$

From Lemma 4, $d^*(3 + \delta)$ either equals to $3 + \delta$ if $\delta \in (0, 1]$, or equals to $(3 + \delta) \cdot \frac{6 \cdot 1 + 4}{3 \cdot 1 + 5} = \frac{5(3 + \delta)}{4}$ if $\delta \in (1, x_2 - x_0]$. This is because $0 \leq m^*(3 + \delta) \leq m^*(6) = 1$. Then, for $\delta \in (0, 1]$,

$$F_2(G_2, \delta) = \frac{7 + \delta}{3 + \delta}, \text{ hence } \sup_{\delta \in (0, 1]} F_2(G_2, \delta) = \frac{7}{3}. \quad (13)$$

For $\delta \in (1, x_2 - x_0]$,

$$F_2(G_2, \delta) = \frac{7 + \delta}{\frac{5(3 + \delta)}{4}} = \frac{28 + 4\delta}{15 + 5\delta}, \text{ hence } \sup_{\delta \in (1, x_2 - x_0]} F_2(G_2, \delta) = \frac{28}{15}. \quad (14)$$

Combining (13) and (14) yields

$$\sup_{\delta \in (0, x_2 - x_0]} F_2(G_2, \delta) = \frac{7}{3}. \quad (15)$$

For given $i \geq 3$, and $\delta \in (0, x_i - x_{i-2}]$, we have

$$F_i(G_2, \delta) = \frac{2T_{i-1} + x_{i-2} + \delta}{d^*(x_{i-1} + x_{i-2} + \delta)} = \frac{2^{i+1} - 2 + 2^{i-2} + \delta}{d^*(2^{i-1} + 2^{i-2} + \delta)} = \frac{9 \cdot 2^{i-2} - 2 + \delta}{d^*(2^{i-1} + 2^{i-2} + \delta)}.$$

Moreover, from Lemma 4 we have that

$$d^*(2^{i-1} + 2^{i-2} + \delta) = (2^{i-1} + 2^{i-2} + \delta) \cdot \frac{6m^*(2^{i-1} + 2^{i-2} + \delta) + 4}{3m^*(2^{i-1} + 2^{i-2} + \delta) + 5}.$$

and

$$m^*(x_{i-1} + x_{i-2} + \delta) \geq m^*(x_{i-1} + x_{i-2}) = m^*(3 \cdot 2^{i-2}).$$

For $i \in \{3, 4\}$ and $\delta \in (0, x_i - x_{i-2}]$, $m^*(x_{i-1} + x_{i-2} + \delta) \geq m^*(3 \cdot 2^{i-2}) \geq 1$, then

$$d^*(2^{i-1} + 2^{i-2} + \delta) \geq (2^{i-1} + 2^{i-2} + \delta) \cdot \frac{6 \cdot 1 + 4}{3 \cdot 1 + 5} = \frac{15 \cdot 2^{i-2} + 5\delta}{4},$$

hence, for $i = 3$,

$$F_3(G_2, \delta) = \frac{9 \cdot 2^{i-2} - 2 + \delta}{d^*(2^{i-1} + 2^{i-2} + \delta)} \leq \frac{16 + \delta}{\frac{30 + 5\delta}{4}} = \frac{64 + 4\delta}{30 + 5\delta}.$$

We obtain that

$$\sup_{\delta \in (0, x_3 - x_1]} F_3(G_2, \delta) \leq \frac{64}{30}. \quad (16)$$

For $i = 4$,

$$F_4(G_2, \delta) = \frac{9 \cdot 2^{i-2} - 2 + \delta}{d^*(2^{i-1} + 2^{i-2} + \delta)} \leq \frac{32 + \delta}{\frac{60 + 5\delta}{4}} = \frac{128 + 4\delta}{60 + 5\delta}.$$

We obtain that

$$\sup_{\delta \in (0, x_4 - x_2]} F_4(G_2, \delta) \leq \frac{128}{60}. \quad (17)$$

For $i \geq 5$ and $\delta \in (0, x_i - x_{i-2}]$, $m^*(x_{i-1} + x_{i-2} + \delta) \geq m^*(3 \cdot 2^{i-2}) \geq 2$, then

$$d^*(2^{i-1} + 2^{i-2} + \delta) \geq (2^{i-1} + 2^{i-2} + \delta) \cdot \frac{6 \cdot 2 + 4}{3 \cdot 2 + 5} = \frac{48 \cdot 2^{i-2} + 16\delta}{11},$$

and

$$F_i(G_2, \delta) = \frac{9 \cdot 2^{i-2} - 2 + \delta}{d^*(2^{i-1} + 2^{i-2} + \delta)} \leq \frac{9 \cdot 2^{i-2} - 2 + \delta}{\frac{48 \cdot 2^{i-2} + 16\delta}{11}} = \frac{99 \cdot 2^{i-2} - 22}{48 \cdot 2^{i-2}} \leq \frac{99}{48},$$

hence, for $i \geq 5$,

$$\sup_{\delta \in (0, x_i - x_{i-2}]} F_i(G_2, \delta) \leq \frac{99}{48}. \quad (18)$$

Combining (12), (15), (16), (17) and (18) yields the proof of the theorem. \square

We conclude this section with the discussion of a strategy that is conceptually opposite to the aggressive strategy. We define the strategy **conservative** as the strategy $X = (x_i)_{i=0}^{\infty}$ in \mathcal{S}^* for which x_i is as small as possible, for all $i \in \mathbb{N}$. As shown in Proposition 17 in [6] this strategy is defined by the sequence of the search lengths

$$1, 1, 2, 4, 8, 16 \dots$$

Note that **conservative** differs from **doubling** only in the additional cost of 2 in the discovery time for any length strictly greater than 1. Adapting the proof of Theorem 6 for this additional cost, we observe that the ratio is maximized for a length arbitrary close to 3 from above, which leads to the following corollary.

Corollary 3 The discovery ratio of **conservative** is 3.

Figure 3 illustrates the discovery ratios shown for the various strategies in this section. Note that the discovery ratio indeed orders the strategies in terms of the “aggressiveness” at which they tend to explore new ground over the line.




Fig. 3 Illustration of the discovery ratios discussed in this section.

3.2 On the uniqueness of strategies with optimal discovery ratio

The results of the previous section lead to a natural question. Is **aggressive** the unique strategy of optimal discovery ratio in \mathcal{S}^* ? We first provide evidence

that optimal strategies cannot be radically different from **aggressive**, in that they must mimic it in the first few iterations.

Theorem 7 (optimal strategies coincide with **aggressive** on the first 5 steps) *Strategy* $X = (x_i)_{i=0}^\infty \in \mathcal{S}^*$ has optimal discovery ratio in \mathcal{S}^* only if $x_i = \bar{x}_i$, for $0 \leq i \leq 4$.

Proof We will prove the theorem by induction on i . Recall that the discovery ratio of X is given by Equation (9).

We first show the base case, namely $i = 0$. The base case holds by the argument used in the proof of Theorem 5 which shows that if $x_0 < 4$, then $\text{dr}(X, \mathcal{S}^*) \geq 2$. For the induction step, suppose that X has optimal discovery ratio, and that for all $j \in [0, i]$, $x_j = \bar{x}_j$, with $i < 4$. We will show $x_{i+1} = \bar{x}_{i+1}$ by way of contradiction, hence suppose that $x_{i+1} < \bar{x}_{i+1}$. For sufficiently small $\epsilon > 0$, we have

$$\begin{aligned} m^*(x_{i+1} + x_i + \epsilon) &= m^*(x_{i+1} + \bar{x}_i + \epsilon) && \text{(by induction hypothesis)} \\ &\leq m^*(\bar{x}_{i+1} + \bar{x}_i) \\ &\text{(by monotonicity of } m^* \text{ and the definition of } \mathbf{aggressive}) \\ &= i + 1, && \text{(by definition of } m^*) \end{aligned}$$

which implies by Lemma 4, that

$$\begin{aligned} d^*(x_i + x_{i-1} + \epsilon) &= (x_i + x_{i-1} + \epsilon) \cdot \frac{6 \cdot m^*(x_{i+1} + x_i + \epsilon) + 4}{3 \cdot m^*(x_{i+1} + x_i + \epsilon) + 5} \\ &\leq (x_i + x_{i-1} + \epsilon) \cdot \frac{6 \cdot (i+1) + 4}{3 \cdot (i+1) + 5}. \end{aligned} \quad (19)$$

Therefore

$$\begin{aligned} F_{i+2}(X, \epsilon) &= \frac{2 \cdot \sum_{j=0}^{i+1} x_j + x_i + \epsilon}{d^*(x_{i+1} + x_i + \epsilon)} \\ &= \frac{2T_i(\mathbf{aggressive}) + 2x_{i+1} + \bar{x}_i + \epsilon}{d^*(x_{i+1} + \bar{x}_i + \epsilon)} && \text{(by ind. hyp.)} \\ &\geq \frac{2T_i(\mathbf{aggressive}) + 2x_{i+1} + \bar{x}_i + \epsilon}{(x_{i+1} + \bar{x}_i + \epsilon) \cdot \frac{6 \cdot (i+1) + 4}{3 \cdot (i+1) + 5}} && \text{(by (19))} \\ &= \frac{(i+1)2^{i+3} + (i+2)2^{i+1} + 2x_{i+1} + \epsilon}{(x_{i+1} + (i+2)2^{i+1} + \epsilon) \cdot \frac{6 \cdot (i+1) + 4}{3 \cdot (i+1) + 5}} && \text{(by Lemma 3)} \\ &\geq \frac{(i+1)2^{i+3} + (i+2)2^{i+1} + (i+3)2^{i+3} + \epsilon}{(i+3)2^{i+2} + (i+2)2^{i+1} + \epsilon} \cdot \frac{3i+8}{6i+10}. && \text{(monoton. on } x_{i+1}) \end{aligned}$$

Hence

$$\sup_{\delta \in (0, x_{i+2} - x_i]} F_{i+2}(X, \delta) \geq \frac{(i+1)2^{i+3} + (i+2)2^{i+1} + (i+3)2^{i+3}}{(i+3)2^{i+2} + (i+2)2^{i+1}} \cdot \frac{3i+8}{6i+10} = \frac{9i+18}{6i+10},$$

which is greater than $\frac{8}{5}$ if $i \leq 3$. We conclude, from (9), that $\text{dr}(X, \mathcal{S}^*) > 8/5$, which is a contradiction. \square

Theorem 7 leaves open the possibility that there exists some strategy $X \in \mathcal{S}^*$ with optimal discovery ratio, and for which x_5 differs from \bar{x}_5 . In particular, one may ask whether by slightly decreasing the value of x_5 , and remaining aggressive for x_i with $i \geq 6$, whether we may obtain a strategy that is in \mathcal{S}^* and has optimal discovery ratio? We will answer this question affirmatively. Formally, we define for an arbitrarily fixed constant $0 < \epsilon \leq 1$ the strategy $Y = (y_0, y_1, \dots)$, in which $y_i = \bar{x}_i$, for $0 \leq i \leq 4$, $y_5 = \bar{x}_5 - 4\epsilon$ and $y_i = 3y_{i-1} - T_{i-2}(Y)$, for $i \geq 6$.

Lemma 5 (closed expression of Y) The strategy Y satisfies $y_i = \bar{x}_i - \epsilon\bar{x}_{i-5}$, where we denote for convenience $\bar{x}_i = 0$ for all $i < 0$. It follows, $T_n(Y) = T_n(\bar{X}) - \epsilon T_{n-5}(\bar{X})$.

Proof We show this by induction on i . The base case holds for $0 \leq i \leq 5$ by definition of Y . We have for $i \geq 6$

$$\begin{aligned} y_i &= 3y_{i-1} - T_{i-2}(Y) && \text{(by definition of } y_i) \\ &= 3\bar{x}_{i-1} - 3\epsilon\bar{x}_{i-6} - T_{i-2}(\bar{X}) + \epsilon T_{i-7}(\bar{X}) && \text{(by induction hypothesis)} \\ &= 3\bar{x}_i - \epsilon\bar{x}_{i-5}. && \text{(by definition of } \bar{x}) \end{aligned}$$

□

Strategy Y is competitively optimal, since it satisfies by construction the conditions of Lemma 2. We will show that it also has the best-possible discovery ratio. This proves that there is an infinite number of strategies with optimal discovery ratio in \mathcal{S}^* .

Theorem 8 (optimal discovery ratio of Y) $dr(Y, \mathcal{S}^*) = \frac{8}{5}$.

Proof We rely on (9) to express the discovery ratio. Since Y mimics **aggressive** in the first five iterations, for $n \in [0, 4]$ we have that

$$\sup_{n \in [0, 4]} \sup_{\delta \in (0, y_n - y_{n-2})} F_n(Y, \delta) = \frac{8}{5}.$$

For $n = 5$ and $\delta \in (0, y_5 - y_3]$, we have $y_4 + y_3 + \delta \in (y_4 + y_3, y_4 + y_5]$, which implies that $m^*(y_4 + y_3 + \delta) = 5$. We obtain that

$$\begin{aligned} F_5(Y, \delta) &= \frac{2T_4(Y) + y_3 + \delta}{d^*(y_4 + y_3 + \delta)} && \text{(by Equation (9))} \\ &= \frac{2T_4(Y) + y_3 + \delta}{(y_4 + y_3 + \delta) \frac{6m^*(y_4 + y_3 + \delta) + 4}{3m^*(y_4 + y_3 + \delta) + 5}} && \text{(by Lemma 4)} \\ &= \frac{2T_4(Y) + y_3 + \delta}{(y_4 + y_3 + \delta)} \cdot \frac{20}{34} && \text{(Subst. of } m^*(x'_4 + x'_3 + \delta)) \\ &= \frac{2 \cdot (4 + 1)2^{4+2} + (3 + 2)2^{3+1} + \delta}{(4 + 2)2^{4+1} + (3 + 2)2^{3+1} + \delta} \cdot \frac{20}{34} && \text{(by Lemma 5)} \end{aligned}$$

$$\begin{aligned}
&= \frac{720 + \delta}{272 + \delta} \cdot \frac{20}{34} && \text{(arranging terms)} \\
&\leq \frac{720 \cdot 20}{272 \cdot 34} && \text{(monotonicity on } \delta) \\
&= \frac{450}{289} < \frac{8}{5}.
\end{aligned}$$

For $n = 6$ and $\delta \in (0, y_6 - y_4]$, we have $y_5 + y_4 + \delta \in (y_5 + y_4, y_6 + y_5]$, which implies that $m^*(y_5 + y_4 + \delta) \in [5, 6]$. Then

$$\begin{aligned}
F_6(Y, \delta) &= \frac{2T_5(Y) + y_4 + \delta}{d^*(y_5 + y_4 + \delta)} && \text{(by Equation (9))} \\
&= \frac{2T_5(Y) + y_4 + \delta}{(y_5 + y_4 + \delta) \frac{6m^*(y_5 + y_4 + \delta) + 4}{3m^*(y_5 + y_4 + \delta) + 5}} && \text{(by Lemma 4)} \\
&\leq \frac{2T_5(Y) + y_4 + \delta}{(y_5 + y_4 + \delta)} \cdot \frac{3 \cdot 5 + 5}{6 \cdot 5 + 4} && \text{(by } m^*(x'_{n-1} + x'_{n-2} + \delta) \geq 5) \\
&= \frac{2 \cdot (5 + 1)2^{5+2} - 8\epsilon + (4 + 2)2^{4+1} + \delta}{(5 + 2)2^{5+1} - 4\epsilon + (4 + 2)2^{4+1} + \delta} \cdot \frac{20}{34} && \text{(by Lemma 5)} \\
&= \frac{1728 - 8\epsilon + \delta}{640 - 4\epsilon + \delta} \cdot \frac{20}{34} && \text{(arranging terms)} \\
&\leq \frac{1720 \cdot 20}{636 \cdot 34} && \text{(monotonicity on } \delta \text{ and } \epsilon) \\
&= \frac{34400}{21624} < \frac{8}{5}.
\end{aligned}$$

For $n \geq 7$ and $\delta \in (0, y_n - y_{n-2}]$, we have

$$\begin{aligned}
y_{n-1} + y_{n-2} + \delta &> y_{n-1} + y_{n-2} && \text{(by } \delta > 0) \\
&= \bar{x}_{n-1} - \epsilon \bar{x}_{n-6} + \bar{x}_{n-2} - \epsilon \bar{x}_{n-7} \\
&= (4 + (2n - 1))2^{n-1} + (4 + 2(n - 2))2^{n-2} \\
&\quad - \epsilon(4 + 2(n - 6))2^{n-6} - \epsilon(4 + 2(n - 7))2^{n-7} \\
&= (2n + 2)2^{n-1} + 2n2^{n-2} - \epsilon(2n - 8)2^{n-6} - \epsilon(2n - 10)2^{n-7} \\
&= (192n - 128 - \epsilon(6n + 7))2^{n-7} \\
&= (n + 1)2^n + n2^{n-1} - \epsilon(n - 4)2^{n-5} - \epsilon(n - 5)2^{n-6} \\
&= (96n + 64 - \epsilon(3n - 13))2^{n-6} \\
&\geq (93n + 77)2^{n-6} && \text{(by } \epsilon \leq 1) \\
&\geq (48n - 11)2^{n-6} && \text{(by } n \geq 7) \\
&= (3(n - 2) + 5)2^{n-2}.
\end{aligned}$$

It follows that $m^*(y_{n-1} + y_{n-2} + \delta) \geq n - 1$. Therefore

$$\begin{aligned}
F_n(Y, \delta) &= \frac{2T_{n-1}(Y) + y_{n-2} + \delta}{d^*(y_{n-1} + y_{n-2} + \delta)} && \text{(by Equation (9))} \\
&= \frac{2T_{n-1}(Y) + y_{n-2} + \delta}{(y_{n-1} + y_{n-2} + \delta) \frac{6m^*(y_{n-1} + y_{n-2} + \delta) + 4}{3m^*(y_{n-1} + y_{n-2} + \delta) + 5}} && \text{(by Lemma 4)}
\end{aligned}$$

$$\begin{aligned}
&\leq \frac{2T_{n-1}(Y) + y_{n-2} + \delta}{(y_{n-1} + y_{n-2} + \delta)} \cdot \frac{3(n-1) + 5}{6(n-1) + 4} \\
&\hspace{15em} \text{(by } m^*(y_{n-1} + y_{n-2} + \delta) \geq n-1) \\
&= \frac{2n2^{n+1} - \epsilon 2(n-4)2^{n-3} + n2^{n-1} - \epsilon(2n-5)2^{n-6} + \delta}{(96n + 64 - \epsilon(3n-13))2^{n-6} + \delta} \cdot \frac{3(n-1) + 5}{6(n-1) + 4} \\
&\hspace{15em} \text{(by Lemma 5)} \\
&= \frac{(256n + 32n - \epsilon(16n - 32 + 2n - 5))2^{n-6} + \delta}{(96n + 64 - \epsilon(3n - 13))2^{n-6} + \delta} \cdot \frac{3(n-1) + 5}{6(n-1) + 4} \\
&= \frac{288n - \epsilon(18n - 37) + \delta}{96n + 64 - \epsilon(3n - 13) + \delta} \cdot \frac{3(n-1) + 5}{6(n-1) + 4} \\
&\leq \frac{288n}{96n + 64} \cdot \frac{3(n-1) + 5}{6(n-1) + 4} \hspace{10em} \text{(by monotonicity in } \delta \text{ and } \epsilon) \\
&< \frac{8}{5}. \hspace{15em} \text{(by } n \geq 7)
\end{aligned}$$

□

4 Connections between the discovery and the bijective ratios

In this last section we establish a connection between the discovery and the bijective ratios. Bijective analysis was introduced in [7] in the context of online computation, assuming that each request is drawn from a discrete, finite set. For instance, in the paging problem (discussed in Section 1), each request belongs to the set of all pages. Let \mathcal{I}_n denote the set of all requests of size n . For a cost minimization problem Π with discrete, finite requests, let $\pi : \mathcal{I}_n \rightarrow \mathcal{I}_n$ denote a bijection over \mathcal{I}_n . Given two online algorithms A and B for Π , the bijective ratio of A against B is defined as

$$\text{br}(A, B) = \min_{\pi: \mathcal{I}_n \rightarrow \mathcal{I}_n} \sup_{\mathcal{S} \in \mathcal{I}_n} \frac{A(\mathcal{S})}{B(\pi(\mathcal{S}))}, \text{ for all } n \geq n_0,$$

where $A(\mathcal{S})$ denotes the cost of A on request sequence \mathcal{S} .

Assuming \mathcal{I}_n is finite, an equivalent definition of $\text{br}(A, B)$ is as follows. Let $A(i, n)$ denote the i -th least costly request sequence for A among request sequences in \mathcal{I}_n . Then

$$\text{br}(A, B) = \sup_n \max_i \frac{A(i, n)}{B(i, n)}.$$

Consider, in contrast, the linear search problem. Here, there is only one request: the unknown position of the hider (i.e., $n = 1$). However, the set of all requests is not only infinite, but uncountable. Thus, the above definitions do not carry over to our setting, and we need to seek alternative definitions. One possibility is to discretize the set of all requests (as in [8]). Namely, we may assume that

the hider can hide only at integral distances from the origin. Then given strategies S_1, S_2 , one could define the bijective ratio of S_1 against S_2 as $\sup_i \frac{S_1(i)}{S_2(i)}$, where $S(i)$ denotes the i -th least costly request (hider position) in strategy S .

While the latter definition may indeed be valid, it is still not a faithful representation of the continuous setting. For instance, for hiding positions “close” to the origin, the discretization adds overheads that should not be present, and skews the expressions of the ratios. For this reason, we need to adapt the definition so as to reflect the continuous nature of the problem. Specifically, note that while the concept “the cost of the i -th least costly request in S ” is not well-defined in the continuous setting, the related concept of “the cost for discovering a total length equal to ℓ in S ” is, in fact, well defined, and is precisely the value $D(S, \ell)$. We can thus define the bijective ratio of S_1 against S_2 as

$$\text{br}(S_1, S_2) = \sup_{\ell} \frac{D(S_1, \ell)}{D(S_2, \ell)},$$

which is the same as the definition of the discovery ratio (Definition 1).

5 Conclusion

In this work we addressed the issue of separating strategies for a well-studied search optimization problem, namely searching on the line. The starting point was the observation that there is an infinite number of strategies of optimal competitive ratio, and in particular a strategy that explores the line aggressively has the same optimal competitive ratio as the more conservative strategy that is based on doubling. We introduced a new measure called discovery ratio, which informally captures the eagerness of the strategy to explore new ground. We then applied synthetically both the discovery and the competitive ratio in the analysis of search strategies for linear search. Our main result established the strict separation of the two strategies, which, to our knowledge, is the first result of this type in the context of search problems. Future work should consider similar issues in more complex search environments, such as the *m-ray search* problem [10, 25], in which the environment consists of $m \geq 2$ infinite rays with a common origin.

6 Declarations

This work was partially funded by the grant ANR-19-CE48-0016 from the French National Research Agency (ANR), DIM RFSI DACM and Labex Mathématique Hadamard. This research also benefited from the support of the FMJH Program PGMO and from the support of EDF-Thales-Orange.

The authors do not have any conflicts of interest.

References

- [1] S. Alpern and S. Gal. *The theory of search games and rendezvous*. Kluwer Academic Publishers, 2003.

- [2] S. Angelopoulos, , D. Arsénio, C. Dürr, and A. López-Ortiz. Multi-processor search and scheduling problems with setup cost. *Theory of Computing Systems*, pages 1–34, 2016.
- [3] S. Angelopoulos. Further connections between contract-scheduling and ray-searching problems. In *Proceedings of the 24th International Joint Conference on Artificial Intelligence (IJCAI)*, pages 1516–1522, 2015.
- [4] S. Angelopoulos. Online search with a hint. In *Proceedings of the 12th Innovations in Theoretical Computer Science Conference (ITCS)*, pages 51:1–51:16, 2021.
- [5] S. Angelopoulos, D. Arsénio, and C. Dürr. Infinite linear programming and online searching with turn cost. *Theoretical Computer Science*, 670:11–22, 2017.
- [6] S. Angelopoulos, D. Arsénio, and S. Kamali. Competitive sequencing with noisy advice. *arXiv*, abs/2111.05281, 2021.
- [7] S. Angelopoulos, R. Dorrigiv, and A. López-Ortiz. On the separation and equivalence of paging strategies. In *Proceedings of the 18th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 229–237, 2007.
- [8] S. Angelopoulos, M. P. Renault, and P. Schweitzer. Stochastic dominance and the bijective ratio of online algorithms. *Algorithmica*, 82(5):1101–1135, 2020.
- [9] S. Angelopoulos and P. Schweitzer. Paging and list update under bijective analysis. *Journal of the ACM*, 60(2):7:1–7:18, 2013.
- [10] R. Baeza-Yates, J. Culberson, and G. Rawlins. Searching in the plane. *Information and Computation*, 106:234–244, 1993.
- [11] A. Beck. On the linear search problem. *Naval Research Logistics*, 2:221–228, 1964.
- [12] A. Beck and D.J. Newman. Yet more on the linear search problem. *Israel Journal of Mathematics*, 8(4):419–429, 1970.
- [13] R. Bellman. An optimal search problem. *SIAM Review*, 5:274, 1963.
- [14] P. Berman. *Online Algorithms: The State of the Art*, chapter Online searching and navigation, pages 232–241. Springer, 1998.
- [15] D.S. Bernstein, L. Finkelstein, and S. Zilberstein. Contract algorithms and robots on rays: unifying two scheduling problems. In *Proceedings of the 18th International Joint Conference on Artificial Intelligence (IJCAI)*,

pages 1211–1217, 2003.

- [16] P. Bose, J. De Carufel, and S. Durocher. Searching on a line: A complete characterization of the optimal solution. *Theoretical Computer Science*, 569:24–42, 2015.
- [17] J. Boyar, S. Irani, and K. S. Larsen. A comparison of performance measures for online algorithms. *Algorithmica*, 72(4):969–994, 2015.
- [18] J. Boyar, K.S. Larsen, and A. Maiti. A comparison of performance measures via online search. *Theoretical Computer Science*, 532:2–13, 2014.
- [19] A. Condon, A. Deshpande, L. Hellerstein, and N. Wu. Algorithms for distributional and adversarial pipelined filter ordering problems. *ACM Transactions on Algorithms*, 5(2):24:1–24:34, 2009.
- [20] J. Czyzowicz, K. Georgiou, E. Kranakis, D. Krizanc, L. Narayanan, J. Opatrny, and S. Shende. Search on a Line by Byzantine Robots. In *Proceedings of the 27th International Symposium on Algorithms and Computation (ISAAC 2016)*, pages 27:1–27:12, 2016.
- [21] E.D. Demaine, S.P. Fekete, and S. Gal. Online searching with turn cost. *Theoretical Computer Science*, 361:342–355, 2006.
- [22] R. Dorrigiv and A. López-Ortiz. A survey of performance measures for on-line algorithms. *SIGACT News*, 36(3):67–81, 2005.
- [23] S. Gal. A general search game. *Israel Journal of Mathematics*, 12:32–45, 1972.
- [24] S. Gal. Minimax solutions for linear search problems. *SIAM Journal on Applied Mathematics*, 27:17–30, 1974.
- [25] S. Gal. *Search Games*. Academic Press, 1980.
- [26] C. Hipke, C. Icking, R. Klein, and E. Langetepe. How to find a point in the line within a fixed distance. *Discrete Applied Mathematics*, 93:67–73, 1999.
- [27] P. Jaillet and M. Stafford. Online searching. *Operations Research*, 49:234–244, 1993.
- [28] M-Y. Kao and M.L. Littman. Algorithms for informed cows. In *Proceedings of the AAAI 1997 Workshop on Online Search*, 1997.
- [29] M-Y. Kao, J.H. Reif, and S.R. Tate. Searching in an unknown environment: an optimal randomized algorithm for the cow-path problem. *Information and Computation*, 131(1):63–80, 1996.

- [30] A. R. Karlin and E. Koutsoupias. Beyond competitive analysis. In *Beyond the Worst-Case Analysis of Algorithms*, pages 529–546. Cambridge University Press, 2020.
- [31] D. G. Kirkpatrick. Hyperbolic dovetailing. In *Proceedings of the 17th Annual European Symposium on Algorithms (ESA)*, pages 616–627, 2009.
- [32] E. Koutsoupias, C.H. Papadimitriou, and M. Yannakakis. Searching a fixed graph. In *Proceedings of the 23rd International Colloquium on Automata, Languages and Programming (ICALP)*, pages 280–289, 1996.
- [33] Andrey Kupavskii and Emo Welzl. Lower bounds for searching robots, some faulty. In *Proceedings of the 2018 ACM Symposium on Principles of Distributed Computing*, pages 447–453, 2018.
- [34] A. López-Ortiz and S. Schuierer. On-line parallel heuristics, processor scheduling and robot searching under the competitive framework. *Theoretical Computer Science*, 310(1–3):527–537, 2004.
- [35] A. McGregor, K. Onak, and R. Panigrahy. The oil searching problem. In *Proceedings of the 17th European Symposium on Algorithms (ESA)*, pages 504–515, 2009.
- [36] S. Schuierer. Lower bounds in online geometric searching. *Computational Geometry: Theory and Applications*, 18(1):37–53, 2001.