



HAL
open science

A construction of the Shephard-Todd group G_{32} through the Weyl group of type E_6

Cédric Bonnafé

► **To cite this version:**

Cédric Bonnafé. A construction of the Shephard-Todd group G_{32} through the Weyl group of type E_6 . *Portugaliae Mathematica*, In press. hal-04248841

HAL Id: hal-04248841

<https://hal.science/hal-04248841v1>

Submitted on 18 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A CONSTRUCTION OF THE SHEPHARD-TODD GROUP G_{32} THROUGH THE WEYL GROUP OF TYPE E_6

by

CÉDRIC BONNAFÉ

Abstract. — It is well-known that the quotient of the derived subgroup of the Shephard-Todd complex reflection group G_{32} (which has rank 4) by its center is isomorphic to the derived subgroup of the Weyl group of type E_6 . We show that this isomorphism can be realized through the second exterior power, and take the opportunity to propose an alternative construction of the group G_{32} .

Let G_{32} denote the complex reflection group constructed by Shephard-Todd [ShTo] and let E_6 be a Weyl group of type E_6 (which is denoted by G_{35} in the Shephard-Todd classification). Let $\mathbf{Sp}_4(\mathbb{F}_3)$ (resp. $\mathbf{SO}_5(\mathbb{F}_3)$) denote the symplectic (resp. orthogonal) group of dimension 4 (resp. 5) over the finite field with three elements \mathbb{F}_3 . Let $\Omega_5(\mathbb{F}_3)$ be the image of $\mathbf{Sp}_4(\mathbb{F}_3)$ in $\mathbf{SO}_5(\mathbb{F}_3)$ through the natural morphism $\mathbf{Sp}_4(\mathbb{F}_3) \rightarrow \mathbf{SO}_5(\mathbb{F}_3)$: this is the normal subgroup of index 2 of $\mathbf{SO}_5(\mathbb{F}_3)$. Finally, if G is a group, let $D(G)$ and $Z(G)$ denote respectively its derived subgroup and its center and, if d is a non-zero natural number, let μ_d be the group of d -th roots of unity in \mathbb{C} .

It is shown in [LeTa, théo. 8.43 et 8.54] that $G_{32} \simeq \mu_3 \times \mathbf{Sp}_4(\mathbb{F}_3)$ and that $E_6 \simeq \mathbf{SO}_5(\mathbb{F}_3)$. In particular,

$$(*) \quad D(G_{32})/\mu_2 \simeq \Omega_5(\mathbb{F}_3) \simeq D(E_6).$$

The purpose of this note is to present a direct elementary explanation of the isomorphism $D(G_{32})/\mu_2 \simeq D(E_6)$, which in fact allows us to construct the complex reflection group G_{32} from the rational reflection group E_6 . This construction uses the classical morphism $\mathbf{SL}_4(\mathbb{C}) \rightarrow \mathbf{SO}_6(\mathbb{C})$, and follows the same lines as our previous paper [Bon] (in which we constructed the complex reflection group G_{31} from the Weyl group of type B_6).

This note does not pretend to prove a deep result: it is just a nice example of the application of the classical theory of reflection groups (invariants, Springer theory,...).

Remark. It is shown in [LeTa, Theo. 8.53] that $G_{33} \simeq \mu_2 \times \Omega_5(\mathbb{F}_3)$. In particular, this gives an indirect isomorphism $D(G_{33}) \simeq D(E_6)$. However, we do not know of any construction of this isomorphism which would be in the same spirit as above.

1. The morphism $\mathbf{SL}_4(\mathbb{C}) \rightarrow \mathbf{SO}_6(\mathbb{C})$

We recall here the construction of this morphism, using some notation from [Bon]. Let us fix a complex vector space V of dimension 4 and let

$$\begin{aligned} \Lambda : \mathbf{GL}(V) &\longrightarrow \mathbf{GL}(\wedge^2 V) \\ g &\longmapsto \wedge^2 g \end{aligned}$$

be the natural morphism of algebraic groups. Note that $\wedge^2 V$ has dimension 6 and that

$$(1.1) \quad \text{Ker } \Lambda \simeq \mu_2 = \{\pm \text{Id}_V\}.$$

The lists of eigenvalues of elements of $\mathbf{GL}(V)$ or $\mathbf{GL}(\wedge^2 V)$ will always be given with multiplicities. If $g \in \mathbf{GL}(V)$ admits a, b, c and d as eigenvalues, then

$$(1.2) \quad \Lambda(g) \text{ admits } ab, ac, ad, bc, bd \text{ and } cd \text{ as eigenvalues.}$$

In particular,

$$(1.3) \quad \det \Lambda(g) = (\det g)^3.$$

Let us fix now a generator ε of the one-dimensional vector space $\wedge^4 V$. The choice of this generator allows to identify \mathbb{C} and $\wedge^4 V$ and to define a bilinear form

$$\begin{aligned} \beta_\wedge : \wedge^2 V \times \wedge^2 V &\longrightarrow \mathbb{C} \\ (x, y) &\longmapsto x \wedge y. \end{aligned}$$

This bilinear form is symmetric and non-degenerate. By definition of the determinant,

$$(1.4) \quad \beta_\wedge(\Lambda(g)(x), \Lambda(g)(y)) = (\det g)\beta_\wedge(x, y)$$

for all $g \in \mathbf{GL}(V)$ and $x, y \in \wedge^2 V$. For dimension and connectedness reasons, the image of $\mathbf{GL}(V)$ through Λ is the neutral component $\mathbf{CO}(\wedge^2 V)^\circ$ of the conformal orthogonal group $\mathbf{CO}(\wedge^2 V) = \mathbf{CO}(\wedge^2 V, \beta_\wedge)$ and Λ induces an isomorphism of algebraic groups

$$(1.5) \quad \mathbf{SL}(V)/\mu_2 \simeq \mathbf{SO}(\wedge^2 V).$$

The next elementary lemma will be useful (in this paper, we denote by j a primitive third root of unity).

Lemma 1.6. — *Let $g \in \mathbf{SO}(\wedge^2 V)$ having j, j, j, j^2, j^2 and j^2 as eigenvalues. Then there exists a unique element $\tilde{g} \in \mathbf{SL}(V)$, having $1, j, j$ and j as eigenvalues, and such that*

$$\Lambda^{-1}(\{g, g^{-1}\}) = \{\pm \tilde{g}, \pm \tilde{g}^{-1}\}.$$

Proof. — Let $h \in \mathbf{SL}(V)$ be such that $\Lambda(h) = g$. Then $\Lambda^{-1}(\{g, g^{-1}\}) = \{\pm h, \pm h^{-1}\}$. Let a, b, c and d be the eigenvalues of h . By (1.2), by reordering if necessary the eigenvalues of h , we may assume that $ab = ac = j$. In particular, $b = c$. So $bd = cd$, which implies that $bd = cd = j^2$ (for otherwise j would be an eigenvalue of g with multiplicity ≥ 4). Still by (1.2), $(ad, bc) = (j, j^2)$ or $(ad, bc) = (j^2, j)$.

If $(ad, bc) = (j, j^2)$, then $b = c = d$ and $b^2 = j^2$, so $b = c = d = \eta j$ and $a = \eta$, for some $\eta \in \{\pm 1\}$. By replacing h by $-h$ if necessary, the eigenvalues of h are then $1, j, j, j$ and so h is indeed the unique element of $\Lambda^{-1}(\{g, g^{-1}\})$ admitting this list of eigenvalues.

If $(ad, bc) = (j^2, j)$, then $d = jb = jc$ and $b^2 = j$, which implies that $a = b = c = \eta j^2$ and $d = \eta$, for some $\eta \in \{\pm 1\}$. By replacing h by $-h$ if necessary, the eigenvalues of h are then $1, j^2, j^2, j^2$ and so h^{-1} is indeed the unique element of $\Lambda^{-1}(\{g, g^{-1}\})$ admitting $1, j, j, j$ as list of eigenvalues. \square

Corollary 1.7. — *Let $g \in \mathbf{SO}(\wedge^2 V)$ having j, j, j, j^2, j^2 and j^2 as eigenvalues. Then g and g^{-1} are not conjugate in $\mathbf{SO}(\wedge^2 V)$.*

Proof. — Assume that we have found $w \in \mathbf{SO}(\wedge^2 V)$ such that $wgw^{-1} = g^{-1}$. Let ω and γ be respective preimages of w and g in $\mathbf{SL}(V)$. Then $\Lambda(\omega\gamma\omega^{-1}) = g^{-1} = \Lambda(\gamma^{-1})$. This shows that γ^{-1} is conjugate to γ or to $-\gamma$, which is impossible by examining the possible lists of eigenvalues of γ obtained in the proof of Lemma 1.6. \square

2. Construction of G_{32}

Let us see E_6 as a finite subgroup of $\mathbf{O}(\wedge^2 V)$ generated by reflections. Set

$$\mathbf{jSO}(\wedge^2 V) = \langle \mathbf{SO}(\wedge^2 V), j \text{Id}_{\wedge^2 V} \rangle$$

and $\sqrt[3]{\mathbf{SL}}(V) = \{g \in \mathbf{GL}(V) \mid \det(g)^3 = 1\} = \langle \mathbf{SL}(V), j \text{Id}_V \rangle$.

Then

$$(2.1) \quad \Lambda^{-1}(\mathbf{jSO}(\wedge^2 V)) = \sqrt[3]{\mathbf{SL}}(V).$$

We then define

$$W = \Lambda^{-1}(\langle \mathbf{D}(E_6), j \text{Id}_{\wedge^2 V} \rangle).$$

It is a subgroup of $\sqrt[3]{\mathbf{SL}}(V)$. The aim of this note is to show that W is isomorphic to the complex reflection group G_{32} of Shephard-Todd. Note first that

$$(2.2) \quad \mu_6 \subset W.$$

2.A. Reflections in W . — The list of degrees E_6 is 2, 5, 6, 8, 9, 12 while its list of codegrees is 0, 3, 4, 6, 7, 10 (see [Bro, Table A.3]). In particular, exactly 3 of the degrees are divisible by 3, which shows [Spr, Theo. 3.4] that E_6 contains an element w_3 admitting the eigenvalue j with multiplicity 3. We will denote by C_3 the conjugacy class of w_3 in W . Since also exactly 3 of the codegrees are divisible by 3, this implies, for instance by [LeMi, Theo. 1.2], that w_3 is regular in the sense of Springer [Spr, §4] (that is, admits an eigenvector for the eigenvalue j whose stabilizer in E_6 is trivial). Since E_6 is a rational group, w_3 also admits j^2 as an eigenvalue with multiplicity 3. Hence, the eigenvalues of w_3 are j, j, j, j^2, j^2 and j^2 . In particular, $\det w_3 = 1$ and so

$$w_3 \in \mathbf{D}(E_6) = E_6 \cap \mathbf{SO}(\wedge^2 V).$$

By [Spr, Theo. 4.2(iii)], the centralizer of w_3 in E_6 has order $6 \cdot 9 \cdot 12$, which shows that

$$(2.3) \quad |C_3| = 2 \cdot 5 \cdot 8 = 80.$$

Moreover [Spr, Theo. 4.2(iii)], if $w \in E_6$, then

$$(2.4) \quad w \in C_3 \text{ if and only if } \dim \text{Ker}(w - j \text{Id}_V) = 3.$$

Hence, if $w \in C_3$, then $w^{-1} \in C_3$ but w^{-1} is not conjugate to w in $\mathbf{D}(E_6)$ (by Corollary 1.7).

Now, let $\text{Ref}(W)$ denote the set of reflections of W . If $s \in \text{Ref}(W)$, the eigenvalues $\Lambda(s)$ are $1, 1, 1, \det(s), \det(s), \det(s)$. Since $\det(s) \in \{j, j^2\}$, the eigenvalues of $\det(s)\Lambda(s)$ are then j, j, j, j^2, j^2, j^2 . It follows from (2.4) that $\det(s)\Lambda(s) \in C_3$. This defines a map

$$\begin{aligned} \lambda : \text{Ref}(W) &\longrightarrow C_3 \\ s &\longmapsto \det(s)\Lambda(s). \end{aligned}$$

The next result will be very useful:

Lemma 2.5. — *The map λ is bijective.*

Proof. — First, if $\lambda(s) = \lambda(s')$, then there exists $\xi \in \mathbb{C}^\times$ such that $s' = \xi s$. Since s and s' are reflections, this is possible only if $\xi = 1$, and so $s = s'$. This shows that λ is injective.

Let us now show the surjectivity. Let $w \in C_3$. By Corollary 1.7, there exists a unique $\tilde{w} \in \Lambda^{-1}(\{w, w^{-1}\})$ admitting $1, j, j, j$ as eigenvalues. Then $j^2\tilde{w}$ and $j\tilde{w}^{-1}$ are reflections satisfying

$$\lambda(j^2\tilde{w}) = \det(j^2\tilde{w})\Lambda(j^2\tilde{w}) = j^8 \cdot j^4\Lambda(\tilde{w}) = \Lambda(\tilde{w})$$

and
$$\lambda(j\tilde{w}^{-1}) = \det(j\tilde{w}^{-1})\Lambda(j\tilde{w}^{-1}) = j^4 \cdot j^2\Lambda(\tilde{w}^{-1}) = \Lambda(\tilde{w})^{-1}.$$

So $w \in \{\lambda(j^2\tilde{w}), \lambda(j\tilde{w}^{-1})\}$, which shows that λ is surjective. \square

We then deduce from (2.3) and Lemma 2.5 that

$$(2.6) \quad |\text{Ref}(W)| = 80.$$

2.B. Structure of W . — Our main result is Theorem 2.7 below: the proof we propose here uses neither known properties of the group G_{32} nor the classification of complex reflection groups and so might be viewed as an alternative construction of G_{32} starting from E_6 (however, note that we use properties of E_6).

Theorem 2.7. — *The group W :*

- (a) *has order 155 520;*
- (b) *is generated by reflections of order 3;*
- (c) *is irreducible and primitive;*
- (d) *admits 12, 18, 24, 30 as list of degrees.*

Proof. — Set $E_6^\# = \langle D(E_6), j \text{Id}_{\wedge^2 V} \rangle$ and $W^+ = W \cap \mathbf{SL}(V)$. Recall that $|E_6| = 51\,840$. So

$$(2.8) \quad |D(E_6)| = 25\,920, \quad |E_6^\#| = 77\,760, \quad |W| = 155\,520 \quad \text{and} \quad |W^+| = 51\,840.$$

This shows (a). Moreover,

$$(2.9) \quad Z(W) = \mu_6 \quad \text{and} \quad W/\mu_6 \simeq D(E_6).$$

Let $\mathcal{R} = \{\det(s)^{-1}s \mid s \in \text{Ref}(W)\}$. Set

$$G = \langle \text{Ref}(W) \rangle \quad \text{and} \quad H = \langle \mathcal{R} \rangle.$$

The statement (b) is equivalent to the following one

$$(\#) \quad W = G.$$

First, $\Lambda(\mathcal{R}) = C_3$ and so $\Lambda(H) = D(E_6)$ (as this last group is simple and C_3 is a conjugacy class). In particular, $W = H \cdot \mu_6$ and so, since $H \subset G \cdot \mu_3$, we obtain

$$W = G \cdot \mu_6,$$

which is almost the expected result (#). Before showing (#), note that, since $\Lambda(H) = D(E_6)$, we have that H (and so G) acts irreducibly on V . Moreover, if G is not primitive, then G (and so H) would be monomial [LeTa, Lem. 2.12], which would imply that $\Lambda(H) = D(E_6)$ is monomial, which is false. So

(♣) G is irreducible and primitive.

Statement (c) follows.

Let us conclude this proof by showing simultaneously (b) et (d). Let d_1, d_2, d_3, d_4 be the degrees of G . It follows from (2.6) and for instance from [Bro, Theo. 4.1] that

$$(\diamond) \quad \begin{cases} d_1 d_2 d_3 d_4 = |G|, \\ d_1 + d_2 + d_3 + d_4 = 84. \end{cases}$$

The morphism $\det : G \rightarrow \mu_3$ is surjective (since $\det(s) \in \{j, j^2\}$ for any $s \in \text{Ref}(W)$), and this implies that $\mu_3 \subset G$ (because $G/(G \cap \mu_6) \simeq D(E_6)$ is simple). It remains to show that $|G| \neq |W|/2 = 77\,760$.

So assume that $|G| = 77\,760$. Since $\mu_3 \subset G$, all the d_i 's are divisible by 3 and, since $\mu_6 \not\subset G$, at least one of them (say d_4) is not divisible by 6. Write $e_i = d_i/3$. Then

$$\begin{cases} e_1 e_2 e_3 e_4 = 960 = 2^6 \cdot 3 \cdot 5, \\ e_1 + e_2 + e_3 + e_4 = 28, \\ e_4 \text{ is odd.} \end{cases}$$

From the second equality, we deduce that at least one of the e_i 's (say e_3) is odd. From the first equality, we deduce that at least one of the e_i 's (say e_2) is even and so e_1 is also even. The first equality shows that e_1 or e_2 (say e_2) is divisible by 8. So $e_2 \in \{8, 16, 24\}$. A quick inspection of the possibilities shows that $e_1 = 4, e_2 = 16$ and $\{e_3, e_4\} = \{3, 5\}$. The degrees of G are then 9, 12, 15 and 48. Since 16 divides one of the degrees, it then follows from [Spr, théo. 3.4(i)] that $G = H \times \mu_3$ contains an element of order 16 and so H contains an element of order 16. Therefore, $D(E_6) = \Lambda(H)$ contains an element of order 8, which is impossible (see Remark 2.11 below for a proof of this fact based only on Springer theory). This contradicts the fact that $|G| = 77\,760$. So we have shown (#), that is,

$$W = \langle \text{Ref}(W) \rangle = G.$$

This is statement (b). In particular, $\mu_6 \subset W$ and so all the d_i 's are divisible by 6. Set $a_i = d_i/6$. Then (\diamond) implies that

$$\begin{cases} a_1 a_2 a_3 a_4 = 120 = 2^3 \cdot 3 \cdot 5, \\ a_1 + a_2 + a_3 + a_4 = 14, \end{cases}$$

By the same argument as before, since $\mu_{12} \not\subset W$, we may assume that a_3 and a_4 are odd and that a_1 and a_2 are even. A quick inspection of the possibilities shows that $\{a_1, a_2\} = \{2, 4\}$ and $\{e_3, e_4\} = \{3, 5\}$. This concludes the proof of (d). \square

Corollary 2.10. — *The group W is isomorphic to the reflection group G_{32} of Shephard-Todd.*

Proof. — This follows from Theorem 2.7 and from the classification of complex reflection groups [ShTo]. \square

Hence, Corollary 2.10 gives an explanation for the fact, mentioned in the introduction, that $D(G_{32})/\mu_2 \simeq D(E_6)$: the isomorphism is realized by Λ .

Remarque 2.11. — In the proof of Theorem 2.7, we have used the fact that $D(E_6)$ does not contain any element of order 8. This fact can be easily obtained by a computer calculation for instance, but we propose here a proof using only Springer theory. Let $w \in E_6$ be an element of order 8. Then w necessarily admits an eigenvalue which is a primitive 8-th root of unity ζ . As only one of the degrees of E_6 and only one of the codegrees of E_6 is divisible by 8, this implies that w is a regular element in the sense of Springer [LeMi, Theo. 1.2]. Then [Spr, Theo. 4.2(v)] the list of eigenvalues of w is $\zeta^{-1}, \zeta^{-4}, \zeta^{-5}, \zeta^{-7}, \zeta^{-8}, \zeta^{-11}$, and so $\det(w) = \zeta^{-36} = -1$. So $w \notin D(E_6)$.

References

- [Bon] C. BONNAFÉ, *Une (nouvelle?) construction du groupe de réflexions complexe G_{31}* , Bull. Math. Soc. Sci. Math. Roumanie **45** (2003), 133-143.
- [Bro] M. BROUÉ, *Introduction to complex reflection groups and their braid groups*, Lecture Notes in Math. **1988**, Springer-Verlag, Berlin, 2010. xii+138 pp.
- [LeMi] G. I. LEHRER & J. MICHEL, *Invariant theory and eigenspaces for unitary reflection groups*, Comptes Rendus Acad. Sci. **336** (2003), 795-800.
- [LeTa] G. I. LEHRER & D. E. TAYLOR, *Unitary reflection groups*, Australian Mathematical Society Lecture Series **20**, Cambridge University Press, Cambridge, 2009.
- [Spr] T.A. SPRINGER, *Regular elements of finite reflection groups*, Invent. Math. **25** (1974), 159-198.
- [ShTo] G. C. SHEPHARD & J. A. TODD, *Finite unitary reflection groups*, Canad. J. Math. **6** (1954), 274-304.

October 18, 2023

CÉDRIC BONNAFÉ, IMAG, Université de Montpellier, CNRS, Montpellier, France
E-mail : cedric.bonnafe@umontpellier.fr