



HAL
open science

Discriminating between tsunamis and tropical cyclones in the sedimentary record using X-ray tomography

Maude Biguenet, Éric Chaumillon, Pierre Sabatier, Raphael Paris, Pierre
Vacher, Nathalie Feuillet

► **To cite this version:**

Maude Biguenet, Éric Chaumillon, Pierre Sabatier, Raphael Paris, Pierre Vacher, et al.. Discriminating between tsunamis and tropical cyclones in the sedimentary record using X-ray tomography. *Marine Geology*, 2022, 450, pp.106864. 10.1016/j.margeo.2022.106864 . hal-04254917

HAL Id: hal-04254917

<https://hal.science/hal-04254917v1>

Submitted on 24 Jan 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Discriminating between tsunamis and tropical cyclones in the sedimentary record using X-ray tomography

M. Biguenet^{ab}, E. Chaumillon^a, P. Sabatier^b, R. Paris^c, P. Vacher^d, N. Feuillet^e.

^a Université de la Rochelle, UMR 7266 LIENSs, 17042 La Rochelle cedex, France.

^b Université Savoie Mont Blanc, CNRS, UMR 5204 EDYTEM, 73376 Le Bourget du Lac cedex, France.

^c Université Clermont Auvergne, CNRS, IRD, OPGC, Laboratoire Magmas et Volcans, F-63000 Clermont-Ferrand, France.

^d Laboratoire SYMME, domaine Universitaire BP 80439, 74944 Annecy le Vieux Cedex, France.

^e Institut de Physique du Globe de Paris, Université de Paris, CNRS, 75005 Paris, France.

1 Abstract

2 Scrub Island lagoon (Anguilla, Caribbean) provides the opportunity to display both tropical cyclones,
3 near-field and far-field tsunami deposits within the same sediment core. Here, we use X-ray
4 tomography to highlight the sedimentary fabric and link it to processes that occurred during these
5 events. Overall, the tsunami sedimentary fabric remains highly homogenous, and display fabric
6 trends that can be associated with variations in sediment concentration and /or flow strength. The
7 near-field tsunami deposit displays fabric characteristics of intense and highly-concentrated flow,
8 whereas the far-field tsunami fabric is less defined which may illustrate a less intense event.
9 Contrastingly, the sediment fabric of the tropical cyclone (hurricane) deposit is more variable with a
10 succession of different dipping-angles and orientations of the grains that can be interpreted as the
11 record of successive waves, likely infragravity waves. Observed differences in sedimentary fabrics
12 between tsunami and tropical cyclone-related deposits are an encouragement to generalize
13 sediment fabric parameters as criteria to distinguish them in coastal sedimentary archives.

14

15 **Keywords:** tsunamis, tropical cyclones, Lesser Antilles, lagoons, X-ray tomography

16

17 **1. Introduction**

18 Geological records constitute key-archives to reconstruct long-term series of marine flooding to
19 extend our knowledge well beyond the observational and historical records. They allow defining
20 return periods of tsunami induced by large earthquake (e.g. Minoura et al., 2001; Nanayama et al.,
21 2003; Monecke et al., 2008) or tropical cyclone/storm activity related to climate variability (e.g. Liu
22 and Fearn, 1993, 2000; Donnelly and Woodruff, 2007; Sabatier et al., 2012; Goslin and Clemmensen,
23 2017; Wallace et al., 2021). Therefore, it is crucial to differentiate tropical cyclones and tsunamis-
24 related deposits in sedimentary archives in order to better reconstruct return time periods of those
25 dramatic events in densely populated coastal areas. Unfortunately, there are still no unequivocal
26 methods for differentiating between tsunami- and tropical cyclone- generated deposits. Moreover,
27 criteria to distinguish those deposits seem to be site dependent and there are few locations where
28 both types of deposits were clearly identified (e.g. Nanayama et al., 2000; Goff et al., 2004; Tuttle et
29 al., 2004; Kortekaas and Dawson, 2007; Biguenet et al., 2021). The most widely recognized
30 sedimentary signatures for tsunami deposits are sequences of normal (fining-up) or inversely
31 (coarsening-up) graded beds, nongraded (massive) or multiple graded sandsheets, internal mud
32 laminae, intraclasts (rip-up clasts), truncated flame structures, or backflow induced structures and
33 erosional lower contact (e.g. Nanayama et al., 2000; Goff et al., 2004; Tuttle et al., 2004; Kortekaas
34 and Dawson, 2007; Biguenet et al., 2021). Storm deposits tend to have many parallel, cross
35 laminations and climbing ripples (e.g., Morton et al., 2007; Baumann, 2017), with a highly variable
36 grain-size distribution and a sharp, non-erosional lower contact (e.g., Goff et al., 2004; Soria et al.,
37 2017; 2018). Here, we use X-ray tomography on previously identified tropical cyclone (hurricane) and
38 tsunami deposits recorded in a coastal lagoon located in Scrub Island (Anguilla, Biguenet et al., 2021)
39 to present new criteria based on the sedimentary fabric. X-ray tomography enables characterization
40 of the three-dimensional structure and texture of rocks and sediments (Cnudde and Boone, 2013).
41 Thus, an increasing number of sedimentological studies use X-ray tomography to observe internal

42 sedimentary structures (e.g. Renter, 1989; Orsi et al., 1994; Flisch and Becker, 2003). Among these
43 studies, the use of X-ray tomography to analyze tsunamis and or tropical cyclones deposits is
44 relatively new (e.g. Cuven et al., 2013; May et al., 2016; Falvard and Paris, 2017; Paris et al., 2020).
45 We propose in this study new diagnostic parameters based on X-ray tomography using a method
46 developed by Falvard and Paris (2017) and Paris et al. (2020) to identify tropical cyclone, near-field
47 and far field tsunamis deposits in coastal archives.

48

49 **2. Study area and available data**

50 Scrub Island is located in the northern part of the Lesser Antilles Arc, southeast of the Anegada
51 Passage (Fig. 1AB). This island is 1.5 km wide from north to south and about 3.5 km long from east to
52 west (Fig. 1C). It consists of low-relief limestones (highest point about 25 m above mean sea-level,
53 a.s.l.). In this study, we focus on a 400 m-long and 130 m-wide very shallow (maximum water depth
54 of 80 cm) closed lagoon, located along the western coast of the island (Fig. 1D). This saltwater lagoon
55 is separated from the ocean by a 2 to 5 m-high and about 50 m-wide sandy barrier, which is covered
56 by small bushes at the top. A 110 cm-long sediment core (SCR1-18-01) was collected in March 2018
57 in this lagoon using an Uwitec gravity corer with hammer. Twenty-five sandy layers resulting from
58 high-energy-marine floods were identified after sedimentological, geochemical, microfaunal and
59 chronological analyses. Two of these layers are interpreted as historical tsunami deposits and the 23
60 remaining sandy layers were interpreted as tropical cyclone (hurricane) deposits, with the three most
61 recent layers being correlated with historical tropical cyclones (Biguenet et al., 2021).

62 In order to compare tsunami and tropical cyclones sedimentary signature with X-ray tomography, we
63 selected two tsunami deposits and one tropical cyclone deposit in the same core SCR1-18-01. Those
64 deposits are all composed of fine-to-coarse bioclastic sand (mean grain-size of samples between 122
65 and 765 μm) interstratified with the silty organic-rich lagoonal sediment-fill. A short description of
66 those deposits is given below (for further details, please refer to Biguenet et al., 2021). The most
67 recent tsunami deposit is dated between 1652 and 1810 cal. CE. It is interpreted to result from the

68 Lisbon transoceanic tsunami that occurred in 1755 CE and also observed and identified in the
69 sedimentary record of a large part of the Lesser Antilles (Engel et al., 2016; NGDC/WDS, 2021; Paris
70 et al., 2021). This sediment record consists of a 2 cm-thick very poorly to poorly sorted sandy layer,
71 with a rip-up clast and topped by a 1.9 cm-thick mud layer with a large amount of wood and plant
72 remains (33.9 to 30 cm-depth in core). The older tsunami deposit is dated between 1364 and 1469
73 cal. CE and is interpreted to record the Pre-Columbian event also identified in St-Thomas (US Virgin
74 Islands, Fig 1A; Fuentes et al., 2017) and in Anegada (British Virgin Islands, Fig 1A; Atwater et al.,
75 2017). This sediment record is a 16.2 cm thick massive poorly to well-sorted sandy bed without any
76 major grain-size variations (63 to 46.8 cm-depth in core). Two internal mud laminae were found at its
77 base. The selected tropical cyclone deposit is interpreted as recording the Category 3 Hurricane
78 Donna in 1960 CE (NOAA, 2020, Historical Hurricane Tracks). This sediment record corresponds to a
79 4.75 cm-thick, poorly sorted, medium to coarse sandy deposit (5 to 0.25 cm-depth in core). We
80 choose to focus on the Hurricane Donna deposit because it is the thickest and it was more accurately
81 dated than the other tropical cyclone deposits found in SCR1-18-01 core.

82

83 **3. Material and methods**

84 The three event deposits were scanned at the SYMME laboratory (Annecy, France) using an X-ray
85 tomography system (Easytom XL 150 from RX SOLUTION company) with a 50 μm resolution (voxel
86 size). After reconstruction of the original projections, the deposits are imaged as stacks of horizontal
87 cross-sections along the core (16 bits Tiff image files). The tsunami deposit dated between 1652 and
88 1810 cal. CE. that we interpret to be formed by the Lisbon tsunami (Biguenet et al., 2021) (299 images
89 from base to top) and Donna deposit (757 images from base to top) were scanned at 150 kV and 70
90 μa , whereas the Pre-Columbian tsunami deposit (2360 images from base to top) was scanned at 150
91 Kv and 200 μa . These parameters were adapted for each deposit to obtain the best image quality.
92 The reconstruction of the sedimentary fabric in 3D and associated numerical values was made
93 following the method of Falvard and Paris (2017), later updated in Paris et al. (2020). The sand grains

94 were semi-automatically separated using the morphological segmentation tool from MorpholibJ
95 collection (Legland et al., 2016) included in the Fiji distribution of ImageJ (Schindelin et al., 2012). All
96 sand grains were then counted and characterised in terms of size, shape, orientation, and dipping-
97 angle, using the Blob3D program (Ketcham, 2005). Sand grains with a measured maximum axis
98 length shorter than 250 μm were excluded to remove artifacts and avoid potential over-
99 segmentation of larger grains. Furthermore, sand grains with an aspect ratio (longest axis / smallest
100 axis of the ellipsoid) higher than 3 were also excluded to avoid overly elongated particles resulting
101 from artifacts. Resolution and image quality allowed sedimentary fabric to be characterized (spatial
102 arrangement of the grains) at 5 mm intervals down core. In every 5-mm subunit, the sedimentary
103 fabric was characterized for three different grain-size fractions (medium sand 250-500 μm ; coarse
104 sand 500-1000 μm ; very coarse sand > 1000 μm). Projections of the orientations (azimuth) and
105 dipping-angle (plunge) of the longest grain axes (sedimentary fabric data) were plotted on
106 stereographic lower hemisphere equal-area projection, using StereoNet. The eigenvalues (the
107 normalized vector magnitudes) S1 (longest axis), S2 (intermediate axis) and S3 (shortest axis) of the
108 ellipsoid fabric were also calculated for the three grain fractions in every subunit of 5 mm. This
109 ellipsoid corresponds to the preferential orientations of the particles extracted from the sample and
110 they were also represented in equilateral ternary diagrams (Benn, 1994). These diagrams describe
111 sedimentary fabrics based on fabric isotropy ($I = S3/S1$) and elongation ($E = 1-(S2/S1)$) of each
112 sample. The three poles of the diagram correspond to an isotropic fabric ($S1 = S2 = S3$; $I = 1$ and $E =$
113 0), a planar girdle-shaped fabric ($S1 = S2 > S3$; $I = E = 0$, oblate ellipsoid), and a cluster shaped fabric
114 ($S1 > S2 > S3$; $I = 0$, $E = 1$, a prolate ellipsoid). Finally, the detailed grain-size parameters of each
115 deposit previously determined in Biguenet et al. (2021), were described here to enable a precise
116 comparison with the sedimentary fabric. The grain-size parameters result from the grain-size
117 analyses that were carried out at the EDYTEM laboratory using a Beckman Coulter LS 13320 XR
118 particle size analyser using sonication, with a range size between 0.010 μm and 3000 μm . The
119 sampling step was approximately 5 mm through core SCR1-18-01 ($n = 126$) respecting facies

120 boundaries. The mean grain size curve and the grain size contour plot were produced using MATLAB
121 software (Biguenet et al., 2021).

122

123 **4. Results**

124 The use of high-resolution X-ray tomography allowed us to identify more sand layers than by eye and
125 thus revise the boundaries of the selected tsunami and tropical cyclone-related layers to obtain more
126 precise deposits delimitations. The sandy part of the deposit interpreted as being formed by the
127 Lisbon Tsunami has been restricted to extend from 33.25 to 31.75 cm (Fig. 2). Thin sand layers
128 alternating with mud at the base of the Pre-Columbian deposit have been interpreted as recording
129 older events. Thus, the Pre-Columbian deposit is restricted to extend from 60.5 to 47.25 cm (Fig. 2).
130 The sand layer (5-3.5 cm, Fig.2) separated from the base of Hurricane Donna deposit by a mud layer
131 has been interpreted as recording and older tropical cyclone event. Thus, Hurricane Donna deposit is
132 restricted to extend from 3.5 to 0.25 cm (Fig. 2). X-ray tomography is hence a promising method to
133 precisely define facies boundaries at a high resolution.

134

135 The study of sedimentary fabric from the three deposits has led us to use the same criteria
136 distinguishing sedimentary fabrics as Paris et al. (2020), to facilitate the comparisons. Thus, all the
137 studied stereograms were listed in the four categories defined by Paris et al. (2020) (Fig. 2).

- 138 – Fabric A: moderate-dipping-angle (15 to 35°) to high-dipping-angle (>35°) fabric with one
139 well-defined orientation, typically flow-parallel.
- 140 – Fabric B: bimodal low-dipping-angle (<15°) fabric (two orientations with opposite dipping-
141 angles), either flow parallel or flow oblique.
- 142 – Fabric C: low-to-moderate dipping-angle fabric with at least two different orientations.
- 143 – Fabric D: dispersed fabric.

144

145 **4.1 Pre-Columbian Tsunami**

146 Fourteen grain-size samples were collected in this deposit from 60.25 to 47.75 cm, a 1-cm interval
147 (Fig. 2). The top of the deposit (56.75 to 47.75 cm) is a unimodal coarse sand with mode at 718 μm .
148 The sand is well sorted and displays symmetrical skewness from 56.75 to 51.75 cm, and is
149 moderately well sorted and finely skewed from 51.75 to 47.75 cm. The base of the deposit (60.25 to
150 56.75 cm) displays unimodal to trimodal, poorly sorted, mostly very fine skewed, fine to coarse sand.
151 The major modes are at 718 μm (coarse sand), 122 μm (fine sand), 596 μm (coarse sand), 134 μm
152 (fine sand), and 53 μm (coarse silt). The 53 μm and 596 μm modes are only present in the sample at
153 60.25 cm, the base of the deposit.

154 As seen on the stereographic projections (Fig. 2), the Pre-Columbian tsunami deposit displays a
155 bimodal low—dipping-angle (mean ellipsoid plunge $<23^\circ$ and often $<10^\circ$) fabric (Fabric B) with two
156 well-defined orientations of opposite dipping-angles through almost the entire deposit and in the
157 three sand fractions (Fig. 2). However, the base of this deposit (60.5 to 59 cm) displays a progressive
158 transition from a poorly- to well-defined fabric, transitional between a dispersed fabric (FD) and
159 bimodal fabric (FB) with a slight variation of orientation (50° rotation), especially in the medium sand
160 (MS) and coarse sand (CS) fractions (Fig. 2).

161 The mean ellipsoid plunge varies between 0.2° and 22.4° (Fig. 3). The mean ellipsoid plunge is 8.7° for
162 the MS fraction, 5.8° for the CS fraction and 4.3° for the VCS fraction. Thus, the MS fraction displays
163 the highest plunges. The ellipsoid degree of elongation of this tsunami deposit is typically larger than
164 0.15 (Fig. 3 & 4), with a broad range of values from 0.07 to 0.53. Compared to the other deposits, the
165 Pre-Columbian tsunami displays a relatively higher degree of elongation. The mean ellipsoid
166 elongation is 0.17 (low elongation) for the MS fraction, 0.25 (low to moderate elongation) for the CS
167 fraction and 0.38 (moderate elongation) for the VCS fraction. Thus, the degree of elongation is
168 highest for the VCS fraction.

169 The ellipsoid degree of isotropy for this deposit ranges between 0.24 and 0.78 (Fig. 4). The mean
170 ellipsoid isotropy is 0.74 (medium to high isotropy) for the medium sand fraction (MS), 0.55 (medium
171 isotropy) for the coarse sand fraction (CS) and 0.36 (moderate isotropy) for the very coarse sand

172 fraction (VCS; Fig. 4). Thus, the degree of isotropy is highest for the MS fraction and lowest for the
173 VCS fraction.

174 For each grain-size fraction, there is no major variation of isotropy and elongation degree, and mean
175 plunge from the base to the top of the deposit (Standard deviation inferior to 0.1 for both isotropy
176 and elongation degree, and inferior to 6° for the mean plunge).

177

178 **4.2 Lisbon Tsunami**

179 Three grain-size samples were analysed for the deposit interpreted as being formed by the Lisbon
180 Tsunami (33.05 to 31.9 cm; Fig. 2). The grain-size sample at 33.05 cm is a trimodal, very poorly sorted
181 symmetrical fine sand. The sample has a mode at 718 µm (coarse sand), a second at 122 µm (fine
182 sand), and a third at 53 µm (coarse silt). The grain-size sample at 32.55 cm shows bimodal, poorly
183 sorted, very fine skewed medium sand. The sample has a mode at 718 µm (coarse sand) and a
184 second at 111 µm (fine sand). The grain-size sample at 31.9 cm is a trimodal, poorly sorted, very fine
185 skewed medium sand. The sample has a mode at 718 µm (coarse sand), a second at 111 µm (fine
186 sand) and a third mode at 48 µm (coarse silt).

187 In this deposit, the MS and CS fractions have dispersed fabrics (FD), with no identifiable orientation
188 (FD, Fig. 2). However, the VCS fraction displays a less-dispersed fabric, with a base-to-top evolution
189 from a transitional FB-FC fabric (low-to-moderate dipping-angle fabric with at least two different
190 orientations) to poorly-defined fabrics (FC to FD).

191 The Lisbon tsunami deposit is characterized by low ellipsoid degree of elongation (Fig. 3), with a
192 range of values between 0.01 and 0.26. The mean ellipsoid elongation is 0.05 for the MS fraction,
193 0.07 for the CS fraction and 0.18 for the VCS fraction. Thus, the degree of elongation is slightly higher
194 for the VCS fraction.

195 The mean ellipsoid plunge is low and varies between 1.1° and 19° (Fig. 3). The mean ellipsoid plunge
196 is 6.0° for the MS fraction, 6.3° for the CS fraction and 11.0° for the VCS fraction. Unlike the Pre-
197 Colombian tsunami deposit, the VCS fraction displays here the highest plunge values (Fig. 3).

198 For each fraction, there is no major variation of isotropy and elongation degree, and mean plunge
199 from the base to the top of the deposit (Standard deviation inferior to 0.14 for both isotropy and
200 elongation degree, and inferior to 9° for the mean plunge).

201 The ellipsoid degree of isotropy for the Lisbon deposit varies from 0.33 to 0.64 (Fig. 4). The mean
202 ellipsoid isotropy is 0.60 (medium isotropy) for the MS fraction, 0.55 (medium isotropy) for the CS
203 fraction and 0.48 (moderate to medium isotropy) for the VCS fraction (Fig. 4). Thus, the degree of
204 isotropy is slightly lower for the VCS fraction.

205

206 **4.3 Hurricane Donna**

207 Three grain-size samples were analysed for the Hurricane Donna deposit (2.8 to 1.05 cm; Fig. 2). They
208 are all poorly sorted, very fine skewed medium to coarse sand. The distribution is bimodal at 2.8 and
209 2.1 cm with a mode at 718 µm (coarse sand) and a second at respectively 101 and 111 µm (fine
210 sand), whereas the sample at 1.05 cm displays one unique mode at 788 µm (coarse sand). Thus,
211 there is a coarsening of the grain-size from bottom to top of the deposit.

212 Hurricane Donna deposit displays vertical variations both in terms of fabric orientation and plunge,
213 as well as variations between the different grain-size fractions analysed. Small variations of the
214 dipping-angle of fabric are visible from base to top of the deposit (Fig. 2). The base of the deposit (3.5
215 to 2 cm) has a bimodal low dipping-angle (<15°) fabric with a moderately well-defined orientation for
216 all the sand fractions, corresponding to FB (Fig. 2). The upper part of the deposit (2 to 0.25 cm) shows
217 significant differences between the different sand fractions. The VCS still define a FB-type fabric,
218 whereas, finer fractions (MS and CS) have higher dipping-angles (>15° and up to 62.4°) with one
219 dominant orientation, this defining a FB to FA-type fabric.

220 The ellipsoid elongations calculated for the Hurricane Donna deposit vary between 0.08 and 0.39
221 (Fig. 3); a range that is slightly less dispersed than for the Pre-Colombian tsunami deposit (Fig. 3). The
222 mean ellipsoid elongation is 0.16 (low elongation) for the MS fraction, 0.17 (low to moderate

223 elongation) for the CS fraction and 0.29 (moderate elongation) for the VCS fraction. Thus, the degree
224 of elongation is highest for the VCS fraction.

225 Donna's deposit has a larger range of ellipsoid plunges (0.5° to 62.4°) compared to the two other
226 deposits (Fig. 3). The mean ellipsoid plunge is 35.0° for the MS fraction, 11.3° for the CS fraction and
227 5.7° for the VCS fraction. Highest plunge values are found in the MS fraction.

228 The ellipsoid degree of isotropy for this deposit ranges between 0.29 and 0.83 (Fig. 4). The mean
229 ellipsoid isotropy is 0.76 (medium to high isotropy) for the MS fraction, 0.66 (medium isotropy) for
230 the CS fraction and 0.41 (moderate isotropy) for the VCS fraction (Fig. 4). These values are almost
231 identical to those in the Pre-Colombian tsunami deposit. The degree of isotropy is also highest for the
232 MS fraction and lowest for the VCS fraction in this deposit.

233 For each fraction, there is no major variation of isotropy and elongation degree from the base to the
234 top of the deposit (Standard deviation inferior to 0.11 for both isotropy and elongation degree).
235 However, the mean plunge values display a large broad of values and varies from base to top of the
236 deposit (between 1.9° and 62.4°), especially in the MS fraction, where the standard deviation reaches
237 24.1° .

238

239 **5. Discussion**

240 **5.1 Grain-size influence on the sedimentary fabric interpretation**

241 The separation of the grains from the matrix and their counting is dependent of the image resolution
242 ($50\ \mu\text{m}$ here) because the larger grains are then better defined. The largest grain-size fraction (VCS
243 fraction, $>1000\ \mu\text{m}$) had a higher degree of anisotropy of the ellipsoid of grain orientation, and it is
244 thus more suitable for identifying the primary fabric. Indeed, a more anisotropic ellipsoid of grain
245 orientation indicates that the grains are preferentially aligned in a given orientation (Fig. 4). The
246 smallest grain-size fraction (MS fraction, $250\text{-}500\ \mu\text{m}$) displays the lowest degree of anisotropy of the
247 ellipsoid of grain-orientation, indicating a lesser defined grain orientation (Fig. 4). Thus, the
248 stereographic projection of the smallest grain-size fraction must be interpreted with caution.

249 However, the differences in the anisotropy degree between the grain-size fractions may also be the
250 result of sedimentary processes such as interactions of coarse sand with finer-grained fractions, and
251 the characteristics of the flow itself (Paris et al., 2020). For instance, the primary fabric is better
252 preserved if fine-grained particles (silts and clays) are mixed with sands, as it was the case for the
253 high plunges of the smallest grain-size (MS fraction, 250-500 μm) at the top of Hurricane Donna
254 deposit (Fig. 3). Thus, these high-plunges could represent a secondary orientation because of
255 hammering during coring or bioturbation for instance, but most likely a primary fabric resulting from
256 sedimentary processes.

257

258 **5.2 Near-field vs far-field tsunami**

259 The Pre-Columbian tsunami deposit is a massive sandy bed and it is thicker than all other high-energy
260 event deposits recorded over the last 1600 years in Scrub Island. The absence of visible grading and
261 laminations (except for the two mud laminae at the base) in this massive coarse sand unit could
262 suggest a rapid and powerful event, maybe the strongest that hit the lagoon over this period. Thus,
263 Morton et al. (2007) suggested that extremely rapid deposition occurs when flow decelerates
264 between the uprush and backwash phases of inundation, often resulting in single-layer,
265 homogeneous and structureless tsunami deposits, where inverse grading is rare. Alternatively, a
266 deposit can also be massive if the source consists of very well-sorted sediments, or if it results from
267 sediment transport convergence (deposition over time as more sediment comes into an area than
268 leaves it) (Jaffe et al., 2012). However, the erosional base of the Pre-Columbian Tsunami deposit is
269 typical of a rapid and powerful event, which remains coherent with the massive nature of the
270 deposit. This type of tsunami deposit was already reported by Switzer et al. (2012). They found a
271 mixture of upper flow regime planar beds and homogenous massive sand in the powerful 2004
272 Indian Ocean tsunami deposit. There are no historical observations for the Pre-Colombian tsunami,
273 and its source is still debated. Because the sediment records associated with this event were found
274 (for now) in the northern part of the Antilles, it is assumed that its source may be close to the Puerto

275 Rico Trench. Cordrie et al. (2022), showed that only a near-field tsunami, originated either from an
276 Mw 8.5-8.7 mega-thrust earthquake on the subduction interface North to the Anegada Island or
277 from a Mw 8.0 outer-rise earthquake along the trench, was able to reach the location of the deposits
278 observed in Saint-Thomas, Anegada, Anguilla and create ~6-10 m height waves on Scrub Island coast
279 leading to the flood of the lagoon.

280 In this study, the Pre-Columbian deposit sedimentary fabric remains highly homogenous (FB-type
281 fabric) through all its thickness and grain-size fractions. FB is either flow-parallel or flow-oblique
282 depending on the deposition rate, sediment concentration, and bed shear stress (Paris et al., 2020).
283 Flow-parallel fabric is common for sands transported at high sediment concentration and rapidly
284 deposited (e.g. Rees, 1968, 1983; Hiscott and Middleton, 1980; Gupta et al., 1987; Baas et al., 2007).
285 Such fabric can be explained by the reduction of the distance between colliding grains, thus creating
286 transfers of angular momentum (Rees, 1983). The massive texture of the Pre-Columbian sand unit
287 suggests a high depositional rate, and it is then assumed that the FB orientation observed here is
288 aligned to the main flow during marine flooding. Given the north-south orientation of the barrier
289 facing the Caribbean Sea to the west of the lagoon (Fig 1D), it is very likely that the flow propagated
290 from the West to East in a cross-shore direction relative to the sediment barrier. Consequently, the
291 main recorded flow/current orientation (FB orientation) during marine flooding should be
292 approximately West-Est. The grain-size parameters are coherent with this observation. Thus, the top
293 of deposit (56.75 to 47.75 cm) remains homogenous: unimodal well sorted sand (718 μm) with some
294 little vertical variations of sorting and skewness. This sediment may then illustrate a source of mature
295 sediment such as the sandy barrier and/or beach with a mode at 718 μm . Finally, the flow-parallel
296 fabric maintains a low-dipping-angle because the sediment is quite well-sorted, as demonstrated by
297 Paris et al. (2020).

298 The less-defined orientation at the base of the deposit, which is more or less sub-parallel to FB
299 orientation illustrates a flow-oblique fabric. Although the base displays a different fabric compared to
300 the rest of the deposit, there is no significant differences in terms of mean plunge, isotropy and

301 elongation of the ellipsoids. Thus, the presence of flow-oblique fabric may be explained by others
302 parameters. Unlike the rest of the deposit, the base (60.25 to 56.75 cm) is poorly sorted with vertical
303 variations in term of skewness and grain sizes (modes at 718 μm , 122 μm , 596 μm , 53 μm), which
304 tend to show several sediment sources. The modes at 53 μm and 596 μm are only present in sample
305 at the base of the deposit. It is thus likely that these sample illustrates a mix of different sources of
306 sediment. The mode at 53 μm (coarse silt) reveals for example silty lagoonal sediment that was
307 incorporated with the sand during the erosion part of the deposition. Moreover, the oblique
308 orientation may be the result of the bed shear stress that occurred at the bottom of the flow during
309 the event.

310 Paris et al. (2020) proposed that vertical variations of the fabric orientation could thus reflect
311 changes in sediment concentration and bed shear stress. The transition from a flow-oblique fabric to
312 a flow-parallel fabric could also illustrate an acceleration of the flow and an increase of the sediment
313 concentration.

314

315 The deposit interpreted as being formed by the Lisbon Tsunami (Biguenet et al., 2021) is composed
316 of a 1.5 cm-thick coarse-sandy layer with rip-up clast, topped by a 1.9 cm-thick silty layer with a large
317 amount of plant remains and wood originated from the sandy barrier or the interior of the island
318 (Biguenet et al., 2021; Fig. 5). Historical archives and sedimentological records (Atwater et al., 2017;
319 Fuentes et al., 2017; Paris et al., 2021) suggested that the Lisbon Transoceanic Tsunami reached the
320 Lesser Antilles, and numerical modelling (Roger et al., 2011) showed that it induced moderate
321 tsunami waves in the Lesser Antilles region. The tsunami has been observed at 18 locations across
322 the north-eastern Caribbean from Bridgetown (Barbados) to Santiago de Cuba (Cuba)
323 (NGDC/WDS,2021), with a maximum wave height of 6.4 m at Saba Island (Fuentes et al., 2017) and a
324 4.5 m-high run-up observed in Saint-Martin, close to Scrub Island (Affleck, 1755).

325 In the studied samples, the MS and CS fractions show a dispersed fabric (FD), whereas the VCS
326 fraction a- base-to-top evolution from a transitional FC fabric to a poorly defined fabric (FC to FD).

327 Falvard and Paris (2017) explained that collision between clasts during a tsunami may produce
328 isotropic ellipsoids in the resulting deposits. In some cases, a flow with a very high concentration of
329 sediment may produce dispersed fabric (Rees, 1983; Paris et al., 2020). The Lisbon tsunami deposit
330 displays no significant differences in terms of mean plunge, isotropy and elongation of the ellipsoids
331 through all the deposit. The entire sandy part of the deposit is composed of multimodal very poorly
332 to poorly sorted sand without any major grain-size variation. The mode at 718 μm , already major in
333 the Pre-Columbian deposit, is also part of all the grain-size samples of this deposit. This may again
334 indicate a contribution of the sandy barrier and/or beach. However, the sorting and multimodal
335 distribution of the Lisbon deposit grain-size samples illustrate also other sources of sediment. The
336 basal sample displays the same grain-size modes (122 μm and 53 μm) as the Pre-Columbian deposit,
337 indicating an incorporation of lagoonal sediment during the erosion phase of the deposition. The two
338 others grain-size samples (middle and top of the deposit) display almost identical grain-size
339 parameters. The only difference is the presence of a mode at 48 μm (coarse silt) in the top deposit.
340 This is most likely because of the presence of rip-up clasts and/or the 1.9 cm-thick silty layer
341 originated from the barrier or the interior of the island that is topped over the sandy part of the
342 deposit. Thus, the grain-size parameters are relatively homogeneous and cannot explain the VCS
343 fraction fabrics. The poorly-defined fabric at the base of the deposit, turning to less-defined,
344 dispersed fabric in the VCS fraction may illustrate a variation in flow regime during the event. The
345 scenario of inundation could be similar to a simulation produced for the same event on the
346 Martinique's island shore (Anse Meunier lagoon, Paris et al., 2021). The water level rises
347 progressively and it can last several hours until the barrier breaks locally, causing a washover that
348 floods the lagoon. The deposits of the Lisbon tsunami in Scrub Island record a similar washover
349 causing dispersed fabric. This hypothesis is in accordance with far-field tsunami inundation.

350

351 **5.3 Tropical cyclone deposit**

352 The sediment record of Hurricane Donna is a 3.25 cm thick poorly-sorted very fine skewed laminated
353 medium to coarse sandy deposit. Donna was a Category 3 tropical cyclone (hurricane) that hit Scrub
354 Island in September 1960. The tropical cyclone passed 21 km South-West with respect to Scrub
355 Island. Winds were at 110 kt and pressure at 952 mb (NOAA, 2020, Historical Hurricane Tracks).

356 There is no significant difference in the isotropy and elongation of the ellipsoids throughout the
357 deposit of Hurricane Donna (with values in the same range than those of the Pre-Columbian deposit),
358 but there are important variations of the fabric from base to top, and also between the different
359 grain size fractions. The lower part (3.5-2 cm depth) of the deposit is characterized by a dominant
360 moderately well-defined FB fabric, with variations of grain orientations. Here, FB fabric is less defined
361 than the one developed in the Pre-Columbian tsunami deposit. The grain-size mode at 718 μm
362 probably illustrates again the arrival of barrier and/or beach sand. The base does not show silty
363 lagoonal sediment which may indicate a less erosive event than the two tsunami deposits. Thus, the
364 base of the deposit tends to show a less intense and/or concentrated initial flow than for the tsunami
365 deposits. The absence of an erosional surface at the base of storm deposits was also observed in
366 washover deposits related to the 1991 Halloween storm in the northeastern U.S. coastline by Switzer
367 and Jones (2008) and in Typhoon Haiyan overwash sediments from Leyte Gulf coastlines by Soria et
368 al. (2018). This was related to a lower inundation velocity compared to tsunami events. This is
369 consistent with a less intense and/or concentrated initial flow during storms compared to tsunamis.

370 The upper part (2-0.25 cm depth) is clearly different, with FB to FA fabric in CS and FB fabric in VCS
371 (Fig. 2). There is then a transition to FB/FA moderate-to-high dipping-angle fabric, typical of a rapidly
372 deposited highly concentrated flow. Very high-dipping-angle fabric seen in the MS fraction should be
373 considered with caution, since it mostly illustrates the fact that MS acts as a matrix between the
374 coarser grains (CS and VCS). This observation is restricted to the tropical cyclone deposit, as both the
375 Lisbon and Pre-Columbian tsunami deposits have very little vertical variations of the mean plunge.
376 The primary fabric is here preserved thanks to a finely-skewed grain size distribution (i.e. the
377 presence of a fine fraction, here the medium sand, intercalated between the coarser fractions). The

378 lamination (upper plane beds) observed in the deposit is consistent with the deposition under a near-
379 supercritical flow regime. Fabric in upper plane beds is usually flow-parallel, which is consistent with
380 the orientation of the fabric throughout the deposit. Moreover, the coarsening at the top of the
381 deposit (788 μm mode at 1.05 cm) tends to show an intensification of the flow. It could also record a
382 more energetic flow entering the lagoon related to the lowering or opening of the sandy barrier.
383 Inverse grading may also be the result of dispersive pressures due to grains collision, that causes
384 larger grains to move toward the zone of lower pressure at the top of the sediment layer (Bagnold,
385 1954). Inverse grading may also result from kinetic sieving, which implies downward migration of
386 smaller grains and upward moving of larger grains during deposition (Middleton, 1970). Leatherman
387 and Williams (1983) concluded that both these mechanisms may occur during the flow and are then
388 part of the primary fabric of the deposit. Overall, the lamination (upper plane beds), and the
389 coarsening of the deposit at the top of Hurricane Donna deposit is consistent with other storms
390 deposits such as the Haiyan overwash deposit (Soria et al., 2017), which were interpreted as
391 indicating upper flow regime conditions of a unidirectional, turbulent, high-velocity flow.

392 Moreover, the small variations of dipping-angle of fabric through the deposit are reminiscent of
393 sediment impulses related to flow variations creating laminations. Plane or laminated bedding in
394 coastal deposits is generally attributed to repeated inundation by storm waves (Switzer et al., 2012;
395 Tuttle and al., 2004). Here, the fabric orientation and dipping-angle variations tend to show
396 successive waves arrival, which is typical of infragravity waves (Baumann et al., 2017). Infragravity
397 waves correspond to surface ocean gravity waves with periods in the range 20–200s and can increase
398 rapidly to over 1 m in nearshore water depths (Baumann et al., 2017). Thus, infragravity waves are
399 known to dominate the hydro-sedimentary dynamics of dissipative beaches, particularly under storm
400 conditions (Russell, 1993). These waves have significant contribution to wave runup and overwash
401 formed during storms (Bertin et al., 2018) and may have brought the sand deposit of Hurricane
402 Donna in Scrub Island lagoon.

403

404 **6. Conclusion**

405 This work is based on an exceptional 500 year-long sediment record located in a small coastal lagoon
406 (Anguilla, Caribbean) including three contrasted high-energy events: a near-field tsunami, a far-field
407 tsunami and a tropical cyclone. Thus, this sediment record gave us the opportunity to explore
408 differentiation between tsunami and tropical cyclone deposits, using X-ray tomography. We found
409 that X-ray tomography is a promising tool to distinguish tsunami from tropical cyclone sandy deposits
410 because it is able to document important sediment fabric variations in those three deposits. Each
411 tsunami deposit displays mostly similar fabric parameters from base to top, but their fabric are
412 contrasted. The near-field Pre-Colombian tsunami deposit displays mostly bimodal low-dipping-angle
413 flow-aligned fabrics from base to top. The homogenous low-dipping-angle fabric of the Pre-
414 Colombian tsunami deposit is both related to its well sorting and probably to an extremely rapid and
415 intense character of tsunami waves generated by the near-field earthquake (Cordrie et al., 2022).
416 The thick and coarse Pre-Colombian tsunami deposit is obviously related to massive erosion of the
417 sandy barrier after this near-field event. The more dispersed fabric of the Lisbon tsunami deposit is
418 both related to its bimodal grain size distribution and to lower energy far-field tsunami waves. The
419 far-field Lisbon tsunami is characterized by a more or less dispersed fabric from base to top.
420 Contrastingly, the tropical cyclone deposit displays vertical variations in medium and coarse sand
421 fabric, with two orientations and opposite low dipping-angle fabric at base and one dominant
422 orientation with moderate to high dipping-angle at top. The more homogenous fabric of the two
423 tsunami deposits from base to top is interpreted as the result of rapid sedimentation induced by the
424 arrival of few inundation waves in the lagoon over a period of minutes to hours, which is typical
425 during coastal tsunamis (e.g., Tuttle et al., 2004). The vertical variations in orientation and dipping-
426 angle of the tropical cyclone deposit are interpreted as resulting from a larger number of waves that
427 occurred during a period of hours to days. These waves can be attributed to infragravity waves,
428 frequently observed during storms. Our observations are promising but still too scarce to be
429 generalized as general criteria to distinguish tsunami and tropical cyclone deposits. Nevertheless,

430 they are consistent and complementary with the results of Falvard and Paris (2017) and Paris et al.
431 (2020). Similar studies of comparable deposits should be conducted in order to obtain reproducible
432 observations and generalize sediment fabric parameters as criteria to distinguish tsunami and
433 tropical cyclone deposits. It appears that these parameters are subtle, leading to the idea that
434 potentially many past high-energy events have been misidentified.

435

436 **Data availability**

437 All data generated or analyzed during this study are included in this article or available from the
438 corresponding author on reasonable request.

439

440 **Acknowledgments**

441 This work is part of the CARQUAKES project (ANR-17-CE03-0006). The objective of the CARQUAKES
442 project is to improve the catalog of large earthquakes and tsunamis in the Lesser Antilles and
443 characterize the related hazards by applying an innovative and novel multidisciplinary approach
444 combining several state-of-the-art methods of offshore and onshore paleoseismology and tsunami
445 modelling. We thank Simon Falvard for helping us in images processing and sediment fabric analyses.
446 Thanks are due to the reviewers and the editor Edward Anthony for their constructive comments
447 that helped improve the manuscript.

448

449 **References cited**

450 Affleck, B., 1755. An account of the agitation of the sea at Antigua, Nov. 1, 1755. By Capt. Affleck of
451 the Advice Man of War. Communicated by Charles Gray, E/q; F.R.S in a Letter to William
452 Watson, F.R.S.
453 Atwater, B.F., ten Brink, U.S., Cescon, A.L., Feuillet, N., Fuentes, Z., Halley, R.B., Nuñez, C., Reinhardt,
454 E.G., Roger, J.H., Sawai, Y., Spiske, M., Tuttle, M.P., Wei, Y., Weil-Accardo, J., 2017. Extreme

455 waves in the British Virgin Islands during the last centuries before 1500 CE. *Geosphere* 13,
456 301–368. <https://doi.org/10.1130/GES01356.1>

457 Baas, J.H., Hailwood, E.A., McCaffrey, W.D., Kay, M., Jones, R., 2007. Directional petrological
458 characterisation of deep-marine sandstones using grain fabric and permeability anisotropy:
459 Methodologies, theory, application and suggestions for integration. *Earth-Science Reviews*
460 82, 101–142. <https://doi.org/10.1016/j.earscirev.2007.02.003>

461 Bagnold, 1954. Experiments on a gravity-free dispersion of large solid spheres in a Newtonian fluid
462 under shear. *Proc. R. Soc. Lond. A* 225, 49–63. <https://doi.org/10.1098/rspa.1954.0186>

463 Baumann, J., Chaumillon, E., Bertin, X., Schneider, J.-L., Guillot, B., Schmutz, M., 2017. Importance of
464 infragravity waves for the generation of washover deposits. *Marine Geology* 391, 20–35.
465 <https://doi.org/10.1016/j.margeo.2017.07.013>

466 Benn, D.I., 1994. Fabric Shape and the Interpretation of Sedimentary Fabric Data. *SEPM JSR Vol. 64A*.
467 <https://doi.org/10.1306/D4267F05-2B26-11D7-8648000102C1865D>

468 Bertin, X., de Bakker, A., van Dongeren, A., Coco, G., André, G., Ardhuin, F., Bonneton, P., Bouchette,
469 F., Castelle, B., Crawford, W.C., Davidson, M., Deen, M., Dodet, G., Guérin, T., Inch, K.,
470 Leckler, F., McCall, R., Muller, H., Olabarrieta, M., Roelvink, D., Ruessink, G., Sous, D.,
471 Stutzmann, É., Tissier, M., 2018. Infragravity waves: From driving mechanisms to impacts.
472 *Earth-Science Reviews* 177, 774–799. <https://doi.org/10.1016/j.earscirev.2018.01.002>

473 Biguenet, M., Sabatier, P., Chaumillon, E., Chagué, C., Arnaud, F., Jorissen, F., Coulombier, T., Geba,
474 E., Cordrie, L., Vacher, P., Develle, A.L., Chalmin, E., Soufi, F., Feuillet, N., 2021. A 1600 year-
475 long sedimentary record of tsunamis and hurricanes in the Lesser Antilles (Scrub Island,
476 Anguilla). *Sedimentary Geology* 412, 105806. <https://doi.org/10.1016/j.sedgeo.2020.105806>

477 Cnudde, V., Boone, M.N., 2013. High-resolution X-ray computed tomography in geosciences: A
478 review of the current technology and applications. *Earth-Science Reviews* 123, 1–17.
479 <https://doi.org/10.1016/j.earscirev.2013.04.003>

480 Cordrie, L., Feuillet, N., Gailler, A., Biguenet, M., Chaumillon, E., Sabatier, P., 2022. A Megathrust
481 earthquake as source of a Pre-Colombian tsunami in Lesser Antilles: Insight from sediment
482 deposits and tsunami modeling. *Earth-Science Reviews* 228, 104018.
483 <https://doi.org/10.1016/j.earscirev.2022.104018>

484 Cuven, S., Paris, R., Falvard, S., Miot-Noirault, E., Benbakkar, M., Schneider, J.-L., Billy, I., 2013. High-
485 resolution analysis of a tsunami deposit: Case-study from the 1755 Lisbon tsunami in
486 southwestern Spain. *Marine Geology* 337, 98–111.
487 <https://doi.org/10.1016/j.margeo.2013.02.002>

488 Donnelly, J.P., Woodruff, J.D., 2007. Intense hurricane activity over the past 5,000 years controlled by
489 El Niño and the West African monsoon. *Nature* 447, 465–468.
490 <https://doi.org/10.1038/nature05834>

491 Engel, M., Oetjen, J., May, S.M., Brückner, H., 2016. Tsunami deposits of the Caribbean – Towards an
492 improved coastal hazard assessment. *Earth-Science Reviews* 163, 260–296.
493 <https://doi.org/10.1016/j.earscirev.2016.10.010>

494 Falvard, S., Paris, R., 2017. X-ray tomography of tsunami deposits: Towards a new depositional model
495 of tsunami deposits. *Sedimentology* 64, 453–477. <https://doi.org/10.1111/sed.12310>

496 Feuillet, N., Manighetti, I., Tapponnier, P., Jacques, E., 2002. Arc parallel extension and localization of
497 volcanic complexes in Guadeloupe, Lesser Antilles. *Journal of Geophysical Research: Solid*
498 *Earth* 107, 2331.

499 Flisch, A., Becker, A., 2003. Industrial X-ray computed tomography studies of lake sediment drill
500 cores. *Geological Society, London, Special Publications* 215, 205–212.
501 <https://doi.org/10.1144/GSL.SP.2003.215.01.19>

502 Fuentes, Z., Tuttle, M.P., Schmidt, W.E., 2017. Sand Scripts of Past Tsunamis in Coastal Ponds of St.
503 Thomas, U.S. Virgin Islands. *Seismological Research Letters* 88, 1516–1526.
504 <https://doi.org/10.1785/0220170038>

505 Goff, J., McFadgen, B.G., Chagué-Goff, C., 2004. Sedimentary differences between the 2002 Easter
506 storm and the 15th-century Okoropunga tsunami, southeastern North Island, New Zealand.
507 Marine Geology 204, 235–250. [https://doi.org/10.1016/S0025-3227\(03\)00352-9](https://doi.org/10.1016/S0025-3227(03)00352-9)

508 Goslin, J., Clemmensen, L.B., 2017. Proxy records of Holocene storm events in coastal barrier
509 systems: Storm-wave induced markers. Quaternary Science Reviews 174, 80–119.
510 <https://doi.org/10.1016/j.quascirev.2017.08.026>

511 Gupta, G.D., Parkash, B., Garde, R.J., 1987. An experimental study of fabric development in plane-bed
512 phases. Sedimentary Geology 53, 101-122.

513 Hiscott, R.N., Middleton, G.V., 1980. Fabric of Coarse Deep-water Sandstones Tourelle Formation,
514 Quebec, Canada. Journal of Sedimentary Petrology 50, No 3, 703-722.

515 Jaffe, B.E., Goto, K., Sugawara, D., Richmond, B., Fujino, S., and Nishimura, Y., 2012. Flow speed
516 estimated by inverse modeling of sandy tsunami deposits: results from the 11 March 2011
517 tsunami on the coastal plain near the Sendai Airport, Honshu, Japan, Sedimentary Geology
518 282, p. 90-109, doi.org/10.1016/j.sedgeo.2012.09.002.

519 Ketcham, R.A., 2005. Three-dimensional grain fabric measurements using high-resolution X-ray
520 computed tomography. Journal of Structural Geology 27, 1217–1228.
521 <https://doi.org/10.1016/j.jsg.2005.02.006>

522 Kortekaas, S., Dawson, A.G., 2007. Distinguishing tsunami and storm deposits: An example from
523 Martinhal, SW Portugal. Sedimentary Geology 200, 208–221.
524 <https://doi.org/10.1016/j.sedgeo.2007.01.004>

525 Leatherman, S.P., Williams, A.T., 1983. Vertical sedimentation units in a barrier island washover fan.
526 Earth Surf. Process. Landforms 8, 141–150. <https://doi.org/10.1002/esp.3290080205>

527 Legland, D., Arganda-Carreras, I., Andrey, P., 2016. MorphoLibJ: integrated library and plugins for
528 mathematical morphology with ImageJ. Bioinformatics btw413.
529 <https://doi.org/10.1093/bioinformatics/btw413>

530 Liu, K., Fearn, M.L., 1993. Lake-sediment record of late Holocene hurricane activities from coastal
531 Alabama. *Geology* 21, 793–796.

532 Liu, K., Fearn, M.L., 2000. Reconstruction of Prehistoric Landfall Frequencies of Catastrophic
533 Hurricanes in Northwestern Florida from Lake Sediment Records. *Quaternary Research* 54,
534 238–245. <https://doi.org/10.1006/qres.2000.2166>

535 May, S.M., Falvard, S., Norpoth, M., Pint, A., Brill, D., Engel, M., Scheffers, A., Dierick, M., Paris, R.,
536 Squire, P., Brückner, H., 2016. A mid-Holocene candidate tsunami deposit from the NW Cape
537 (Western Australia). *Sedimentary Geology* 332, 40–50.
538 <https://doi.org/10.1016/j.sedgeo.2015.11.010>

539 Middleton, G. V., 1970. Experimental studies related to problems of flysch sedimentation. Lajoie, J.,
540 ed., *Flysch Sedimentology in North America*. Geol. Assoc. Canada Spec. Paper 7, 253-272.

541 Minoura, K., Imamura, F., Sugawara, D., Kono, Y., Iwashita, T., 2001. The 869 Jogan tsunami deposit
542 and recurrence interval of large-scale tsunami on the Pacific coast of northeast Japan.
543 *Journal of Natural Disaster Science* 23, 83–88.

544 Monecke, K., Finger, W., Klarer, D., Kongko, W., McAdoo, B.G., Moore, A.L., Sudrajat, S.U., 2008. A
545 1,000-year sediment record of tsunami recurrence in northern Sumatra. *Nature* 455, 1232–
546 1234. <https://doi.org/10.1038/nature07374>

547 Morton, R.A., Gelfenbaum, G., and Jaffe, B.E., 2007, Physical criteria for distinguishing sandy tsunami
548 and storm deposits using modern examples. *Sedimentary Geology*, v. 200, p. 184-207,
549 [doi:10.1016/j.sedgeo.2007.01.003](https://doi.org/10.1016/j.sedgeo.2007.01.003).

550 Nanayama, F., Shigeno, K., Satake, K., Shimokawa, K., Koitabashi, S., Miyasaka, S., Ishii, M., 2000.
551 Sedimentary differences between the 1993 Hokkaido-nansei-oki tsunami and the 1959
552 Miyakojima typhoon at Taisei, southwestern Hokkaido, northern Japan. *Sedimentary Geology*
553 135, 255–264. [https://doi.org/10.1016/S0037-0738\(00\)00076-2](https://doi.org/10.1016/S0037-0738(00)00076-2)

554 Nanayama, F., Satake, K., Furukawa, R., Shimokawa, K., Atwater, B.F., Shigeno, K., Yamaki, S., 2003.
555 Unusually large earthquakes inferred from tsunami deposits along the Kuril trench. *Nature*
556 424, 660–663. <https://doi.org/10.1038/nature01864>

557 National Geophysical Data Center/World Data Service: NGDC/WDS Global Historical Tsunami
558 Database. NOAA National Centers for Environmental Information. doi:10.7289/V5PN93H7.
559 (Accessed 25 October 2021).

560 NOAA (National Oceanic and Atmospheric Administration), 2020. Historical Hurricane Tracks.
561 <https://coast.noaa.gov/hurricanes/#map=4/32/-80>. (Accessed 6 November 2020).

562 Orsi, T.H., Edwards, C.M., Anderson, A.L., 1994. X-ray computed tomography; a nondestructive
563 method for quantitative analysis of sediment cores. *Journal of Sedimentary Research* 64,
564 690–693. <https://doi.org/10.1306/D4267E74-2B26-11D7-8648000102C1865D>

565 Paris, R., Sabatier, P., Biguenet, M., Bougouin, A., André, G., Roger, J., 2021. A tsunami deposit at
566 Anse Meunier, Martinique Island: Evidence of the 1755 CE Lisbon tsunami and implication for
567 hazard assessment. *Marine Geology* 439, 106561.
568 <https://doi.org/10.1016/j.margeo.2021.106561>

569 Paris, R., Falvard, S., Chagué, C., Goff, J., Etienne, S., Doumalin, P., 2020. Sedimentary fabric
570 characterized by X-ray tomography: A case-study from tsunami deposits on the Marquesas
571 Islands, French Polynesia. *Sedimentology* 67, 1207–1229. <https://doi.org/10.1111/sed.12582>

572 Pinet, C., 2011. New Aerial Survey of Anguilla. European R3 Regional Risk Reduction initiative- United
573 Nations Development Program, contract BAR/UNDP/PSC/2011/ 003, IGN/IMAO report for
574 the Government of Anguilla.

575 Rees, A.I., 1968. The Production of Preferred Orientation in a Concentrated Dispersion of Elongated
576 and Flattened Grains. *The Journal of Geology* 76, 457–465. <https://doi.org/10.1086/627343>

577 Rees, A.I., 1983. Experiments on the production of transverse grain alignment in a sheared
578 dispersion. *Sedimentology* 30, 437–448. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-3091.1983.tb00682.x)
579 [3091.1983.tb00682.x](https://doi.org/10.1111/j.1365-3091.1983.tb00682.x)

580 Renter, J.A.M., 1989. Applications of computerized tomography in sedimentology. *Marine*
581 *Geotechnology* 8, 201–211. <https://doi.org/10.1080/10641198909379868>

582 Roger, J., Baptista, M.A., Sahal, A., Accary, F., Allgeyer, S., Hébert, H., 2011. The Transoceanic 1755
583 Lisbon Tsunami in Martinique. *Pure and Applied Geophysics* 168, 1015–1031.
584 <https://doi.org/10.1007/s00024-010-0216-8>

585 Russell, P.E., 1993. Mechanisms for beach erosion during storms. *Continental Shelf Research* 13,
586 1243–1265. [https://doi.org/10.1016/0278-4343\(93\)90051-X](https://doi.org/10.1016/0278-4343(93)90051-X)

587 Sabatier, P., Dezileau, L., Colin, C., Briqueu, L., Bouchette, F., Martinez, P., Siani, G., Raynal, O., Von
588 Grafenstein, U., 2012. 7000 years of paleostorm activity in the NW Mediterranean Sea in
589 response to Holocene climate events. *Quaternary Research* 77, 1–11.
590 <https://doi.org/10.1016/j.yqres.2011.09.002>

591 Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden,
592 C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J., Hartenstein, V., Eliceiri, K., Tomancak,
593 P., Cardona, A., 2012. Fiji: an open-source platform for biological-image analysis. *Nature*
594 *Methods* 9, 676–682. <https://doi.org/10.1038/nmeth.2019>

595 SHOM: Service Hydrographique et Océanographique de la Marine, 2018. MNT bathymétrie de
596 façade de Saint-Martin et Saint-Barthélemy (Projet Homonim).
597 http://dx.doi.org/10.17183/MNT_ANTN100m_HOMONIM_WGS84.

598 Soria, J.L.A., Switzer, A.D., Pilarczyk, J.E., Siringan, F.P., Khan, N.S., Fritz, H.M., 2017. Typhoon Haiyan
599 overwash sediments from Leyte Gulf coastlines show local spatial variations with hybrid
600 storm and tsunami signatures. *Sedimentary Geology* 358, 121–138.
601 <https://doi.org/10.1016/j.sedgeo.2017.06.006>

602 Soria, J.L.A., Switzer, A.D., Pilarczyk, J.E., Tang, H., Weiss, R., Siringan, F., Manglicmot, M., Gallentes,
603 A., Lau, A.Y.A., Cheong, A.Y.L., Koh, T.W.L., 2018. Surf beat-induced overwash during Typhoon
604 Haiyan deposited two distinct sediment assemblages on the carbonate coast of Hernani,

605 Samar, central Philippines. *Marine Geology, Geological Records of Extreme Wave Events* 396,
606 215–230. <https://doi.org/10.1016/j.margeo.2017.08.016>

607 Switzer, A.D., Jones, B.G., 2008. Large-scale washover sedimentation in a freshwater lagoon from the
608 southeast Australian coast: sea-level change, tsunami or exceptionally large storm? *The*
609 *Holocene* 18, 787–803. <https://doi.org/10.1177/0959683608089214>

610 Switzer, A.D., Srinivasalu, S., Thangadurai, N., Mohan, V.R., 2012. Bedding structures in Indian
611 tsunami deposits that provide clues to the dynamics of tsunami inundation. *Geological*
612 *Society, London, Special Publications* 361, 61–77. <https://doi.org/10.1144/SP361.6>

613 Tuttle, M.P., Ruffman, A., Anderson, T., Jeter, H., 2004. Distinguishing Tsunami from Storm Deposits
614 in Eastern North America: The 1929 Grand Banks Tsunami versus the 1991 Halloween Storm.
615 *Seismological Research Letters* 75, 15.

616 Wallace, E.J., Coats, S., Emanuel, K., Donnelly, J.P., 2021. Centennial-Scale Shifts in Storm Frequency
617 Captured in Paleohurricane Records From The Bahamas Arise Predominantly From Random
618 Variability. *Geophys. Res. Lett.* 48. <https://doi.org/10.1029/2020GL091145>

619

620 **Figure captions**

621 Figure 1. (A) The Antilles Arc with main faults and islands (PR: Puerto Rico, VI: British and US Virgin
622 Islands, S: Saba, StE: Sint Eustatius, StK: Saint Kitts, N: Nevis, B: Barbuda, Ant: Antigua, Mo:
623 Montserrat, G: Guadeloupe, D: Dominica, Ma: Martinique, modified from Feuillet et al., 2002, and
624 references therein). The bathymetry and topographic DEM (digital elevation model) are derived from
625 the SHOM (Service hydrographique et océanographique de la Marine) website (SHOM, 2018). The
626 small yellow square corresponds to the Anguilla-Saint-Barthélemy platform shown in panel B. (B) The
627 Anguilla-Saint-Barthélemy platform with its islands. (C) Scrub Island with topographic data obtained
628 by LIDAR (Pinet, 2011) and the bathymetry from SHOM
629 (data.shom.fr/donnees/legend/RASTER_MARINE_3857_WMETS). (D) Satellite image of the
630 westernmost coastal lagoon from Scrub Island and its sandy barrier (Google Earth Pro V 7.3.4.8248.

631 November 4, 2020. Scrub Island, Anguilla. 18°17'04,68''N, 62°57'11,82''W, Eye alt 561 meters. Maxar
632 Technologies 2021 [September 10, 2021]). The white dot corresponds to the location of SCR1-18-01
633 sedimentary core.

634

635 Figure 2. Data for core SCR1-18-01. From left to right: tsunami and tropical cyclone events (in grey),
636 sedimentary log, grain-size contour plot, X-ray photography and their associated projections of the
637 orientations (azimuth) and dipping-angle (plunge) of the longest grain axes for Hurricane Donna,
638 Lisbon Tsunami and Pre-Columbian Tsunami. The grey rectangles in the X-ray photography of the
639 Pre-Columbian Tsunami indicate area with no data because some sedimentary levels were sampled
640 before this study (Biguenet et al., 2021). The FA to FD stereographic representations are relative to
641 the facies observed in the studied deposits.

642

643 Figure 3. Elongation of the ellipsoids compared to the plunges for all grain-size fractions in the three
644 studied events. Examples of ellipsoid shapes depending on their elongation value are displayed
645 below the horizontal axis (for a plunge of 0°). Examples of ellipsoid dipping-angle depending on their
646 plunge value are displayed left to the vertical axis (for a degree of elongation of 0.3).

647

648 Figure 4. Anisotropy of the ellipsoids of grain orientation for different types of fabric (Benn, 1994),
649 subdivided by grain-size fraction for every studied deposit. The shape of the ellipsoid is represented
650 at each end of the central plot: isotropic (top), oblate (low left) and prolate (low right).

651

652 Figure 5. Three-dimensions visualisation of the Lisbon tsunami deposit. (A) Entire deposit; (B)
653 Observation of the sediment fabric in the centre of the deposit and of its erosive base.