



HAL
open science

Toric sheaves and flips

Andrew Clarke, Achim Napame, Carl Tipler

► **To cite this version:**

| Andrew Clarke, Achim Napame, Carl Tipler. Toric sheaves and flips. 2023. hal-04255427v1

HAL Id: hal-04255427

<https://hal.science/hal-04255427v1>

Preprint submitted on 24 Oct 2023 (v1), last revised 27 Jun 2024 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

TORIC SHEAVES AND FLIPS

ANDREW CLARKE, ACHIM NAPAME, AND CARL TIPLER

ABSTRACT. Any toric flip naturally induces an equivalence between the associated categories of equivariant reflexive sheaves, and we investigate how slope stability behaves through this functor. On one hand, for a fixed toric sheaf, and natural polarisations that make the exceptional loci small, we provide a simple numerical criterion that characterizes when slope stability is preserved through the flip. On the other hand, for a given flip, we introduce full subcategories of logarithmic toric sheaves and characterize when polystability is preserved for all toric sheaves in those subcategories at once.

1. INTRODUCTION

Introduced by Mumford [16] and generalized by Takemoto [23], slope stability of vector bundles, and more generally of torsion-free coherent sheaves, can be used as a device to produce moduli spaces. While slope stability is not a GIT notion in higher dimension, it behaves well with respect to tensor products and restrictions, and has a differential geometric interpretation in gauge theory through the Hitchin–Kobayashi correspondence (see e.g. [14] and reference therein). In particular, stable bundles, and more generally stable reflexive sheaves ([1]), are of particular interest in gauge theory and mathematical physics (see e.g. [13] for a survey on stable sheaves on toric varieties addressed to the mathematical physics community).

Despite its usefulness, checking stability in practice remains a difficult problem. Our goal is to add to the list of known methods to produce stable sheaves via transformations of the underlying polarised manifold. In the equivariant context of toric geometry, the behaviour of slope stability through descent under GIT quotients was studied in [4], while the problem of pulling back stable sheaves on fibrations was considered in [18]. In this paper, we study how slope stability is affected through a toric flip between polarised (simplicial) toric varieties. Those transformations are of particular interest in several aspects. From the complex geometry point of view, they form building blocks for toric Minimal Model Program (see [6, Chapter 15] and references therein), and together with fibrations and blow-ups addressed in [18], our results complete the study of slope stability through any type of extremal contraction arising in toric MMP. From the mathematical physics point of view, toric flips can be seen as singular transitions between toric varieties. Those transitions

ANDREW CLARKE, INSTITUTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DO RIO DE JANEIRO, AV. ATHOS DA SILVEIRA RAMOS 149, RIO DE JANEIRO, RJ, 21941-909, BRAZIL

ACHIM NAPAME, INSTITUTO DE MATEMÁTICA, ESTATÍSTICA E COMPUTAÇÃO CIENTÍFICA - UNICAMP, RUA SÉRGIO BUARQUE DE HOLANDA 651, 13083-970 CAMPINAS-SP, BRAZIL

CARL TIPLER, UNIV BREST, UMR CNRS 6205, LABORATOIRE DE MATHÉMATIQUES DE BRETAGNE ATLANTIQUE, FRANCE

E-mail addresses: `andrew@im.ufrj.br`, `achim@unicamp.br`, `Carl.Tipler@univ-brest.fr`.

2010 Mathematics Subject Classification. Primary: 14M25, Secondary: 14J60.

Key words and phrases. Toric flips, equivariant sheaves, slope-stability.

are of particular importance given the construction of Calabi–Yau hypersurfaces in Fano reflexive toric varieties ([2]) and the connections between various Calabi–Yau vacua through conifold transitions ([3, 21]). Our results then provide a toy model in the study of stable sheaves through singular transitions between toric varieties (see [5] for a differential geometric approach to stability of the tangent bundle through conifold transitions).

Consider a toric flip (see Section 2 for precise the definitions):

$$\begin{array}{ccc} X & \overset{\psi}{\dashrightarrow} & X' \\ & \searrow \phi & \swarrow \phi' \\ & X_0 & \end{array}$$

between two simplicial toric varieties X and X' . There is a \mathbb{Q} -Cartier divisor $D_+ \subset X$ naturally attached to the flip, such that $-D_+$ is ϕ -ample and restricts to the anticanonical divisor of the ϕ -contracted fibers (see Section 2.2). By abuse of notation, we will still denote D_+ the divisor $\psi_*(D_+) \subset X'$. Then, for some ample Cartier divisor L_0 on X_0 , there exists $\varepsilon_0 > 0$ such that the divisors

$$L_{-\varepsilon} := \phi^* L_0 - \varepsilon D_+ \subset X$$

and

$$L_\varepsilon := (\phi')^* L_0 + \varepsilon D_+ \subset X'$$

define ample \mathbb{Q} -Cartier divisors for $\varepsilon \in (0, \varepsilon_0)$. Then, our first result (Theorem 4.4) relates slope stability of toric sheaves on (X_0, L_0) , $(X, L_{-\varepsilon})$ and (X', L_ε) :

Theorem 1.1. *Let \mathcal{E} be a torus equivariant reflexive sheaf on X . Then, up to shrinking ε_0 , we have for all $\varepsilon \in (0, \varepsilon_0)$:*

- (i) *If $\phi_* \mathcal{E}$ is L_0 -stable, then \mathcal{E} (resp. $\psi_* \mathcal{E}$) is $L_{-\varepsilon}$ -stable on X (resp. L_ε -stable on X').*
- (ii) *If $\phi_* \mathcal{E}$ is L_0 -unstable, then \mathcal{E} (resp. $\psi_* \mathcal{E}$) is $L_{-\varepsilon}$ -unstable on X (resp. L_ε -unstable on X').*
- (iii) *If $\phi_* \mathcal{E}$ is L_0 -semistable, let \mathfrak{E} be the finite set of equivariant and saturated reflexive subsheaves $\mathcal{F} \subseteq \phi_* \mathcal{E}$ arising in a Jordan–Hölder filtration of $\phi_* \mathcal{E}$. If for any $\mathcal{F} \in \mathfrak{E}$,*

$$(1.1) \quad \frac{c_1(\mathcal{E}) \cdot D_+ \cdot (\phi^* L_0)^{n-2}}{\text{rank}(\mathcal{E})} < \frac{c_1((\phi^* \mathcal{F})^{\vee\vee}) \cdot D_+ \cdot (\phi^* L_0)^{n-2}}{\text{rank}(\mathcal{F})}$$

then \mathcal{E} is $L_{-\varepsilon}$ -stable on X .

A similar statement as (iii) holds for $\psi_* \mathcal{E}$ with the reverse inequalities in (1.1) (see Theorem 4.4). The intersection number that appears in (1.1) is the first order term in the ε -expansion of the $L_{-\varepsilon}$ -slope. As Theorem 1.1 shows, if $\phi_* \mathcal{E}$ is strictly semistable, this term will never allow for both \mathcal{E} and $\psi_* \mathcal{E}$ to be stable at the same time, for the considered polarisations. As \mathfrak{E} is finite, the numerical criterion in (iii) can be used in practice to produce examples of toric sheaves that go from being unstable to stable through a toric flip (see Section 4.3).

Remark 1.2. Theorem 1.1 actually holds uniformly for flat families of toric sheaves with fixed characteristic function (see the discussion at the end of Section 4.2).

While our first result focuses on specific flat families of sheaves for a given flip, our second result describes toric flips that preserve slope polystability for *all* equivariant reflexive sheaves at once, in some adapted full subcategories. Denote by $\mathfrak{R}\mathfrak{e}\mathfrak{f}^T(X)$ the category of torus equivariant reflexive sheaves on X . For any given torus invariant divisor $D \subset X$, we introduce in Section 3.3 a full subcategory $\mathfrak{R}\mathfrak{e}\mathfrak{f}^T(X, D)$ of *logarithmic toric sheaves*. Our terminology is inspired by the fact that the logarithmic tangent sheaf $\mathcal{T}_X(-\log D)$ belongs to $\text{Obj}(\mathfrak{R}\mathfrak{e}\mathfrak{f}^T(X, D))$. Setting $D' = \psi_*D$, the birational transformation $\psi : X \dashrightarrow X'$ induces an equivalence of categories (still denoted ψ_*) between $\mathfrak{R}\mathfrak{e}\mathfrak{f}^T(X, D)$ and $\mathfrak{R}\mathfrak{e}\mathfrak{f}^T(X', D')$. We will say that the functor ψ_* preserves polystability for a pair of ample classes $(\alpha, \alpha') \in \text{Pic}(X)_{\mathbb{Q}} \times \text{Pic}(X')_{\mathbb{Q}}$ if for any $\mathcal{E} \in \text{Obj}(\mathfrak{R}\mathfrak{e}\mathfrak{f}^T(X, D))$, \mathcal{E} is polystable on (X, α) if and only if $\psi_*\mathcal{E}$ is polystable on (X', α') . Then, our result is as follows (see Theorem 5.1) :

Theorem 1.3. *Let Σ be the fan of X , and let D be the torus invariant divisor*

$$D := \sum_{\rho \in \Delta} D_{\rho} \subset X,$$

for $\Delta \subset \Sigma(1)$. Then, the following assertions are equivalent, for a pair of ample classes $(\alpha, \alpha') \in \text{Pic}(X)_{\mathbb{Q}} \times \text{Pic}(X')_{\mathbb{Q}}$.

- (i) The functor $\psi_* : \mathfrak{R}\mathfrak{e}\mathfrak{f}^T(X, D) \rightarrow \mathfrak{R}\mathfrak{e}\mathfrak{f}^T(X', D)$ preserves polystability for (α, α') .
- (ii) There is $c \in \mathbb{Q}_{>0}$ such that for all $\mathcal{E} \in \mathfrak{R}\mathfrak{e}\mathfrak{f}^T(X, D)$,

$$\mu_{\alpha}(\mathcal{E}) = c \mu_{\alpha'}(\psi_*\mathcal{E}).$$

- (iii) There is $c \in \mathbb{Q}_{>0}$ such that for all $\rho \notin \Delta$,

$$\deg_{\alpha} D_{\rho} = c \deg_{\alpha'} D'_{\rho}.$$

In the above statement, D_{ρ} stands for the torus invariant divisor associated to a ray $\rho \in \Sigma(1)$, and μ_{α} stands for the slope on (X, α) . We should point out that condition (iii) becomes very restrictive when Δ is small, while $\mathfrak{R}\mathfrak{e}\mathfrak{f}^T(X, D)$ becomes smaller for larger Δ 's. Nevertheless, Theorem 1.3 provides a simple numerical criterion for pairs of classes on X and X' to preserve polystability of specific toric sheaves through the toric flip.

Acknowledgments. The authors are partially supported by the grant BRIDGES ANR-FAPESP ANR-21-CE40-0017. AN and CT also benefited from the grant MARGE ANR-21-CE40-0011. AC is also partially supported by the grant Projeto CAPES - PrInt UFRJ 88887.311615/2018-00.

Notations and conventions. All varieties we will consider will be normal toric varieties over the complex numbers. For such a variety X , we denote by T its torus, N its lattice of one parameter subgroups, M its lattice of characters. For \mathbb{K} a field of characteristic zero, and a lattice W , we denote $W_{\mathbb{K}} := W \otimes_{\mathbb{Z}} \mathbb{K}$. The fan of X will be denoted Σ_X , or simply Σ when the situation is clear enough, and we will also use the notation X_{Σ} for the variety associated to the fan Σ . For a given cone $\sigma \in \Sigma$, we let $U_{\sigma} = \text{Specm}(\mathbb{C}[\sigma^{\vee} \cap M])$ the affine chart in X , $O(\sigma) \subset X$ the orbit associated to σ by the orbit-cone correspondence, and $V(\sigma)$ the closure in X of $O(\sigma)$. Finally, for a fan Σ and a subset $S \subset N_{\mathbb{R}}$, we denote by $\Sigma_S = \{\sigma \in \Sigma \mid \sigma \subset S\}$.

2. BACKGROUND ON TORIC FLIPS

2.1. Toric flips. We recall in this section the basics on toric flips (and refer the reader to [6, Section 3.3] for the definition of toric morphisms and their fan description). While our presentation differs slightly from [6, Chapter 15], we will keep most of the notations from this book, and the properties that we list here can be recovered from [6, Lemma 15.3.7, Lemma 15.3.11 and Theorem 15.3.13]. Let N be a rank n lattice, with $n \geq 3$.

Definition 2.1. A full dimensional strictly convex cone $\sigma_0 \subset N_{\mathbb{R}}$ will be called a *flipping cone* if there exist primitive elements $\{\nu_1, \dots, \nu_{n+1}\} \in N^{n+1}$ such that

- (1) The cone σ_0 is spanned by the ν_i 's :

$$\sigma_0 = \text{Cone}(\nu_1, \dots, \nu_{n+1}),$$

- (2) there exists $(b_1, \dots, b_{n+1}) \in \mathbb{Z}^{n+1}$ such that

$$\sum_{i=0}^{n+1} b_i \nu_i = 0,$$

- (3) the sets $J_- = \{i \mid b_i < 0\}$ and $J_+ = \{i \mid b_i > 0\}$ both contain at least 2 elements.

For a given flipping cone $\sigma_0 \subset N_{\mathbb{R}}$ as in Definition 2.1, we will denote

$$J_0 = \{i \mid b_i = 0\}.$$

We also introduce the notation, for any $J \subset \{1, \dots, n+1\}$,

$$\sigma_J = \text{Cone}(\nu_i \mid i \in J)$$

together with the fans

$$\Sigma_- = \{\sigma_J \mid J_+ \not\subseteq J\}$$

and

$$\Sigma_+ = \{\sigma_J \mid J_- \not\subseteq J\}.$$

Identifying σ_0 with the fan of its faces, we see that Σ_- and Σ_+ provide refinements of σ_0 . Those refinements induce toric morphisms $\phi_{\pm} : X_{\Sigma_{\pm}} \rightarrow U_{\sigma_0}$, whose properties are listed below (see [6, Lemma 15.3.11.(c)]) :

Lemma 2.2. *The morphisms $\phi_{\pm} : X_{\Sigma_{\pm}} \rightarrow U_{\sigma_0}$ are surjective and birational. Their exceptional loci are given by $V(\sigma_{J_{\pm}}) \subset X_{\Sigma_{\pm}}$ and satisfy $\phi_{\pm}(V(\sigma_{J_{\pm}})) = V(\sigma_{J_- \cup J_+})$ and $\dim V(\sigma_{J_{\pm}}) = n - |J_{\pm}|$, while $\dim V(\sigma_{J_- \cup J_+}) = |J_0|$.*

We now introduce the notion of toric flips that we will use in this paper.

Definition 2.3. Let X and X' be two n -dimensional simplicial toric varieties with fans Σ and Σ' and common lattice of one-parameter subgroups N . We will say that they are related by a *toric flip* if there exists a normal toric variety X_0 with fan Σ_0 containing a flipping cone $\sigma_0 \in \Sigma_0$ such that

$$\Sigma|_{\sigma_0} = \Sigma_+, \Sigma'|_{\sigma_0} = \Sigma_- \quad \text{and} \quad \Sigma|_{N_{\mathbb{R}} \setminus \sigma_0} = \Sigma'|_{N_{\mathbb{R}} \setminus \sigma_0} = (\Sigma_0)|_{N_{\mathbb{R}} \setminus \sigma_0}.$$

In that situation, the refinements Σ and Σ' of Σ_0 induce toric morphisms $\phi : X \rightarrow X_0$ and $\phi' : X' \rightarrow X_0$, the latter being called the *flip* of the former.

In Definition 2.3, the fans Σ_{\pm} are the one associated to the flipping cone σ_0 as described above. Note that the definition, together with Lemma 2.2, imply that X and X' are birational and isomorphic in codimension 2, and that $X_0 \setminus (\sigma_{J_- \cup J_+})$ is simplicial. The situation of Definition 2.3 can be better summarized in the following commutative diagram :

$$\begin{array}{ccccc}
 & & \psi & & \\
 & & \dashrightarrow & & \\
 X & & & & X' \\
 \uparrow \iota_+ & \searrow \phi & & \swarrow \phi' & \uparrow \iota_- \\
 V(\sigma_{J_+}) & & X_0 & & V(\sigma_{J_-}) \\
 \searrow \phi_+ & & \uparrow \iota_0 & & \swarrow \phi_- \\
 & & V(\sigma_{J_+ \cup J_-}) & &
 \end{array}$$

The maps ϕ and ϕ' are the toric morphisms induced by the refinements Σ and Σ' of Σ_0 , while the maps ι_+ , ι_- and ι_0 denote inclusions. Finally,

$$\psi = (\phi')^{-1} \circ \phi : X \dashrightarrow X'$$

is the birational morphism that is an isomorphism away from $V(\sigma_{J_{\pm}})$. From now on, we fix X and X' two simplicial toric varieties related by a toric flip ψ , and retain the notation from the previous diagram.

2.2. The exceptional loci and relatively ample divisors. We will be interested in stability later on, whose definition requires that the varieties be polarised. We hence turn to the description of the fibers of ϕ_{\pm} , and describe some ϕ -ample (and ϕ' -ample) divisors. For a cone $\sigma \subset N_{\mathbb{R}}$, we denote $N_{\sigma} \subset N$ the sublattice

$$N_{\sigma} = \text{Span}(\sigma) \cap N,$$

and the quotient map

$$\pi_{\sigma} : N \rightarrow N(\sigma) := N/N_{\sigma}.$$

Recall that from the orbit-cone correspondence (see [6, Section 3.2, Proposition 3.2.7]), the toric variety $V(\sigma)$ can be obtained as the toric variety associated to the fan of cones in $(N/N_{\sigma})_{\mathbb{R}}$:

$$\text{Star}(\sigma) = \{\pi_{\sigma}(\tau) \mid \sigma \preceq \tau\}.$$

In particular, the lattices of one parameter subgroups of $V(\sigma_{J_-})$ and $V(\sigma_{J_- \cup J_+})$ are respectively $N/N_{\sigma_{J_-}}$ and $N/N_{\sigma_{J_- \cup J_+}}$. One can show that the projection map

$$N/N_{\sigma_{J_-}} \rightarrow N/N_{\sigma_{J_- \cup J_+}}$$

is compatible with the fans $\text{Star}(\sigma_{J_-})$ and $\text{Star}(\sigma_{J_- \cup J_+})$ and induces the toric morphism

$$\phi_- : V(\sigma_{J_-}) \rightarrow V(\sigma_{J_- \cup J_+}).$$

Those lattices fit naturally in the sequence

$$0 \rightarrow N_{\sigma_{J_- \cup J_+}}/N_{\sigma_{J_-}} \rightarrow N/N_{\sigma_{J_-}} \rightarrow N/N_{\sigma_{J_- \cup J_+}} \rightarrow 0.$$

As we are interested in the fibers of ϕ_- , we introduce the quotient lattice

$$N_{\mathcal{R}} := N_{\sigma_{J_- \cup J_+}}/N_{\sigma_{J_-}},$$

and denote the projection $N_{\sigma_{J_- \cup J_+}} \rightarrow N_{\mathcal{R}}$, and its \mathbb{R} -linear extension, by $u \mapsto \bar{u}$. We finally introduce the fan

$$\Sigma_{\mathcal{R}} := \{\bar{\sigma}_J \mid J \subsetneq J_+\},$$

and the associated toric variety $X_{\mathcal{R}}$.

Remark 2.4. We keep the notation $X_{\mathcal{R}}$ to be consistent with [6], where the \mathcal{R} stands for an *extremal ray* being responsible for the flip in the context of toric MMP.

For the following, see [6, Lemma 15.4.2 and Proposition 15.4.5.(c)].

Proposition 2.5. *The fibers of ϕ_- are isomorphic to the \mathbb{Q} -Fano toric variety $X_{\mathcal{R}}$. Moreover, $X_{\mathcal{R}}$ has dimension $|J_+| - 1$ and Picard rank one.*

From the above, we deduce that the anticanonical divisor of $X_{\mathcal{R}}$

$$-K_{\mathcal{R}} = \sum_{\rho \in \Sigma_{\mathcal{R}}(1)} D_{\rho}$$

is \mathbb{Q} -Cartier and ample. Note that by construction it can be written

$$-K_{\mathcal{R}} = \sum_{i \in J_+} D_{\mathbb{R}_+ \cdot \nu_i} = \sum_{i \in J_+} D_{\bar{\rho}_i}$$

where we set

$$\rho_i = \mathbb{R}_+ \cdot \nu_i,$$

and D_{ρ} stands for the torus-invariant divisor associated to the ray ρ . An easy exercise, using the orbit-cone correspondence, together with [6, Proposition 15.5.1], shows that

Proposition 2.6. *The \mathbb{Q} -Cartier torus invariant divisor*

$$(2.1) \quad -D_{J_+} := -\sum_{i \in J_+} D_{\rho_i} \in \text{Div}(X)$$

is ϕ -ample, while the \mathbb{Q} -Cartier invariant divisor

$$(2.2) \quad D'_{J_+} := \psi_*(D_{J_+}) = \sum_{i \in J_+} D'_{\rho_i} \in \text{Div}(X')$$

is ϕ' -ample.

Remark 2.7. In the above proposition, we use a superscript to D' to indicate that the torus invariant divisor D' is taken as the orbit closure of some ray in X' . As $\psi : X \dashrightarrow X'$ is an isomorphism in codimension 2, it induces an isomorphism ψ_* between the groups of torus invariant \mathbb{Q} -Cartier divisors on X and X' , which can be written on basis elements $D_{\rho} \mapsto D'_{\rho}$ for any $\rho \in \Sigma(1) = \Sigma'(1)$. Then, to ease notation later on, we will omit the superscript to D'_{ρ} , the context being clear enough whether D_{ρ} is considered as a divisor on X or X' . We will also simply denote D_{J_+} and D'_{J_+} by

$$D_+ = \sum_{i \in J_+} D_{\rho_i},$$

so that the conclusion of Proposition 2.6 is that $-D_+$ is ϕ -ample on X and D_+ is ϕ' -ample on X' .

Finally, we note that we could have considered the divisor

$$D_- = \sum_{i \in J_-} D_{\rho_i},$$

which is ϕ -ample (while $-D_-$ is ϕ' -ample). However, the *wall relation*

$$\sum_{i \in J_-} b_i \nu_i + \sum_{i \in J_+} b_i \nu_i = 0$$

from Definition 2.1 implies that from the intersection theory point of view, computing slopes with D_+ or D_- will produce the same results regarding stability notions (see e.g. [6, Section 6.4, Proposition 6.4.4]).

3. THE FLIP FUNCTOR

3.1. Equivariant sheaves and Klyachko's equivalence. We now turn to the description of the flip functor, which requires first introducing the categories of torus equivariant reflexive sheaves. For a given toric variety X with fan Σ , a *torus equivariant reflexive sheaf* is a reflexive sheaf \mathcal{E} on X together with an isomorphism

$$\varphi : \alpha^* \mathcal{E} \rightarrow \pi_2^* \mathcal{E}$$

satisfying certain cocycle conditions, where $\alpha : T \times X \rightarrow X$ and $\pi_2 : T_N \times X \rightarrow X$ stand for the torus action and the projection on X respectively (see for example [20, Section 5]).

Definition 3.1. A *toric sheaf* is a torus equivariant reflexive sheaf.

Klyachko has shown (see [12] for locally free sheaves and [20] in general) that any toric sheaf is uniquely described by a *family of filtrations*, denoted

$$(E, E^\rho(i))_{\rho \in \Sigma(1), i \in \mathbb{Z}}.$$

Here, E stands for a finite dimensional complex vector space of dimension $\text{rank}(\mathcal{E})$, and for each ray $\rho \in \Sigma(1)$, $(E^\rho(i))_{i \in \mathbb{Z}}$ is a bounded *increasing* filtration of E (we will use increasing filtrations as in [20], rather than decreasing ones as in [12]). Then, the equivariant reflexive sheaf \mathcal{E} is recovered from the formula, for $\sigma \in \Sigma$:

$$\Gamma(U_\sigma, \mathcal{E}) := \bigoplus_{m \in M} \bigcap_{\rho \in \sigma(1)} E^\rho(\langle m, u_\rho \rangle) \otimes \chi^m$$

where $u_\rho \in N$ is the primitive generator of ρ , $\langle \cdot, \cdot \rangle$ the duality pairing and χ^m the weight m character function.

A morphism between two families of filtrations

$$b : (E_1, E_1^\rho(i))_{\rho \in \Sigma(1), i \in \mathbb{Z}} \rightarrow (E_2, E_2^\rho(i))_{\rho \in \Sigma(1), i \in \mathbb{Z}}$$

is a linear map $b : E_1 \rightarrow E_2$ that satisfies

$$b(E_1^\rho(i)) \subset E_2^\rho(i)$$

for any $\rho \in \Sigma(1)$ and $i \in \mathbb{Z}$. Any such morphism corresponds uniquely to a morphism between the associated reflexive sheaves, and a fundamental result of Klyachko and Perling ([12, 20]) asserts that the categories of families of filtrations and of toric sheaves are equivalent. For our purposes, it seems more natural to slightly restrict the definition of morphisms. We will consider morphisms between toric sheaves \mathcal{E}_i to be equivariant morphisms of coherent sheaves $\beta : \mathcal{E}_1 \rightarrow \mathcal{E}_2$

that satisfy that $\text{Im}(\beta)$ is a saturated reflexive subsheaf of \mathcal{E}_2 . Those morphisms correspond through Klyachko's equivalence to linear maps $b : E_1 \rightarrow E_2$ such that

$$b(E_1^\rho(i)) = b(E_1) \cap E_2^\rho(i)$$

for any (ρ, i) , as can be seen via [18, Lemma 2.15] for example. We denote by $\mathfrak{R}ef^T(X)$ on one hand and by $\mathfrak{Filt}(X)$ on the other, the categories of toric sheaves and of families of filtrations, endowed with those classes of morphisms. We will denote by

$$\mathfrak{Rl} : \mathfrak{Filt}(X) \rightarrow \mathfrak{R}ef^T(X)$$

Klyachko's functor as described above. Then, Klyachko and Perling's work readily implies the following :

Theorem 3.2 ([12, 20]). *The functor \mathfrak{Rl} is an equivalence of categories.*

Remark 3.3. A nice feature of the categories $\mathfrak{Filt}(X)$ and $\mathfrak{R}ef^T(X)$ is that they are abelian. This is no longer true when we consider all morphisms of reflexive sheaves, as for example the quotient of $\mathcal{O}_{\mathbb{P}^1}$ by the subsheaf $\mathcal{O}_{\mathbb{P}^1}(1)$ is torsion, and hence not reflexive.

3.2. Flip functor. Assume now that $\phi' : X' \rightarrow X_0$ is the flip of $\phi : X \rightarrow X_0$ as in the previous section. As ϕ and ϕ' are isomorphisms in codimension 2, we have

$$\Sigma(1) = \Sigma_0(1) = \Sigma'(1).$$

We deduce that there are equivariant injections i (resp. i_0 and i') of the T -invariant Zariski open set

$$U := \bigcup_{\tau \in \Sigma(0) \cup \Sigma(1)} U_\tau$$

into X (resp. X_0 and X'). Then, from [10, Proposition 1.6], we deduce that for any toric sheaf $\mathcal{E} \in \text{Obj}(\mathfrak{R}ef^T(X))$ (resp. $\mathcal{F} \in \text{Obj}(\mathfrak{R}ef^T(X_0))$ and $\mathcal{G} \in \text{Obj}(\mathfrak{R}ef^T(X'))$), we have

$$i_*(\mathcal{E}|_U) \simeq \mathcal{E}$$

(resp. $(i_0)_*(\mathcal{F}|_U) \simeq \mathcal{F}$ and $i'_*(\mathcal{G}|_U) \simeq \mathcal{G}$). As reflexive sheaves are normal, meaning their sections extend over codimension 2 Zariski closed sets, we have :

Proposition 3.4. *The pushforward i_* (resp. i'_* and $(i_0)_*$) induces an equivalence of categories*

$$i_* : \mathfrak{R}ef^T(U) \rightarrow \mathfrak{R}ef^T(X)$$

(resp. $\mathfrak{R}ef^T(U) \simeq \mathfrak{R}ef^T(X_0)$ and $\mathfrak{R}ef^T(U) \simeq \mathfrak{R}ef^T(X')$). Hence, we have equivalences

$$\mathfrak{R}ef^T(X) \simeq \mathfrak{R}ef^T(X_0) \simeq \mathfrak{R}ef^T(X').$$

It is straightforward to check that the equivalence

$$\mathfrak{R}ef^T(X) \simeq \mathfrak{R}ef^T(X_0)$$

is induced by the pushforward ϕ_* while the equivalence

$$\mathfrak{R}ef^T(X') \simeq \mathfrak{R}ef^T(X_0)$$

is induced by ϕ'_* . Moreover, the categories of families of filtrations on U , X , X' and X_0 are readily the same, and the above equivalence of categories simply correspond to the self-equivalence of $\mathfrak{Filt}(X)$ induced by the identity on objects and morphisms.

Definition 3.5. We define the *flip functor*

$$\psi_* : \mathfrak{R}ef^T(X) \rightarrow \mathfrak{R}ef^T(X')$$

to be the composition of functors induced by $((\phi')^{-1})_*$ and ϕ_* .

We conclude this section by noting that the flip functor sends the tangent sheaf \mathcal{T}_X of X to the tangent sheaf $\mathcal{T}_{X'}$ of X' :

Lemma 3.6. *We have*

$$\phi_* \mathcal{T}_X = \mathcal{T}_{X_0} = \phi'_* \mathcal{T}_{X'}.$$

Proof. It follows directly from the description of the family of filtrations associated to the tangent sheaf, that only depends on N , $\Sigma(1)$ and the primitive generators of elements $\rho \in \Sigma(1)$ (see [12, Example 2.3(5) on page 350]). As this data is the same for X , X' and X_0 , and as the functors induced by ϕ_* and ϕ'_* correspond to the identity functor on

$$\mathfrak{Filt}(X) = \mathfrak{Filt}(X_0) = \mathfrak{Filt}(X'),$$

the result follows. \square

3.3. The logarithmic subcategories. In Section 5, we will be interested in specific subcategories of $\mathfrak{R}ef^T(X)$ and $\mathfrak{R}ef^T(X')$. For any $\Delta \subset \Sigma(1)$, we introduce the torus invariant divisor

$$D_\Delta := \sum_{\rho \in \Delta} D_\rho$$

and the full subcategory $\mathfrak{R}ef^T(X, D_\Delta)$ of $\mathfrak{R}ef^T(X)$ whose objects are the toric sheaves on X whose associated families of filtrations $(E, E^\rho(i))_{\rho \in \Sigma(1), i \in \mathbb{Z}}$ satisfy for all $\rho \in \Delta$:

$$(3.1) \quad E^\rho(i) = \begin{cases} 0 & \text{if } i < 0 \\ E & \text{if } i \geq 0. \end{cases}$$

Remark 3.7. From [17, Theorem 1.1], the logarithmic tangent sheaf $\mathcal{T}_X(-\log D_\Delta)$ belongs to $\text{Obj}(\mathfrak{R}ef^T(X, D_\Delta))$, which justify our choice of terminology.

It is then straightforward to see that the flip functor ψ_* induces an equivalence between $\mathfrak{R}ef^T(X, D_\Delta)$ and $\mathfrak{R}ef^T(X', D'_\Delta)$, where we use D'_Δ to denote $\sum_{\rho \in \Delta} D'_\rho$. Note also that ψ_* sends $\mathcal{T}_X(-\log D_\Delta)$ to $\mathcal{T}_{X'}(-\log D'_\Delta)$.

4. FLIPS AND STABILITY FOR A GIVEN SHEAF

4.1. Slope stability of toric sheaves. Let (X, L) be a polarised complex variety. Recall that a reflexive sheaf \mathcal{E} on (X, L) is said to be *slope stable* (resp. *slope semistable*), or simply *stable* (resp. *semistable*) for short, if for any coherent and saturated subsheaf $\mathcal{F} \subset \mathcal{E}$ of strictly smaller rank, one has

$$\mu_L(\mathcal{F}) < \mu_L(\mathcal{E}),$$

(resp. $\mu_L(\mathcal{F}) \leq \mu_L(\mathcal{E})$), where for any coherent torsion-free sheaf \mathcal{F} , the *slope* $\mu_L(\mathcal{F})$ is defined by

$$\mu_L(\mathcal{F}) = \frac{c_1(\mathcal{F}) \cdot L^{n-1}}{\text{rank}(\mathcal{F})} \in \mathbb{Q}.$$

A *polystable sheaf* is a direct sum of stable ones with the same slope. A sheaf will be called *unstable* if it is *not semistable*.

Remark 4.1. When referring to a specific polarisation L used to define stability notions, we will use the terminology L -stable (resp. L -unstable, L -semistable, etc).

A remarkable fact, proved by Kool ([15, Proposition 4.13]), is that if we assume X and \mathcal{E} to be toric, to check stability for \mathcal{E} , it is enough to compare slopes with equivariant and saturated reflexive subsheaves, that is sub objects of \mathcal{E} in $\mathfrak{Rcf}^T(X)$ (note that this was proved in the smooth case by Kool, but it was noted in [4] that the proof extends in the normal case). If $(E, E^\rho(\bullet))_{\rho \in \Sigma(1)}$ stands for the family of filtrations of \mathcal{E} , any saturated equivariant reflexive subsheaf of \mathcal{E} is associated to a family of filtrations of the form $(F, F \cap E^\rho(i))_{\rho \in \Sigma(1), i \in \mathbb{Z}}$ for some vector subspace $F \subsetneq E$ (see [18, Lemma 2.15]). Moreover, Klyachko's formula for the slope of a toric sheaf ([4, Corollary 2.18]) is

$$(4.1) \quad \mu_L(\mathcal{E}) = -\frac{1}{\text{rank}(\mathcal{E})} \sum_{\rho \in \Sigma(1)} \iota_\rho(\mathcal{E}) \deg_L(D_\rho),$$

where $\deg_L(D_\rho)$ is the degree with respect to L , and

$$\iota_\rho(\mathcal{E}) := \sum_{i \in \mathbb{Z}} i (\dim(E^\rho(i)) - \dim(E^\rho(i-1))).$$

To sum up, we have

Proposition 4.2. *The toric sheaf associated to $(E, E^\rho(i))_{\rho \in \Sigma(1), i \in \mathbb{Z}}$ is stable if and only if for any subspace $F \subsetneq E$, we have*

$$\frac{1}{\dim(F)} \sum_{\rho \in \Sigma(1)} \iota_\rho(F) \deg_L(D_\rho) > \frac{1}{\dim(E)} \sum_{\rho \in \Sigma(1)} \iota_\rho(\mathcal{E}) \deg_L(D_\rho),$$

where

$$\iota_\rho(F) := \sum_{i \in \mathbb{Z}} i (\dim(F \cap E^\rho(i)) - \dim(F \cap E^\rho(i-1))).$$

The similar statement holds for semistability by replacing the strict inequality by a large inequality.

Remark 4.3. As observed in Section 3.2, for a given flip as in Section 2.1, the families of filtrations for $\mathcal{E} \in \text{Obj}(\mathfrak{Rcf}^T(X))$, $\psi_*\mathcal{E}$ and $\phi_*\mathcal{E}$ are the same. We thus have the equalities $\iota_\rho(\mathcal{E}) = \iota_\rho(\phi_*\mathcal{E}) = \iota_\rho(\psi_*\mathcal{E})$. Then, to compare slopes on X , X_0 and X' , only the terms coming from the degrees of the invariant divisors D_ρ 's will vary according to the polarisations on each variety.

4.2. Main result and its proof. Consider now $\phi : X \rightarrow X_0$ and its toric flip $\phi' : X' \rightarrow X_0$ as defined in Section 2. From Proposition 2.6 (recall also Remark 2.7), for any ample Cartier divisor L_0 on X_0 , there exists $\varepsilon_0 > 0$ such that the divisors

$$L_{-\varepsilon} := \phi^*L_0 - \varepsilon D_+$$

on X and

$$L_\varepsilon := (\phi')^*L_0 + \varepsilon D_+$$

on X' define ample \mathbb{Q} -Cartier divisors for $\varepsilon \in (0, \varepsilon_0)$. We will then be interested in the behaviour of stability for toric sheaves related by the flip functor on $(X, L_{-\varepsilon})$ and (X', L_ε) , for $0 < \varepsilon < \varepsilon_0$. Note that a necessary condition for stability of an element $\mathcal{E} \in \text{Obj}(\mathfrak{Rcf}^T(X))$ under those polarisations is L_0 -semistability of $\phi_*\mathcal{E}$.

Conversely, if the sheaf $\phi_*\mathcal{E}$ is L_0 -semistable, it then admits a Jordan-Hölder filtration

$$0 = \mathcal{E}_1 \subseteq \mathcal{E}_2 \subseteq \dots \subseteq \mathcal{E}_\ell = \phi_*\mathcal{E}$$

by slope semistable coherent and saturated subsheaves with stable quotients of same slope as $\phi_*\mathcal{E}$ (see e.g. [11]). We denote by

$$\mathrm{Gr}(\phi_*\mathcal{E}) := \bigoplus_{i=1}^{\ell-1} \mathcal{E}_{i+1}/\mathcal{E}_i$$

the graded object of $\phi_*\mathcal{E}$ and by \mathfrak{E} the finite set of equivariant and saturated reflexive subsheaves $\mathcal{F} \subseteq \phi_*\mathcal{E}$ arising in a Jordan-Hölder filtration of $\phi_*\mathcal{E}$. Note that by Proposition 3.4, for any $\mathcal{F} \in \mathfrak{E}$, $(\phi^*\mathcal{F})^{\vee\vee}$ (resp. $((\phi')^*\mathcal{F})^{\vee\vee}$) is saturated in \mathcal{E} (resp. in $\psi_*\mathcal{E}$).

Theorem 4.4. *Let \mathcal{E} be a toric sheaf on X . Then, up to shrinking ε_0 , we have for all $\varepsilon \in (0, \varepsilon_0)$:*

- (i) *If $\phi_*\mathcal{E}$ is L_0 -stable, then \mathcal{E} (resp. $\psi_*\mathcal{E}$) is $L_{-\varepsilon}$ -stable on X (resp. L_ε -stable on X').*
- (ii) *If $\phi_*\mathcal{E}$ is L_0 -unstable, then \mathcal{E} (resp. $\psi_*\mathcal{E}$) is $L_{-\varepsilon}$ -unstable on X (resp. L_ε -unstable on X').*
- (iii) *If $\phi_*\mathcal{E}$ is L_0 -semistable, and if for any $\mathcal{F} \in \mathfrak{E}$,*

$$\frac{c_1(\mathcal{E}) \cdot D_+ \cdot (\phi^*L_0)^{n-2}}{\mathrm{rank}(\mathcal{E})} < \frac{c_1((\phi^*\mathcal{F})^{\vee\vee}) \cdot D_+ \cdot (\phi^*L_0)^{n-2}}{\mathrm{rank}(\mathcal{F})}$$

then \mathcal{E} (resp. $\psi_\mathcal{E}$) is $L_{-\varepsilon}$ -stable on X (resp. L_ε -unstable on X').*

- (iv) *If $\phi_*\mathcal{E}$ is L_0 -semistable, and if for any $\mathcal{F} \in \mathfrak{E}$,*

$$\frac{c_1(\mathcal{E}) \cdot D_+ \cdot (\phi^*L_0)^{n-2}}{\mathrm{rank}(\mathcal{E})} > \frac{c_1((\phi^*\mathcal{F})^{\vee\vee}) \cdot D_+ \cdot (\phi^*L_0)^{n-2}}{\mathrm{rank}(\mathcal{F})}$$

then \mathcal{E} (resp. $\psi_\mathcal{E}$) is $L_{-\varepsilon}$ -unstable on X (resp. L_ε -stable on X').*

Remark 4.5. Note that, given the semistability of $\phi_*\mathcal{E}$, the numerical criterion in points (iii) and (iv) only requires testing a *finite* number of inequalities, as \mathfrak{E} is finite. This makes this criterion useful in practice.

Before proving this theorem, we first recall some facts on intersection products in toric varieties that will be used. Let $\{u_1, \dots, u_k\}$ be a set of primitive elements of N such that $\sigma = \mathrm{Cone}(u_1, \dots, u_k)$ is simplicial. We define $\mathrm{mult}(\sigma)$ as the index of the sublattice $\mathbb{Z}u_1 + \dots + \mathbb{Z}u_k$ in $N_\sigma = \mathrm{Span}(\sigma) \cap N$. If Σ is simplicial, according to [9, Section 5.1], one can consider intersections of cycles or cycle classes with *rational* coefficients. The Chow group

$$A^\bullet(X)_\mathbb{Q} = \bigoplus_{p=0}^n A^p(X) \otimes \mathbb{Q} = \bigoplus_{p=0}^n A_{n-p}(X) \otimes \mathbb{Q}$$

has the structure of a graded \mathbb{Q} -algebra and by [6, Lemma 12.5.2], if $\rho_1, \dots, \rho_d \in \Sigma(1)$ are distinct, then in $A^\bullet(X)_\mathbb{Q}$, we have

$$(4.2) \quad [D_{\rho_1}] \cdot [D_{\rho_2}] \cdots [D_{\rho_d}] = \begin{cases} \frac{1}{\mathrm{mult}(\sigma)} [V(\sigma)] & \text{if } \sigma = \rho_1 + \dots + \rho_d \in \Sigma \\ 0 & \text{otherwise.} \end{cases} .$$

If χ^m is the weighted m character function on X_Σ , then by [6, Proposition 4.1.2 and (12.5.4)], the divisor of χ^m is given by

$$(4.3) \quad \operatorname{div}(\chi^m) = \sum_{\rho \in \Sigma(1)} \langle m, u_\rho \rangle D_\rho$$

and $\operatorname{div}(\chi^m) \sim_{\text{lin}} 0$ in $A^1(X_\Sigma)$.

Proof of Theorem 4.4. We first prove that for any $\rho \in \Sigma(1)$,

$$(4.4) \quad \psi_*(D_\rho \cdot D_+) = D'_\rho \cdot D'_+.$$

We use the notation of Section 2.1 for the toric flip, and set

$$\Delta = \{\operatorname{Cone}(\nu_i) : i \in J_+\}.$$

Recall from Definition 2.1 that J_+ and J_- have at least two elements. It then follows from the definition of Σ_\pm that for any $\rho \in \sigma_0(1) \setminus \Delta$, and any $j \in J_+$, $\rho + \operatorname{Cone}(\nu_j)$ is a two-dimensional cone of Σ_+ and Σ_- . Together with the definition of ψ , we deduce that in the Chow ring $A^\bullet(X')_{\mathbb{Q}}$,

$$\psi_*(D_\rho \cdot D_{\rho_j}) = D'_\rho \cdot D'_{\rho_j}.$$

If $\rho \in \Sigma(1) \setminus \sigma_0(1)$, then for any $j \in J_+$,

$$\rho + \operatorname{Cone}(\nu_j) \notin \{\tau : \tau \preceq \sigma_0\}.$$

As by Definition 2.3,

$$\Sigma|_{N_{\mathbb{R}} \setminus \sigma_0} = \Sigma'|_{N_{\mathbb{R}} \setminus \sigma_0} = (\Sigma_0)|_{N_{\mathbb{R}} \setminus \sigma_0},$$

we deduce that:

- either $\rho + \operatorname{Cone}(\nu_j) \in \Sigma_0 \setminus \{\tau : \tau \preceq \sigma_0\}$, and then $\psi_*(D_\rho \cdot D_{\rho_j}) = D'_\rho \cdot D'_{\rho_j}$ in $A^\bullet(X')_{\mathbb{Q}}$;
- or $\rho + \operatorname{Cone}(\nu_j) \notin \Sigma_0 \setminus \{\tau : \tau \preceq \sigma_0\}$, in which case $D_\rho \cdot D_{\rho_j} = 0$ in $A^\bullet(X)_{\mathbb{Q}}$ and $D'_\rho \cdot D'_{\rho_j} = 0$ in $A^\bullet(X')_{\mathbb{Q}}$.

This proves that for any $\rho \in \Sigma(1) \setminus \Delta$ and any $j \in J_+$,

$$\psi_*(D_\rho \cdot D_{\rho_j}) = D'_\rho \cdot D'_{\rho_j}$$

and then by linearity and definition of D_+ :

$$\psi_*(D_\rho \cdot D_+) = D'_\rho \cdot D'_+.$$

We now assume that $\rho \in \Delta$. By Lemma 2.2, one has $\dim V(\sigma_{J_+}) = n - |J_+|$; therefore σ_{J_+} is a simplicial cone and then $\{\nu_j : j \in J_+\}$ form a part of a \mathbb{Q} -basis of $N \otimes_{\mathbb{Z}} \mathbb{Q}$. Let $\{\nu_j^* : j \in J_+\}$ be a part of a \mathbb{Q} -basis of $M \otimes_{\mathbb{Z}} \mathbb{Q}$ such that for any $i, j \in J_+$,

$$\langle \nu_j^*, \nu_i \rangle = \begin{cases} 0 & \text{if } i \neq j \\ 1 & \text{if } i = j \end{cases}.$$

For any $j \in J_+$, there is $a_j \in \mathbb{N}^*$ such that $a_j \nu_j^* \in M$. By using (4.3) with $m = a_j \nu_j^*$, we get

$$a_j D_{\rho_j} \sim_{\text{lin}} - \sum_{\rho \in \Sigma(1) \setminus \Delta} \langle a_j \nu_j^*, u_\rho \rangle D_\rho.$$

on X and X' . By the first cases, we deduce that for any $j \in J_+$,

$$\psi_*(D_{\rho_j} \cdot D_+) = -\psi_* \left(\sum_{\rho \in \Sigma(1) \setminus \Delta} \langle \nu_j^*, u_\rho \rangle D_\rho \cdot D_+ \right)$$

$$\begin{aligned}
&= - \sum_{\rho \in \Sigma(1) \setminus \Delta} \langle \nu_j^*, u_\rho \rangle D'_\rho \cdot D'_+ \\
&= D'_{\rho_j} \cdot D'_+
\end{aligned}$$

in $A^\bullet(X')_{\mathbb{Q}}$. This concludes the proof of (4.4).

We can now compute the slopes. We have

$$\begin{aligned}
(L_{-\varepsilon})^{n-1} &= (\phi^* L_0)^{n-1} - (n-1)\varepsilon D_+ \cdot (\phi^* L_0)^{n-2} + o(\varepsilon) \quad \text{and} \\
(L_\varepsilon)^{n-1} &= ((\phi')^* L_0)^{n-1} + (n-1)\varepsilon D_+ \cdot ((\phi')^* L_0)^{n-2} + o(\varepsilon).
\end{aligned}$$

By the Projection formula [8, Proposition 2.3], for any $\rho \in \Sigma(1)$, one has

$$D_\rho \cdot (\phi^* L_0)^{n-1} = (\phi_* D_\rho) \cdot (L_0)^{n-1} = \deg_{L_0}(D_\rho)$$

and

$$D_\rho \cdot ((\phi')^* L_0)^{n-1} = (\phi'_* D_\rho) \cdot (L_0)^{n-1} = \deg_{L_0}(D_\rho).$$

By (4.4), we get

$$D_\rho \cdot D_+ \cdot (\phi^* L_0)^{n-2} = D_\rho \cdot D_+ \cdot ((\phi')^* L_0)^{n-2}.$$

Hence, for any coherent sheaf \mathcal{E} on X ,

$$\mu_{L_{-\varepsilon}}(\mathcal{E}) = \mu_{L_0}(\phi_* \mathcal{E}) - \frac{c_1(\mathcal{E}) \cdot D_+ \cdot (\phi^* L_0)^{n-2}}{\text{rank}(\mathcal{E})} (n-1)\varepsilon + o(\varepsilon)$$

and

$$\mu_{L_\varepsilon}(\psi_* \mathcal{E}) = \mu_{L_0}(\phi_* \mathcal{E}) + \frac{c_1(\mathcal{E}) \cdot D_+ \cdot (\phi^* L_0)^{n-2}}{\text{rank}(\mathcal{E})} (n-1)\varepsilon + o(\varepsilon).$$

Assume now $\mathcal{E} \in \text{Obj}(\mathfrak{Rcf}^T(X))$ is given by the family of filtrations $(E, E^\rho(j))$. By Proposition 4.2, to check the stability of \mathcal{E} , it is enough to compare the slope of \mathcal{E} with the slope of any equivariant reflexive sheaf \mathcal{F} given by the family of filtrations $(F, F \cap E^\rho(j))$ for $F \subsetneq E$ a subspace. As the set of vector subspaces $F \subset E$ that are necessary to test slopes on is actually finite (see [18, Lemma 2.17]), we deduce from the above ε -expansions for the slopes that there is $\varepsilon_0 > 0$ such that for all $\varepsilon \in (0, \varepsilon_0)$:

- if $\phi_* \mathcal{E}$ is L_0 -stable, then \mathcal{E} (resp. $\psi_* \mathcal{E}$) is $L_{-\varepsilon}$ -stable on X (resp. L_ε -stable on X');
- and if $\phi_* \mathcal{E}$ is L_0 -unstable, then \mathcal{E} (resp. $\psi_* \mathcal{E}$) is $L_{-\varepsilon}$ -unstable on X (resp. L_ε -unstable on X').

We now consider the case where $\phi_* \mathcal{E}$ is L_0 -semistable. We first observe, as in the (un)stable case, that there is $\varepsilon_1 > 0$, such that for all $\varepsilon \in (0, \varepsilon_1)$ and for all $\mathcal{F} \in \text{Obj}(\mathfrak{Rcf}^T(X_0))$ with $\mathcal{F} \subsetneq \phi_* \mathcal{E}$ such that $\mu_{L_0}(\mathcal{F}) < \mu_{L_0}(\phi_* \mathcal{E})$ one has

$$\mu_{L_{-\varepsilon}}((\phi^* \mathcal{F})^{\vee\vee}) < \mu_{L_{-\varepsilon}}(\mathcal{E}) \quad \text{and} \quad \mu_{L_\varepsilon}(((\phi')^* \mathcal{F})^{\vee\vee}) < \mu_{L_\varepsilon}(\psi_* \mathcal{E}).$$

If $\mathcal{F} \subsetneq \phi_* \mathcal{E}$ is a sub object such that $\mu_{L_0}(\mathcal{F}) = \mu_{L_0}(\phi_* \mathcal{E})$, then there is a Jordan-Hölder filtration

$$0 = \mathcal{E}_1 \subseteq \dots \subseteq \mathcal{E}_l = \phi_* \mathcal{E}$$

with $l \geq 1$ such that $\mathcal{F} = \mathcal{E}_i$ for some $i \in \{1, \dots, l\}$ (see [11, Section 1.6]) and we deduce that $\mathcal{F} \in \mathfrak{E}$. Therefore, from the expansions of the slopes, to get the points (iii) and (iv) of the theorem, it suffices to compare

$$\frac{c_1(\mathcal{E}) \cdot D_+ \cdot (\phi^* L_0)^{n-2}}{\text{rank}(\mathcal{E})} \quad \text{and} \quad \frac{c_1((\phi^* \mathcal{F})^{\vee\vee}) \cdot D_+ \cdot (\phi^* L_0)^{n-2}}{\text{rank}(\mathcal{F})}$$

for any $\mathcal{F} \in \mathfrak{E}$. By uniqueness of the reflexive hull of the graded object of a Jordan–Hölder filtration ([11, Theorem 1.6.7]), \mathfrak{E} is finite, and the result follows. \square

Remark 4.6. In the proof, we have shown that for any $\rho \in \Sigma(1)$,

$$D_\rho \cdot D_+ \cdot (\phi^* L_0)^{n-2} = D'_\rho \cdot D'_+ \cdot ((\phi')^* L_0)^{n-2}$$

If $\ell \geq 2$, the equality

$$D_\rho \cdot (D_+)^{\ell} \cdot (\phi^* L_0)^{n-1-\ell} = D'_\rho \cdot (D'_+)^{\ell} \cdot ((\phi')^* L_0)^{n-1-\ell}$$

is no longer true in general (see for example Equations (4.5) and (4.6) in Section 4.3). The arguments used to prove Theorem 4.4 are very close to those used in [18]. One should be careful though that the results from [18] do not imply directly Theorem 4.4, as X_0 is not \mathbb{Q} -factorial.

Remark 4.7. While the case when $\phi_* \mathcal{E}$ is semistable on X_0 is not fully covered by Theorem 4.4, items (iii) and (iv), one can easily adapt the numerical criterion of [18, Theorem 1.3] to take into account higher order terms in the ε -expansions of the $L_{-\varepsilon}$ and L_ε slopes, and obtain a full description of the stability behaviour of \mathcal{E} in terms of that of $\phi_* \mathcal{E}$, for the considered polarisations.

Actually, Theorem 4.4 holds for some specific flat families of toric sheaves. We recall that the *characteristic function* χ of an equivariant reflexive sheaf \mathcal{F} with family of filtrations $(F, \{F^\rho(j)\})$ is the function

$$\begin{aligned} \chi(\mathcal{F}) : M &\longrightarrow \mathbb{Z}^{\#\Sigma(n)} \\ m &\longmapsto \left(\dim \left(\bigcap_{\rho \in \sigma(1)} F^\rho(\langle m, u_\rho \rangle) \right) \right)_{\sigma \in \Sigma(n)}. \end{aligned}$$

Let S be a scheme of finite type over \mathbb{C} and $\mathcal{E} = (\mathcal{E}_s)_{s \in S}$ be an S -family of equivariant reflexive sheaves over X (see [18, Section 3.5] for more details). We denote by $(E_s, E_s^\rho(i))$ the family of filtrations of \mathcal{E}_s . There is a collection of increasing filtrations of reflexive sheaves

$$(\mathcal{F}, \{\mathcal{F}_m^\rho : m \in M\}_{\rho \in \Sigma(1)})$$

such that for any $s \in S$ and all $m \in M$,

$$E_s = \mathcal{F}(s) \quad \text{and} \quad E_s^\rho(\langle m, u_\rho \rangle) = \mathcal{F}_m^\rho(s)$$

where $\mathcal{F}(s)$ and $\mathcal{F}_m^\rho(s)$ are respectively the fibers of \mathcal{F} and \mathcal{F}_m^ρ at s .

Lemma 4.8. *Let X be a toric variety given by a simplicial fan Σ and let $\mathcal{E} = (\mathcal{E}_s)_{s \in S}$ be an S -family of equivariant reflexive sheaves over X such that:*

- (1) \mathcal{E} is locally free over $X \times S$, or
- (2) the map $s \mapsto \chi(\mathcal{E}_s)$ is constant.

Then, for all $\rho \in \Sigma(1)$ and $j \in \mathbb{Z}$, the map $s \mapsto \dim(E_s^\rho(j))$ is constant.

Proof. If \mathcal{E} is locally free over $X \times S$, by [19, Proposition 3.13] (Klyachko’s compatibility condition for S -families of locally free sheaves), for any $\sigma \in \Sigma(n)$, there is a multiset $A_\sigma \subseteq M$ of size $\text{rank}(\mathcal{E})$ such that for any $m \in M$, \mathcal{F}_m^ρ is a locally free sheaf of rank

$$|\{\alpha \in A_\sigma : \langle \alpha, u_\rho \rangle \leq \langle m, u_\rho \rangle\}|.$$

As for any $s \in S$ and $m \in M$, $\dim(\mathcal{F}_m^\rho(s)) = \text{rank}(\mathcal{F}_m^\rho)$, we deduce that the map

$$s \longmapsto \dim(E_s^\rho(\langle m, u_\rho \rangle))$$

is constant.

We now assume that the map $s \mapsto \chi(\mathcal{E}_s)$ is constant. For any $\rho \in \Sigma(1)$, we denote by i_ρ the smallest integer such that for any $j \geq i_\rho$ and any $s \in S$,

$$E_s^\rho(j) = E_s.$$

Let $\sigma \in \Sigma(n)$. The set $\{u_\rho : \rho \in \sigma(1)\}$ is a \mathbb{Q} -basis of $N \otimes_{\mathbb{Z}} \mathbb{Q}$; we denote by $\{u_\rho^* : \rho \in \sigma(1)\}$ its dual basis. For any $\rho' \in \sigma(1)$, there is $m' \in M$, such that $\langle m', u_{\rho'} \rangle = j$. Let $m \in M$ be given by

$$m = m' + \sum_{\rho \in \sigma(1) \setminus \{\rho'\}} a_\rho u_\rho^*$$

where for any $\rho \in \sigma(1) \setminus \{\rho'\}$, $a_\rho \in \mathbb{Z}$ satisfies $a_\rho u_\rho^* \in M$ and $a_\rho + \langle m', u_\rho \rangle \geq i_\rho$. By construction of m , one has

$$\bigcap_{\rho \in \sigma(1)} E_s^\rho(\langle m, u_\rho \rangle) = E_s^{\rho'}(j).$$

As $s \mapsto \chi(\mathcal{E}_s)$ is constant, we deduce that the map $s \mapsto \dim(E_s^{\rho'}(j))$ is constant for any $\rho' \in \sigma(1)$ and any $j \in \mathbb{Z}$. \square

If \mathcal{E} is an S -family of equivariant reflexive sheaves on X which satisfies the conditions of Lemma 4.8, then by [18, Lemma 3.12], for any ample \mathbb{Q} -Cartier divisor L on X , the set

$$\{\mu_L(\mathcal{G}_s) : s \in S, \mathcal{G}_s \text{ is an equivariant and saturated reflexive subsheaf of } \mathcal{E}_s\}$$

is finite. Therefore, in that case, the ε_0 in Theorem 4.4 can be taken uniformly for $(\mathcal{E}_s)_{s \in S}$.

4.3. An example. We denote by (e_1, e_2, e_3) the standard basis of \mathbb{Z}^3 . Let

$$u_1 = e_1, \quad u_2 = e_1 + e_2 - e_3, \quad u_3 = e_2, \quad u_4 = e_3, \quad u_0 = -(e_1 + e_2 + e_3)$$

and Σ_0 be a fan in \mathbb{R}^3 given by

$$\Sigma_0 = \{\text{Cone}(u_1, u_2, u_3, u_4)\} \cup \bigcup_{i=1}^4 \{\text{Cone}(A) : A \subseteq \{u_0, u_i, u_{i+1}\}\}$$

where $u_5 = u_1$. We denote by σ_0 the flipping cone $\text{Cone}(u_1, u_2, u_3, u_4)$; we have

$$u_2 + u_4 - u_1 - u_3 = 0.$$

Let

$$\Sigma = (\Sigma_0 \setminus \{\sigma_0\}) \cup \Sigma_+ \quad \text{and} \quad \Sigma' = (\Sigma_0 \setminus \{\sigma_0\}) \cup \Sigma_-$$

where

$$\Sigma_+ = \{\text{Cone}(u_j : j \in J) : J \subset \{1, \dots, 4\} \text{ and } \{1, 3\} \not\subseteq J\} \quad \text{and}$$

$$\Sigma_- = \{\text{Cone}(u_j : j \in J) : J \subset \{1, \dots, 4\} \text{ and } \{2, 4\} \not\subseteq J\}.$$

We denote by D_i the torus invariant divisor associated to the ray $\text{Cone}(u_i)$. By using (4.3) with $m \in \{e_1, e_2, e_3\}$, we get the following linear equivalences of divisors on X_0 , X and X' :

$$D_1 \sim_{\text{lin}} D_3 \sim_{\text{lin}} D_0 - D_2 \quad \text{and} \quad D_4 \sim_{\text{lin}} D_0 + D_2.$$

By [6, Theorem 4.2.8.(d)], the divisor D_0 generates the set of invariant Cartier divisors of X_0 . As Σ (resp. Σ') is simplicial, by [6, Proposition 4.2.7] any invariant divisor of X (resp. X') is \mathbb{Q} -Cartier.




FIGURE 1. Fans of varieties given in Section 4.3

$\mathbb{Z}u_0 + \mathbb{Z}u_1 + \mathbb{Z}u_2$	$\mathbb{Z}u_0 + \mathbb{Z}u_2 + \mathbb{Z}u_3$	$\mathbb{Z}u_0 + \mathbb{Z}u_3 + \mathbb{Z}u_4$	$\mathbb{Z}u_0 + \mathbb{Z}u_4 + \mathbb{Z}u_1$
2	2	1	1

$\mathbb{Z}u_2 + \mathbb{Z}u_4 + \mathbb{Z}u_1$	$\mathbb{Z}u_2 + \mathbb{Z}u_4 + \mathbb{Z}u_3$	$\mathbb{Z}u_1 + \mathbb{Z}u_3 + \mathbb{Z}u_2$	$\mathbb{Z}u_1 + \mathbb{Z}u_3 + \mathbb{Z}u_4$
1	1	1	1

TABLE 1. Sublattices and their index in \mathbb{Z}^3

Lemma 4.9 (Intersections of divisors).

(1) On X and X' , we have:

$$\begin{aligned}
 D_1 \cdot D_0^2 &= \frac{1}{2} & D_2 \cdot D_0^2 &= \frac{1}{4} & D_4 \cdot D_0^2 &= 1 & D_0 \cdot D_0^2 &= \frac{3}{4} \\
 D_1 \cdot (D_0 \cdot D_2) &= \frac{1}{2} & D_4 \cdot (D_0 \cdot D_2) &= 0 & D_2 \cdot (D_0 \cdot D_2) &= -\frac{1}{4}.
 \end{aligned}$$

(2) On X , we have

$$D_1 \cdot D_2^2 = \frac{1}{2} \quad D_4 \cdot D_2^2 = -1 \quad D_0 \cdot D_2^2 = \frac{-1}{4} \quad D_2 \cdot D_2^2 = \frac{-3}{4}.$$

(3) On X' , we have

$$D_1 \cdot D_2^2 = \frac{-1}{2} \quad D_4 \cdot D_2^2 = 0 \quad D_0 \cdot D_2^2 = \frac{-1}{4} \quad D_2 \cdot D_2^2 = \frac{1}{4}.$$

Proof. The lemma follows from Formula (4.2) and Table 1. We show the first point to illustrate the computations, the other intersection numbers follow in the same way.

$$\begin{aligned}
 D_1 \cdot D_0^2 &= D_1 \cdot D_0 \cdot (D_2 + D_3) = D_1 \cdot D_0 \cdot D_2 + D_1 \cdot D_0 \cdot D_3 = \frac{1}{2} \\
 D_2 \cdot D_0^2 &= \frac{1}{2} D_2 \cdot D_0 \cdot (D_3 + D_4) = \frac{1}{4} \\
 D_4 \cdot D_0^2 &= D_4 \cdot D_0 \cdot (D_2 + D_3) = 1
 \end{aligned}$$

$$D_0 \cdot D_0^2 = \frac{1}{2}(D_3 + D_4) \cdot D_0^2 = \frac{1}{4} + \frac{1}{2} = \frac{3}{4}.$$

□

We assume that $\mathcal{E} = \mathcal{T}_X$. By [7, Corollary 2.2.17], the family of filtrations of the tangent sheaf \mathcal{T}_X of X is given by:

$$E^p(j) = \begin{cases} 0 & \text{if } j < -1 \\ \text{Span}(u_\rho) & \text{if } j = -1 \\ N \otimes_{\mathbb{Z}} \mathbb{C} & \text{if } j > -1 \end{cases}.$$

In this case, the inequalities of Proposition 4.2 become

$$\frac{1}{\dim(F)} \sum_{u_\rho \in F} \deg_L(D_\rho) \stackrel{(\leq)}{<} \frac{1}{n} \sum_{\rho \in \Sigma(1)} \deg_L(D_\rho).$$

Let $L_0 = D_0$ be an ample Cartier divisor on X_0 . We have

$$\mu_L(\mathcal{E}) = 1 = \frac{1}{\dim(F_1)} \sum_{u_\rho \in F_1} \deg_L(D_\rho) = \frac{1}{\dim(F_2)} \sum_{u_\rho \in F_2} \deg_L(D_\rho)$$

where $F_1 = \text{Span}(u_4)$ and $F_2 = \text{Span}(u_0, u_2, u_4)$. As for any vector subspace $F \subsetneq E$ such that $F \notin \{F_1, F_2\}$, one has

$$\frac{1}{\dim(F)} \sum_{u_\rho \in F} \deg_L(D_\rho) < 1,$$

we deduce that \mathcal{E} is semistable. We denote by \mathcal{F} the subsheaf of \mathcal{E} given by the family of filtrations $(F_1, F_1 \cap E^p(i))$.

Let

$$D_+ = D_2 + D_4.$$

We have

$$D_+ \sim_{\text{lin}} D_0 + 2D_2.$$

So

$$L_{-\varepsilon} = \phi^* L_0 - \varepsilon D_+ \sim_{\text{lin}} (1 - \varepsilon) D_0 - 2\varepsilon D_2 = (1 - \varepsilon) \left(D_0 - \frac{2\varepsilon}{1 - \varepsilon} D_2 \right)$$

and

$$L_\varepsilon = (\phi')^* L_0 + \varepsilon D_+ \sim_{\text{lin}} (1 + \varepsilon) D_0 + 2\varepsilon D_2 = (1 + \varepsilon) \left(D_0 + \frac{2\varepsilon}{1 + \varepsilon} D_2 \right).$$

According to Lemma 4.9, on X we have

$$(4.5) \quad \begin{aligned} \deg_{L_{-\varepsilon}}(D_1) &= \frac{1}{2} - 3\varepsilon + \frac{9}{2}\varepsilon^2 & \deg_{L_{-\varepsilon}}(D_2) &= \frac{1}{4} + \frac{1}{2}\varepsilon - \frac{15}{4}\varepsilon^2 \\ \deg_{L_{-\varepsilon}}(D_4) &= 1 - 2\varepsilon - 3\varepsilon^2 & \deg_{L_{-\varepsilon}}(D_0) &= \frac{3}{4} - \frac{5}{2}\varepsilon + \frac{3}{4}\varepsilon^2; \end{aligned}$$

and on X' we have

$$(4.6) \quad \begin{aligned} \deg_{L_{-\varepsilon}}(D_1) &= \frac{1}{2} + 3\varepsilon + \frac{1}{2}\varepsilon^2 & \deg_{L_{-\varepsilon}}(D_2) &= \frac{1}{4} - \frac{1}{2}\varepsilon + \frac{1}{4}\varepsilon^2 \\ \deg_{L_{-\varepsilon}}(D_4) &= 1 + 2\varepsilon + \varepsilon^2 & \deg_{L_{-\varepsilon}}(D_0) &= \frac{3}{4} + \frac{5}{2}\varepsilon + \frac{3}{4}\varepsilon^2. \end{aligned}$$

By Formulas (4.5), we get

$$\mu_{L_{-\varepsilon}}(\mathcal{E}) = 1 - \frac{10}{3}\varepsilon + \varepsilon^2 \quad \text{and} \quad \mu_{L_{-\varepsilon}}(\mathcal{F}) = 1 - 2\varepsilon - 3\varepsilon^2.$$

Hence, there is ε_0 such that for any $\varepsilon \in (0, \varepsilon_0) \cap \mathbb{Q}$, the tangent sheaf \mathcal{T}_X is $L_{-\varepsilon}$ -unstable. By Formulas (4.6), we have

$$\mu_{L_{-\varepsilon}}(\mathcal{E}) = 1 + \frac{10}{3}\varepsilon + \varepsilon^2 \quad \text{and} \quad \mu_{L_{-\varepsilon}}(\mathcal{F}) = 1 + 2\varepsilon + \varepsilon^2.$$

Hence, there is ε_0 such that for any $\varepsilon \in (0, \varepsilon_0) \cap \mathbb{Q}$, the tangent sheaf $\mathcal{T}_{X'}$ is L_{ε} -stable.

5. FLIPS AND STABILITY FOR LOGARITHMIC SUBCATEGORIES

Consider a toric flip

$$\begin{array}{ccc} X & \overset{\psi}{\dashrightarrow} & X' \\ & \searrow \phi & \swarrow \phi' \\ & X_0 & \end{array}$$

as in Section 2. Fix $\Delta \subset \Sigma(1)$, and introduce the divisor

$$D := \sum_{\rho \in \Delta} D_\rho$$

seen as a divisor on X , X' and X_0 . We consider the equivalent categories $\mathfrak{R}ef^T(X, D)$ and $\mathfrak{R}ef^T(X', D)$ as in Section 3.3. We will say that

$$\psi_* : \mathfrak{R}ef^T(X, D) \rightarrow \mathfrak{R}ef^T(X', D)$$

preserves polystability for the pair of polarisations $(\alpha, \alpha') \in \text{Pic}(X)_{\mathbb{Q}} \times \text{Pic}(X')_{\mathbb{Q}}$ if for any $\mathcal{E} \in \text{Obj}(\mathfrak{R}ef^T(X, D))$, \mathcal{E} is α -polystable if and only if $\psi_*\mathcal{E}$ is α' -polystable.

Theorem 5.1. *The following assertions are equivalent, for a pair of ample classes $(\alpha, \alpha') \in \text{Pic}(X)_{\mathbb{Q}} \times \text{Pic}(X')_{\mathbb{Q}}$.*

- (i) *The functor $\psi_* : \mathfrak{R}ef^T(X, D) \rightarrow \mathfrak{R}ef^T(X', D)$ preserves polystability for (α, α') .*
- (ii) *There is $c \in \mathbb{Q}_{>0}$ such that for all $\mathcal{E} \in \mathfrak{R}ef^T(X, D)$, $\mu_\alpha(\mathcal{E}) = c \mu_{\alpha'}(\psi_*\mathcal{E})$.*
- (iii) *There is $c \in \mathbb{Q}_{>0}$ such that for all $\rho \notin \Delta$, $\deg_\alpha D_\rho = c \deg_{\alpha'} D'_\rho$.*

Proof. Recall the formula

$$\mu_\alpha(\mathcal{E}) = -\frac{1}{\text{rank}(\mathcal{E})} \sum_{\rho \in \Sigma(1)} \iota_\rho(\mathcal{E}) \deg_\alpha(D_\rho)$$

for the slope, with

$$\iota_\rho(\mathcal{E}) := \sum_{i \in \mathbb{Z}} i (\dim(E^\rho(i)) - \dim(E^\rho(i-1)))$$

where $(E, E^\rho(\bullet))_{\rho \in \Sigma(1)}$ stands for the family of filtrations associated to \mathcal{E} . Then, by definition of the logarithmic category $\mathfrak{R}ef^T(X, D)$, for any $\mathcal{E} \in \text{Obj}(\mathfrak{R}ef^T(X, D))$, and any $\rho \in \Delta$, we have

$$\iota_\rho(\mathcal{E}) = 0,$$

so the slope reads

$$\mu_\alpha(\mathcal{E}) = -\frac{1}{\text{rank}(\mathcal{E})} \sum_{\rho \notin \Delta} \iota_\rho(\mathcal{E}) \deg_\alpha(D_\rho).$$

Note also that by construction, for any ρ ,

$$\iota_\rho(\mathcal{E}) = \iota_\rho(\psi_*\mathcal{E}).$$

Then, $(iii) \Rightarrow (ii) \Rightarrow (i)$ is straightforward. To prove $(i) \Rightarrow (iii)$, we argue as in [4, proof of Proposition 4.8], and consider for any pair $(\rho_1, \rho_2) \in \Sigma(1)^2$ the polystable toric sheaf

$$\mathcal{E} = \mathcal{O}_X(d \deg_\alpha(D_{\rho_2})D_{\rho_1}) \oplus \mathcal{O}_X(d \deg_\alpha(D_{\rho_1})D_{\rho_2}),$$

where d is the common denominator of $\deg_\alpha(D_{\rho_2})$ and $\deg_\alpha(D_{\rho_1})$. Its image by ψ_* is

$$\psi_*\mathcal{E} = \mathcal{O}_{X'}(d \deg_\alpha(D_{\rho_2})D'_{\rho_1}) \oplus \mathcal{O}_{X'}(d \deg_\alpha(D_{\rho_1})D'_{\rho_2}).$$

As $\mathcal{E} \in \text{Obj}(\mathfrak{Rcf}^T(X, D))$ (see e.g. [7, Example 2.2.13] for the family of filtrations of rank one toric sheaves), if ψ preserves polystability, we must have

$$\deg_\alpha(D_{\rho_2}) \deg_{\alpha'}(D'_{\rho_1}) = \deg_\alpha(D_{\rho_1}) \deg_{\alpha'}(D'_{\rho_2}).$$

The result follows. \square

Remark 5.2. The reason for considering logarithmic subcategories is the following. If one considers the full $\mathfrak{Rcf}^T(X)$, and assume that ψ_* preserves polystability for all $\mathcal{E} \in \text{Obj}(\mathfrak{Rcf}^T(X, D))$, then by (iii) of Theorem 5.1, for any $\rho \in \Sigma(1)$, one must have

$$\deg_\alpha D_\rho = c \deg_{\alpha'} D'_\rho.$$

Up to scale, we can assume $c = 1$. But then, a result due to Minkowski ([22, p.455]), translated into the toric setting in [4, Proposition 5.3 and Corollary 5.4], implies that the polytope associated to (X, α) equals the polytope associated to (X', α') , which is absurd, as X and X' are not isomorphic.

REFERENCES

- [1] Shigetoshi Bando and Yum-Tong Siu. Stable sheaves and Einstein-Hermitian metrics. *Geometry and analysis on complex manifolds. Festschrift for Professor S. Kobayashi's 60th birthday*, pages 39–59, 1994. [1](#)
- [2] Victor V. Batyrev. Dual polyhedra and mirror symmetry for Calabi-Yau hypersurfaces in toric varieties. *J. Algebr. Geom.*, 3(3):493–535, 1994. [2](#)
- [3] Philip Candelas, Paul S. Green, and Tristan Hübsch. Rolling among Calabi-Yau vacua. *Nuclear Physics B*, 330(1):49–102, 1990. [2](#)
- [4] Andrew Clarke and Carl Tipler. Equivariant stable sheaves and toric GIT. *Proceedings of the Royal Society of Edinburgh: Section A Mathematics*, pages 1–32, 2022. [1](#), [10](#), [19](#)
- [5] Tristan C. Collins, Sebastien Picard, and Shing-Tung Yau. Stability of the tangent bundle through conifold transitions. *Preprint ArXiv 2102.11170*, 2021. [2](#)
- [6] David A. Cox, John B. Little, and Henry K. Schenck. *Toric Varieties*. Graduate Studies in Mathematics. American Mathematical Soc., 2011. [1](#), [4](#), [5](#), [6](#), [7](#), [11](#), [12](#), [15](#)
- [7] J. Dasgupta, A. Dey, and B. Khan. Stability of equivariant vector bundle over toric varieties. *Documenta Mathematica*, 25: 1787–1833, 2020. [17](#), [19](#)
- [8] William Fulton. *Intersection Theory*. Springer New York, 1984. [13](#)
- [9] William Fulton. *Introduction to toric varieties*. Annals of Mathematics Studies. Princeton Univ. Press, Princeton, NJ, 1993. [11](#)
- [10] Robin Hartshorne. Stable reflexive sheaves. *Mathematische Annalen*, 254: 121 – 176, 1980. [8](#)
- [11] Daniel Huybrechts and Manfred Lehn. *The Geometry of Moduli Spaces of Sheaves*. Cambridge Mathematical Library. Cambridge University Press, 2 edition, 2010. [11](#), [13](#), [14](#)
- [12] Alexander Klyachko. Equivariant bundle on toral varieties. *Mathematics of the USSR-Izvestiya*, 35(2): 337 – 375, 1990. [7](#), [8](#), [9](#)
- [13] Allen Knutson and Eric Sharpe. Sheaves on toric varieties for physics. *Adv. Theor. Math. Phys.*, 2(4):873–961, 1998. [1](#)

- [14] Shoshichi Kobayashi. *Differential Geometry of Complex Vector Bundles*. Publications of the Mathematical Society of Japan. Princeton University Press, 1987. [1](#)
- [15] Martijn Kool. Fixed point loci of moduli spaces of sheaves on toric varieties. *Advances in Mathematics*, 227(4): 1700 – 1755, 2011. [10](#)
- [16] David Mumford. Projective invariants of projective structures and applications. *Proc. Internat. Congr. Mathematicians*, pages 526–530, 1963. [1](#)
- [17] Achim Napame. Stability of equivariant logarithmic tangent sheaves on toric varieties of Picard rank two. *To appear in “Annales de la faculté des sciences de Toulouse”*, 2023. [9](#)
- [18] Achim Napame and Carl Tipler. Toric sheaves, stability and fibrations. *Journal of Pure and Applied Algebra*, 228(3):107494, 2024. [1](#), [8](#), [10](#), [13](#), [14](#), [15](#)
- [19] Sam Payne. Moduli of toric vector bundles. *Compositio Mathematica*, 144(5):1199–1213, 2008. [14](#)
- [20] Markus Perling. Graded rings and equivariant sheaves on toric varieties. *Mathematische Nachrichten*, 263/264: 181–197, 2004. [7](#), [8](#)
- [21] Miles Reid. The moduli space of 3-folds with $K = 0$ may nevertheless be irreducible. *Math. Ann.*, 278:329–334, 1987. [2](#)
- [22] Rolf Schneider. *Convex Bodies: the Brunn-Minkowski Theory*, volume 44 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, 2nd. edition, 2013. [19](#)
- [23] Fumio Takemoto. Stable vector bundles on algebraic surfaces. *Nagoya Math. J.*, 47: 29 – 48, 1972. [1](#)