



**HAL**  
open science

# Cobalt-Catalyzed Reductive Cross-Coupling To Construct Csp<sup>3</sup>–Csp<sup>3</sup> Bonds via Csp<sup>3</sup>–S and Csp<sup>3</sup>–X Bonds Activation

Meng-Yu Gao, Corinne Gosmini

► **To cite this version:**

Meng-Yu Gao, Corinne Gosmini. Cobalt-Catalyzed Reductive Cross-Coupling To Construct Csp<sup>3</sup>–Csp<sup>3</sup> Bonds via Csp<sup>3</sup>–S and Csp<sup>3</sup>–X Bonds Activation. *Organic Letters*, 2023, 25 (42), pp.7689-7693. 10.1021/acs.orglett.3c02993 . hal-04262030

**HAL Id: hal-04262030**

**<https://hal.science/hal-04262030>**

Submitted on 27 Oct 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Cobalt-Catalyzed Reductive Cross-Coupling to Construct Csp<sup>3</sup>-Csp<sup>3</sup> Bonds via Csp<sup>3</sup>-S and Csp<sup>3</sup>-X Bonds Activation

Meng-Yu Gao,<sup>a</sup> Corinne Gosmini<sup>a\*</sup>


Laboratoire de Chimie Moléculaire, École Polytechnique, Institut Polytechnique de Paris, CNRS Route de Saclay 91128 Palaiseau cedex, France

Email: corinne.gosmini@polytechnique.edu; [mengyu.gao@polytechnique.edu](mailto:mengyu.gao@polytechnique.edu)

Supporting Information Placeholder


**ABSTRACT:** A cross-electrophilic coupling of benzyl sulfonium salts with alkyl halides forming Csp<sup>3</sup>-Csp<sup>3</sup> bonds is described using a Co-based catalytic system. The activation of the stable Csp<sup>3</sup>-S bond of benzyl sulfonium salts under mild reaction conditions leads to various functionalized alkyl derivatives. Preliminary mechanistic studies suggest the involvement of alkyl radicals both formed from alkyl halides and benzyl sulfoniums through a single electron transfer.

The construction of carbon-carbon bonds is important in synthetic organic chemistry but also in materials and biology. Thus, transition metal-catalyzed cross-coupling reactions have been extensively studied over the past decades and recognized with the Nobel Prize in Chemistry in 2010 for the development of palladium-catalyzed cross-coupling reactions.<sup>1</sup> Most of the cross-coupling reactions imply a nucleophilic organometallic reagent, such as for example Grignard reagents, organozinc, or organoboron derivatives, and an electrophile in the presence of a transition metal.<sup>1a,2</sup> However, to avoid the need or preformation of stoichiometric, reactive, and often sensitive organometallic species, especially when a reactive functional group is present, electrophilic reductive cross-coupling reactions have been developed since the beginning of this century to form a C-C bond in only one step. This interesting procedure exhibits high functional group compatibility, regioselectivity, and chemoselectivity. In most cases, these cross-coupling reactions involve organohalides as electrophiles.<sup>3</sup> However, in order to reduce halogen wastes, other electrophiles such as C-O (alcohol derivatives: sulfonates,<sup>4</sup> esters,<sup>5</sup> carbonates,<sup>5d,6</sup> oxalates<sup>7</sup>, etc.), C-N (amine derivatives, such as pyridinium salts,<sup>8</sup> ammonium salts,<sup>9</sup> etc.), C-S (sulfonium salts<sup>10</sup>), as well as carboxylic acid derivatives (N-hydroxy phthalimide esters<sup>11</sup>) have been more recently involved in Ni, Cu or Co catalyzed cross-couplings. Moreover, although the formation of Csp<sup>2</sup>-Csp<sup>2</sup> bonds between two Csp<sup>2</sup> electrophiles is well established,<sup>3g,12</sup> the Csp<sup>3</sup>-Csp<sup>3</sup> bond formation from two Csp<sup>3</sup> electrophiles remains much more challenging while these bonds are ubiquitous in molecules. Indeed, the unwanted facile  $\beta$ -hydride elimination from alkyl metal complexes is a key impediment to efficient cross-coupling of alkyl electrophiles. In the case of Co-catalyzed cross-couplings involving benzylic substrates such as benzyl chloride<sup>3d</sup>, benzyl zinc<sup>13</sup>, and benzyl mesylates<sup>4b</sup>, Csp<sup>2</sup> partners were always employed. Howev-



er, some electrophilic cross-couplings have been already developed between two different alkyl compounds using Ni or Co as catalysts<sup>3b, c, 14</sup> and notably benzyl sulfonium salts.<sup>15</sup> In particular, Aleman and our group showed recently that these compounds could react by reductive cross-coupling with activated alkyl halides.<sup>16</sup> Aleman reported the cross-coupling with benzyl bromides using a nickel catalyst<sup>16a</sup> (**Scheme 1a**) while allylic acetates or ethers were used by our group under cobalt catalysis<sup>16b</sup>. Nevertheless, compared to the widely used C-O or C-N electrophiles, sulfonium salts still did not receive much attention in cross-coupling reactions since Liebeskind's reports<sup>10a, b</sup>, although S-containing motifs are very present in drugs and materials.<sup>17</sup> These salts are bench-stable and can be synthesized easily and economically from alcohols, organohalides, thiols, or thioethers.<sup>15a, b</sup> They can smoothly undergo oxidative addition with transition metal catalysts and release a neutral sulfide instead of anionic sulfur fragments limiting the risk of catalyst poisoning. In the development and application of sulfonium salts in cross-coupling, different elegant reactions were reported such as C-H arylation<sup>18</sup>, alkylation/ alkenylation<sup>19</sup>, and carbonylation<sup>20</sup>. Hence, considering that no cross-electrophilic coupling was reported between non-activated alkyl halides and benzyl sulfonium salts, regardless of the catalyst, we envisioned the Co-catalyzed reductive cross-coupling of benzyl sulfonium salts with non-activated alkyl iodides and the extension to activated benzyl chlorides (**Scheme 1b**).


**Scheme 1.** Reductive cross-couplings of benzyl sulfonium salts with alkyl halides.



The Co-catalyzed cross electrophilic-coupling of benzyl sulfonium salts (**S1**) with an alkyl iodide such as 1-iodoheptane (**a1**) led to a 41% GC yield of the desired product (1-phenyloctane) **b1** in allylation conditions we previously reported. (see SI for more details).

Then, numerous experiments were conducted to improve this cross-electrophilic coupling (see SI for more details). As shown in **Table 1**, the combination of 10 mol% of CoBr<sub>2</sub>, 2.0 equivalents of Mn activated by 25 mol% of TMSCl, with addition of 75 mol% of pyridine in a mixture of MeCN/ 1,4-dioxane led to a 60% GC yield of the desired product **b1** at 45 °C overnight. With MeCN or 1,4-dioxane as the sole solvent, yields of 1-phenyloctane **b1** decreased to 48% or 11% respectively, but the formation of the homo-coupling from **a1** was significantly inhibited in sole 1,4-dioxane (**Table 1, entries 2, 3**). In the mixture of MeCN and 1,4-dioxane, the quantity of 1,4-dioxane, is importante (**Table 1, entries 4, 5**). With other solvents in place of MeCN, like THF, DMA, and MeOH, the yield of **b1** was lower. (**Table 1, entries 6-8**). Decreasing the amount of CoBr<sub>2</sub> to 5% gave a similar yield of **b1** (62% GC yield, **Table 1, entry 9**). In this condition, decreasing the amount of Mn or using Zn instead of Mn affected the efficiency of the cross-coupling significantly (**Table 1, entries 10, 11**). Moreover, at a higher or lower temperature (60 °C or rt), a low yield of **b1** was obtained together with more dibenzyl or remaining **a1** (**Table 1, entry 12, 13**). However, extending the reaction time to 24 h at rt, the yield of **b1** increased to 73% (**Table 1, entry 14**). Cobalt salt and Mn were necessary for this alkylation reaction but a of **b1** was formed (33% GC yield) even in the absence of pyridine. Scaling up the reaction to 10 mmol led to 78 % GC yield (65% isolated yield). Finally, no reaction occurred upon replacing CoBr<sub>2</sub> by NiBr<sub>2</sub> under the condition (see SI- **Table 3.5**).

**Table 1.** Optimization of Reaction Conditions of benzyl dimethyl sulfonium salt (**S1**) with 1-iodoheptane (**a1**).<sup>a</sup>



Entry	Variation	<b>b1</b> yield/%
1	None	60
2	MeCN (0.5 M)	48
3	1,4-dioxane (0.5 M)	11
4	MeCN/ 1,4-dioxane (3.5 mL: 0.5 mL)	56
5	MeCN/ 1,4-dioxane (2.0 mL: 2.0 mL)	30
6	THF (0.5 M)	36
7	DMA (0.5 M)	Trace
8	MeOH (0.5 M)	25
9	CoBr <sub>2</sub> (5 mol%)	63
11	Mn (1.5 equiv)	11 <sup>b</sup>
11	Zn (2.0 equiv)	20 <sup>b</sup>
12	60 °C	43 <sup>b</sup>
13	rt	40 <sup>b</sup>
14	CoBr <sub>2</sub> (5 mol%), rt, 24 h	73/66 (57 <sup>c</sup> )


<sup>a</sup> Reactions were performed with **S1** (2.0 mmol), **a1** (2.5 mmol), Mn (2.0 equiv), pyridine (75 mol%), TMSCl (25 mol%) in MeCN/1,4-dioxane under N<sub>2</sub>, GC yields using decane as an internal standard. <sup>b</sup> CoBr<sub>2</sub> (5 mol%). <sup>c</sup> Isolated yield.

The nature of the sulfonium partner was modified: benzyl dimethyl sulfonium bromide and 1-benzyl tetrahydro-1H-thiophen-1-ium hexafluorophosphate formed **b1** in respectively 10% and 56% yield under the optimized condition defined in **Table 1**. To compare with other benzylic electrophiles, benzyl chloride gave a similar yield as the benzyl sulfonium salt **S1**, while only a trace amount of **b1** was detected with the more reactive benzyl bromide. Among C-O electrophiles, benzyl carbonate was unreactive, and benzyl sulfonates (mesylate or tosylate) only gave trace or a low yield of the desired coupling product.

With the optimized conditions described in **Table 1** in hand, the scope of benzyl sulfonium tetrafluoroborate salts with various alkyl iodides was studied (**Table 2**). As shown, a large number of substituted non-activated alkyl iodides and functionalized benzyl sulfonium salts reacted under the established conditions with either 25 or 75 mol% of TMSCl. It is obvious that this method is general for primary alkyl iodides, providing a wide range of cross-coupling products in moderate to good isolated yields. For unsubstituted primary linear alkyl iodides, 50% to 69% yields of the cross-coupling products were obtained (**b2-b17**). Alkyl iodides with various functionalities, such as acetate (**b7, b8**), ether (**b9**), phenyl (**b10, b11, b17**), and ester (**b12**), were well tolerated. A functional group could be tolerated both on the benzyl sulfonium salt and the primary alkyl iodide (**b13- b16**). In contrast, cross-couplings of secondary or tertiary alkyl iodides with benzyl sulfonium salt led to only traces or few of products. (**b19-b21**) due to their higher reactivity except for the coupling with cycloalkane such as iodocyclohexane delivering the desired product **b18** in 46% isolated yield.

Although different functionalized groups on the benzyl moiety of sulfonium salts were well tolerated, the coupling of 1-iodoheptane (**a1**) with electron-donating groups substituted benzyl sulfonium salts leading to **b22-b25** was more efficient than with electron-withdrawing groups substituted benzyl sulfonium salts (**b30, b31**) because of the competitive homocoupling of the benzyl sulfonium salt. Halides (F or Cl) also led to cross-coupling products in good yields (**b28, b29**). The steric hindrance on the benzyl sulfonium salt did not affect the cross-coupling (**b22-b24**). Several electron-donating groups on the aromatic ring were also allowed (**b26, b27**).


Finally, we extended the cross-coupling of benzyl sulfonium tetrafluoroborate salts with benzyl chlorides (**Table 2**). In contrast to the work carried out by Aleman<sup>16a</sup>, benzyl chloride showed good and even better efficiency than benzyl bromide in this Co-catalyzed Csp<sup>3</sup>-Csp<sup>3</sup> bond-forming reaction. With a methoxy substituted benzyl chloride, the yield increased significantly compared to the same substituent on the benzyl sulfonium salt (**c1, c2**). These results prompted us to attempt the synthesis of a dihydro resveratrol precursor - a key intermediate in the inhibition of pancreatic oxidative damage<sup>21</sup>. However, the cross-coupling product yields (**c3, c4**) were moderate and similar, regardless of the entity bearing the two methoxy groups.

**Table 2.** Scopes of benzyl dimethyl sulfonium salts (**S**) with alkyl iodides (**a**)/benzyl chlorides. <sup>a b</sup>

<sup>a</sup> Standard condition: **S** (2.0 mmol), **a** (2.5 mmol), CoBr<sub>2</sub> (5 mol%), Mn (2.0 equiv), pyridine (75 mol%), TMSCl (25 or 75 mol%) in MeCN (3.0 mL) and 1,4-dioxane (1.0 mL) at rt, 24 h, N<sub>2</sub>. <sup>b</sup> Isolated yield. <sup>c</sup> GC yield.

To investigate the mechanism of this Co-catalyzed reductive Csp<sup>3</sup>-Csp<sup>3</sup> bond-forming process, a few experiments were conducted. At the outset, the addition of 2 equivalents of 2,2,6,6-tetramethylpiperine-1-oxyl (TEMPO) in the alkylation reaction between **S1** and **a1** inhibited significantly the coupling (**Scheme 2**). Only a trace amount of **b1** was detected by GC-MS, along with trace amounts of the TEMPO-trapped products **1** and **2** (**eq 1**). Moreover, the reaction of either **S1** or **a1** in the presence of TEMPO (**eq 2**), led to traces of 1-(benzyloxy)-2,2,6,6-tetramethyl-piperidin (**1**) or 1-(heptyloxy)-2,2,6,6-tetramethyl-piperidine (**2**) detected by GC-MS as well as either the dimerization product of **S1** (32% GC yield of dibenzyl) or **a1** (16% GC yield of tetradecane) respectively. Likewise, adding glavinoxyl free radical in the reaction between **S1** and **a1** led to a significantly decreased cross-coupling product yield when increasing the amount of free radical reagent (see SI). These results suggested that both benzyl and alkyl radicals were involved in this Co-catalyzed alkylation reaction.


**Scheme 2.** TEMPO trapping experiments with alkyl iodide.



<sup>a</sup> detected by GC-MS. <sup>b</sup> GC yields using decane as an internal standard.

To confirm this hypothesis, iodomethylcyclopropane (**3**) was coupled with **S1** under the optimized alkylation condition (**Scheme 3**, **eq 3**). A mixture of ring-opened products **4** [5-phenylpent-1-ene **4a** and (*E*)-3-pentenyl benzene (**4b**)] was obtained in 33% isolated yield, without unrearranged cross-coupling product (**5**).


**Scheme 3.** Radical clock trapping experiments.



These results prompted us to propose a mechanism (**Scheme 4**) based on our previous reports related to the  $Csp^3-Csp^3$  homocoupling.<sup>22</sup> The cross-coupling of benzyl sulfonium salts and benzyl chlorides suggested a lower reactivity of benzyl sulfonium salts compared to benzyl chlorides. Thus, we proposed that the alkyl radical from the alkyl iodide formed prior to the benzyl radical from sulfonium salt.

First, the catalytic cycle starts with the reduction of the Co(II) pre-catalyst by the manganese into an active low-valent Co species. Then, the oxidative addition of the low-valent Co species with an alkyl radical generated from the alkyl iodide leads to a cobalt complex **A**. After subsequent reduction by Mn, the formed alkyl complex **B** undergoes one electron transfer to benzyl sulfonium salt, delivering Co complex **C**, a benzyl radical and a thioether. Co complex **C** reduced another time by manganese, lead to Co complex **D**. Finally, the benzyl alkyl Co complex **E**, generated by the combination of the benzyl radical and the low-valent Co complex **D**, provides the cross-coupling product by reductive elimination along with the regeneration of the active Co species.

**Scheme 4.** Proposal mechanism



In conclusion, we have demonstrated that benzyl sulfonium salts are suited for the construction of  $Csp^3-Csp^3$  bonds with non-activated alkyl iodides or benzyl chlorides via a Co-catalyzed reductive cross-coupling. Compared to other Co-catalyzed reductive cross-coupling, a mixture of acetonitrile and dioxane as solvent was necessary to obtain best yields. A large variety of functionalized benzyl sulfonium salts reacted with a broad range and small excess of primary or cyclic secondary alkyl iodides in moderate to good yields. To the best of our knowledge, it is the first report using sulfonium salts as electrophiles in metal-catalyzed reductive cross-coupling with non-activated alkyl halides to build  $Csp^3-Csp^3$  bonds. Mechanistic studies suggested both the generations of benzyl and alkyl radicals

from benzyl sulfonium salts and alkyl iodides respectively. Further studies aiming at using other sulfonium salts in cobalt-catalyzed C-C bond formation and gaining detailed mechanistic insights are currently underway in our laboratory.

## ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

brief description (file type, i.e., PDF)

## AUTHOR INFORMATION

### Corresponding Author

\*Corinne Gosmini - Laboratoire de Chimie Moléculaire, École Polytechnique, Institut Polytechnique de Paris, CNRS Route de Saclay 91128 Palaiseau cedex, France.

Email: corinne.gosmini@polytechnique.edu

### Author

Meng-Yu Gao - Laboratoire de Chimie Moléculaire, École Polytechnique, Institut Polytechnique de Paris, CNRS Route de Saclay 91128 Palaiseau cedex, France.

## ACKNOWLEDGMENT

The authors thank Sophie Bourcier from LCM for the HRMS analysis. M.G. also thanks the “China Scholarship Council” for the fellowship. Finally, the authors are very grateful to the CNRS and Institut Polytechnique de Paris, which support this work.

## REFERENCES

- (1) (a) Marsden, J.; Haley, M.; de Meijere, A.; Diederich, F. In *Metal-Catalyzed Cross-Coupling Reactions*, WILEY-VCH: Weinheim, (2004). (b) Knochel, P. In *Handbook of Functionalized Organometallics. Applications in Synthesis*, Wiley-VCH: Weinheim, Germany, (2005). (c) Johansson, Seechurn, C.C.C., Kitching, M.O., Colacot, T.J. Snieckus, V. Palladium-Catalyzed Cross-Coupling: A Historical Contextual Perspective to the 2010 Nobel Prize. *Angew. Chem. Int. Ed.* **2012**, *51*, 5062-5085 (c) Campeau, J.-L.; Hazari, N. Cross-Coupling and Related Reactions: Connecting Past Success to the Development of New Reactions for the Future. *Organometallics* **2019**, *38*, 3-35.
- (2) (a) Liu, C.; Zhang, H.; Shi, W.; Lei, A. Bond Formations between Two Nucleophiles: Transition Metal Catalyzed Oxidative Cross-Coupling Reactions. *Chem. Rev.* **2011**, *111*, 1780-1824. (b) Yu, D. G.; Li, B. J.; Shi, Z. J. Exploration of New C–O Electrophiles in Cross-Coupling Reactions. *Acc. Chem. Res.* **2010**, *43*, 1486–1495. (c) Jana, R.; Pathak, T. P.; Sigman, M. S. Advances in Transition Metal (Pd, Ni, Fe)-Catalyzed Cross-Coupling Reactions Using Alkyl-Organometallics as Reaction Partners. *Chem. Rev.* **2011**, *111*, 1417–1492.
- (3) (a) Lucas, E.; Jarvo, E. Stereospecific and Stereoconvergent Cross-Couplings between Alkyl Electrophiles. *Nat Rev Chem.* **2017**, *1*, 0065. (b) Xu, H.; Zhao, C.; Qian, Q.; Deng, W.; Gong, H. Nickel-Catalyzed Cross-Coupling of Unactivated Alkyl Halides Using Bis (pinacolato) Diboron as Reductant. *Chem. Sci.* **2013**, *4*, 4022-4029. (c) Yu, X.; Yang, T.; Wang, S.; Xu, H.; Gong, H. Nickel-Catalyzed Reductive Cross-Coupling of Unactivated Alkyl Halides. *Org. Lett.* **2011**, *13*, 2138–2141. (d) Pal, S.; Chowdhury, S.; Rozwadowski, E.; Auffrant, A.; Gosmini,

C. Cobalt-Catalyzed Reductive Cross-Coupling Between Benzyl Chlorides and Aryl Halides. *Adv. Synth. Catal.* **2016**, *358*, 2431–2435 (e) Gosmini, C.; Moncomble, A. Cobalt-Catalyzed Cross-Coupling Reactions of Aryl Halides. *Isr. J. Chem.* **2010**, *50*, 568–576. (f) Amatore, M.; Gosmini, C. Direct Method for Carbon–Carbon Bond Formation: The Functional Group Tolerant Cobalt-Catalyzed Alkylation of Aryl Halides. *Chem. Eur. J.*, **2010**, *16*, 5848. (g) Amatore, M.; Gosmini, C. Efficient Cobalt-Catalyzed Formation of Unsymmetrical Biaryl Compounds and Its Application in the Synthesis of a Sartan Intermediate. *Angew. Chem. Int. Ed.* **2008**, *120*, 2119–2122.

(4) Selected reports: (a) Ackerman, L. K. G.; Anka-Lufford, L. L.; Naodovic, M.; Weix, D. J. Cobalt Co-Catalysis for Cross-Electrophile Coupling: Diarylmethanes from Benzyl Mesylates and Aryl Halides. *Chem. Sci.* **2015**, *6*, 1115–1119. (b) Reddy, B. R. P.; Chowdhury, S.; Auffrant, A.; Gosmini, C. Cobalt-Catalyzed Formation of Functionalized Diarylmethanes from Benzylmesylates and Aryl Halides. *Adv. Synth. Catal.* **2018**, *360*, 3026–3029. (c) Komeyama, K.; Michiyuki, T.; Osaka, I. Nickel/Cobalt-Catalyzed C(sp<sup>3</sup>)-C(sp<sup>3</sup>) Cross-Coupling of Alkyl Halides with Alkyl Tosylates. *ACS Catal.* **2019**, *9*, 9285–9291. (e) Liu, J.-H.; Yang C.-T.; Lu X.-Y.; Fu, Y.; Liu, L. Copper-Catalyzed Reductive Cross-Coupling of Nonactivated Alkyl Tosylates and Mesylates with Alkyl and Aryl Bromides. *Chem. Eur. J.* **2014**, *20*, 15334 – 15338. (f) Ma, W.-Y.; Han, G.-Y.; Kang, S.; Pang, X.; Liu, X.-Y.; Shu, X.-Z. Cobalt-Catalyzed Enantiospecific Dynamic Kinetic Cross-Electrophile Vinylation of Allylic Alcohols with Vinyl Triflates. *J. Am. Chem. Soc.* **2021**, *143*, 15930–15935.

(5) (a) Konev, M. O.; Hanna, L. E.; Jarvo, E. R. Intra- and Intermolecular Nickel-Catalyzed Reductive Cross-Electrophile Coupling Reactions of Benzylic Esters with Aryl Halides. *Angew. Chem. Int. Ed.* **2016**, *128*, 6842–6845. (b) Qian, X.; Auffrant, A.; Felouat, A.; Gosmini, C. Cobalt-Catalyzed Reductive Allylation of Alkyl Halides with Allylic Acetates or Carbonates. *Angew. Chem. Int. Ed.* **2011**, *123*, 10586–10589. (c) He, R.-D.; Bai, Y.; Han, G.-Y.; Zhao, Z.-Z.; Pang, X.; Pan, X.; Liu, X.-Y.; Shu, X.-Z. Reductive Alkylation of Alkenyl Acetates with Alkyl Bromides by Nickel Catalysis. *Angew. Chem. Int. Ed.* **2022**, *61*, e202114556. (d) Amatore, M.; Gosmini, C.; Périchon, J. Cobalt-Catalyzed Vinylation of Functionalized Aryl Halides with Vinyl Acetates. *Eur. J. Org. Chem.* **2005**, 989–992.

(6) Chen, H.; Jia, X.; Yu, Y.; Qian, Q.; Gong, H. Nickel-Catalyzed Reductive Allylation of Tertiary Alkyl Halides with Allylic Carbonates. *Angew. Chem. Int. Ed.* **2017**, *56*, 3103–3106.

(7) (a) Yan, X.-B.; Li, C.-L.; Jin, W.-J.; Guo, P.; Shu, X.-Z. Reductive Coupling of Benzyl Oxalates with Highly Functionalized Alkyl Bromides by Nickel Catalysis. *Chem. Sci.* **2018**, *9*, 4529–4534. (b) Ye, Y.; Chen, H.; Sessler, J. L.; Gong, H. Zn-Mediated Fragmentation of Tertiary Alkyl Oxalates Enabling Formation of Alkylated and Arylated Quaternary Carbon Centers. *J. Am. Chem. Soc.* **2019**, *141*, 820–824. (c) Gao, M.; Sun, D.; Gong, H. Ni-Catalyzed Reductive C–O Bond Arylation of Oxalates Derived from  $\alpha$ -Hydroxy Esters with Aryl Halides. *Org. Lett.* **2019**, *21*, 1645–1648. (d) Ye, Y.; Chen, H.; Ma, G.; Gong, H. Iron-Catalyzed Reductive Vinylation of Tertiary Alkyl Oxalates with Activated Vinyl Halides. *Org. Lett.* **2020**, *22*, 2070–2075.

(8) (a) Liao, J.; Basch, C. H.; Hoerrner, M. E.; Talley, M. R.; Boscoe, B. P.; Tucker, J. W.; Garnsey, M. R.; Watson, M. P. Deaminative Reductive Cross-Electrophile Couplings of Alkylpyridinium Salts and Aryl Bromides. *Org. Lett.* **2019**, *21*, 2941–2946. (b) Martin-Montero, R.; Yatham, V. R.; Yin, H.; Davies, J.; Martin, R. Ni-catalyzed Reductive Deaminative Arylation at sp<sup>3</sup> Carbon Centers. *Org. Lett.* **2019**, *21*, 2947–2951. (c) Yue, H.; Zhu, C.; Shen, L.; Geng, Q.; Hock, K. J.; Yuan, T.; Cavallo, L.; Rueping, M. Nickel-Catalyzed C–N Bond Activation: Activated Primary Amines as Alkylating Reagents in Reductive Cross-Coupling. *Chem. Sci.* **2019**, *10*, 4430–4435. (d) Ni, S.; Li, C.-X.; Mao, Y.; Han, J.; Wang, Y.; Yan, H.; Pan, Y. Ni-Catalyzed Deaminative Cross-Electrophile Coupling of Katritzky Salts with Halides via C–N Bond Activation. *Sci. Adv.* **2019**, *5*, eaaw9516.

(9) He, R.-D.; Li, C.-L.; Pan, Q.-Q.; Guo, P.; Liu, X.-Y.; Shu, X.-Z. Reductive Coupling between C–N and C–O Electrophiles. *J. Am. Chem. Soc.* **2019**, *141*, 12481–12486.

(10) (a) Srogl, J.; Allred, G. D.; Liebeskind, L. S. Sulfonium Salts. Participants par Excellence in Metal-Catalyzed Carbon–Carbon Bond-Forming Reactions. *J. Am. Chem. Soc.* **1997**, *119*, 12376–12377. (b) Zhang, S.; Marshall, D.; Liebeskind, L. S. Efficient Pd-Catalyzed Heterobenzylic Cross-Coupling Using Sulfonium Salts as Substrates and (PhO)<sub>3</sub>P as a Supporting Ligand. *J. Org. Chem.* **1999**, *64*, 2796–2804.

(11) (a) Huihui, K. M. M.; Caputo, J. A.; Melchor, Z.; Olivares, A. M.; Spiewak, A. M.; Johnson, K. A.; DiBenedetto, T. A.; Kim, S.; Ackerman, L. K. G.; Weix, D. J. Decarboxylative Cross-Electrophile Coupling of N-Hydroxyphthalimide Esters with Aryl Iodides. *J. Am. Chem. Soc.* **2016**, *138*, 5016–5019. (b) Suzuki, N.; Hofstra, J. L.; Poremba, K. E.; Reisman, S. E. Nickel-Catalyzed Enantioselective Cross-Coupling of N-Hydroxyphthalimide Esters with Vinyl Bromides. *Org. Lett.* **2017**, *19*, 2150–2153.

(12) (a) Xu, C.-L.; Liu, C.-Y.; Pang, X.; Liu, X.-Y.; Shu, X.-Z. Nickel-Catalyzed Cross-Electrophile Vinyl–Vinyl Coupling: An Approach to Structurally Versatile Dienylboronates. *CCS Chem.* **2022**, *4*, 864–871. (b) Pan, F.-F.; Guo, P.; Li, C.-L.; Su, P.; Shu, X.-Z. Enones from Acid Fluorides and Vinyl Triflates by Reductive Nickel Catalysis. *Org. Lett.* **2019**, *21*, 3701–3705. (c) Dorval, C.; Tricoire, M.; Begouin, J.-M.; Gandon, V.; Gosmini, C. Cobalt-Catalyzed C(sp<sup>2</sup>)-CN Bond Activation: Cross-Electrophile Coupling for Biaryl Formation and Mechanistic Insight. *ACS Catal.* **2020**, *10*, 12819–12827.

(13) Benischke, A. D.; Knoll, I.; Rérat, A.; Gosmini, C.; Knochel, P. A Practical Cobalt-Catalyzed Cross-Coupling of Benzylic Zinc Reagents with Aryl and Heteroaryl Bromides or Chlorides. *Chem. Commun.* **2016**, *52*, 3171–3174.

(14) Selected reports: (a) Wang, J. Gong, Y. Sun, D. Gong, H. Nickel-Catalyzed Reductive Benzoylation of Tertiary Alkyl Halides with Benzyl Chlorides and Chloroformate. *Org. Chem. Front.* **2021**, *8*, 2944–2948. (b) Zhao, W.-T.; Meng, H.; Lin, J.-N.; Shu, W. Ligand- Controlled Nickel- Catalyzed Regiodivergent Cross- Electrophile Alkyl- Alkyl Couplings of Alkyl Halides. *Angew. Chem. Int. Ed.* **2023**, *62*, e202215779. (c) Liang, Z.; Xue, W.; Lin, K.; Gong, H. Nickel-Catalyzed Reductive Methylation of Alkyl Halides and Acid Chlorides with Methyl *p*-Tosylate. *Org. Lett.* **2014**, *16*, 21, 5620–5623. (d) Dai, Y.; Wu, F.; Zang, Z.; You, H.; Gong, H. Ni- Catalyzed Reductive Allylation of Unactivated Alkyl Halides with Allylic Carbonates. *Chem. Eur. J.* **2012**, *18*, 808 – 812. (e) Guo, P.; Wang, K.; Jin, W.-J.; Xie, H.; Qi, L.; Liu, X.-Y.; Shu, X.-Z. Dynamic Kinetic Cross-Electrophile Arylation of Benzyl Alcohols by Nickel Catalysis. *J. Am. Chem. Soc.* **2021**, *143*, 513–523. (f) Suga, T.; Takahashi, Y.; Ukaji, Y. One- Shot Radical Cross Coupling Between Benzyl Alcohols and Alkenyl Halides Using Ni/Ti/Mn System. *Adv. Synth. Catal.* **2020**, *362*, 5622–5626.

(15) (a) Péter, Á.; Perry, G. J. P.; Procter, D. J. Radical C–C Bond Formation Using Sulfonium Salts and Light. *Adv. Synth. Catal.* **2020**, *362*, 2135–2142. (b) Kozhushkov, S. I.; Alcarazo, M. Synthetic Applications of Sulfonium Salts. *Eur. J. Inorg. Chem.* **2020**: 2486–2500. (c) Wang, W.; Yao, K.; Wu, F. Nickel-Catalyzed Reductive Cross-Coupling of Benzylic Sulfonium Salts with Aryl Iodides. *Synlett* **2022**, *33*, 361–366.

(16) (a) del Río-Rodríguez, R.; Blanco, L.; Collado, A.; Fernández-Salas, J. A.; Alemán, J. Nickel-Catalysed Cross-Electrophile Coupling of Benzyl Bromides and Sulfonium Salts

towards the Synthesis of Dihydrostilbenes. *Chem. Eur. J.* **2022**, e202201644. (b) Gao, M., Gosmini, C. Co-Catalyzed Reductive Cross-Couplings of Csp<sup>3</sup>-S with C<sub>allyl</sub>-O electrophiles to form Csp<sup>3</sup>-Csp<sup>3</sup> bonds. accepted in *Adv. Synth. Catal.*

(17) (a) He, Y.; Huang, Z.; Ma, J.; Lin, J.; Zhou, Y.-G.; Yu, Z. Transition-Metal-Free Olefinic C-H Azidoalkylthiolation via C(sp<sup>3</sup>)-S Bond Cleavage of Vinylsulfonium Salts. *Adv. Synth. Catal.* **2022**, *364*, 1–13. (b) Gao, J.; Feng, J.; Du, D. Shining Light on C-S Bonds: Recent Advances in C-C Bond Formation Reactions via C-S Bond Cleavage under Photoredox Catalysis. *Chem. Asian J.* **2020**, *15*, 3637–3659. (c) Pan, F.; Shi, Z.-J. Recent Advances in Transition Metal-Catalyzed C-S Activation: From Thioester to (Hetero)aryl Thioether. *ACS Catal.* **2014**, *4*, 280–288. (d) Nambo, M.; Crudden, C. M. Transition Metal-Catalyzed Cross-Couplings of Benzylic Sulfone Derivatives. *Chem. Rec.* **2021**, *21*, 3978–3989.

(18) (a) Vasu D., Yorimitsu, H., Osuka, A. Palladium-Assisted “Aromatic Metamorphosis of Dibenzo-thiophenes into Triphenylenes *Angew. Chem. Int. Ed.* **2015**, *54*, 7162-7166. (b) Aukland, M. H.; Talbot, F. J. T.; Fernández-Salas, J. A.; Ball, M.; Pulis, A. P.; Procter, D. J. An Interrupted Pummerer/Nickel-Catalysed Cross-Coupling Sequence. *Angew. Chem. Int. Ed.* **2018**, *57*, 9785. (c) Berger, F.; Plutschack, M. B.; Riegger, J.; Yu, W.; Speicher, S.; Ho, M.; Frank, N.; Ritter, T. Site-Selective and Versatile Aromatic C-H Functionalization by Thianthrenation. *Nature* **2019**, *567*, 223–228. (d) Chen, J.; Li, J.; Plutschack, M. B.; Berger, F.; Ritter, T.; Regio- and Stereoselective Thianthrenation of Olefins to Access Versatile Alkenyl Electrophiles. *Angew. Chem. Int. Ed.* **2020**, *59*, 5616. (e) Cowper, P.; Jin, Y.; Turton, M. D.; Kociok-Köhn, G.; Lewis, S. E. Azulenesulfonium Salts: Accessible, Stable, and Versatile Reagents for Cross-Coupling. *Angew. Chem. Int. Ed.* **2016**, *55*, 2564.

(19) (a) Minami, H.; Nogi, K.; Yorimitsu, H. Nickel-Catalyzed Negishi-Type Arylation of Trialkylsulfonium Salts. *Synlett.* **2021**, *32*, 1542-1546. (b) Wang, M.; Song, H.-X.; Wang, X.-Y.; Liu, N.; Qin, H.-L.; Zhang, C.-P. Palladium-Catalyzed Mizoroki–Heck-type Reactions of [Ph<sub>2</sub>SR<sub>m</sub>][OTf] with Alkenes at Room Temperature. *Chem. Commun.*, **2016**, *52*, 11893-11896 (c) Uno, D.; Minami, H.; Otsuka, S.; Nogi, K.; Yorimitsu, H. Palladium-Catalyzed Mizoroki-Heck-Type Alkenylation of Monoaryldialkylsulfoniums. *Chem. Asian J.* **2018**, *13*, 2397 – 2400.

(20) (a) Minami, H.; Nogi, K.; Yorimitsu, H. Palladium-Catalyzed Alkoxyacylation of Arylsulfoniums. *Org. Lett.* **2019**, *21*, 8, 2518–2522. (b) Yanagi, T.; Somerville, R. J.; Nogi, K.; Martin, R.; Yorimitsu, H. Ni-Catalyzed Carboxylation of C(sp<sup>2</sup>)-S Bonds with CO<sub>2</sub>: Evidence for the Multifaceted Role of Zn. *ACS Catal.* **2020**, *10*, 2117–2123.

(21) Tsang, S. W.; Guan, Y.-F.; Wang, J.; Bian, Z.-X.; Zhang, H.-J. Inhibition of Pancreatic Oxidative Damage by Stilbene Derivative Dihydro-Resveratrol: Implication for Treatment of Acute Pancreatitis. *Sci. Rep.* **2016**, *6*, 22859.

(22) Cai, Y.; Qian, X.; Gosmini, C. Cobalt-Catalyzed Csp<sup>3</sup>-Csp<sup>3</sup> Homocoupling. *Adv. Synth. Catal.* **2016**, *358*, 2427.