



HAL
open science

Preliminary inventory of intertidal flora in Vigur Island, Iceland

David Pierre Milesi-Gaches

► **To cite this version:**

David Pierre Milesi-Gaches. Preliminary inventory of intertidal flora in Vigur Island, Iceland. Biodiversity Observations, 2023, 13, pp.273-281. 10.15641/bo.1455 . hal-04263081

HAL Id: hal-04263081

<https://hal.science/hal-04263081>

Submitted on 31 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Preliminary inventory of intertidal flora in Vigur Island, Iceland

David Pierre Milesi-Gaches



Milesi-Gaches DP 2023. Preliminary inventory of intertidal flora in Vigur Island, Iceland. Biodiversity Observations 13: 273–281.

26 October 2023

DOI: 10.15641/bo.1455

Fauna & Flora

Preliminary inventory of intertidal flora in Vigur Island, Iceland

David Pierre Milesi-Gaches

Department of Zoology, University of South Bohemia in České Budějovice, Smiřugata 12, 400 Ísafjörður, Iceland

**milesi.david@gmail.com*

Abstract

This study presents the initial findings of a pilot intertidal inventory conducted on Vigur Island, Iceland. The primary focus was to identify the flora species present in the intertidal zone, without delving into their distribution or coverage. Using transect and scanning methods, this inventory provides insights into the biodiversity of this coastal region. This preliminary assessment lays the groundwork for future comprehensive studies on the island's intertidal biodiversity.

Introduction

The intertidal zone, often described as the transitional boundary between terrestrial and marine environments, plays a pivotal role in coastal ecosystems. This zone is characterized by its unique biotic and abiotic factors, influenced by both marine and terrestrial conditions. The ecological significance of intertidal zones is multifaceted. They serve as habitats for a diverse range of organisms, act as buff-

ers against coastal erosion, and play a role in nutrient cycling and energy flow within coastal ecosystems (Tomanek 2002, Spalding et al. 2014, Lv et al. 2020).

One of the primary ecological functions of the intertidal zone is its role in supporting a rich biodiversity. This biodiversity is not only limited to macro-organisms but also includes a plethora of microeukaryotes, which contribute to the overall functioning and resilience of these ecosystems (Brawley et al. 2017, Kong et al. 2019). However, despite their significance, there is a growing concern about the degradation of these habitats. Principally due to climate change anthropogenic activities, such as the introduction of pollutants as polycyclic aromatic hydrocarbons (PAHs) (He & Silliman 2019). These pollutants can influence the distribution and health of the flora and fauna residing in these zones (Lv et al. 2020). Meanwhile in Iceland, climate change was observed responsible for the diminution of intertidal species richness and diversity towards the north, likely influenced predominantly by temperature variations (Espinosa & Guerra-García 2005). Consequently, the intertidal zone remains vulnerable to various anthropogenic pressures, emphasizing the need for continuous monitoring and research in these areas (Lv et al. 2020). Additionally, the interplay between biotic and abiotic factors in these zones can influence community structures, with factors such as sediment composition and water content playing crucial roles (Kong et al. 2019). Hence, an inventory of the intertidal flora is paramount to understand its status, monitor changes, and implement conservation strategies.

Vigur Island, situated in the Westfjords of Iceland just below the Arctic Circle, stands as a beacon in the North Atlantic. Encircled by the clear waters of Ísafjarðardjúp fjord, the intertidal zone of Vigur Island is a testament to the delicate balance between marine elements and the rugged coastline. The development of nearby fish farms adds another layer to the marine ecosystem, highlighting the coexistence of natural and human-induced elements (Hafskipulag 2023). With its rocky coastline and tidal pools, Vigur serves as an invaluable site for the study of intertidal flora and fauna. As both a sought-after tourist destination and a sanctuary for diverse marine and avian species (Hansen 2019, Milesi-Gaches & Lhériaud 2022), Vigur Island encapsulates the essence of Iceland's coastal ecology.

Therefore, understanding the flora of Vigur Island's intertidal zones becomes essential to gauge the health and resilience of its coastal ecosystems. This pilot inventory aims to shed light on the biodiversity of the intertidal flora of the island, providing a foundation for future conservation and management efforts.

Methods

Transect method

To assess the biodiversity of the intertidal zone, four transects were established at random locations (Figure 1), ensuring that each site was safely accessible during low tide. Each transect extended from the land, moving towards the sea. At every 5 m interval along these transects, a quadrat was placed, and the species within it were identified.

It's important to note that the primary objective of this survey was species identification; thus, no data on species coverage or distribution within the quadrats was recorded.

Scanning method

In addition to the transect surveys, a scanning method was employed to further understand the biodiversity of Vigur Island's intertidal zone. Referring to Figure 1, specific sections of the intertidal zone along the coast were selected for scanning. This involved visually inspecting these areas for the presence of various species and subsequently identifying them. Utilising fishing waders enabled a prolonged stay in the water, facilitating the continuation of fieldwork even after the low tide zone became submerged (Figure 2).

The fieldwork was informed and supported by two key reference books, ensuring accurate species identification: Sept (2008) and Preston-Mafham (2010). These reference texts were instrumental in guiding the identification process throughout the study.



Figure 1: Schematic representation of Vigur Island's intertidal zone study areas (Basemap: Loftmyndir ehf).



Figure 2: The observer scans around the last quadrat of a transect in the intertidal zone of Vigur Island, Iceland.

Results and Conclusion

In the course of the investigation on Vigur Island's intertidal zone, three distinct areas were scanned, and four transects were established (Figure 1). Across these transects, a total of 28 quadrats were examined. Within this framework, a total of 26 species were identified:

23 during the transect sessions, and three species while scanning. This diverse assemblage encompassed green, brown, and red algae, as well as lichens. Notably, a single moss species was also documented (Table 1).

Despite the primary focus being on the investigation of intertidal flora without an in-depth analysis of species coverage, the transect observations still yielded valuable insights into species abundance and distribution. Common sea lettuce emerged as the most frequently observed species, appearing in 11 quadrats (Table 2). Knotted wrack and flat rockweed were also commonly found, each appearing in 10 quadrats. Bushy red seaweed was noted in nine records, making it another frequent species. For a comprehensive breakdown of all species occurrences, refer to Figure 3, in which the distribution of each species is visually represented.

The 'scanning' sessions served as an opportunity to identify recurring species across different locations, as well as species that were not observed within the quadrats. Three species were found through this method: *Urospora bangioides*, *Urospora neglecta*, and *Prasiola* sp. (commonly known as short sea lettuce).

While the current sampling plan may not be sufficiently robust to provide a comprehensive assessment of the entire intertidal zone around Vigur Island, it does reveal variations in species richness based on the specific locations where data were collected (as shown in Figure 3). Transect T3, with only eight species, has the lowest plant diversity. This may be attributed to its location within the bay, near the pier, which serves as the island's sole landing point for boats. This relatively sheltered area is frequented by nesting birds, including the Common Eider *Somateria mollissima*, Black Guillemot *Cephus grylle*, and Arctic Tern *Sterna paradisaea*, as noted in the study by Milesi-Gaches & Lh eriau (2022). The primary activities in this area for these bird species are swimming and feeding.

Transects T2 and T4 are located on the outskirts of the tern colony and are primarily visited by ducks and a few Black Guillemots. These transects are also adjacent to the Atlantic Puffin *Fratercula arctica*

Table 1: Overall inventory of the identified species found on the intertidal zone of Vigur Island, Iceland.

	Species	Common name
Green algae	<i>Acrosiphonia</i> sp.	Green rope seaweed
	<i>Blidingia minima</i>	Dwarf green weed
	<i>Prasiola</i> sp.	Short sea lettuce
	<i>Ulva intestinalis</i>	Maiden hair sea lettuce
	<i>Ulva lactuca</i>	Common sea lettuce
	<i>Ulva linza</i>	Flat-tube sea lettuce
	<i>Urospora bangioides</i>	–
	<i>Urospora neglecta</i>	–
	Brown algae	<i>Alaria esculenta</i>
<i>Ascophyllum nodosum</i>		Knotted wrack
<i>Fucus distichus</i>		Flat rockweed
<i>Fucus spiralis</i>		Spiral wrack
<i>Fucus vesiculosus</i>		Northern rockweed
<i>Himanthalia elongata</i>		Thongweed
<i>Laminaria digitata</i>		Oarweed
<i>Leathesia difformis</i>		Sea cauliflower
Red algae		<i>Corallina officinalis</i>
	<i>Cystoclonium purpureum</i>	Bushy red seaweed
	<i>Devaleraea ramentacea</i>	Variable tube seaweed
	<i>Hildenbrandia rubra</i>	Red crust alga
	<i>Palmaria palmata</i>	Dulse
	<i>Porphyra umbilicalis</i>	Purple laver
	<i>Vertebrata lanosa</i>	Wrack fringe tubeweed
	Lichens	<i>Physcia caesia</i>
<i>Verrucaria maura</i>		Black seaside lichen
<i>Xanthoria parietina</i>		Maritime sunburst lichen
Moss	<i>Cladophora rupestris</i>	Tufted sea moss

Table 2: Species frequency of species recorded in the intertidal zone of Vigur Island, Iceland.

Species	Common name	Quadrats
<i>Ulva lactuca</i>	Common sea lettuce	11
<i>Ascophyllum nodosum</i>	Knotted wrack	10
<i>Fucus distichus</i>	Flat rockweed	10
<i>Cystoclonium purpureum</i>	Bushy red seaweed	9
<i>Corallina officinalis</i>	Coral weed	6
<i>Fucus vesiculosus</i>	Northern rockweed	6
<i>Hildenbrandia rubra</i>	Red crust alga	5
<i>Blidingia minima</i>	Dwarf green weed	5
<i>Ulva intestinalis</i>	Maiden hair sea lettuce	4
<i>Verrucaria maura</i>	Black seaside lichen	4
<i>Leathesia difformis</i>	Sea cauliflower	4
<i>Elachista</i> spp.	Tufted fringe	3
<i>Acrosiphonia</i> spp.	Green rope seaweed	2
<i>Cladophora rupestris</i>	Tufted sea moss	2
<i>Palmaria palmata</i>	Dulse	2
<i>Porphyra umbilicalis</i>	Purple laver	2
<i>Ulva linza</i>	Flat-tube sea lettuce	2
<i>Aleria esculenta</i>	Windged kelp	1
<i>Devaleraea ramentacea</i>	Variable tube seaweed	1
<i>Physcia caesia</i>	Blue-gray rosette lichen	1
<i>Xanthoria parietina</i>	Maritime sunburst lichen	1
<i>Laminaria digitata</i>	Oarweed	1
<i>Fucus spiralis</i>	Spiral wrack	1



Figure 3: Distribution of the intertidal flora species found through the transect method on Vigur Island, Iceland.



Figure 4: With seven identified species, the quadrat T3-Q6 was the one with the largest species richness in the intertidal zone of Vigur Island, Iceland.

colony. As for Transect T1, the area is frequented by a variety of bird species, including puffins, ducks, seagulls, and Northern Fulmars *Fulmarus glacialis*, who engage in activities such as feeding and swimming.

The island's coastline exhibits notable variation not solely in the distribution of breeding and nesting species. Beyond the evident geographical exposure, distinct tidal pools and bays present varying current characteristics and tidal exposure. This naturally leads to the an-

tipication of species richness variation throughout the coastline. A mere four transects underscore Vigur's potential to harbour a significant intertidal biodiversity. Such variation is likely extendable to fauna as well. Although not the focal point of this pilot study, several fauna species were identified during fieldwork, including the breadcrumb sponge *Halichondria panicea*, common starfish *Asterias rubens*, blue mussel *Mytilus edulis*, northern rock barnacle *Semibalanus balanoides*, dog whelk *Nucella lapillus*, and small, rough, flat, and smooth periwinkles.

Therefore, I advocate for expanded research within Vigur Island's intertidal zone to include a broader array of factors such as currents, exposure, coastline variations, and the presence of nesting birds. Given the tourism activity and the recent establishment of fish farms in the vicinity (Hafskipulag 2023), it is imperative that decisions pertaining to biodiversity conservation are grounded in rigorous scientific research (Ferðamálastofa 2020). This pilot study has showcased the potential richness of intertidal biodiversity, and proposed a mixed method which could be further developed to monitor the richness and evolution of the identified species.

Acknowledgements


I thank the owners of Vigur Island for accommodating me during the fieldwork phase of this pilot study.

References

- Brawley SH, Blouin NA, Ficko-Blean E, Wheeler GL, Lohr M, Goodson HV, Jenkins JW, Blaby-Haas CE, Helliwell KE, Chan CX, Marriage TN, Bhattacharya D, Klein AS, Badis Y, Brodie J, Cao Y, Collén J, Dittami SM, Gachon CMM, Green BR, Karpowicz SJ, Kim JW, Kudahl UJ, Lin S, Michel G, Mittag M, Olson BJSC, Pangilinan JL, Peng Y, Qiu H, Shu S, Singer JT, Smith AG, Sprecher BN, Wagner V, Wang W, Wang Z-Y, Yan J, Yarish C, Zäuner-Riek S, Zhuang Y, Zou Y, Lindquist EA, Grimwood J, Barry KW, Rokhsar DS, Schmutz J, Stiller JW, Grossman AR, Prochnik SE** 2017. Insights into the red algae and eukaryotic evolution from the genome of *Porphyra umbilicalis* (Bangiophyceae, Rhodophyta). *Proceedings of the National Academy of Sciences* 114: E6361–E6370. Available online at <https://doi.org/10.1073/pnas.1703088114>
- Espinosa F, Guerra-García JM** 2005. Algae, macrofaunal assemblages and temperature: a quantitative approach to intertidal ecosystems of Iceland. *Helgoland Marine Research* 59: 273–285. Available online at <https://doi.org/10.1007/s10152-005-0003-6>
- Ferðamálastofa** 2020. Leading in sustainable development: Icelandic tourism 2030. Available online at https://www.ferdamalastofa.is/static/files/ferdamalastofa/Misc/leading-in-sustainable-development_icelandic-tourism-2030.pdf
- Hafskipulag** 2023. Strandsvæðisskipulag Vestfjarða. Available online at <https://luk.skipulag.is/portal/apps/experiencebuilder/experience/?id=47fd5a7e682743fc8af535f940435dff>
- Hansen ES** 2019. Stofnvöktun lunda - Lokakýrsla 2017–2019. Technical report, South Iceland Nature Research Centre. Available online at https://www.researchgate.net/profile/Erpur-Hansen/publication/338013826_Stofnvoktun_lunda_-_Lokakyrsla_2017-2019/
- He Q, Silliman BR** 2019. Climate change, human impacts, and coastal ecosystems in the Anthropocene. *Current Biology* 29: R1021–R1035. <https://doi.org/10.1016/j.cub.2019.08.042>
- Kong J, Wang Y, Warren A, Huang B, Sun P** 2019. Diversity distribution and assembly mechanisms of planktonic and benthic microeukaryote communities in intertidal zones of southeast Fujian, China. *Frontiers in Microbiology* 10:2640. Available online at <https://www.frontiersin.org/articles/10.3389/fmicb.2019.02640/full>
- Lv M, Luan X, Liao C, Wang D, Liu D, Zhang G, Jiang G, Chen L** 2020. Human impacts on polycyclic aromatic hydrocarbon distribution in Chinese intertidal zones. *Nature Sustainability* 3: 878–884. Available online at <https://doi.org/10.1038/s41893-020-0565-y>
- Milesi-Gaches D, Lhériaux A** 2022. Census of breeding seabirds in Vigur Island, Westfjords, Iceland in 2021. *Bird Census News* 35: 10–20.
- Preston-Mafham K** 2010. *Seashore of Britain and Europe*. London: HarperCollins Publishers.
- Sept JD** 2008. *A photographic guide to seashore life in the North Atlantic: Canada to Cape Cod*. Princeton, New Jersey: Princeton University Press.
- Spalding MD, Ruffo S, Lacambra C, Meliane I, Hale LZ, Shepard CC, Beck MW** 2014. The role of ecosystems in coastal protection: Adapting to climate change and coastal hazards. *Ocean & Coastal Management* 90: 50–57. Available online at <https://doi.org/10.1016/j.ocecoaman.2013.09.007>

Tomanek L 2002. The heat-shock response: Its variation, regulation and ecological importance in intertidal gastropods (genus *Tegula*) 1. Integrative and Comparative Biology 42: 797–807. Available online at <https://doi.org/10.1093/icb/42.4.797>

*Paper edited by Megan Loftie-Eaton
Biodiversity and Development Institute*



Biodiversity Observations is powered by [Open Journal Systems \(OJS\)](#) and is hosted by the [University of Cape Town Libraries](#). OJS is an open source software application for managing and publishing scholarly journals. Developed and released by the [Public Knowledge Project](#) in 2001, it is the most widely used open source journal publishing platform in existence, with over 30,000 journals using it worldwide.

Biodiversity Observations

The scope of Biodiversity Observations includes papers describing observations about biodiversity in general, including animals, plants, algae and fungi. This includes observations of behaviour, breeding and flowering patterns, distributions and range extensions, foraging, food, movement, measurements, habitat and colouration/plumage variations. Biotic interactions such as pollination, fruit dispersal, herbivory and predation fall within the scope, as well as the use of indigenous and exotic species by humans. Observations of naturalised plants and animals will also be considered. Biodiversity Observations will also publish a variety of other interesting or relevant biodiversity material: reports of projects and conferences, annotated checklists for a site or region, specialist bibliographies, book reviews and any other appropriate material. Further details and guidelines to authors are on the journal website (<https://journals.uct.ac.za/index.php/BO/>).

ISSN 2959-3441

Editor: LG Underhill

