

# A global database of dissolved organic matter (DOM) measurements in coastal waters (CoastDOM v1)

Christian Lønborg, Cátia Carreira, Gwenaël Abril, Susana Agustí, Valentina Amaral, Agneta Andersson, Javier Arístegui, Punyasloke Bhadury, Mariana Bernardi Bif, Alberto V Borges, et al.

# ▶ To cite this version:

Christian Lønborg, Cátia Carreira, Gwenaël Abril, Susana Agustí, Valentina Amaral, et al.. A global database of dissolved organic matter (DOM) measurements in coastal waters (CoastDOM v1). Earth System Science Data: Papers in open discussion, 2023, 10.5194/essd-2023-348. hal-04266492

# HAL Id: hal-04266492 https://hal.science/hal-04266492

Submitted on 31 Oct 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





## 1 A global database of dissolved organic matter (DOM) measurements in coastal

## 2 waters (CoastDOM v1)

Christian Lønborg<sup>1\*</sup>, Cátia Carreira<sup>2</sup>, Gwenaël Abril<sup>3</sup>, Susana Agustí<sup>4</sup>, Valentina Amaral<sup>5</sup>, 3 Agneta Andersson<sup>6</sup>, Javier Arístegui<sup>7</sup>, Punyasloke Bhadury<sup>8</sup>, Mariana B. Bif<sup>9</sup>, Alberto V. 4 Borges<sup>10</sup>, Steven Bouillon<sup>11</sup>, Maria LI. Calleja<sup>4,12</sup>, Luiz C. Cotovicz Jr<sup>13,49</sup>, Stefano Cozzi<sup>14</sup>, 5 Maryló Doval<sup>15</sup>, Carlos M. Duarte<sup>4</sup>, Bradley Eyre<sup>16</sup>, Cédric G. Fichot<sup>17</sup>, E. Elena García-6 Martín<sup>18</sup>, Alexandra Garzon-Garcia<sup>19</sup>, Michele Giani<sup>20,21</sup>, Rafael Gonçalves-Araujo<sup>22</sup>, 7 Renee Gruber<sup>23</sup>, Dennis A. Hansell<sup>24</sup>, Fuminori Hashihama<sup>25</sup>, Ding He<sup>26</sup>, Johnna M. 8 Holding<sup>27</sup>, William R. Hunter<sup>28</sup>, J. Severino P. Ibánhez<sup>29</sup>, Valeria Ibello<sup>30</sup>, Shan Jiang<sup>31</sup>, 9 10 Guebuem Kim<sup>32</sup>, Katja Klun<sup>33</sup>, Piotr Kowalczuk<sup>34</sup>, Atsushi Kubo<sup>35</sup>, Choon Weng Lee<sup>36</sup>, Cláudia B. Lopes<sup>37</sup>, Federica Maggioni<sup>38</sup>, Paolo Magni<sup>39</sup>, Celia Marrase<sup>40</sup>, Patrick 11 12 Martin<sup>41</sup>, S. Leigh McCallister<sup>42</sup>, Roisin McCallum<sup>43</sup>, Patricia M Medeiros<sup>44</sup>, Xosé Anxelu G. Morán<sup>4,45</sup>, Frank Muller-Karger<sup>46</sup>, Allison Myers-Pigg<sup>47</sup>, Marit Norli<sup>48</sup>, Joanne M. 13 Oakes<sup>15</sup>, Helena Osterholz<sup>49</sup>, Hyekyung Park<sup>32</sup>, Maria Lund Paulsen<sup>50</sup>, Judith A. 14 Rosentreter<sup>16,51</sup>, Digna Rueda-Roa<sup>46</sup>, Chiara Santinelli<sup>52</sup>, Yuan Shen<sup>53</sup>, Eva Teira<sup>54</sup>, 15 Tinkara Tinta<sup>33</sup>, Guenther Uher<sup>55</sup>, Masahide Wakita<sup>56</sup>, Nicholas Ward<sup>47</sup>, Kenta 16 Watanabe<sup>57</sup>, Yu Xin<sup>58</sup>, Youhei Yamashita<sup>59</sup>, Liyang Yang<sup>60</sup>, Jacob Yeo<sup>16</sup>, Huamao 17 Yuan<sup>61</sup>, Qiang Zheng<sup>53,61</sup>, Xosé Antón Álvarez-Salgado<sup>29</sup> 18

19 <sup>1</sup>Section for Marine Diversity and Experimental Ecology, Department of Ecoscience,

20 Aarhus University, 4000 Roskilde, Denmark.

21 <sup>2</sup>Department of Environmental Science, Aarhus University, 4000 Roskilde, Denmark.

<sup>22</sup> <sup>3</sup>Laboratoire de Biologie des Organismes et Ecosystèmes Aquatiques (BOREA), CNRS,

23 Muséum National d'Histoire Naturelle, 61 rue Buffon, 75005, Paris, France.

<sup>4</sup>King Abdullah University of Science and Technology, Thuwal 23955-6900, Kingdom of

25 Saudi Arabia.





- 26 <sup>5</sup>Departamento Interdisciplinario de Sistemas Costero Marinos, Centro Universitario
- 27 Regional Este, Universidad de la República, Ruta 9 y 15, CP 27000, Rocha, Uruguay.
- 28 <sup>6</sup>Umeå Marine Sciences Centre, Umeå University, Sweden.
- <sup>7</sup>Instituto de Oceanografía y Cambio Global (IOCAG), Universidad de Las Palmas de
- 30 Gran Canaria, Las Palmas, Spain.
- 31 <sup>8</sup>Centre for Climate and Environmental Studies, Indian Institute of Science Education
- 32 and Research Kolkata, Mohanpur, Nadia, West Bengal, India.
- <sup>9</sup>Monterey Bay Aquarium Research Institute, Moss Landing, California, United States.
- <sup>10</sup>University of Liège, Chemical Oceanography Unit, Liège, Belgium.
- 35 <sup>11</sup>KU Leuven, Department of Earth and Environmental Sciences, Leuven, Belgium.
- 36 <sup>12</sup>Marine Ecology and Systematics (MarES), Department of Biology, Universitat de les
- 37 Illes Balears, 07122 Palma de Mallorca, Spain.
- 38 <sup>13</sup>Departamento de Geoquímica, Universidade Federal Fluminense, Outeiro São João
- 39 Batista s/n, 24020015 Niterói, RJ, Brazil.
- 40 <sup>14</sup>Consiglio Nazionale delle Ricerche, Istituto di Scienze Marine (CNR-ISMAR), Strada
- 41 Statale 14, km 163.5, 34149 Trieste, Italy.
- 42 <sup>15</sup>Instituto Tecnolóxico para o Control do Medio Mariño de Galicia, 36611 Vilagarcía de
- 43 Arousa, Spain.
- 44 <sup>16</sup>Centre for Coastal Biogeochemistry, Faculty of Science and Engineering, Southern
- 45 Cross University, Lismore 2480, NSW, Australia.
- <sup>17</sup>Department of Earth and Environment, Boston University, Boston, MA, United States.
- 47 <sup>18</sup>National Oceanography Centre, European Way, Southampton, SO14 3ZH, United
- 48 Kingdom.
- 49 <sup>19</sup>Department of Environment and Science, PO Box 5078, Brisbane, Queensland 4001,
- 50 Australia.
- <sup>20</sup>National Institute of Oceanography and Applied Geophysics (OGS), Trieste, Italy.





- 52 <sup>21</sup>Istituto Centrale per la Ricerca scientifica e tecnologia Applicata al MAre, Chioggia,
- 53 Italy
- 54 <sup>22</sup>National Institute of Aquatic Resources, Technical University of Denmark, Kgs.
- 55 Lyngby, Denmark.
- 56 <sup>23</sup>Australian Institute of Marine Science, PMB 3, Townsville QLD 4810, Australia
- 57 <sup>24</sup>Department of Ocean Sciences, Rosenstiel School of Marine and Atmospheric
- 58 Science, University of Miami, Miami, FL, United States.
- 59 <sup>25</sup>Tokyo University of Marine Science and Technology, Japan.
- 60 <sup>26</sup>Department of Ocean Science and Center for Ocean Research in Hong Kong and
- 61 Macau, The Hong Kong University of Science and Technology, Clear Water Bay, Hong
- 62 Kong, China.
- 63 <sup>27</sup>Department of Ecoscience, Aarhus University, 8000 Aarhus, Denmark.
- 64 <sup>28</sup>Fisheries and Aquatic Ecosystems Branch, Agri-Food and Biosciences Institute,
- 65 Belfast, Northern Ireland, United Kingdom.
- 66 <sup>29</sup>CSIC, Instituto de Investigacións Mariñas, Eduardo Cabello 6, 36208 Vigo, Spain.
- <sup>30</sup>Institute of Marine Sciences, Middle East Technical University, 33731 Erdemli-Mersin,
- 68 Turkey.
- <sup>31</sup>State Key Laboratory of Estuarine and Coastal Research, East China Normal
- 70 University, 200241, Shanghai, China.
- <sup>32</sup>School of Earth and Environmental Sciences, Seoul National University, Seoul 08826,
- 72 Korea.
- <sup>33</sup>Marine Biology Station, National Institute of Biology, Fornače 41, 6330 Piran,
- 74 Slovenia.
- 75 <sup>34</sup>Remote Sensing Laboratory, Institute of Oceanology, Polish Academy of Sciences,
- 76 Sopot, Poland.





- <sup>35</sup>Department of Geosciences, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka,
- 78 422-8529, Japan.
- 79 <sup>36</sup>Laboratory of Microbial Ecology, Institute of Biological Sciences, Institute of Ocean
- 80 and Earth Sciences, Universiti Malaya, 50603 Kuala Lumpur, Malaysia.
- 81 <sup>37</sup>CICECO Aveiro Institute of Materials, Department of Chemistry, University of Aveiro,
- 82 Campus de Santiago, 3810-193 Aveiro, Portugal.
- 83 <sup>38</sup>University of New Caledonia and Institute de recherche pour le development (IRD),
- 84 New Caledonia.
- <sup>39</sup>Consiglio Nazionale delle Ricerche, Istituto per lo studio degli impatti Antropici e
- 86 Sostenibilità in ambiente marino (CNR-IAS), Loc. Sa Mardini, Torregrande, 09170,
- 87 Oristano, Italy.
- <sup>40</sup>Institut de Ciències del Mar (CSIC), Passeig Marítim de la Barceloneta, 37
- 89 08003 Barcelona, Spain.
- <sup>41</sup>Asian School of the Environment, Nanyang Technological University, 639798,
- 91 Singapore.
- <sup>42</sup>Virginia Commonwealth University, Department of Biology, Richmond, Virginia, United
- 93 States.
- <sup>43</sup>Centre for Marine Ecosystems Research, School of Science, Edith Cowan University,
- 95 270 Joondalup Drive, Joondalup 6027 WA, Australia.
- <sup>44</sup>Department of Marine Sciences, University of Georgia, Athens, GA30602, United
- 97 States.
- 98 <sup>45</sup>Centro Oceanográfico de Gijón/Xixón, Instituto Español de Oceanografía, Gijón/Xixón,
- 99 Spain.
- 100 <sup>46</sup>College of Marine Science, University of South Florida, Saint Petersburg, Florida,
- 101 United States.





- 102 <sup>47</sup>Pacific Northwest National Laboratory, Marine and Coastal Research Laboratory,
- 103 Sequim, Washington, United States.
- <sup>48</sup>Norwegian Institute for Water Research, Oslo, Norway.
- 105 <sup>49</sup>Leibniz Institute for Baltic Sea Research Warnemuende, D- 18119 Rostock-
- 106 Warnemuende, Germany.
- <sup>50</sup>Marine microbiology, University of Bergen, Norway.
- 108 <sup>51</sup>Yale School of the Environment, Yale University, New Haven, Connecticut, United
- 109 States.
- <sup>52</sup>Biophysics Institute, CNR, Pisa, Italy.
- 111 <sup>53</sup>State Key Laboratory of Marine Environmental Science (MEL) & College of Ocean
- 112 and Earth Sciences, Xiamen University, China.
- 113 <sup>54</sup>Departamento de Ecología y Biología Animal, Universidade de Vigo, Centro de
- 114 Investigacion Mariña da Universidade de Vigo(CIM-UVigo), Vigo, Spain.
- 115 <sup>55</sup>School of Natural and Environmental Science, Newcastle University, Newcastle upon
- 116 Tyne, United Kingdom.
- <sup>56</sup>Mutsu Institute for Oceanography, Research Institute for Global Change, Japan
- 118 Agency for Marine-Earth Science and Technology, 690 Kitasekine, Sekine, Mutsu,
- 119 Aomori, Japan.
- 120 <sup>57</sup>Coastal and Estuarine Environment Research Group, Port and Airport Research
- 121 Institute, Yokosuka 239-0826, Japan.
- 122 <sup>58</sup>Key Laboratory of Marine Chemistry Theory and Technology, Ministry of Education,
- 123 Institute for Advanced Ocean Study, Ocean University of China, Qingdao, Shandong,
- 124 China.
- <sup>59</sup>Faculty of Environmental Earth Science, Hokkaido University, Hokkaido 060-0810,
- 126 Japan.
- 127 <sup>60</sup>College of Environment and Safety Engineering, Fuzhou University, China.





- 128 <sup>61</sup>Fujian Key Laboratory of Marine Carbon Sequestration, Xiamen University, Xiamen
- 129 361102, China.
- 130
- 131 \*Corresponding author:
- 132 E-mail: <u>c.lonborg@ecos.au.dk</u> or <u>clonborg@gmail.com</u>
- 133
- 134 ORCID nr.:
- 135 Christian Lønborg: 0000-0001-8380-0238
- 136 Cátia Carreira: 0000-0002-1520-9320
- 137 Gwenaël Abril: 0000-0002-4914-086X
- 138 Susana Agustí: 0000-0003-0536-7293
- 139 Valentina Amaral: 0000-0002-1088-1484
- 140 Agneta Andersson: 0000-0001-7819-9038
- 141 Javier Arístegui: 0000-0002-7526-7741
- 142 Punyasloke Bhadury: 0000-0001-8714-7475
- 143 Mariana Bernardi Bif: 0000-0002-2148-4556
- 144 Alberto V. Borges: 0000-0002-5434-2247
- 145 Steven Bouillon: 0000-0001-7669-2929
- 146 Maria Ll. Calleja: 0000-0002-5992-2013
- 147 Luiz C. Cotovicz Jr: 0000-0002-3914-8155
- 148 Stefano Cozzi: 0000-0003-0116-742X
- 149 Maryló Doval: 0000-0002-8565-8703
- 150 Carlos M. Duarte: 0000-0002-1213-1361
- 151 Bradley Eyre: 0000-0001-5502-0680
- 152 Cédric G. Fichot: 0000-0002-1099-5764
- 153 E. Elena García-Martín: 0000-0003-4807-3287
- 154 Alexandra Garzon-Garcia: 0000-0002-6804-8890
- 155 Michele Giani: 0000-0002-3306-7725
- 156 Rafael Gonçalves-Araujo: 0000-0001-8344-8326
- 157 Renee Gruber: 0000-0002-8788-6910
- 158 Dennis A. Hansell: 0000-0001-9275-3445
- 159 Fuminori Hashihama: 0000-0003-3835-7681
- 160 Ding He: 0000-0001-9620-6115
- 161 Johnna M. Holding: 0000-0002-7364-0055
- 162 William R. Hunter: 0000-0001-8801-7947
- 163 J. Severino P. Ibánhez: 0000-0001-6093-3054
- 164 Valeria Ibello: 0000-0002-1067-0425
- 165 Shan Jiang: 0000-0002-1121-6080
- 166 Guebuem Kim: 0000-0002-5119-0241
- 167 Katja Klun: 0000-0001-6111-1650
- 168 Piotr Kowalczuk: 0000-0001-6016-0610
- 169 Atsushi Kubo: 0000-0002-6457-5386
- 170 Choon Weng Lee: 0000-0001-9805-9980
- 171 Cláudia Lopes: 0000-0001-7378-8677
- 172 Federica Maggioni: 0000-0002-7109-4257





- 173 Paolo Magni: 0000-0001-5955-6829
- 174 Celia Marrase: 0000-0002-5097-4829
- Patrick Martin: 0000-0001-8008-5558 175
- S. Leigh McCallister: 0000-0002-9041 176
- 177 Roisin McCallum: 0000-0002-0358-2371
- Patricia M Medeiros: 0000-0001-6818-2603 178
- Xosé Anxelu G. Morán: 0000-002-9823-5339 179
- Frank Muller-Karger: 0000-0003-3159-5011 180
- 181 Allison Myers-Pigg: 0000-0002-6905-6841
- Marit Norli: 0000-0001-7472-1562 182
- Joanne M. Oakes: 0000-0002-9287-2652 183
- Helena Osterholz: 0000-0002-2858-9799 184
- 185 Hyekyung Park: 0000-0002-4743-5883
- 186 Maria Lund Paulsen: 0000-0002-1474-7258
- 187 Judith A. Rosentreter: 0000-0001-5787-5682
- Digna Rueda-Roa: 000-0003-4621-009X 188 Chiara Santinelli: 0000-0002-8921-275X 189
- 190 Yuan Shen: 0000-0001-6618-4226
- Eva Teira: 0000-0002-4333-0101
- 191
- 192 Tinkara Tinta: 0000-0001-6740-8973
- Guenther Uher: 0000-0001-5105-4445 193
- Masahide Wakita: 0000-0002-3333-0546 194
- 195 Nicholas Ward: 0000-0001-6174-5581
- 196 Kenta Watanabe: 0000-0002-0106-3623
- 197 Yu Xin: 0000-0002-5328-7717
- 198 Youhei Yamashita: 0000-0002-9415-8743
- Livang Yang: 0000-0001-8767-8698 199
- 200 Jacob Yeo: 0000-0003-2443-5378
- 201 Huamao Yuan: 0000-0003-2014-619X
- 202 Qiang Zheng: 0000-0002-6836-2310
- 203 Xosé Antón Álvarez-Salgado: 0000-0002-2387-9201





#### 204 Abstract

205 The measurements of dissolved organic carbon (DOC), nitrogen (DON), and phosphorus 206 (DOP) are used to characterize the dissolved organic matter (DOM) pool and are 207 important components of biogeochemical cycling in the coastal ocean. Here, we present 208 the first edition global database (CoastDOM v1; of а available at 209 https://figshare.com/s/512289eb43c4f8e8eaef) compiling previously published and 210 unpublished measurements of DOC, DON, and DOP collected in coastal waters. These 211 data are complemented by hydrographic data such as temperature and salinity and, to 212 the extent possible, other biogeochemical variables (e.g., Chlorophyll-a, inorganic 213 nutrients) and the inorganic carbon system (e.g., dissolved inorganic carbon and total 214 alkalinity). Overall, CoastDOM v1 includes observations from all continents however, 215 most data were collected in the Northern Hemisphere, with a clear gap in coastal water 216 DOM measurements from the Southern Hemisphere. The data included were collected 217 from 1978 to 2022 and consist of 62339 data points for DOC, 20360 for DON and 13440 218 for DOP. The number of measurements decreases progressively in the sequence DOC 219 > DON > DOP, reflecting both differences in the maturity of the analytical methods and 220 the greater focus on carbon cycling by the aquatic science community. The global 221 database shows that the average DOC concentration in coastal waters (average 222 (standard deviation; SD): 182 (314) µmol C L-1; median: 103 µmol C L-1), is 13-fold greater 223 than the average coastal DON concentrations (average (SD): 13.6 (30.4) µmol N L<sup>-1</sup>; 224 median: 8.0 µmol N L<sup>-1</sup>), which was itself 39-fold greater than the average coastal DOP 225 concentrations (average (SD): 0.34 ± 1.11 µmol P L<sup>-1</sup>; median: 0.18 µmol P L<sup>-1</sup>). This 226 dataset will be useful to identify global spatial and temporal patterns in DOM and to 227 facilitate reuse of DOC, DON and DOP data in studies aimed at better characterising local 228 biogeochemical processes, closing nutrient budgets, estimating carbon, nitrogen and





- 229 phosphorous pools, as well as identifying a baseline for modelling future changes in
- 230 coastal waters.
- 231
- 232 Keywords: Dissolved organic matter, Dissolved organic carbon, Dissolved organic
- 233 nitrogen, Dissolved organic phosphorus, Coastal waters, Global database.





#### 234 **1. Introduction**

Coastal waters are the most biogeochemical dynamic areas of the ocean, exhibiting the highest standing stocks, process rates and transport fluxes of carbon (C), nitrogen (N), and phosphorus (P) per unit area (Bauer et al., 2013; Mackenzie et al., 2011). In these areas, organic matter plays a critical role in numerous biogeochemical processes, serving as both a C, N and P reservoir and substrate (Carreira et al., 2021).

240 Organic matter found in the marine environment is commonly distinguished by its size; 241 material retained on a filter with a pore size typically between 0.2 and 0.7 µm is classified 242 as particulate organic matter (POM), whereas organic matter that passes through the filter 243 is referred to as dissolved organic matter (DOM). This partitioning is operational but has 244 implications for biogeochemical cycling: POM can be suspended in the water column or 245 sink to the sediments controlled by its size, shape and density (Laurenceau-Cornec et al., 246 2015), whereas DOM is a solute that mostly remains in the water column. In most coastal 247 waters, the DOM concentrations are greater than POM, with the POM fraction being less 248 degraded and more bioavailable (Boudreau and Ruddick, 1991; Lønborg et al., 2018).

249 The DOM pool consists mainly of C (DOC), N (DON), and P (DOP) but it also includes 250 other elements such as oxygen, sulphur and trace elements (Lønborg et al., 2020). In 251 coastal waters, DOM originates from multiple sources. Internal, or autochthonous, sources include planktonic organisms (Lønborg et al., 2009; Carlson and Hansell, 2015), 252 253 benthic microalgae, macrophytes, and sediment porewater (Burdige and Komada, 2014; 254 Wada et al., 2008). On the other hand, DOM from external, or allochthonous, sources, 255 has mainly terrestrial origins, including wetlands, river and surface runoff, groundwater 256 discharges, and atmospheric deposition (lavorivska et al., 2016; Raymond and Spencer, 257 2015; Taniguchi et al., 2019; Santos et al., 2021). The main sinks for DOM from the water 258 column in coastal waters are: 1) bubble coagulation and abiotic flocculation (Kerner et al., 259 2003) or sorption to particles (Chin et al., 1998); 2) sunlight mediated photodegradation





260 (Mopper et al., 2015); and 3) microbial degradation by mainly heterotrophic prokaryotes

261 (Lønborg and Álvarez-Salgado, 2012).

262 Given the importance of DOM as a source of nutrients and for coastal biogeochemical 263 cycling in general, numerous studies have measured the C, N and P content of the DOM 264 pool over the last few decades (e.g., García-Martín et al., 2021; Cauwet, 2002; Osterholz 265 et al., 2021). Most data, however, are often unavailable or stored in an inaccessible 266 manner, making it difficult to e.g., analyse global spatial and temporal patterns effectively. 267 A global open ocean DOM data compilation already exists, but it contains few coastal 268 samples (< 200m) (Hansell et al., 2021). Hence, there is a clear need for a comprehensive 269 global and integrated database of DOC, DON and DOP measurements for coastal waters. 270 To address this need, we have prepared the first edition of a coastal DOM database 271 (named CoastDOM v1), by compiling both previously reported as well as unpublished 272 data. These data have been obtained from authors of the original studies or extracted 273 directly from the original studies. In order to allow the DOM measurements to be 274 interpreted across larger scales, and to better understand their relationship with local 275 environmental conditions, we have included concurrently collected ancillary data (such 276 as physical and/or chemical seawater properties) whenever available. The objective of 277 this database is multifaceted. Firstly, we aimed to compile all available coastal DOM data 278 into a single repository. Secondly, our intention was to make these data easily accessible 279 to the research community and thirdly, we sought to achieve long-term consistency of the 280 measurements, to enable data intercomparison, and establish a robust baseline for 281 assessing, for example, the impacts of climate change and land use changes.

282

#### 283 2. Methods

#### 284 2.1. Data compilation





285 The measurements included in CoastDOM v1 were obtained either directly from 286 authors of previously published studies, online databases, or scientific papers. An 287 extensive search of published reports, Ph.D. theses, and peer-reviewed literature was 288 performed to identify studies dealing with DOM in coastal waters. First, a formal search 289 was performed using Google Scholar in January 2022 using the search terms "dissolved 290 organic carbon", "dissolved organic nitrogen", and "dissolved organic phosphorus" in 291 connection with "marine" or "ocean", which yielded a total of 897 articles (after filtering 292 the query by searching content in the title and abstract and excluding non-coastal 293 articles). When data could not be obtained directly from the corresponding authors, 294 relevant data were extracted. Further searches for relevant datasets were conducted 295 using the reference lists of the identified scientific papers as well as databases and 296 repositories to capture as many datasets as possible. Additionally, research groups that 297 were invited to participate in this effort were also encouraged to submit unpublished data 298 to CoastDOM v1.

299

#### 300 2.2. Dissolved organic matter analysis

301 The DOC concentrations included in CoastDOM v1 were commonly measured using a 302 total organic carbon (TOC) high temperature catalytic oxidation (HTCO) analyser (81% 303 of samples). Some were measured by a combined wet chemical oxidation (WCO) step 304 and/or UV digestion, after which the carbon dioxide generated was guantified (19% of 305 samples). Similarly, concentrations of total dissolved nitrogen (TDN) were determined 306 using either a nitric oxide chemiluminescence detector connected in series with the HTCO 307 analyser used for DOC analyses (31% of the samples), or by employing a UV and/or 308 chemical oxidation step (69%). In the latter approach, both organic and inorganic N 309 compounds were oxidised to nitrate, which was subsequently quantified through a 310 colorimetric method to determine the concentration of inorganic N (Valderrama, 1981;





311 Álvarez-Salgado et al., 2023; Halewood et al., 2022; Foreman et al., 2019). The reported 312 DON concentrations were calculated as the difference between TDN and dissolved inorganic nitrogen (DIN; sum of ammonium (NH<sub>4</sub><sup>+</sup>) and nitrate/nitrite (NO<sub>3</sub><sup>-</sup> + NO<sub>2</sub><sup>-</sup>); DON 313 314 = TDN - DIN) (Álvarez-Salgado et al., 2023). Analyses of total dissolved phosphorus 315 (TDP) were determined by UV (4%) or wet chemical oxidation (66%), or a combination of 316 these (30%), and subsequently were analysed for inorganic phosphorus by a colorimetric 317 method (Álvarez-Salgado et al., 2023). The DOP concentrations were calculated as the difference between TDP and soluble reactive phosphorus (SRP: HPO42-) (DOP = TDP -318 SRP) (Álvarez-Salgado et al., 2023). 319

320

#### 321 3. Description of the dataset

322 The data compiled in CoastDOM v1 were collected, analysed and processed by different 323 laboratories, however, all data included have undergone quality control measures, either 324 by using reference samples or internal quality assurance procedures. While many of the 325 included DOC and TDN data have been systematically compared against consensus 326 reference material (CRM) mainly provided by the University of Miami's CRM program 327 (Hansell, 2005), there is a limitation in CoastDOM v1 regarding the intercalibration across 328 different measurement systems used for both DOP and DON determination. While the 329 CRM could be used for DOC, DON and DOP measurements, this has not yet been 330 attempted for DOP and measurement uncertainties increase in the sequence DOC > 331 DON > DOP. Although some of the reported measurements have quantified the DOP 332 recovery based on commercially available DOP compounds such as Adenosine 333 triphosphate (ATP), it is not known if these were conducted systematically in all cases. 334 Therefore, we strongly recommend undertaking further intercalibrations across 335 laboratories for future measurements of TDP, as has been done for DOC and TDN





336	measurements (e.g., Sharp et al., 2002). Since additional quality control is not possible
337	in retrospect, we assessed the quality of CoastDOM v1 based on its internal consistency.
338	In CoastDOM v1, we defined "coastal water" as encompassing estuaries (salinity >
339	0.1) to the continental shelf break (water depth < 200 m). However, some locations, such
340	as deep fjords which are close to the coast cannot be classed as coastal due to
341	bathymetry (deeper than > 200 m). Therefore, we evaluated the inclusion of some
342	datasets on a case-by-case basis. For inclusion in the database, each DOM
343	measurement needed at a minimum to contain the following information (if reported in the
344	original publication or otherwise available):
345	- Country where samples were collected
346	- Latitude of measurement (in decimal units)
347	- Longitude of measurement (in decimal units)
348	- Year of sampling
349	- Month of sampling
350	- Sampling day (when available)
351	- Depth (m) at which the discrete sample were collected
352	- Temperature (°C) of the sample
353	- Salinity of the sample
354	- Dissolved organic carbon (DOC) concentration (µmol L <sup>-1</sup> )
355	- Method used to measure DOC concentration
356	- DOC - QA flag: Quality flag for DOC measurement
357	- Dissolved organic nitrogen (DON) concentration (µmol L <sup>-1</sup> )
358	- Total dissolved nitrogen (TDN) concentration (µmol L <sup>-1</sup> )
359	- Method used to measure TDN concentration
360	- TDN - QA flag: Quality flag for TDN measurement

- 360 TDN QA flag: Quality flag for TDN measurement
- 361 Dissolved organic phosphorus (DOP) concentration (µmol L<sup>-1</sup>)





- 362 Total dissolved phosphorus (TDP) concentration (µmol L<sup>-1</sup>)
- 363 Method used to measure TDP concentration
- 364 TDP QA flag: Quality flag for TDP measurement
- 365 Responsible person
- 366 Originator institution
- 367 Contact of data originator
- 368 It should be noted that in all entries, at least DOC, DON or DOP should have been
- 369 measured. In addition, we also included other relevant data, when available, in the
- 370 CoastDOM v1 dataset:
- Depth at the station where the sample was collected (Bottom depth, m).
- Total suspended solids (TSS) concentration (mg L<sup>-1</sup>)
- 373 Chlorophyll-*a* (Chl *a*) concentration (μg L<sup>-1</sup>)
- Chl a QA flag: Quality flag for chlorophyll-a measurement
- Sum of nitrate and nitrite (NO<sub>3</sub><sup>-</sup>+NO<sub>2</sub><sup>-</sup>) concentration (µmol L<sup>-1</sup>)
- 376 NO<sub>3</sub><sup>-</sup> + NO<sub>2</sub><sup>-</sup> QA flag: Quality flag for NO<sub>3</sub><sup>-</sup> + NO<sub>2</sub><sup>-</sup> measurement
- 377 Ammonium (NH<sub>4</sub><sup>+</sup>) concentration (μmol L<sup>-1</sup>)
- 378 NH<sub>4</sub><sup>+</sup> QA flag: Quality flag for NH<sub>4</sub><sup>+</sup> measurement
- 379 Soluble reactive phosphorus (HPO<sub>4</sub><sup>2-</sup>) concentration (μmol L<sup>-1</sup>)
- 380 HPO<sub>4</sub><sup>2-</sup> QA flag: Quality flag for HPO<sub>4</sub><sup>2-</sup> measurement
- 381 Particulate organic carbon (POC) concentration (μmol L<sup>-1</sup>)
- 382 Method used to measure POC concentration
- 383 POC QA flag: Quality flag for POC measurement
- Particulate nitrogen (PN) concentration (μmol L<sup>-1</sup>)
- 385 Method used to measure PN concentration
- 386 PN QA flag: Quality flag for PN measurement
- 387 Particulate phosphorus (PP) concentration (µmol L<sup>-1</sup>)





- 388 Method used to measure PP concentration
- 389 PP QA flag: Quality flag for PP measurement
- Dissolved inorganic carbon (DIC) concentration (µmol kg<sup>-1</sup>)
- 391 DIC QA flag: Quality flag for DIC measurement
- Total alkalinity (TA) concentration (μmol kg<sup>-1</sup>)
- 393 TA QA flag: Quality flag for TA measurement
- 394

395 Quality control of large datasets is crucial to ensure their reliability and usefulness. 396 Thus, we have not included data that were deemed compromised, such as records that 397 had not gone through quality control by the data originators. We also accepted a certain 398 degree of measurement error since multiple groups have been involved in the collection, 399 analysis, and/or compilation of the information. Some of these errors were corrected (e.g., 400 when a value was placed in a wrong column, or clearly inaccurate locations were 401 reallocated for consistency with the place of study), while others could not be rectified 402 (e.g., values showing clear signs of contamination) and were consequently excluded from 403 CoastDOM v1. It should also be noted that differences in analytical capabilities between 404 laboratories and individual measurement campaigns likely caused additional uncertainty. 405 Outliers, arising for example from contamination, were removed from the dataset. The 406 data were moreover screened for zero values (i.e., concentrations below the detection 407 limit or absence of data). In cases where concentrations were below the detection limit, 408 the zero values were replaced with half the value of the limit-of-detection. To ensure the 409 inclusion of only high-quality data, we only accepted entries with specific World Ocean 410 Circulation Experiment (WOCE) quality codes: "2- Acceptable measurement" and "6-411 Mean of replicate measurements". In our quality control assessments, we carefully 412 avoided overly strict criteria, known as "data grooming", which could potentially overlook 413 genuine patterns and changes in the dataset that may be significant over longer temporal 16





and/or wider spatial scales. Coastal waters are known to exhibit a wide range of
environmental concentrations, influenced by factors such as seasonality and local
anthropogenic activities. Consequently, these data points may encompass a wide
concentration range.

418

#### 419 **3.1 Summary of dissolved organic carbon (DOC) observations**

420 Measurements of DOC concentrations were conducted between 1978 to 2022, with a 421 total of 62339 individual data points (Table 1). The DOC concentrations ranged from 17 422 to 30327 µmol C L<sup>-1</sup> (average (Standard Deviation; SD): 182 (314) µmol C L<sup>-1</sup>; median: 103 µmol C L<sup>-1</sup>; Table 1). The majority (53%) of the concentrations fell within the range of 423 424 60 to 120 µmol C L<sup>-1</sup> (Fig. 1). A large number of DOC observations (17%) ranged between 425 300 and 600 µmol C L<sup>-1</sup>, which were predominantly collected in eutrophic and river-426 influenced coastal waters of the Northern Hemisphere, such as the Baltic Sea (Fig. 1). It 427 was observed that 75% of the DOC concentrations were higher than 77 µmol C L<sup>-1</sup>, while 428 25% of the measurements surpassed 228 µmol C L<sup>-1</sup> (Table 1).

429 Coastal environments that experience minimal continental runoff, such as Palmer 430 Station in Antarctica, typically exhibit low DOC concentrations. On the other hand, coastal 431 waters heavily influenced by humic-rich terrigenous inputs, such as the Sarawak region 432 in Malaysia, tended to have high DOC concentrations. In addition some extreme high 433 DOC concentrations were measured in the Derwent River in Australia which is impacted 434 by paper mill effluents. There has been a large increase in the number of DOC 435 observations after 1992 (Fig. 2), and those measurements were from a wide range of 436 locations. However, these observations were not evenly distributed, with the Southern 437 Hemisphere being relatively under-sampled, especially in the African, South American 438 and Antarctic continents (Fig. 2, 3).





#### 440 **3.2. Summary of dissolved organic nitrogen (DON) observations**

- The DON measurements were collected between 1990 and 2021, with a total of 20357 data points (Table 1). Concentrations of DON ranged from < 0.1 to 2095.3  $\mu$ mol N L<sup>-1</sup> (average (SD): 13.6 (30.4)  $\mu$ mol N L<sup>-1</sup>; median: 8.0  $\mu$ mol N L<sup>-1</sup>; Table 1), with the most common range (42%) for DON concentrations between 4 to 8  $\mu$ mol N L<sup>-1</sup> (Fig. 1). Overall 75% of DON concentrations were above 5.5  $\mu$ mol N L<sup>-1</sup>, while 25% were above 15.8  $\mu$ mol N L<sup>-1</sup> (Table 1).
- The lowest DON concentrations were recorded in Young Sound, Greenland, which receives direct run-off from the Greenland Ice Sheet, whereas the highest concentrations were detected during a flood event in the Richmond River Estuary, Australia. Since 1995, there has been a large increase in the number of DON measurements conducted in coastal waters globally (Fig. 2); however, the majority of those measurements have been in the Northern Hemisphere, mostly in Europe and the United States (Fig. 2, 3).
- 453

### 454 3.3. Summary of dissolved organic phosphorus (DOP) observations

- 455 CoastDOM v1 includes a total of 13534 DOP measurements, collected between 1990 456 and 2021 (Table 1). Overall, DOP concentrations ranged from < 0.10 to 84.27  $\mu$ mol P L<sup>-</sup> 457 <sup>1</sup> (average (SD): 0.34 (1.11)  $\mu$ mol P L<sup>-1</sup>; median: 0.18  $\mu$ mol P L<sup>-1</sup>; Table 2). The majority 458 (74%) of DOP concentrations were below 0.30  $\mu$ mol P L<sup>-1</sup> (Fig. 1). Analysis of the DOP 459 dataset revealed that 75% of the concentrations were above 0.11  $\mu$ mol P L<sup>-1</sup>, while 25% 460 were above 0.30  $\mu$ mol P L<sup>-1</sup> (Table 1).
- The lowest DOP concentrations were measured off the Kimberley Coast in Australia, while the highest concentrations were found in the Vasse-Wonnerup Estuary in the South west region of Australia. Similarly to DOC and DON, most of the DOP measurements have been conducted from the 1990s onwards, with a predominant focus in the Northern Hemisphere, particularly in Europe and the United States (Fig. 2, 3).





#### 466

#### 467 3.4. Summary of dissolved organic matter (DOM) observations

468 In CoastDOM v1 the number of measurements decreases progressively in the 469 sequence DOC > DON > DOP, reflecting both differences in the maturity of the analytical 470 methods and the greater focus on carbon cycling by the aquatic science community. In 471 addition the average DOC concentration in coastal waters (182 (314) µmol C L<sup>-1</sup>), was 472 13-fold greater than the average coastal DON concentrations 13.6 (30.4) µmol N L<sup>-1</sup>), 473 which was itself 39-fold greater than the average coastal DOP concentrations (0.34 (1.11) 474 µmol P L<sup>-1</sup>) (Table 1). Interestingly the coefficient of variation (C.V.) increased from DOC (173%) to DON (224%) and DOP (326%), which is related to that the % contribution of 475 476 refractory organic material decreases in the same sequence (Table 1).

477

#### 478 **3.5. Potential use of the dataset**

479 The use of the CoastDOM v1 dataset should be accompanied by the citation of this 480 paper and the inclusion of the correct doi-reference. CoastDOM v1 will be available in full 481 open access on the PANGEA homepage after acceptance of the manuscript, where it will 482 be available as a \*.csv file. The dataset includes a brief description of the metadata and 483 methods employed, with emphasis on measurement techniques and data units. We 484 chose the terminology most familiar to the ocean science community. It is important to 485 note that all data included in CoastDOM v1, as well as this manuscript, are considered 486 public domain; as such, a subset of this global dataset may also be present in previous 487 data compilations (e.g., Hansell et al., 2021). The list of citations and links referenced in 488 CoastDOM v1 also provide users with information as to how these data has been 489 previously used in publications or databases.

490

#### 491 3.6. Recommendations and conclusions





492 In CoastDOM v1, we have compiled available coastal DOM data in a single repository, 493 making it freely available to the research community. This compilation has established a 494 consistent global dataset, serving as a valuable information source to investigate a variety 495 of environmental questions and to explore spatial and temporal trends. We suggest a set 496 of recommendations for the future expansion of this global dataset. Firstly, our analysis 497 highlights a spatial bias, with a concentration of sampling efforts and/or data availability 498 predominantly concentrated in the Northern Hemisphere. The data gap in coastal DOM 499 measurements in the Southern Hemisphere needs to be addressed to provide a more 500 representative global understanding of the role of DOM in coastal water biogeochemistry. 501 Additionally, increased sampling efforts especially in the African and South American 502 continents are warranted due to the vulnerability of many coastal areas to climate change 503 and intensifying human activities, which will undoubtedly impact DOM biogeochemistry. 504 Further it is also worth noting that there is comparatively few data from coastal waters 505 affected by river discharge into the tropics, e.g., Amazon, Indian and Indonesian rivers 506 that together dominate freshwater inputs to the coastal ocean. Secondly, there is a need 507 for more comprehensive temporal and spatial datasets to capture the variability of DOM 508 levels in highly dynamic and productive coastal systems. Focused efforts should be made 509 to resolve these temporal and spatial changes. Thirdly, it is also important to collect and 510 report ancillary data, such as temperature, salinity, nutrient measurements, and 511 particulate components, to provide context and better understand the underlying 512 processes driving the observed DOM levels. Lastly, we strongly recommend that the 513 DOM research community conducts regular inter-calibration exercises to establish 514 standardised and interoperable methods, particularly for DON and DOP measurements. 515 This will ensure the comparability and reliability of data across different studies and 516 enhance our understanding of DON and DOP dynamics in coastal waters.





517 In light of ongoing global environmental changes, the mobilisation and open sharing of 518 existing data of important biogeochemical variables, such as the DOM pool, are crucial 519 for establishing baselines and determining global trends and changes in coastal waters. 520 The aim is to publish an updated version of the database periodically to determine global 521 trends of DOM levels in coastal waters, and we therefore encourage researchers to 522 submit new data to the corresponding author. The CoastDOM v1 dataset was developed 523 according to the FAIR principles regarding Findability, Accessibility, Interoperability and 524 Reusability of data. Thus, CoastDOM v1 will serve as a reliable open-source information 525 resource, enabling in-depth analyses and providing quality-controlled input data for large 526 scale ecosystem models.

527

#### 528 4. Data availability

529 The dataset available the is for review process Figshare at 530 https://figshare.com/s/512289eb43c4f8e8eaef). The dataset is furthermore submitted to 531 the PANGEA database and is currently waiting to be assigned a Doi number (Lønborg et 532 al., 2023). The file will be available as a \*.csv merged file and will be available in full open 533 access in the PANGEA database after acceptance of the manuscript.

534

#### 535 Competing interests

- 536 The authors declare no competing interests.
- 537 Author Contribution
- 538 C.L., C.C., and X.A.A-S started the initiative and finalised the data compilation. All co-
- 539 authors contributed data. C.L. wrote the manuscript with input from all co-authors.
- 540 Acknowledgement





541 Qi Chen is acknowledged for his very skilful help with Figure 3.

#### 542 Funding

543 During the drafting of the manuscript C.L. received funding from the Independent Research Fund Denmark Grant No. 1127-00033B. The monitoring data obtained from 544 545 Bermuda received funding from the Bermuda Government Department of Environment 546 and Natural Resources. A subset of the data obtained from UK coastal estuaries received 547 funding from the Natural Environment Research Council (Grant NE/N018087/1). Data 548 retrieved from the Palmer LTER data were collected with support from the Office of Polar 549 Programs, US National Science Foundation. Data obtained from the Great Barrier Reef 550 Marine Monitoring Program for Inshore Water Quality, which is a partnership between the 551 Great Barrier Reef Marine Park Authority, the Australian Institute of Marine Science, 552 James Cook University, and the Cape York Water Partnership. The contribution by P.K. 553 was supported by DiSeDOM project contract no. UMO-2019/33/B/ST10/01232 funded by 554 the NCN - National Science Centre, Poland. N.W. and A.M.-P. participated in this 555 synthesis effort with funding provided by the U.S. Department of Energy funded 556 COMPASS-FME project; the provided data was collected with funding from the PREMIS 557 Initiative, conducted under the Laboratory Directed Research and Development Program 558 at Pacific Northwest National Laboratory. The data obtained from the Levantine Sea (Med 559 Sea) received funding from the Scientific and Technological Research Council of Türkiye 560 (TÜBİTAK, 1001 program, Grant 115Y629).

561

#### 562 References

- Álvarez-Salgado, X. A., Nieto-Cid, M., and Rossel, P. E.: Dissolved Organic Matter, in:
   Marine Analytical Chemistry, edited by: Blasco, J., and Tovar-Sánchez, A., Springer
   International Publishing, Cham, 39-102, 10.1007/978-3-031-14486-8\_2, 2023.
- Bauer, J. E., Cai, W. J., Raymond, P. A., Bianchi, T. S., Hopkinson, C. S., and Regnier,
  P. A.: The changing carbon cycle of the coastal ocean, Nature, 504, 61-70,
- 568 10.1038/nature12857, 2013.
- 569 Boudreau, B. P. and Ruddick, B. R.: On a reactive continuum representation of organic 570 matter diagenesis, American Journal of Science, 291, 507-538, 1991.





571	Burdige, D. J. and Komada, T.: Sediment pore waters, in: Biogeochemistry of marine
572	dissolved organic matter, edited by: Hansen, D. A., and Carlson, C. A., Elsevier, 535-
573	577, 2014.
574	Carlson, C. A. and Hansell, D. A.: DOM Sources, Sinks, Reactivity, and Budgets, in:
575	Biogeochemistry of Marine Dissolved Organic Matter, edited by: Carlson, C. A., and
576	Hansell, D. A., Elsevier Science & Technology, 65-126, 10.1016/b978-0-12-405940-
577	5.00003-0, 2015.
578	Carreira, C., Talbot, S., and Lønborg, C.: Bacterial consumption of total and dissolved
579	organic carbon in the Great Barrier Reef, Biogeochemistry, 10.1007/s10533-021-
580	00802-x, 2021.
581	Cauwet, G.: DOM in coastal areas, in: Biogeochemistry of Dissolved organic matter,
582	edited by: Hansell, D., and Carlson, C. A., Academic Press, London, 579-609, 2002.
583	Chin, W. C., Orellana, M. V., and Verdugo, P.: Spontaneous assembly of marine
584	dissolved organic matter into polymer gels, Nature, 391, 568-572, 1998.
585	Foreman, R. K., Björkman, K. M., Carlson, C. A., Opalk, K., and Karl, D. M.: Improved
586	ultraviolet photo-oxidation system yields estimates for deep-sea dissolved organic
587	nitrogen and phosphorus, Limnology and Oceanography: Methods, 17, 277-291,
588	https://doi.org/10.1002/lom3.10312, 2019.
589	García-Martín, E. E., Sanders, R., Evans, C. D., Kitidis, V., Lapworth, D. J., Rees, A. P.,
590	Spears, B. M., Tye, A., Williamson, J. L., Balfour, C., Best, M., Bowes, M., Breimann,
591	S., Brown, I. J., Burden, A., Callaghan, N., Felgate, S. L., Fishwick, J., Fraser, M.,
592	Gibb, S. W., Gilbert, P. J., Godsell, N., Gomez-Castillo, A. P., Hargreaves, G., Jones,
593	O., Kennedy, P., Lichtschlag, A., Martin, A., May, R., Mawji, E., Mounteney, I.,
594	Nightingale, P. D., Olszewska, J. P., Painter, S. C., Pearce, C. R., Pereira, M. G.,
595	Peel, K., Pickard, A., Stephens, J. A., Stinchcombe, M., Williams, P., Woodward, E.
596	M. S., Yarrow, D., and Mayor, D. J.: Contrasting Estuarine Processing of Dissolved
597	Organic Matter Derived From Natural and Human-Impacted Landscapes, Global
598	Biogeochemical Cycles, 35, e2021GB007023,
599	https://doi.org/10.1029/2021GB007023, 2021.
600	Halewood, E., Opalk, K., Custals, L., Carey, M., Hansell, D. A., and Carlson, C. A.:
601	Determination of dissolved organic carbon and total dissolved nitrogen in seawater
602	using High Temperature Combustion Analysis, Frontiers in Marine Science, 9,
603	10.3389/fmars.2022.1061646, 2022.
604	Hansell, D. A.: Dissolved Organic Carbon Reference Material Program, Eos,
605 606	Transactions American Geophysical Union, 86, 318-318,
606 607	https://doi.org/10.1029/2005EO350003, 2005. Hansell, D. A., Carlson, C. A., Amon, R. M. W., Álvarez-Salgado, X. A., Yamashita, Y.,
608	Romera-Castillo, C., and Bif, M. B.: Compilation of dissolved organic matter (DOM)
608 609	data obtained from global ocean observations from 1994 to 2021. Version 2. (NCEI
610	Accession 0227166) [dataset], doi.org/10.25921/s4f4-ye35, 2021.
611	lavorivska, L., Boyer, E. W., and DeWalle, D. R.: Atmospheric deposition of organic
612	carbon via precipitation, Atmospheric Environment, 146, 153-163,
613	https://doi.org/10.1016/j.atmosenv.2016.06.006, 2016.
614	Kerner, M., Hohenberg, H., Ertl, S., Reckermann, M., and Spitzy, A.: Self-organization
615	of dissolved organic matter tomicelle-like microparticles in river water, Nature, 422,
616	150-154, 2003.
617	Laruelle, G. G., Dürr, H. H., Lauerwald, R., Hartmann, J., Slomp, C. P., Goossens, N.,
618	and Regnier, P. A. G.: Global multi-scale segmentation of continental and coastal
619	waters from the watersheds to the continental margins, Hydrol. Earth Syst. Sci., 17,
620	2029-2051, 10.5194/hess-17-2029-2013, 2013.





621	Laurenceau-Cornec, E. C., Trull, T. W., Davies, D. M., De La Rocha, C. L., and Blain,
622	S.: Phytoplankton morphology controls on marine snow sinking velocity, Marine
623	Ecology Progress Series, 520, 35-56, 2015.
624	Lønborg, C. and Álvarez-Salgado, X. A.: Recycling versus export of bioavailable
625	dissolved organic matter in the coastal ocean and efficiency of the continental shelf
626	pump, Global biogeochemical cycles 26, GB3018, 10.1029/2012GB004353, 2012.
627	Lønborg, C., Álvarez-Salgado, X. A., Davidson, K., and Miller, A. E. J.: Production of
628	bioavailable and refractory dissolved organic matter by coastal heterotrophic
629	microbial populations, Estuarine, Coastal and Shelf science, 82, 682–688,
630	10.1016/j.ecss.2009.02.026, 2009.
631	Lønborg, C., Álvarez-Salgado, X. A., Duggan, S., and Carreira, C.: Organic matter
632	bioavailability in tropical coastal waters: The Great Barrier Reef, Limnology and
633	Oceanography, 63, 1015-1035, 10.1002/lno.10717, 2018.
634	Lønborg, C., Carreira, C., Jickells, T., and Álvarez-Salgado, X. A.: Impacts of Global
635	Change on Ocean Dissolved Organic Carbon (DOC) Cycling, Frontiers in Marine
636	Science, 7, 466, 10.3389/fmars.2020.00466, 2020.
637	Lønborg, C., Carreira, C., Abril, G., Agustí, S., Amaral, V., Andersson, A., Arístegui, J.,
638	Bhadury, P., Bif, M. B., Borges, A. V., Bouillon, S., Calleja, M. L., Cotovicz Jr, L. C.,
639	Cozzi, S., Doval, M. D., Duarte, C. M., Eyre, B. D., Fichot, C. G., García-Martín, E.
640	E., Garzon-Garcia, A., Giani, M., Gonçalves-Araujo, R., Gruber, R., Hansell, D. A.,
641	Hashihama, F., He, D., Holding, J. M., Hunter, W. R., Ibánhez, J. S., Ibello, V., Jiang,
642	S., Kim, G., Klun, K., Kowalczuk, P., Kubo, A., Weng Lee, C., Lopes, C., Maggioni,
643	F., Magni, P., Marrase, C., Martin, P., McCallister, S. L., McCallum, R., Medeiros, P.
644	M., Morán, X. A. G., Muller-Karger, F., Myers-Pigg, A., Norli, M., Oakes, J. M.,
645	Osterholz, H., Park, H., Paulsen, M. L., Rosentreter, J. A., Rueda-Roa, D., Santinelli,
646	C., Shen, Y., Teira, E., Tinta, T., Uher, G., Wakita, M., Ward, N., Watanabe, K., Xin,
647	Y., Yamashita, Y., Yang, L., Yeo, J., Yuan, H., Zheng, Q., and Alvarez-Salgado, X.
648	A.: A global database of dissolved organic matter (DOM) measurements in coastal
649	waters (CoastDOM v1) [dataset], 2023.
650	Mackenzie, F. T., De Carlo, E. H., and Lerman, A.: 5.10 - Coupled C, N, P, and O
651	Biogeochemical Cycling at the Land–Ocean Interface, in: Treatise on Estuarine and
652	Coastal Science, edited by: Wolanski, E., and McLusky, D., Academic Press,
653	Waltham, 317-342, https://doi.org/10.1016/B978-0-12-374711-2.00512-X, 2011.
654	Mopper, K., Kieber, D. J., and Stubbins, A.: Marine Photochemistry of Organic Matter,
655	in: Biogeochemistry of Marine Dissolved Organic Matter, edited by: Carlson, C. A.,
656	and Hansell, D. A., 389-450, 10.1016/b978-0-12-405940-5.00008-x, 2015.
657	Osterholz, H., Burmeister, C., Busch, S., Dierken, M., Frazão, H. C., Hansen, R.,
658	Jeschek, J., Kremp, A., Kreuzer, L., and Sadkowiak, B.: Nearshore dissolved and
659	particulate organic matter dynamics in the southwestern Baltic Sea: environmental
660	drivers and time series analysis (2010–2020), Frontiers in Marine Science, 8,
661	795028, 2021.
662	Raymond, P. A. and Spencer, R. G. M.: Riverine DOM, in: Biogeochemistry of Marine
663	Dissolved Organic Matter, edited by: Hansell, D. A., and Carlson, C. A., Elsevier, Amsterdam, 509-533, 10.1016/b978-0-12-405940-5.00011-x, 2015.
664 665	
665 666	Santos, I. R., Burdige, D. J., Jennerjahn, T. C., Bouillon, S., Cabral, A., Serrano, O.,
666 667	Wernberg, T., Filbee-Dexter, K., Guimond, J. A., and Tamborski, J. J.: The renaissance of Odum's outwelling hypothesis in 'Blue Carbon' science, Estuarine,
668	Coastal and Shelf Science, 255, 10.1016/j.ecss.2021.107361, 2021.
669	Sharp, J. H., Rinker, K. R., Savidge, K. B., Abell, J., Benaim, J. Y., Bronk, D., Burdige,
670	D. J., Cauwet, G., Chen, W., Doval, M. D., Hansell, D., Hopkinson, C., Kattner, G.,
671	Koumover N. McClethene K. L. Marting, L. Martin, N. Nagal, K. Ogove, H.

Kaumeyer, N., McGlathery, K. J., Merriam, J., Morley, N., Nagel, K., Ogawa, H.,





- Pollard, C., Pujo-Pay, M., Raimbault, P., Sambrotto, R., Seitzinger, S., Spyres, G.,
- Tirendi, F., Walsh, T. W., and Wong, C. S.: A preliminary methods comparison for
- 674 measurement of dissolved organic nitrogen in seawater, Marine Chemistry, 78, 171-675 184, 2002.
- 75 184, 2002.
- Taniguchi, M., Dulai, H., Burnett, K. M., Santos, I. R., Sugimoto, R., Stieglitz, T., Kim,
- 677 G., Moosdorf, N., and Burnett, W. C.: Submarine Groundwater Discharge: Updates
- on Its Measurement Techniques, Geophysical Drivers, Magnitudes, and Effects,
  Frontiers in Environmental Science, 7, 10.3389/fenvs.2019.00141, 2019.
- Valderrama, J. C.: The simultaneous analysis of total nitrogen and total phosphorus in
   natural waters, Marine Chemistry, 10, 109-122, 1981.
- Wada, S., Aoki, M. N., Mikami, A., Komatsu, T., Tsuchiya, Y., Sato, T., Shinagawa, H.,
- and Hama, T.: Bioavailability of macroalgal dissolved organic matter in seawater,
- 684 Marine Ecology Progress Series, 370, 33-44, 2008





685	Figure	legends
-----	--------	---------

- Figure 1. Histograms showing the distribution of observations for a) dissolved organic
   carbon (DOC), b) nitrogen (DON) and c) phosphorus (DOP), within defined
   concentration ranges in the coastal ocean. Note that the concentration ranges are not
- 689 uniform in all cases due to the large difference in concentration levels.
- 690 Figure 2. a) Cumulative number of observations for dissolved organic carbon (DOC),
- 691 nitrogen (DON), and phosphorus (DOP). Number of observations shown as a function
  692 of b) latitude, and c) longitude, grouped into bins of 10° latitude or longitude.
- Figure 3. Global distribution of observations included in CoastDOM v1 for a) dissolved organic carbon (DOC), b) nitrogen (DON), and c) phosphorus (DOP). The black dots on the map represent the reported data that are included in the CoastDOM v1 database. Histograms show the distribution of observations for DOC, DON and DOP within defined concentration ranges in the continents where measurements are available. Maps were created using the GIS shape file obtained from Laurelle et al.
- 699 (2013)






**Table 1.** Descriptive statistics for the dissolved organic carbon (DOC), dissolved organic
nitrogen (DON), and dissolved organic phosphorus (DOP) measurements included in the
CoastDOM v1 dataset. The minimum (Min), maximum (Max), average values (Avg.) and
standard deviation (SD), coefficient of variation (CV %), median, 25th and 75th
percentiles (perc.) and number of samples (N) for each variable are shown.

	DOC	DON	DOP
	µmol C L⁻¹	µmol N L⁻¹	µmol P L⁻¹
Min	17	< 0.1	< 0.01
Max	30327	2095.3	84.27
Avg. (SD)	182 (314)	13.6 (30.4)	0.34 (1.11)
Median	103	8.0	0.18
CV %	173	224	326
25th perc.	77	5.5	0.11
75th perc.	228	15.8	0.30
Ν	62339	20357	13534
















709 Figure 2.





