



HAL
open science

Area preserving homeomorphisms of surfaces with rational rotational direction

Pierre-Antoine Guihéneuf, Patrice Le Calvez, Alejandro Passeggi

► **To cite this version:**

Pierre-Antoine Guihéneuf, Patrice Le Calvez, Alejandro Passeggi. Area preserving homeomorphisms of surfaces with rational rotational direction. 2023. hal-04267469

HAL Id: hal-04267469

<https://hal.science/hal-04267469>

Preprint submitted on 1 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

AREA PRESERVING HOMEOMORPHISMS OF SURFACES WITH RATIONAL ROTATIONAL DIRECTION

PIERRE-ANTOINE GUIHÉNEUF, PATRICE LE CALVEZ,
AND ALEJANDRO PASSEGGI

ABSTRACT. Let S be a closed surface of genus $g \geq 2$, furnished with a Borel probability measure λ with total support. We show that if f is a λ -preserving homeomorphism isotopic to the identity such that the rotation vector $\text{rot}_f(\lambda) \in H_1(S, \mathbb{R})$ is a multiple of an element of $H_1(S, \mathbb{Z})$, then f has infinitely many periodic orbits.

Moreover, these periodic orbits can be supposed to have their rotation vectors arbitrarily close to the rotation vector of any fixed ergodic Borel probability measure.

Keywords: Rotation vector, maximal isotopy, transverse foliation

MSC 2020: 37C25 37E30, 37E45

CONTENTS

1. Introduction	2
1.1. Rotation vector	2
1.2. The main theorem	3
1.3. Idea of the proof	7
1.4. Acknowledgements	8
2. Definitions, notations and preliminaries	8
2.1. Loops and paths	8
2.2. Poincaré-Birkhoff theorem	8
2.3. Homeomorphisms of hyperbolic surfaces	9
2.4. Caratheodory theory of prime ends	10
2.5. Rotational topological horseshoes	10
3. Foliations on surfaces	11
3.1. \mathcal{F} -transverse intersections	11
3.2. Recurrence, equivalence and accumulation	12
3.3. Strips	13
3.4. More about the accumulation property	14
4. Forcing theory	20
4.1. Maximal isotopies and transverse foliations	20
4.2. Forcing theory in the annular covering space	25
5. Proof of the main theorem	31
References	37

1. INTRODUCTION

1.1. Rotation vector. If S is a smooth compact boundaryless oriented surface of genus g , we denote $\text{Homeo}(S)$ the space of homeomorphisms of S furnished with the C^0 -topology. This topology coincides with the uniform topology because S is compact. The path-connected component of the identity map Id , usually called the space of homeomorphisms *isotopic to the identity*, will be denoted $\text{Homeo}_*(S)$. A continuous path $I = (f_t)_{t \in [0,1]}$ joining the identity to a map $f \in \text{Homeo}_*(S)$ is called an *identity isotopy* of f . We call *trajectory* of a point $z \in S$ defined by I the path $I(z) : t \mapsto f_t(z)$ joining z to $f(z)$.

By compactness of S , one knows by Krylov-Bogolioubov's theorem that the set $\mathcal{M}(f)$ of f -invariant Borel probability measures is not empty. More precisely it is a non empty compact convex subset of the space \mathcal{M} of Borel probability measures furnished with the weak* topology. Remind that the *support* of μ , denoted $\text{supp}(\mu)$, is the smallest closed set of μ -measure 1.

Let us recall the definition of the *rotation vector* of a measure $\mu \in \mathcal{M}(f)$ (see [Mats1], [Pol] or [Sc]). Let $I = (f_t)_{t \in [0,1]}$ be an identity isotopy of f . Fix $z \in S$. The homotopy class of $I(z)$, relative to the endpoints, contains a smooth path γ joining z to $f(z)$. If α is a closed 1-form, the quantity $\int_\gamma \alpha$ does not depend on the choice of γ and we denote it $\int_{I(z)} \alpha$. It is equal to $h(f(z)) - h(z)$ if α is exact and h is a primitive of α . One gets a real valued morphism $\alpha \mapsto \int_S \left(\int_{I(z)} \alpha \right) d\mu(z)$ defined on the space of closed 1-forms, that vanishes on the space of exact 1-forms because μ is invariant by f . So, it induces a natural linear form on the first cohomology group $H^1(S, \mathbb{R})$. Hence, there exists a homology class $\text{rot}_I(\mu) \in H_1(S, \mathbb{R})$, uniquely defined by the equation

$$\langle [\alpha], \text{rot}_I(\mu) \rangle = \int_S \left(\int_{I(z)} \alpha \right) d\mu(z),$$

where α is any closed 1-form, $[\alpha] \in H^1(S, \mathbb{R})$ its cohomology class and

$$\langle \cdot, \cdot \rangle : H^1(S, \mathbb{R}) \times H_1(S, \mathbb{R}) \rightarrow \mathbb{R}$$

the natural bilinear form. By definition $\text{rot}_I(\mu) \in H_1(S, \mathbb{R})$ is the rotation vector of μ (for the isotopy I). It is well known that two identity isotopies of f are homotopic relative to the ends if the genus of S is larger than 1 (see [H]). In that case, $\int_{I(z)} \alpha$ does not depend on I and one can write

$$\text{rot}_f(\mu) = \text{rot}_I(\mu).$$

If O is a periodic orbit of f , one can define the rotation vector $\text{rot}_I(O)$ of O (or $\text{rot}_f(O)$ if the genus of S is larger than 1) as being equal to the rotation vector of μ_O , where μ_O is the probability measure equidistributed on O . In particular we have $\text{rot}_I(O) = 0$ if O is a contractible periodic orbit, which means that the loop $I^q(z)$ is homotopic to zero, if $z \in O$.

Let us give an equivalent definition. Furnish S with a Riemannian metric and for every points z, z' in S , choose a path $\gamma_{z,z'}$ joining z to z' in such a way that the lengths of the paths $\gamma_{z,z'}$ are uniformly bounded. For every

$z \in S$, and every $n \geq 1$, consider the path

$$I^n(z) = I(z)I(f(z)) \cdots I(f^{n-1}(z))$$

defined by concatenation, and the loop

$$\Gamma_n(z) = I^n(z)\gamma_{f^n(z),z}.$$

One can prove that there exists a μ -integrable function $\text{rot}_f : S \rightarrow H_1(S, \mathbb{R})$ such that for μ -almost every point $z \in S$, the sequence $[\Gamma_n(z)]/n$ converges to $\text{rot}_f(z)$. This allows to define

$$\text{rot}_I(\mu) = \int \text{rot}_f(z) d\mu(z).$$

Let us give a last definition that will be used in this article. In the whole text we will write $[\Gamma] \in H_1(S, \mathbb{Z})$ for the homology class of an oriented loop $\Gamma \subset S$. Let $U \subset S$ be a topological open disk (meaning a simply connected domain) such that $\mu(U) \neq 0$. Write $\varphi_U : U \rightarrow U$ for the first return map of f and $\tau_U : U \rightarrow \mathbb{N} \setminus \{0\}$ for the time of first return map. These maps are defined μ -almost everywhere on U . Kac's Lemma [K] tells us that φ_U preserves the measure $\mu|_U$ and that τ_U is $\mu|_U$ -integrable, and that moreover

$$\int_U \tau_U d\mu = \mu \left(\bigcup_{k \geq 0} f^k(U) \right) = \mu \left(\bigcup_{k \in \mathbb{Z}} f^k(U) \right).$$

We also denote by μ_U the normalized probability measure $\mu|_U/\mu(U)$. One can construct a map $\rho_U : U \rightarrow H_1(S, \mathbb{Z})$ defined μ_U -almost everywhere as follows: if $\varphi_U(z)$ is well defined, one closes the trajectory $I^{\tau_U(z)-1}(z)$ with a path γ contained in U that joins $\varphi_U(z)$ to z , and set $\rho_U(z) = [I^{\tau_U(z)-1}(z)\gamma]$, noting that $[I^{\tau_U(z)-1}(z)\gamma]$ is independent of the choice of γ . If the genus of S is bigger than 1 (what we suppose from now), then this map does not depend on the choice of I . It is easy to prove that the map ρ_U/τ_U is uniformly bounded on U and consequently that ρ_U is μ_U -integrable. So, by Birkhoff's theorem, there exist μ_U -integrable functions $\rho_U^* : U \rightarrow H_1(S, \mathbb{R})$ and $\tau_U^* : U \rightarrow \mathbb{R}$ such that for μ_U -almost every point z it holds that

$$(1) \quad \lim_{n \rightarrow +\infty} \frac{1}{n} \sum_{k=0}^{n-1} \rho_U(\varphi_U^k(z)) = \rho_U^*(z), \quad \lim_{n \rightarrow +\infty} \frac{1}{n} \sum_{k=0}^{n-1} \tau_U(\varphi_U^k(z)) = \tau_U^*(z).$$

These quantities are related to the rotation number by the fact that for μ_U -almost every point z , we have $\text{rot}_f(z) = \rho_U^*(z)/\tau_U^*(z)$.

1.2. The main theorem. Let us begin this section by introducing the notion of *homotopical interval of rotation*. If S is an oriented closed surface, denote $\mathcal{FHL}(S)$ the free homotopy loop space of S . For every $\kappa \in \mathcal{FHL}(S)$ and every $\Gamma \in \kappa$, the homology class $[\Gamma] \in H_1(S, \mathbb{Z})$ does not depend on the choice of Γ , we denote it $[\kappa]$. If $\Gamma : \mathbb{R}/\mathbb{Z} \rightarrow S$ is a loop and k an integer, we can define the loop $\Gamma^k : t \mapsto \Gamma(kt)$. For every $\kappa \in \mathcal{FHL}(S)$, every $\Gamma \in \kappa$ and every $k \in \mathbb{Z}$, the free homotopy class of Γ^k does not depend on the choice of Γ , we denote it κ^k . A homotopical interval of rotation of $f \in \text{Homeo}_*(S)$ is a couple (κ, r) , where $\kappa \in \mathcal{FHL}(S)$ and r is a positive integer, that satisfies the following: there exists an integer $s > 0$ such that for every $p/q \in [0, 1] \cap \mathbb{Q}$, one can find a point $z \in S$ of period at least q/s ,

such that the loop naturally defined by $I^{rq}(z)$ belongs to κ^p . In particular, we have $\text{rot}_f(z) = p/(rq)[\kappa]$.

Let us state the main result of the article.

Theorem A. *Let S be an oriented closed surface of genus $g \geq 2$. If $f \in \text{Homeo}_*(S)$ preserves a Borel probability measure λ such that $\text{supp}(\lambda) = S$ and $\text{rot}_f(\lambda) \in \mathbb{R}H_1(S, \mathbb{Z})$, then f has infinitely many periodic points.*

More precisely, for every ergodic measure $\nu \in \mathcal{M}(f)$ that is not a Dirac measure at a contractible fixed point and every neighborhood \mathcal{U} of $\text{rot}_f(\nu)$ in $H_1(S, \mathbb{R})$, there exists a homotopical interval of rotation (κ, r) such that $[\kappa]/r \in \mathcal{U}$.

Note that if f satisfies the hypotheses of the theorem and is different from identity, then by ergodic decomposition it has an ergodic invariant probability measure ν that is not supported on a fixed point. Theorem A applies and implies the existence of a homotopical interval of rotation; in particular f has an infinite number of periodic points, of arbitrarily large period, and of rotation vector arbitrarily close to 0. If $\text{rot}_f(\lambda) \neq 0$, the measure ν can be chosen such that $\text{rot}_f(\nu) \neq 0$ and consequently, f has periodic orbits of arbitrary large period and with non zero rotation vector. In any case, any ergodic Borel probability measure, supported on a contractible fixed point or not, has its rotation vector approximated by rotation vectors of an infinite number of periodic points. Remark that this property is also true for f equal to the identity.

Before explaining what are the two different sources of creation of homotopical interval of rotation in Paragraph 1.3, let us comment Theorem A. We start by giving a direct application. If ω is a smooth area form on S , denote $\text{Diff}_\omega^r(S)$, $1 \leq r \leq \infty$, the space of C^r diffeomorphisms of S preserving ω , endowed with the C^r -topology, and $\text{Diff}_{\omega,*}^r(S)$ the connected component of $\text{Diff}_\omega^r(S)$ that contains the identity. It is a classical fact that $\text{Diff}_{\omega,*}^r(S) = \text{Diff}_\omega^r(S) \cap \text{Homeo}_*(S)$.

Corollary 1.1. *Suppose that $g \geq 2$. Then, for any $1 \leq r \leq \infty$, the set of maps $f \in \text{Diff}_{\omega,*}^r(S)$ that have infinitely many periodic points is dense in $\text{Diff}_{\omega,*}^r(S)$.*

Proof. There is no loss of generality by supposing that the measure μ_ω naturally defined by ω is a probability measure. Note that the map $f \mapsto \text{rot}_f(\mu_\omega)$ is a morphism defined on $\text{Diff}_\omega^r(S)$. One can find a family of simple loops $(\Gamma_i)_{1 \leq i \leq 2g}$ in S such that the family $([\Gamma_i])_{1 \leq i \leq 2g}$ generates $H_1(S, \mathbb{R})$. For every $i \in \{1, \dots, 2g\}$ consider a closed tubular neighborhood W_i of Γ_i . It is easy to construct a divergence free smooth vector field ζ_i supported on W_i with an induced flow $(h_i^t)_{t \in \mathbb{R}}$ satisfying $\text{rot}_{h_i^t}(\mu_\omega) = t[\Gamma_i]$. For every $t = (t_1, \dots, t_{2g}) \in \mathbb{R}^{2g}$, define $f^t = h_1^{t_1} \circ \dots \circ h_{2g}^{t_{2g}} \circ f$. We have

$$\text{rot}_{f^t}(\mu_\omega) = \text{rot}_f(\mu_\omega) + \sum_{i=1}^{2g} \text{rot}_{h_i^{t_i}}(\mu_\omega) = \text{rot}_f(\mu_\omega) + \sum_{i=1}^{2g} t_i[\Gamma_i].$$

So, we can find t “arbitrarily small” such that $\text{rot}_{f^t}(\mu_\omega) \in H_1(S, \mathbb{Q})$. \square

Remark. A very close version of the theorem has been proved independently by Rohil Prasad. A very strong recent result of Cristofer-Prasad-Zhang [CPrZ], whose proof uses Periodic Floer Homology theory, asserts that if ω is a smooth area form on S , then for every $k \in \mathbb{N} \cup \{\infty\}$, the set of maps $f \in \text{Diff}_\omega^k(S)$ that have a dense set of periodic points is dense in $\text{Diff}_\omega^k(S)$ (which of course implies that Corollary 1.1 holds in the smooth category, see also [EH] and [CPoPrZ]). The following result is used in their proof: in the case where $f \in \text{Diff}_{\omega,*}^\infty(S)$ and $\text{rot}_f(\mu_\omega) \in H_1(S, \mathbb{Q}) \setminus \{0\}$, the map f has a periodic orbit with non zero rotation vector. Moreover they find an explicit upper bound of the period related to $\text{rot}(\mu_\omega)$ and to the genus of S . As explained by Prasad [Pr] in a recent note, a simple approximation process permits to extend this result to the case where $f \in \text{Homeo}_*(S)$ preserves μ_ω and satisfies $\text{rot}_f(\mu_\omega) \in H_1(S, \mathbb{Q}) \setminus \{0\}$. Moreover a blow-up argument allows to extend the result in the case where $\text{rot}_f(\mu_\omega) \in \mathbb{R}H_1(S, \mathbb{Z}) \setminus \{0\}$. Consequently it holds that f has infinitely many periodic orbits of period arbitrarily large. This last point is a consequence of previous works where area preserving homeomorphisms with finitely many periodic points are characterized ([AdT] in the case of the torus, [Lec3] in the case of surfaces with higher genus). Using Oxtoby-Ulam theorem [OxU] and the fact that every invariant probability measure is the barycenter of two invariant probability measures, the first one atomic and the second one with no atom, the measure μ_ω can be replaced with any probability measure with total support. In the present article, we give some precisions about the structure of the periodic points.

Remark. The theorem is untrue in the sphere and in the torus. Indeed, suppose that $\alpha \in \mathbb{R} \setminus \mathbb{Q}$.

The diffeomorphism f_α of the Riemann sphere \mathbb{S}^2 defined as follows

$$f_\alpha(z) = \begin{cases} \infty & \text{if } z = \infty, \\ e^{2i\pi\alpha} z & \text{if } z \in \mathbb{C}, \end{cases}$$

preserves a probability measure μ_ω associated to an area form and has no periodic point but 0 and ∞ . If I is an identity isotopy of f , then $\text{rot}_I(\mu_\omega) = 0$ because $H_1(\mathbb{S}^2, \mathbb{R}) = 0$.

The diffeomorphism

$$g_\alpha : \mathbb{R}^2/\mathbb{Z}^2 \longrightarrow \mathbb{R}^2/\mathbb{Z}^2 \\ (x, y) \longmapsto (x + (\alpha + \mathbb{Z}), y)$$

preserves the area form $\omega = dx \wedge dy$ and has no periodic orbit. If $I = (R_{t\alpha})_{t \in [0,1]}$, then we have $\text{rot}_I(\mu_\omega) = \alpha(1, 0) \in \mathbb{R}H_1(\mathbb{T}^2, \mathbb{Z})$.

Remark. In particular, the theorem asserts that if $\text{rot}_f(\lambda) = 0$, then there exists infinitely many periodic orbits. Moreover the set of periods is infinite if f is not the identity because there exist ergodic invariant measures that are not Dirac measures at a fixed point. This result, that admits a version for the case $g = 1$, was already known (see [Lec2]). It is a generalization of a result stated in the differential setting (see [FH]) which itself is the two dimensional version of what is called Conley conjecture, later proved in any

dimension (see [G]). Note that in [Lec2] it is proved that if f has finitely many fixed points, then there are infinitely many contractible periodic orbits.

Remark. The theorem was well known for the time one map of a conservative flow. Indeed, let X be a (time independent) vector field of class C^1 whose flow preserves ω . The equalities $0 = L_X\omega = i_X d\omega + di_X\omega$ tell us that the 1-form $\beta = i_X\omega$ is closed. Moreover it is invariant by the flow of X because $L_X\beta = i_X d\beta + di_X(i_X\omega) = 0$. If f is the time one map of the flow $(f^t)_{t \in \mathbb{R}}$ of X , then, denoting $I = (f^t)_{t \in [0,1]}$, we know that for every closed 1-form α , we have

$$\begin{aligned} \langle [\alpha], \text{rot}_I(\mu_\omega) \rangle &= \int_S \left(\int_{I(z)} \alpha \right) d\mu_\omega(z) \\ &= \int_S \left(\int_0^1 \alpha(X(f_t(z))) dt \right) d\mu_\omega(z) \\ &= \int_0^1 \left(\int_S \alpha(X(f_t(z))) d\mu_\omega(z) \right) dt \\ &= \int_S \alpha(X(z)) d\mu_\omega(z) \end{aligned}$$

Noting that $0 = i_X(\alpha \wedge \omega) = i_X\alpha \wedge \omega - \alpha \wedge i_X\omega$ we deduce that

$$\langle [\alpha], \text{rot}_I(\mu_\omega) \rangle = \int_S \alpha \wedge \beta.$$

The fact that $\text{rot}_I(\mu_\omega) \in \mathbb{R}H_1(S, \mathbb{Z})$ implies that $[\beta] \in \mathbb{R}H^1(S, \mathbb{Z})$. Suppose for instance that $[\beta] \in H^1(S, \mathbb{Z})$. Then there exists a function $H : S \rightarrow \mathbb{R}/\mathbb{Z}$ of class C^2 such that $\beta = dH$. Indeed, let us fix $z_0 \in S$. For every point $z \in M$, the value modulo 1, denoted $H(z)$, of $\int_\gamma \beta$ does not depend on the C^1 path γ joining z_0 to z . We get in that way a function $H : S \rightarrow \mathbb{R}/\mathbb{Z}$ of class C^2 such that $\beta = dH$. It is invariant by the flow of X because

$$L_X H = i_X dH + di_X H = i_X \beta = i_X(i_X\omega) = 0.$$

Denote $\text{sing}(X)$ the set of singular points of X . Remind that the α -limit set $\alpha(z)$ and the ω -limit set $\omega(z)$ of a point $z \in S$ are the sets of subsequential limits of the sequences $(f^{-n}(z))_{n \geq 0}$ and $(f^n(z))_{n \geq 0}$ respectively. If z is not singular, either the orbit of z is periodic, or its limit sets $\alpha(z)$ and $\omega(z)$ are contained in $\text{sing}(X)$. In particular the ergodic invariant probability measures that are non supported on a singular point are supported on a periodic orbit of f lying on a periodic orbit of the flow with rational period, or supported on a whole periodic orbit of the flow with irrational period. The union W of periodic orbits of the flow is non empty (by Sard's theorem) and open. Moreover every connected component V of W is annular (meaning homeomorphic to $\mathbb{R}/\mathbb{Z} \times \mathbb{R}$). The genus being at least two, there exist singular points. Furthermore S is not a sphere. It implies that there exists at least one end of V such that for every sequence $(z_n)_{n \geq 0}$ in V converging to this end, the period of z (for the flow) converges to $+\infty$. So the period is not constant on V . It implies that f has periodic points of arbitrarily large period. More precisely, the loops Γ that appear in the Theorem are the simple loops contained in such a component V that are non homotopic to zero in V and suitably oriented. Note that if $\text{rot}_I(\mu_\omega) \neq 0$, there exists

at least one connected component V of W such that $i_*(H_1(V, \mathbb{Z})) \neq \{0\}$, where $i_* : H_1(V, \mathbb{Z}) \rightarrow H_1(S, \mathbb{Z})$ is the morphism naturally defined by the inclusion map $i : V \rightarrow S$, meaning that the periodic points in V have non zero rotation vector.

Remark. The hypothesis $\text{rot}_f(\lambda) \in \mathbb{R}H_1(S, \mathbb{Z})$ is necessary to get the theorem. Indeed one can find smooth vector fields with finitely many singular points, whose flows preserves an area form ω and such that every orbit is dense if not reduced to a singular point. The time one map of this flow f has no periodic points but the singular points. Of course it holds that $\text{rot}_f(\lambda) \notin \mathbb{R}H_1(S, \mathbb{Z})$. Classical examples are given by translation flows in a minimal direction.

Remark. Corollary 1.1 was already known. In fact we have a much stronger result: the set of maps $f \in \text{Diff}_\omega^r(S)$ that have a hyperbolic periodic point with transverse homoclinic intersection, is an open and dense subset of $\text{Diff}_\omega^r(S)$ (see [LecSa]). This result has been known for a long time in the case where $g \leq 1$ (see [Ad], [AdT], [D], [OI1], [OI2], [Pi], [R]). A difficult step in the proof of the case $g \geq 2$ is to show that the set of maps $f \in \text{Diff}_{\omega,*}^r(S)$ having at least $2g - 1$ periodic points is dense in $\text{Diff}_{\omega,*}^r(S)$.

1.3. Idea of the proof. The main tool of the proof is the forcing theory developed in [LecT1, LecT2, Lel], which we introduce in Paragraphs 3.1 and 4.1. Using this tool, we analyse the possible configurations that can occur under the hypotheses of Theorem A. In most of the cases, we will find a rotational horseshoe (defined in Paragraph 2.5), which will allow us to get the conclusion of the theorem. In only one case we will not be able to find such a horseshoe and indeed, there are some examples of homeomorphisms satisfying the hypotheses of Theorem A and without topological horseshoe, for example time one maps of area preserving flows. The conclusion will be obtained using an improved version of Poincaré-Birkhoff Theorem 2.1 in a suitable annulus. Caratheodory's theory of prime ends (see Paragraph 2.4) will be used in this last case.

More precisely, one can find a suitable identity isotopy I of f and a singular oriented foliation \mathcal{F} on S whose regular set coincide with the set $\text{dom}(I)$ of points with non trivial trajectory under the isotopy, that satisfy the following fundamental property: every non trivial trajectory $I(z)$ is homotopic in $\text{dom}(I)$ to a path transverse to \mathcal{F} . Given an f -invariant ergodic probability measure ν such that $\nu(\text{dom}(I)) = 1$, the proof starts by building an *approximation* of a typical orbit for ν (Lemma 5.1): it is an oriented loop Γ_* transverse to \mathcal{F} , such that $[\Gamma_*]$ is close to $\text{rot}_f(\nu)$, and such that, for ν -almost every point z , the transverse path defined naturally by the whole orbit of z draws this loop. We will consider an annular covering space $\widehat{\text{dom}}(I)$ of $\text{dom}(I)$ where Γ_* is lifted to a non contractible simple loop $\hat{\Gamma}_*$. The isotopy $I|_{\text{dom}(I)}$ and the foliation \mathcal{F} can be lifted to $\widehat{\text{dom}}(I)$. The union of leaves that meet $\hat{\Gamma}_*$ is an open annulus \tilde{B} . Depending of the properties of the trajectories of typical points for the measure ν with respect to this annulus \tilde{B} , we get different conclusions: if they cross or visit this annulus (see Paragraph 3.3 for definitions), then we are able to find a topological rotational horseshoe, by means of the forcing theory results

proved in Paragraph 4.2; if they stay forever in this annulus then we prove that Poincaré-Birkhoff Theorem 2.1 applies and implies the existence of an infinite number of periodic orbits.

We strongly use, or develop, the results proved by Gabriel Lellouch in his PhD thesis [Lel]. In particular we will need the main result of [Lel], where \wedge denotes the natural intersection form on $H_1(S, \mathbb{R})$ (see Paragraph 4.1): if μ and μ' are two invariant probability measures such that $\text{rot}_f(\mu) \wedge \text{rot}_f(\mu') \neq 0$, then f has a rotational horseshoe. The hypothesis $\text{rot}_f(\lambda) \in \mathbb{R}H_1(S, \mathbb{Z})$ will be used once: with the help of Atkinson's theorem [At], it will permit us to assume that $[\Gamma_*] \wedge \text{rot}_f(\lambda) = 0$.

1.4. Acknowledgements. We would like to thank Sobhan Seyfaddini for suggesting us this problem. While ending this article we received the recent note of Rohil Prasad. We would like to thank him for his useful comments.

2. DEFINITIONS, NOTATIONS AND PRELIMINARIES

In the sequel, the letter S will refer to a closed surface while the letter Σ will refer to any surface (not necessarily compact, not necessarily connected). If f is a surface homeomorphism, μ will refer to any f -invariant measure, λ to an f -invariant measure with total support, and ν to an f -invariant ergodic measure.

2.1. Loops and paths. Let Σ be an oriented surface (not necessarily closed, not necessarily boundaryless, not necessarily connected). A *loop* of Σ is a continuous map $\Gamma : \mathbb{T} \rightarrow \Sigma$, where $\mathbb{T} = \mathbb{R}/\mathbb{Z}$. It will be called *essential* if it is not homotopic to a constant loop. A *path* of Σ is a continuous map $\gamma : I \rightarrow \Sigma$ where $I \subset \mathbb{R}$ is an interval. A loop or a path will be called *simple* if it is injective. The *natural lift* of a loop $\Gamma : \mathbb{T} \rightarrow \Sigma$ is the path $\gamma : \mathbb{R} \rightarrow \Sigma$ such that $\gamma(t) = \Gamma(t + \mathbb{Z})$. A *segment* is a simple path $\sigma : [a, b] \rightarrow \Sigma$, where $a < b$. The points $\sigma(a)$ and $\sigma(b)$ are the *endpoints* of σ . We will say that σ *joins* $\sigma(a)$ to $\sigma(b)$. More generally if A and B are disjoint, we will say that σ *joins* A to B , if $\sigma(a) \in A$ and $\sigma(b) \in B$. A *line* is a proper simple path $\lambda : \mathbb{R} \rightarrow \Sigma$. As it is usually done we will use the same name and the same notation to refer to the image of a loop or a path γ .

Note that a simple loop or a simple path is naturally oriented. Let Γ be a simple loop of Σ , and denote Σ' the connected component of Σ it belongs to. If $\Sigma' \setminus \Gamma$ has two connected components, we say that Γ *separates* Σ ; in this case the connected component that is located on the right of Γ will be denoted $R(\Gamma)$ and the other one $L(\Gamma)$. We will use the same notations $R(\lambda)$, $L(\lambda)$ for a line λ that separates the connected component it belongs to.

Let f be an orientation preserving homeomorphism of Σ . A *Brouwer line* of f is a line λ that separates Σ such that $f(\lambda) \subset L(\lambda)$ and $f^{-1}(\lambda) \subset R(\lambda)$. Equivalently it means that $f(\overline{L(\lambda)}) \subset L(\lambda)$ or that $f^{-1}(\overline{R(\lambda)}) \subset R(\lambda)$.

2.2. Poincaré-Birkhoff theorem. Let us consider the annulus $\mathbb{A} = \mathbb{T} \times I$, where $(0, 1) \subset I \subset [0, 1]$, and its universal covering space $\tilde{\mathbb{A}} = \mathbb{R} \times I$. We define the covering projection $\tilde{\pi} : (x, y) \mapsto (x + \mathbb{Z}, y)$ and the generating covering automorphism $T : (x, y) \mapsto (x + 1, y)$. We denote $\tilde{p}_1 : \tilde{\mathbb{A}} \rightarrow \mathbb{R}$ the projection on the first factor.

Let f be a homeomorphism of \mathbb{A} isotopic to the identity (meaning orientation preserving and fixing the boundary circles or ends) and \tilde{f} a lift of f to $\tilde{\mathbb{A}}$. The map $p_1 \circ \tilde{f} - p_1$ lifts a continuous function $\psi_{\tilde{f}} : \mathbb{A} \rightarrow \mathbb{R}$ because \tilde{f} and T commute. In particular, for every $z \in \mathbb{A}$, for every lift $\tilde{z} \in \tilde{\mathbb{A}}$ of z and every $n \geq 1$, we have

$$\sum_{i=0}^{n-1} \psi_{\tilde{f}}(f^i(z)) = p_1(\tilde{f}^n(\tilde{z})) - p_1(\tilde{z}).$$

Let z be a positively recurrent point. Say that f has $\text{rot}_{\tilde{f}}(z) \in \mathbb{R}$ as a rotation number if for every subsequence $(f^{n_k}(z))_{k \geq 0}$ of $(f^n(z))_{n \geq 0}$ that converges to z , we have

$$\lim_{k \rightarrow +\infty} \frac{1}{n_k} \sum_{i=0}^{n_k-1} \psi_{\tilde{f}}(f^i(z)) = \text{rot}_{\tilde{f}}(z).$$

If O is a periodic point of f of period q , then there exists $p \in \mathbb{Z}$ such that for every $\tilde{z} \in \tilde{\pi}^{-1}(O)$ we have $\tilde{f}^q(\tilde{z}) = T^p(\tilde{z})$. In this case, p/q is the rotation number of O for the lift \tilde{f} . We will use the following extension of the classical Poincaré-Birkhoff Theorem (see for example [Lec1]):

Theorem 2.1. *Let f be a homeomorphism of \mathbb{A} isotopic to the identity and \tilde{f} a lift of f to $\tilde{\mathbb{A}}$. We suppose that there exist two positively recurrent points z_1 and z_2 , such that $\text{rot}_{\tilde{f}}(z_1) < \text{rot}_{\tilde{f}}(z_2)$. Then:*

- *either, for every rational number $p/q \in (\text{rot}_{\tilde{f}}(z_1), \text{rot}_{\tilde{f}}(z_2))$, written in an irreducible way, there exists a periodic orbit O of f of period q and rotation number p/q for \tilde{f} ;*
- *or there exists an essential simple loop $\Gamma \subset \mathbb{T} \times (0, 1)$ such that $f(\Gamma) \cap \Gamma = \emptyset$.*

Of course, we have a similar result in an abstract annulus, meaning a topological space homeomorphic to \mathbb{A} .

2.3. Homeomorphisms of hyperbolic surfaces. Let Σ be a connected oriented hyperbolic surface without boundary, meaning different from the sphere, the plane, the open annulus or the torus. One can furnish Σ with a complete Riemannian metric of constant negative curvature -1 . The universal covering space of Σ is the disk $\mathbb{D} = \{z \in \mathbb{C} \mid |z| < 1\}$ and the group of covering transformations, denoted \mathcal{G} , is composed of Möbius automorphisms of \mathbb{D} . One can suppose that the metric is of first type, meaning that the closure in \mathbb{C} of every \mathcal{G} -orbit contains $\mathbb{S}_1 = \{z \in \mathbb{C} \mid |z| = 1\}$ (see [Mats2] for instance). Every hyperbolic element $T \in \mathcal{G}$ can be extended to a homeomorphism of $\overline{\mathbb{D}}$ having two fixed points on the boundary: a repelling fixed point $\alpha(T)$ and an attracting fixed point $\omega(T)$. For every $z \in \overline{\mathbb{D}} \setminus \{\alpha(T), \omega(T)\}$, it holds that

$$\lim_{k \rightarrow -\infty} T^k z = \alpha(T), \quad \lim_{k \rightarrow +\infty} T^k z = \omega(T).$$

The metric being of first type, the set of points $\alpha(T)$ and the set of points $\omega(T)$, T among all hyperbolic automorphism, is dense in \mathbb{S}_1 . Every parabolic

element $T \in \mathcal{G}$ can be extended to a homeomorphism of $\overline{\mathbb{D}}$ having one fixed point $\alpha\omega(T)$ on the boundary. For every $z \in \overline{\mathbb{D}} \setminus \{\alpha\omega(T)\}$, it holds that

$$\lim_{k \rightarrow \pm\infty} T^k z = \alpha\omega(T).$$

A homeomorphism f of Σ isotopic to the identity has a unique lift \tilde{f} to \mathbb{D} that commutes with the covering automorphisms. We will call it the *canonical lift* of f . It is well known that \tilde{f} extends to a homeomorphism $\overline{\tilde{f}}$ of $\overline{\mathbb{D}}$ that fixes every point of \mathbb{S}_1 . If $T \in \mathcal{G}$ is hyperbolic, then \tilde{f} lifts a homeomorphism \hat{f} of $\hat{\Sigma} = \tilde{\Sigma}/T$. Moreover \hat{f} extends to a homeomorphism of the compact annulus $\overline{\hat{\Sigma}}$ obtained by adding the two circles $\hat{J} = \tilde{J}/T$ and $\hat{J}' = \tilde{J}'/T$, where \tilde{J} and \tilde{J}' are the two connected components of $\mathbb{S}_1 \setminus \{\alpha(T), \omega(T)\}$. Note that every point of $\hat{J} \cup \hat{J}'$ is fixed, with a rotation number equal to zero for the lift $\overline{\tilde{f}}|_{\overline{\mathbb{D}} \setminus \{\alpha(T), \omega(T)\}}$. Similarly, if $T \in \mathcal{G}$ is parabolic, then \tilde{f} lifts a homeomorphism \hat{f} of $\hat{\Sigma} = \tilde{\Sigma}/T$ that extends to a homeomorphism of $\overline{\hat{\Sigma}}$ obtained by adding the circle $(\mathbb{S}_1 \setminus \{\alpha\omega(T)\})/T$ at one end of $\hat{\Sigma}$. Every point of this circle is fixed, with a rotation number equal to zero for the lift $\overline{\tilde{f}}|_{\overline{\mathbb{D}} \setminus \{\alpha\omega(T)\}}$.

2.4. Caratheodory theory of prime ends. In this small subsection we state a result that will be used once in the article, consequence of what is called *prime end theory* (see [Math] for instance). Let S be a closed surface of genus ≥ 1 and U an open annulus of S . Say that an end e of U is *singular* if there exists a point $z \in S$ and a neighborhood of e in U that is a punctured neighborhood of z in S . Otherwise say that e is *regular*. There is at least one regular end because S is not the 2-sphere. Suppose that U is invariant by an orientation preserving homeomorphism f . Then the homeomorphism $f|_U$ extends to a homeomorphism \overline{f}_U of a larger annulus \overline{U}_{pe} obtained by blowing up each regular end of U and replace it with the associated circle of prime ends. Moreover if U is a connected component of the complement of a closed subset X of $\text{fix}(f)$, then the extended map fixes each point of the circles of prime ends. More precisely, suppose that $I = (f_t)_{t \in [0,1]}$ is an identity isotopy of f , such that $f_t(U) = U$ and $X \subset \text{fix}(f_t)$ for every $t \in [0,1]$. Then, the rotation number of the points on the added circles (they are fixed) is equal to 0, for the lift of \overline{f}_U to the universal covering space of \overline{U}_{pe} , that extends the lift of $f|_U$ to the universal covering space of U , naturally defined by $I|_U$.

2.5. Rotational topological horseshoes. Let Σ be a connected oriented surface. Say that $Y \subset S$ is a *topological horseshoe* of $f \in \text{Homeo}_*(S)$ if Y is closed, invariant by a power f^r of f , and if $f^r|_Y$ admits a finite extension $g : Z \rightarrow Z$ on a Hausdorff compact space Z such that:

- g is an extension of the Bernoulli shift $\sigma : \{1, \dots, m\}^{\mathbb{Z}} \rightarrow \{1, \dots, m\}^{\mathbb{Z}}$, where $m \geq 2$;
- the preimage of every s -periodic sequence of $\{1, \dots, m\}^{\mathbb{Z}}$ by the factor map contains at least one s -periodic point of g .

It means that g is a homeomorphism of Z that is semi-conjugated to $f^r|_Y$ and that the fibers of the factor map are all finite with an uniform bound

M in their cardinality. Note that, if $h(f)$ denotes the topological entropy of f , then it holds that

$$rh(f) = h(f^r) \geq h(f^r|_Y) = h(g) \geq h(\sigma) = \log q,$$

and that f^r has at least q^n/M fixed points for every $n \geq 1$.

Suppose now that S is a connected closed oriented surface. Say that a topological horseshoe Y of $f \in \text{Homeo}_*(S)$ is a *rotational topological horseshoe of type* (κ, r) , where $\kappa \in \mathcal{FHL}(S)$ and r is a positive integer, if there exists a positive integer s such that for every $p/q \in [0, 1] \cap \mathbb{Q}$, there exists a point $z \in Y$ of period at least q/s , such that the loop naturally defined by $I^{r^q}(z)$ belongs to κ^p . In particular the horseshoe defines a homotopical interval of rotation. The rotational topological horseshoes that appear in the present article will be constructed in an annular covering of an invariant open set, satisfying the geometric definition given in [PaPotSa].

3. FOLIATIONS ON SURFACES

In this section we will consider an oriented boundaryless surface Σ , not necessarily closed, not necessarily connected, and a non singular oriented topological foliation \mathcal{F} on Σ . We will consider:

- the universal covering space $\tilde{\Sigma}$ of Σ ;
- the covering projection $\tilde{\pi} : \tilde{\Sigma} \rightarrow \Sigma$;
- the group \mathcal{G} of covering automorphisms;
- the lifted foliation $\tilde{\mathcal{F}}$ on $\tilde{\Sigma}$.

For every point $z \in \Sigma$, we denote ϕ_z the leaf of \mathcal{F} that contains z . If $\phi_z : \mathbb{R} \rightarrow \Sigma$ is a parametrization of ϕ_z inducing the orientation, such that $\phi_z(0) = z$, we set $\phi_z^+ = \phi_z|_{[0, +\infty)}$ and $\phi_z^- = \phi_z|_{(-\infty, 0]}$. Similarly, for every point $\tilde{z} \in \tilde{\Sigma}$, we denote $\tilde{\phi}_{\tilde{z}}$ the leaf of $\tilde{\mathcal{F}}$ that contains \tilde{z} and we define in the same way $\tilde{\phi}_{\tilde{z}}^+$ and $\tilde{\phi}_{\tilde{z}}^-$.

3.1. \mathcal{F} -transverse intersections. A path $\gamma : J \rightarrow \Sigma$ is *positively transverse*¹ to \mathcal{F} if it locally crosses each leaf of \mathcal{F} from the right to the left. Observe that every lift $\tilde{\gamma} : J \rightarrow \tilde{\Sigma}$ of γ is positively transverse to $\tilde{\mathcal{F}}$ and that for every $a < b$ in J :


- $\tilde{\gamma}|_{[a, b]}$ meets once every leaf $\tilde{\phi}$ such that $R(\tilde{\phi}_{\tilde{\gamma}(a)}) \subset R(\tilde{\phi}) \subset R(\tilde{\phi}_{\tilde{\gamma}(b)})$;
- $\tilde{\gamma}|_{[a, b]}$ does not meet any other leaf.

Two transverse paths $\tilde{\gamma}_1 : J_1 \rightarrow \tilde{\Sigma}$ and $\tilde{\gamma}_2 : J_2 \rightarrow \tilde{\Sigma}$ are said *equivalent* if they meet the same leaves of $\tilde{\mathcal{F}}$. Two transverse paths $\gamma_1 : J_1 \rightarrow \Sigma$ and $\gamma_2 : J_2 \rightarrow \Sigma$ are *equivalent* if there exists a lift $\tilde{\gamma}_1 : J_1 \rightarrow \tilde{\Sigma}$ of γ_1 and a lift $\tilde{\gamma}_2 : J_2 \rightarrow \tilde{\Sigma}$ of γ_2 that are equivalent.

Let $\tilde{\gamma}_1 : J_1 \rightarrow \tilde{\Sigma}$ and $\tilde{\gamma}_2 : J_2 \rightarrow \tilde{\Sigma}$ be two transverse paths such that there exist $t_1 \in J_1$ and $t_2 \in J_2$ satisfying $\tilde{\gamma}_1(t_1) = \tilde{\gamma}_2(t_2)$. We will say that $\tilde{\gamma}_1$ and $\tilde{\gamma}_2$ have a *$\tilde{\mathcal{F}}$ -transverse intersection* at $\tilde{\gamma}_1(t_1) = \tilde{\gamma}_2(t_2)$ if there exist $a_1, b_1 \in J_1$ satisfying $a_1 < t_1 < b_1$ and $a_2, b_2 \in J_2$ satisfying $a_2 < t_2 < b_2$ such that:

- $\tilde{\phi}_{\tilde{\gamma}_1(a_1)} \subset L(\tilde{\phi}_{\tilde{\gamma}_2(a_2)})$, $\tilde{\phi}_{\tilde{\gamma}_2(a_2)} \subset L(\tilde{\phi}_{\tilde{\gamma}_1(a_1)})$;
- $\tilde{\phi}_{\tilde{\gamma}_1(b_1)} \subset R(\tilde{\phi}_{\tilde{\gamma}_2(b_2)})$, $\tilde{\phi}_{\tilde{\gamma}_2(b_2)} \subset R(\tilde{\phi}_{\tilde{\gamma}_1(b_1)})$;

¹In the whole text, “transverse” will mean “positively transverse”.

FIGURE 1. Example of $\tilde{\mathcal{F}}$ -transverse intersection.

- every path joining $\tilde{\phi}_{\tilde{\gamma}_1(a_1)}$ to $\tilde{\phi}_{\tilde{\gamma}_1(b_1)}$ and every path joining $\tilde{\phi}_{\tilde{\gamma}_2(a_2)}$ to $\tilde{\phi}_{\tilde{\gamma}_2(b_2)}$ must intersect.

It means that there is a “crossing” between the two paths naturally defined by $\tilde{\gamma}_1$ and $\tilde{\gamma}_2$ in the space of leaves of $\tilde{\mathcal{F}}$, which is a one-dimensional topological manifold, usually non Hausdorff (see Figure 1).

Now, let $\gamma_1 : J_1 \rightarrow \Sigma$ and $\gamma_2 : J_2 \rightarrow \Sigma$ be two transverse paths such that there exist $t_1 \in J_1$ and $t_2 \in J_2$ satisfying $\gamma_1(t_1) = \gamma_2(t_2)$. Say that γ_1 and γ_2 have a \mathcal{F} -transverse intersection at $\gamma_1(t_1) = \gamma_2(t_2)$ if $\tilde{\gamma}_1$ and $\tilde{\gamma}_2$ have a $\tilde{\mathcal{F}}$ -transverse intersection at $\tilde{\gamma}_1(t_1) = \tilde{\gamma}_2(t_2)$, where $\tilde{\gamma}_1 : J_1 \rightarrow \tilde{\Sigma}$ and $\tilde{\gamma}_2 : J_2 \rightarrow \tilde{\Sigma}$ are lifts of γ_1 and γ_2 such that $\tilde{\gamma}_1(t_1) = \tilde{\gamma}_2(t_2)$. If $\gamma_1 = \gamma_2$ one speaks of a \mathcal{F} -transverse self-intersection. This means that if $\tilde{\gamma}_1$ is a lift of γ_1 , there exists $T \in \mathcal{G}$ such that $\tilde{\gamma}_1$ and $T\tilde{\gamma}_1$ have a $\tilde{\mathcal{F}}$ -transverse intersection at $\tilde{\gamma}_1(t_1) = T\tilde{\gamma}_1(t_2)$.

3.2. Recurrence, equivalence and accumulation. A transverse path $\gamma : \mathbb{R} \rightarrow \Sigma$ is *positively recurrent* if, for every $a < b$, there exist $c < d$, with $b < c$, such that $\gamma|_{[a,b]}$ and $\gamma|_{[c,d]}$ are equivalent. Similarly γ is *negatively recurrent* if, for every $a < b$, there exist $c < d$, with $d < a$, such that $\gamma|_{[a,b]}$ and $\gamma|_{[c,d]}$ are equivalent. Finally γ is *recurrent* if it is both positively and negatively recurrent.

Two transverse paths $\gamma_1 : \mathbb{R} \rightarrow \Sigma$ and $\gamma_2 : \mathbb{R} \rightarrow \Sigma$ are *equivalent at $+\infty$* if there exists a_1 and a_2 in \mathbb{R} such that $\gamma_1|_{[a_1,+\infty)}$ and $\gamma_2|_{[a_2,+\infty)}$ are equivalent. Similarly γ_1 and γ_2 are *equivalent at $-\infty$* if there exists b_1 and b_2 in \mathbb{R} such that $\gamma_1|_{(-\infty,b_1]}$ and $\gamma_2|_{(-\infty,b_2]}$ are equivalent.

A transverse path $\gamma_1 : \mathbb{R} \rightarrow \Sigma$ *accumulates positively* on the transverse path $\gamma_2 : \mathbb{R} \rightarrow \Sigma$ if there exist real numbers a_1 and $a_2 < b_2$ such that $\gamma_1|_{[a_1,+\infty)}$ and $\gamma_2|_{[a_2,b_2)}$ are equivalent. Similarly, γ_1 *accumulates negatively* on γ_2 if there exist real numbers b_1 and $a_2 < b_2$ such that $\gamma_1|_{(-\infty,b_1]}$ and $\gamma_2|_{(a_2,b_2]}$ are equivalent. Finally γ_1 *accumulates* on γ_2 if it accumulates positively or negatively on γ_2 .

3.3. Strips. We fix $T \in \mathcal{G} \setminus \{0\}$ and consider

- the annulus $\hat{\Sigma} = \tilde{\Sigma}/T$;
- the covering projections $\pi : \tilde{\Sigma} \rightarrow \hat{\Sigma}$ and $\hat{\pi} : \hat{\Sigma} \rightarrow \Sigma$;
- the foliation $\hat{\mathcal{F}}$ on $\hat{\Sigma}$ induced by $\tilde{\mathcal{F}}$.

Suppose that $\hat{\Gamma}_*$ is a simple loop transverse to $\hat{\mathcal{F}}$. Then, $\hat{\Gamma}_*$ is essential and $\tilde{\gamma}_* = \pi^{-1}(\hat{\Gamma}_*)$ is an oriented line of $\tilde{\Sigma}$, invariant by T and transverse to $\tilde{\mathcal{F}}$. The set

$$\hat{B} = \{\hat{z} \in \hat{\Sigma} \mid \hat{\phi}_{\hat{z}} \cap \hat{\Gamma}_* \neq \emptyset\}$$

is an open annulus which is $\hat{\mathcal{F}}$ -saturated, meaning that it is a union of leaves. Similarly

$$\tilde{B} = \pi^{-1}(\hat{B}) = \{\tilde{z} \in \tilde{\Sigma} \mid \tilde{\phi}_{\tilde{z}} \cap \tilde{\gamma}_* \neq \emptyset\}$$

is an $\tilde{\mathcal{F}}$ -saturated plane invariant by T . We will call such a set a *strip* or a *T-strip* if we want to be more precise. The frontier of \tilde{B} , denoted $\partial\tilde{B}$, is a union of leaves (possibly empty) and can be written $\partial\tilde{B} = \partial\tilde{B}^R \sqcup \partial\tilde{B}^L$, where

$$\partial\tilde{B}^R = \partial\tilde{B} \cap R(\tilde{\gamma}_*), \quad \partial\tilde{B}^L = \partial\tilde{B} \cap L(\tilde{\gamma}_*).$$

Let us state some facts that can be proven easily (see [LecT2] or [Lel]). Note first that:

- if there is a leaf $\tilde{\phi} \subset \partial\tilde{B}$ that is invariant by T , then the set $\partial\tilde{B}^R$ or $\partial\tilde{B}^L$ that contains $\tilde{\phi}$ is reduced to this leaf;
- if $\tilde{\gamma} : \mathbb{R} \rightarrow \tilde{\Sigma}$ is transverse to $\tilde{\mathcal{F}}$, then the set of real numbers t such that $\gamma(t) \in \tilde{B}$ is an interval (possibly empty).

Suppose now that $\tilde{\gamma} : \mathbb{R} \rightarrow \tilde{\Sigma}$ is transverse to $\tilde{\mathcal{F}}$ and that

$$\{t \in \mathbb{R} \mid \gamma(t) \in \tilde{B}\} = (a, b),$$

where $-\infty \leq a < b \leq \infty$. Say that

- $\tilde{\gamma}$ *draws* \tilde{B} if there exist $t < t'$ in (a, b) such that $\tilde{\phi}_{\tilde{\gamma}(t')} = T\tilde{\phi}_{\tilde{\gamma}(t)}$.

If, moreover, we suppose that $-\infty < a < b < +\infty$, say that:

- $\tilde{\gamma}$ *crosses* \tilde{B} *from the right to the left* if $\tilde{\gamma}(a) \in \partial\tilde{B}^R$ and $\tilde{\gamma}(b) \in \partial\tilde{B}^L$;
- $\tilde{\gamma}$ *crosses* \tilde{B} *from the left to the right* if $\tilde{\gamma}(a) \in \partial\tilde{B}^L$ and $\tilde{\gamma}(b) \in \partial\tilde{B}^R$;
- $\tilde{\gamma}$ *visits* \tilde{B} *on the right* if $\tilde{\gamma}(a) \in \partial\tilde{B}^R$ and $\tilde{\gamma}(b) \in \partial\tilde{B}^R$;
- $\tilde{\gamma}$ *visits* \tilde{B} *on the left* if $\tilde{\gamma}(a) \in \partial\tilde{B}^L$ and $\tilde{\gamma}(b) \in \partial\tilde{B}^L$.

We will say that $\tilde{\gamma}$ *crosses* \tilde{B} if it crosses it from the right to the left or from the left to the right. Similarly, we will say that $\tilde{\gamma}$ *visits* \tilde{B} if it visits it on the right or on the left. Note that $T(\tilde{\gamma})$ satisfies the same properties as $\tilde{\gamma}$. Note also that if $\tilde{\gamma}$ visits \tilde{B} on the right, then $\partial\tilde{B}^R$ is not reduced to a T -invariant leaf. An analogous property holds if $\tilde{\gamma}$ visits \tilde{B} on the left. Finally, observe that at least one of the following situations occurs (the two last assertions are not incompatible):

- $\tilde{\gamma}$ crosses \tilde{B} ;
- $\tilde{\gamma}$ visits \tilde{B} ;
- $\tilde{\gamma}$ is equivalent to $\tilde{\gamma}_*$ at $+\infty$ or at $-\infty$;
- $\tilde{\gamma}$ accumulates on $\tilde{\gamma}_*$ positively or negatively.

Let us conclude this list of properties by the following ones (see [Lel, Section 2.1.2.c]):




FIGURE 2. An example where Proposition 3.2 holds.

Proposition 3.1. *We have the following results:*

- If $\tilde{\gamma}$ visits and draws \tilde{B} , then $\tilde{\gamma}$ and $T(\tilde{\gamma})$ have an $\tilde{\mathcal{F}}$ -transverse intersection and so $\gamma = \tilde{\pi} \circ \tilde{\gamma}$ has an \mathcal{F} -transverse self intersection.
- If $\tilde{\gamma}_1$ crosses \tilde{B} from the right to the left, if $\tilde{\gamma}_2$ crosses \tilde{B} from the right to the left and at least one of the paths $\tilde{\gamma}_1$ or $\tilde{\gamma}_2$ draws \tilde{B} , then there exists $k \in \mathbb{Z}$ such that $\tilde{\gamma}_1$ and $T^k(\tilde{\gamma}_2)$ have an $\tilde{\mathcal{F}}$ -transverse intersection, and so $\gamma_1 = \tilde{\pi} \circ \tilde{\gamma}_1$ and $\gamma_2 = \tilde{\pi} \circ \tilde{\gamma}_2$ have a \mathcal{F} -transverse intersection.

3.4. More about the accumulation property. In this final paragraph, we will suppose moreover than Σ is connected and that $\Sigma \neq \mathbb{R}^2/\mathbb{Z}^2$. The goal is to prove the following result that has its own interest and will be used in the sequel to prove Theorem A. This statement is stronger than some results of [Lel, Section 2.1.1].

Proposition 3.2. *Suppose that $\gamma_1 : \mathbb{R} \rightarrow \Sigma$ is a positively recurrent transverse path that accumulates positively on a transverse path $\gamma_2 : \mathbb{R} \rightarrow \Sigma$. Then, there exists a transverse simple loop $\Gamma_* \subset \Sigma$ with the following properties.*

- (1) *The set B of leaves met by Γ_* is an open annulus of Σ .*
- (2) *The path γ_1 stays in B and is equivalent to the natural lift of Γ_* .*
- (3) *If $\tilde{\gamma}_1, \tilde{\gamma}_2$ are lifts of γ_1, γ_2 to the universal covering space $\tilde{\Sigma}$ such that $\tilde{\gamma}_1|_{[a_1, +\infty)}$ is equivalent to $\tilde{\gamma}_2|_{[a_2, b_2)}$ and if \tilde{B} is the lift of B that contains $\tilde{\gamma}_1$, then one of the inclusions $\tilde{\phi}_{\tilde{\gamma}_2(b_2)} \subset \partial \tilde{B}^R, \tilde{\phi}_{\tilde{\gamma}_1(a_1)} \subset \partial \tilde{B}^L$ holds. In the first case, we have $\tilde{B} \subset L(\tilde{\phi})$ for every $\tilde{\phi} \subset \partial \tilde{B}^R$ and in the second case, we have $\tilde{B} \subset R(\tilde{\phi})$ for every $\tilde{\phi} \subset \partial \tilde{B}^L$.*

An example of a situation where Proposition 3.2 holds is depicted in Figure 2.

In Proposition 4.17 we will get additional properties when the paths are supposed to be trajectories that are typical for some ergodic f -invariant measures.

Proof. Let us start with a lemma.




FIGURE 3. Local configuration of the path Γ and the foliation \mathcal{F} (in red) around the point $0 = \Psi_z(z)$.

Lemma 3.3. *Let $\Gamma : \mathbb{T} \rightarrow \Sigma$ be a transverse loop, $\tilde{\gamma} : \mathbb{R} \rightarrow \widetilde{\text{dom}}(I)$ a lift of Γ and \tilde{B} the strip that contains $\tilde{\gamma}$. Let $T \in \mathcal{G}$ be the deck transformation associated to \tilde{B} . Suppose that there exists a deck transformation $R \in \mathcal{G}$ and $a \in \mathbb{R}$ such that $\tilde{\gamma}|_{[a, a+1]}$ is equivalent to a subpath of $R\tilde{\gamma}$. Then $\tilde{\gamma}|_{[a, a+1]} \cap R\tilde{\gamma} \neq \emptyset$.*

Note that if moreover Γ is a simple path, then the conclusion of the lemma implies that $R \in \langle T \rangle$.

This lemma can be reduced easily to the following fact.

Sub-lemma 3.4. *Let \mathcal{F} be a singular foliation on Σ , and $\Gamma : \mathbb{T} \rightarrow \Sigma$ a loop of Σ that is transverse to \mathcal{F} . Then, there exists $z \in \Gamma$ such that ϕ_z^+ does not meet Γ but at the end point.*

Proof of Lemma 3.3. By Sub-lemma 3.4, there exist z, z' in Γ (possibly equal) such that ϕ_z^+ and $\phi_{z'}^-$ do not meet Γ but at their end point. Denote \tilde{z}, \tilde{z}' the respective lifts of z, z' that belong to $\tilde{\gamma}|_{[a, a+1]}$. We know that $\tilde{\phi}_{\tilde{z}}^+ \cap R\tilde{\gamma} = \emptyset$, and that $\tilde{\phi}_{\tilde{z}'}^- \cap R\tilde{\gamma} \neq \emptyset$. We deduce that $R\tilde{\gamma} \cap \tilde{\gamma}|_{[a, a+1]} \neq \emptyset$. \square

Proof of Sub-lemma 3.4. Fix $z \in \Gamma$. The loop Γ being transverse to \mathcal{F} , there are finitely many parameters $t \in \mathbb{T}$ such that $z = \Gamma(t)$. Consequently, there exists a compact neighborhood W_z of z , a homeomorphism $\Phi_z : W_z \rightarrow [-1, 1]^2$ and a finite set I_z such that:

- Φ_z sends z onto $(0, 0)$;
- Φ_z sends $\mathcal{F}|_{W_z}$ onto the vertical foliation oriented upward;
- we have $\Phi_z(\Gamma \cap W_z) = \bigcup_{i \in I_z} \text{gr}(\psi_{i,z})$, where $\psi_{i,z} : [-1, 1] \rightarrow [-1, 1]$ is a continuous function satisfying $\psi_{i,z}(0) = 0$.

Here the notation $\text{gr}(\psi)$ denotes the graph of $\psi : [-1, 1] \rightarrow [-1, 1]$ oriented from the right to the left. See Figure 3 for an example of such a configuration.

Consider the two continuous functions

$$\psi_z^- = \min_{i \in I_z} \psi_{i,z}, \quad \psi_z^+ = \max_{i \in I_z} \psi_{i,z}$$

and define

$$\gamma_z^- = \Phi_z^{-1}(\text{gr}(\psi_z^-)), \quad \gamma_z^+ = \Phi_z^{-1}(\text{gr}(\psi_z^+)).$$

We will argue by contradiction by supposing that for any $z \in \Gamma$, the path ϕ_z^+ meets Γ in a point that is not the end point. In that case, for every $z \in \Gamma$, there exists a sub-path $\delta_z : [0, 1] \rightarrow \Sigma$ of ϕ_z^+ such that

$$\delta_z(0) = z, \quad \delta_z(1) \in \Gamma, \quad \delta_z((0, 1)) \cap \Gamma = \emptyset.$$

In particular we can define a first return map $\theta : \Gamma \rightarrow \Gamma$ by setting $\theta(z) = \delta_z(1)$. We will prove that $X = \bigcup_{z \in \Gamma} \delta_z([0, 1])$ is a compact sub-surface with boundary. Note that for every $z \in \Gamma$, the function θ induces a homeomorphism from a compact neighborhood α_z of z in γ_z^+ to a compact neighborhood ω_z of $\theta(z)$ in $\gamma_{\theta(z)}^-$ and consequently that every point $\delta_z(t)$, $t \in (0, 1)$, belongs to the interior of X . Note also that for every $z \in \Gamma$, the set $\Phi_z^-(\{(x, y) \mid y \geq \psi_z^-(x)\})$ is included in X . By compactness, one can cover Γ with finitely many α_z , $z \in \Gamma$. We deduce that the image of θ , denoted $\text{im}(\theta)$, is the union of finitely many compact subsets (the corresponding ω_z) and therefore is compact. We deduce also that X is compact because for every $z \in \Gamma$, the set $\bigcup_{z' \in \alpha_z} \delta_{z'}([0, 1])$ is compact. Now, observe that for every $z \in \Gamma$ and every $z' \in \gamma_z^-$, the sets $\gamma_{z'}^-$ and γ_z^- coincide in a neighborhood of z' . It implies that $\text{im}(\theta) \cap \gamma_z^-$ is an open subset of γ_z^- . By connectedness of γ_z^- , either γ_z^- is contained in $\text{im}(\theta)$ or it is disjoint from $\text{im}(\theta)$. In the first case, W_z is contained in X , in the second case $W_z \cap X = \Phi_z^{-1}(\{(x, y) \mid y \geq \psi_z^-(x)\})$: we have proved that X is a compact sub-surface of Σ (possibly with boundary). Note that for every $z \in \partial X$ it holds that $\phi_z^+ \setminus \{z\} \subset \text{int}(X)$ (in other terms the foliation is pointing inward on the boundary).

By hypothesis, Σ is connected and different from $\mathbb{R}^2/\mathbb{Z}^2$. So, it does not bear a non-singular foliation. We deduce that X is a surface with boundary. More precisely it is homeomorphic to the closed annulus because it bears a non singular foliation. Let $\Psi : X \rightarrow \mathbb{S}^2$ be a topological embedding compatible with the usual orientations. The loop $\Psi(\Gamma)$ is homologous to 0 in \mathbb{S}^2 and one can define a dual function $\delta : \mathbb{S}^2 \setminus \Psi(\Gamma) \rightarrow \mathbb{Z}$. Such a function is defined by the following property: for every z, z' in $\mathbb{S}^2 \setminus \Psi(\Gamma)$ and every path β joining z to z' , the algebraic intersection number $\Psi(\Gamma) \wedge \beta$ is equal to $\delta(z') - \delta(z)$. Let U be a connected component of $\mathbb{S}^2 \setminus \Psi(\Gamma)$ where δ reaches its maximum. The set $\Psi(\Gamma)$ being connected, the closure of U is a topological disk. Moreover the fact that δ reaches its maximum in U implies that for every $z \in \partial U$ it holds that $\phi_z^+ \setminus \{z\} \subset U$. So U is not a connected component of $\mathbb{S}^2 \setminus \Psi(X)$ and it holds that $\bar{U} \subset \psi(X)$. Summarizing, we have found a closed topological disk bearing a non-singular foliation pointing inward on the boundary. We have got a contradiction. \square

Let us explain how to construct the simple loop Γ_* that appears in Proposition 3.2. As γ_1 is positively recurrent, there exist two numbers $c_1 < c'_1$, with $c_1 > a_1$, such that $\phi_{\gamma_1(c_1)} = \phi_{\gamma_1(c'_1)}$ (see Figure 4 for these different points). It implies that $\gamma_1|_{[c_1, c'_1]}$ is equivalent to a transverse path $\gamma_* : [c_1, c'_1] \rightarrow \Sigma$ such that $\gamma_*(c_1) = \gamma_*(c'_1)$. The set

$$X = \{(t, t') \in [c_1, c'_1]^2 \mid t < t' \text{ and } \gamma_*(t) = \gamma_*(t')\}$$




FIGURE 4. The different objects appearing in the proof of Proposition 3.2, Lemma 3.5 and Claim 3.6. The leaves are in orange.

is non empty (because it contains (c_1, c'_1)) and compact. Indeed, it is closed in $\{(t, t') \in [c_1, c'_1]^2 \mid t < t'\}$, and its closure in the compact set $\{(t, t') \in [c_1, c'_1]^2 \mid t \leq t'\}$ does not contain any couple (t, t) . The function $(t, t') \mapsto t' - t$ being continuous and positive on X , reaches its minimum at a couple (c''_1, c'''_1) . So, replacing (c_1, c'_1) with (c''_1, c'''_1) if necessary, one can always suppose that the loop Γ_* naturally defined by γ_* is simple. We denote B the union of leaves met by Γ_* .

By hypothesis there exist two lifts $\tilde{\gamma}_1$ and $\tilde{\gamma}_2$ of respectively γ_1 and γ_2 to $\tilde{\Sigma}$ such that $\tilde{\gamma}_1|_{[a_1, +\infty)}$ and $\tilde{\gamma}_2|_{[a_2, b_2)}$ are equivalent. We denote \tilde{B} the strip that lifts B and contains $\tilde{\gamma}_1|_{[c_1, c'_1]}$. We denote $\tilde{\gamma}_*$ a lift of Γ_* that lies inside \tilde{B} and $T \in \mathcal{G}$ the primitive deck transformation associated to \tilde{B} (chosen accordingly to the orientation of $\tilde{\gamma}_*$).

Lemma 3.5. *The path $\tilde{\gamma}_1|_{[c_1, +\infty)}$ is included in \tilde{B} .*

Proof. We will argue by contradiction and suppose it is not. Then there exists $d_1 > c'_1$, uniquely defined, such that $\tilde{\gamma}_1(d_1) \notin \tilde{B}$ and $\tilde{\gamma}_1|_{[c_1, d_1]} \subset \tilde{B}$.

Claim 3.6. *There exists a deck transformation $R \in \mathcal{G}$ and real numbers $e_1 < e'_1$, with $e_1 \geq a_1$, such that either $R\tilde{\gamma}_1|_{[e_1, e'_1]}$ draws and crosses \tilde{B} , or it draws and visits \tilde{B} .*

Proof. Note that to prove this claim one has to find $R \in \mathcal{G}$ and $e_1 < e'_1$ such that $R\tilde{\gamma}_1|_{[e_1, e'_1]}$ draws \tilde{B} and both $R\tilde{\gamma}_1(e_1)$ and $R\tilde{\gamma}_1(e'_1)$ do not belong to \tilde{B} .

As γ_1 is positively recurrent, there exist real numbers $e''_1 < e'_1$, with $e''_1 > d_1$, and a deck transformation $R \in \mathcal{G}$ such that $R\tilde{\gamma}_1|_{[e''_1, e'_1]}$ is equivalent to $\tilde{\gamma}_1|_{[c_1, d_1]}$; in particular:

- $\tilde{\gamma}_1|_{[c_1, c'_1]}$ is equivalent to a subpath of $R\tilde{\gamma}_1|_{[e''_1, e'_1]}$;
- $R\tilde{\gamma}_1([e''_1, e'_1]) \subset \tilde{B}$ and $R\tilde{\gamma}_1(e'_1) \notin \tilde{B}$.

To prove the claim, it is sufficient to show that $R\tilde{\gamma}_1([a_1, e'_1]) \not\subset \tilde{B}$, because in that case there exists $e_1 \in [a_1, e''_1]$ such that $R\tilde{\gamma}_1([e_1, e'_1]) \subset \tilde{B}$ and $R\tilde{\gamma}_1(e_1) \notin \tilde{B}$.

We argue by contradiction. Suppose that $R\tilde{\gamma}_1([a_1, e'_1])$ is contained in \tilde{B} . Then $\tilde{\gamma}_1([a_1, e'_1])$ is contained in $R^{-1}(\tilde{B})$. Recall that there exists t such that $\tilde{\gamma}_*|_{[t, t+1]}$ is equivalent to $\tilde{\gamma}_1|_{[c_1, c'_1]}$ which is a subpath of $\tilde{\gamma}_1|_{[a_1, e'_1]}$. It implies that $\tilde{\gamma}_*|_{[t, t+1]}$ is equivalent to a subpath of $R^{-1}\tilde{\gamma}_*$ because $\tilde{\gamma}_1([a_1, e'_1])$ is contained in $R^{-1}(\tilde{B})$. Lemma 3.3 applies and ensures that $R^{-1} \in \langle T \rangle$. As \tilde{B} is invariant by T , the condition $R\tilde{\gamma}_1([a_1, e'_1]) \subset \tilde{B}$ gives $\tilde{\gamma}_1([a_1, e'_1]) \subset \tilde{B}$. This contradicts the condition $\tilde{\gamma}_1(d_1) \notin \tilde{B}$, because $a_1 < d_1 < e'_1$. \square

As γ_1 is positively recurrent, there exist sequences $(e_{1,n})_{n \geq 0}$ and $(e'_{1,n})_{n \geq 0}$ with $a_1 < e_{1,n} < e'_{1,n} < e_{1,n+1}$, and a sequence $(R_n)_{n \geq 0}$ of deck transformations, such that $R_n\tilde{\gamma}_1|_{[e_{1,n}, e'_{1,n}]}$ is equivalent to $R\tilde{\gamma}_1|_{[e_1, e'_1]}$. As $\tilde{\gamma}_1$ accumulates on $\tilde{\gamma}_2$, a similar statement holds for $\tilde{\gamma}_2$: there exist sequences $(e_{2,n})_{n \geq 0}$ and $(e'_{2,n})_{n \geq 0}$ with $a_2 < e_{2,n} < e'_{2,n} < e_{2,n+1} < b_2$ such that $R_n\tilde{\gamma}_2|_{[e_{2,n}, e'_{2,n}]}$ is equivalent to $R\tilde{\gamma}_2|_{[a_2, b_2]}$. Note that the R_n are all different because every leaf of $\tilde{\mathcal{F}}$ intersects $\tilde{\gamma}_2([a_2, b_2])$ at most once.

We have two possibilities given by Claim 3.6: either $R\tilde{\gamma}_1|_{[e_1, e'_1]}$ draws and crosses \tilde{B} , or it draws and visits \tilde{B} .

Suppose that $R\tilde{\gamma}_1|_{[e_1, e'_1]}$ draws and crosses \tilde{B} . In this case, for any $n \in \mathbb{N}$, the path $R_n\tilde{\gamma}_2|_{[e_{2,n}, e'_{2,n}]}$ intersects $\tilde{\gamma}_*$. Replacing R_n with $T^{k_N} \circ R_n$ for a certain $k_N \in \mathbb{Z}$ if necessary, one can suppose that $R_n\tilde{\gamma}_2|_{[e_{2,n}, e'_{2,n}]}$ intersects $\tilde{\gamma}_*|_{[t, t+1]}$ and so $R_n^{-1}(\tilde{\gamma}_*|_{[t, t+1]})$ intersects $\tilde{\gamma}_2([a_2, b_2])$. It contradicts the fact that the action of \mathcal{G} on compact subsets is proper.

Suppose now that $R\tilde{\gamma}_1|_{[e_1, e'_1]}$ draws and visits \tilde{B} . Then $R\tilde{\gamma}_1|_{[e_1, e'_1]}$ and $TR\tilde{\gamma}_1|_{[e_1, e'_1]}$ have an $\tilde{\mathcal{F}}$ -transverse intersection. One deduces that for any $n \in \mathbb{N}$, one has $R_n\tilde{\gamma}_2|_{[e_{2,n}, e'_{2,n}]}$ and $TR\tilde{\gamma}_1|_{[e_1, e'_1]}$ have an $\tilde{\mathcal{F}}$ -transverse intersection because $R_n\tilde{\gamma}_2|_{[e_{2,n}, e'_{2,n}]}$ and $R\tilde{\gamma}_1|_{[e_1, e'_1]}$ are equivalent. Consequently, it holds that $R_n\tilde{\gamma}_2|_{[e_{2,n}, e'_{2,n}]} \cap TR\tilde{\gamma}_1|_{[e_1, e'_1]} \neq \emptyset$ and so that $R_n\tilde{\gamma}_2|_{[a_2, b_2]} \cap TR\tilde{\gamma}_1|_{[e_1, e'_1]} \neq \emptyset$. It contradicts once again the fact that the action of \mathcal{G} on compact subsets is proper. This finishes the proof of Lemma 3.5. \square

By Lemma 3.5, we know that $\tilde{\gamma}_1|_{[c_1, +\infty)}$ stays in \tilde{B} . We first prove that $\tilde{\gamma}_1$ cannot accumulate in $\tilde{\gamma}_*$.

Indeed, otherwise, as γ_1 is positively recurrent, there exist deck transformations $(R_n)_{n \geq 0} \in \mathcal{G}$ and parameters $d_n < d'_n$ both going to $+\infty$ such that $\tilde{\gamma}_1|_{[d_n, d'_n]}$ is equivalent to $R_n\tilde{\gamma}_1|_{[c_1, c'_1]}$, which is itself equivalent to $R_n\tilde{\gamma}_*|_{[t, t+1]}$. The fact that $\tilde{\gamma}_1$ accumulates in $\tilde{\gamma}_*$ implies that $R_n \notin \langle T \rangle$ eventually. Recall that for any n , the path $\tilde{\gamma}_1|_{[d_n, d'_n]}$ is equivalent to a subpath of $\tilde{\gamma}_*$; this allows to apply Lemma 3.3 to the simple path Γ_* , which implies that $R_n \in \langle T \rangle$, a contradiction.

Hence, there exists $t_1 \in \mathbb{R}$ such that $\tilde{\gamma}_1|_{[c_1, +\infty)}$ is equivalent to $\tilde{\gamma}_*|_{[t_1, +\infty)}$. Moreover it is equivalent to $\tilde{\gamma}_2|_{[c_2, b_2]}$, where $c_2 \in [a_2, b_2]$. It implies that $\tilde{\phi}_{\tilde{\gamma}_2(b_2)} \subset \partial\tilde{B}$. We do not lose generality by supposing that $\tilde{\phi}_{\tilde{\gamma}_2(b_2)} \subset \partial\tilde{B}^L$. We choose $a'_2 \in [c_2, b_2]$ such that $\tilde{\gamma}_2([a'_2, b_2]) \in L(\tilde{\gamma}_*)$.

Lemma 3.7. *For every leaf $\tilde{\phi} \subset \partial\tilde{B}^L$ it holds that $\tilde{B} \subset R(\tilde{\phi})$.*




FIGURE 5. The configuration of the proof of Lemma 3.7.

Proof. See Figure 5 for an example of configuration of the proof. Suppose that there exists a leaf $\tilde{\phi}_0 \subset \partial \tilde{B}^L$ such that $\tilde{B} \subset L(\tilde{\phi}_0)$. One can find a transverse path $\tilde{\gamma}_3 : [a_3, b_3] \rightarrow \tilde{\Sigma}$ such that $\tilde{\gamma}_3(a_3) \in \tilde{\phi}_0$ and $\tilde{\gamma}_3((a_3, b_3]) \subset \tilde{B}$. Such a path enters in \tilde{B} by the left. By taking a smaller b_3 if necessary, we can suppose moreover than $\tilde{\gamma}_3([a_3, b_3]) \subset L(\tilde{\gamma}_*)$. We will prove that it prevents $\tilde{\gamma}_1$ accumulating positively in $\tilde{\gamma}_2$. If $\tilde{\lambda}$ is an oriented line of \tilde{B} , denote $R_{\tilde{B}}(\tilde{\lambda})$ the connected component of $\tilde{B} \setminus \tilde{\lambda}$ located on the right of $\tilde{\lambda}$ and $L_{\tilde{B}}(\tilde{\lambda})$ the connected component of $\tilde{B} \setminus \tilde{\lambda}$ located on the left of $\tilde{\lambda}$. One defines two oriented lines $\tilde{\lambda}_2, \tilde{\lambda}_3$ of \tilde{B} by setting

$$\tilde{\lambda}_2 = (\tilde{\gamma}_2|_{[a'_2, b_2]})^{-1} \tilde{\phi}_{\tilde{\gamma}_2(a'_2)}^+, \quad \tilde{\lambda}_3 = \tilde{\gamma}_3|_{(a_3, b_3]} \tilde{\phi}_{\tilde{\gamma}_3(b_3)}^+.$$

The line $\tilde{\gamma}_*$ intersects $\tilde{\phi}_{\tilde{\gamma}_2(a'_2)}^+$ in a unique point \tilde{z}_2 and we have $\tilde{z}_2 \in \tilde{\phi}_{\tilde{\gamma}_2(a'_2)}^+$. Similarly, $\tilde{\gamma}_*$ intersects $\tilde{\phi}_{\tilde{\gamma}_3(b_3)}^+$ in a unique point \tilde{z}_3 and we have $\tilde{z}_3 \in \tilde{\phi}_{\tilde{\gamma}_3(b_3)}^+$. Denote $\tilde{\sigma}_2 \subset \tilde{\phi}_{\tilde{\gamma}_2(a'_2)}$ the segment that joins $\tilde{\gamma}_2(a'_2)$ to \tilde{z}_2 and $\tilde{\sigma}_3 \subset \tilde{\phi}_{\tilde{\gamma}_3(b_3)}$ the segment that joins $\tilde{\gamma}_3(b_3)$ to \tilde{z}_3 . By compactness of all segments, if n is large enough, then we have

$$T^n (\tilde{\gamma}_3([a_3, b_3]) \cup \tilde{\sigma}_3) \cap (\tilde{\gamma}_2([a'_2, b_2]) \cup \tilde{\sigma}_2) = \emptyset.$$

Moreover, one can suppose that

$$T^n \tilde{\phi}_{\tilde{\gamma}_3(b_3)}^+ \subset L(\tilde{\phi}_{\tilde{\gamma}_2(a'_2)}^+).$$

The fact that $\tilde{\gamma}_2([a'_2, b_2])$ and $\tilde{\gamma}_3([a_3, b_3])$ are included in $L(\tilde{\gamma}_*)$ while $\tilde{\phi}_{\tilde{\gamma}_2(a'_2)}^+$ and $\tilde{\phi}_{\tilde{\gamma}_3(b_3)}^+$ are included in $\overline{R(\tilde{\gamma}_*)}$ tells us that

$$T^n \tilde{\phi}_{\tilde{\gamma}_3(b_3)}^+ \cap (\tilde{\gamma}_2([a'_2, b_2]) \cup \tilde{\sigma}_2) = \emptyset.$$

We deduce that the lines $\tilde{\lambda}_2$ and $T^n \tilde{\lambda}_3$ are disjoint.

The sub-path of $\tilde{\gamma}_*$ that joins \tilde{z}_2 to $T^n \tilde{z}_3$ is disjoint from $\tilde{\lambda}_2$ and $T^n \tilde{\lambda}_3$ but at the endpoints, entering in $L_{\tilde{B}}(\tilde{\lambda}_2)$ at \tilde{z}_2 and leaving $R_{\tilde{B}}(T^n \tilde{\lambda}_3)$ at $T^n \tilde{z}_3$. Consequently the following inclusion $\overline{L_{\tilde{B}}(T^n \tilde{\lambda}_3)} \subset L_{\tilde{B}}(\tilde{\lambda}_2)$ holds. Every leaf $\tilde{\phi} \subset L_{\tilde{B}}(\tilde{\phi}_{T^n \tilde{z}_3})$ is disjoint from $T^n \tilde{\lambda}_3$. It is contained in $L(T^n \tilde{\lambda}_3)$ because the sub-path of $\tilde{\gamma}_*$ that joins $\tilde{\phi}_{T^n(\tilde{z}_3)}$ to $\tilde{\phi}$ is disjoint from $T^n \tilde{\lambda}_3$ but at $T^n \tilde{z}_3$ and enters in $L_{\tilde{B}}(T^n \tilde{\lambda}_3)$ at $T^n \tilde{z}_3$. The contradiction comes from the fact that $\tilde{\phi}$ must intersect $\tilde{\gamma}_2|_{[a'_2, b_2]}$ because $\tilde{\phi} \subset L_{\tilde{B}}(\tilde{\phi}_{z_2})$. \square

Lemma 3.8. *The set B is an open annulus of Σ .*

Proof. Suppose it is not. Then there exists a deck transformation $R \notin \langle T \rangle$ of $\tilde{\Sigma}$ such that $R\tilde{B} \cap \tilde{B} \neq \emptyset$. As \tilde{B} is the set of leaves met by $\tilde{\gamma}_*$, it implies the existence of $t \in \mathbb{R}$ such that $R\tilde{\gamma}_*(t) \in \tilde{B}$. The line $\tilde{\gamma}_*$ lifts the simple loop Γ_* and so we have $R\tilde{\gamma}_* \cap \tilde{\gamma}_* = \emptyset$. Moreover, there is at least one leaf of $\tilde{\mathcal{F}}$ that is met both by $\tilde{\gamma}_*$ and $R\tilde{\gamma}_*$. Consequently, one of the following inclusions $\overline{L(R\tilde{\gamma}_*)} \subset L(\tilde{\gamma}_*)$, $\overline{L(\tilde{\gamma}_*)} \subset L(R\tilde{\gamma}_*)$ holds. Replacing R by R^{-1} if necessary, one can suppose that the first inclusion holds, which implies that $R\tilde{\gamma}_* \subset L(\tilde{\gamma}_*)$.

Note that $R\tilde{\gamma}_*$ cannot accumulate on $\tilde{\gamma}_*$ (neither positively nor negatively) because the natural lift γ_* of Γ_* is recurrent and so, by Lemma 3.3, cannot accumulate on itself. Moreover it cannot be equivalent to $\tilde{\gamma}_*$ neither at $+\infty$ nor at $-\infty$ (by using Lemma 3.3). It cannot cross \tilde{B} because $R\tilde{\gamma}_* \cap \tilde{\gamma}_* = \emptyset$. It remains to prove that it cannot visit \tilde{B} .

Using the fact that $R\tilde{\gamma}_* \subset L(\tilde{\gamma}_*)$, the line $R\tilde{\gamma}_*$ must visit \tilde{B} by the left if it visits \tilde{B} . This contradicts Lemma 3.7: no transverse trajectory enters in \tilde{B} by the left side. \square

To prove Proposition 3.2, it remains to prove that γ_1 is entirely contained in B (which will imply that $\tilde{\gamma}_1$ is entirely contained in \tilde{B}). But this is implied by the facts that $\gamma_1|_{[a_1, +\infty)}$ is contained in B and that γ_1 is recurrent. This finishes the proof of Proposition 3.2. \square

The following results (and others related to the accumulation property) were already stated by Lellouch in [Lel, Section 2.1.1]. Using the precise description given here, we get them as a trivial corollary.

Corollary 3.9. *Suppose that $\gamma_1 : \mathbb{R} \rightarrow \Sigma$ is a positively recurrent transverse path that accumulates positively on a transverse path $\gamma_2 : \mathbb{R} \rightarrow \Sigma$. Then there is no positively or negatively recurrent transverse path $\gamma_0 : \mathbb{R} \rightarrow \Sigma$ that accumulates positively or negatively on γ_1 . In particular a positively recurrent transverse path does not accumulate on itself. Also, the accumulated leaf $\phi_{\gamma_2(b_2)}$ is not met by γ_1 .*

Proof. To prove the first point, it suffices to note that by Proposition 3.2, the function $t \mapsto \phi_{\gamma_1(t)}$ is locally injective. The last point comes from the fact that γ_1 is contained in B while $\phi_{\gamma_2(b_2)}$ is contained in the frontier of B . \square

4. FORCING THEORY

4.1. Maximal isotopies and transverse foliations. Let Σ be an oriented boundaryless surface, not necessarily closed, not necessarily connected and

f a homeomorphism isotopic to the identity. Recall that if $I = (f_t)_{t \in [0,1]}$ is an identity isotopy of f , the trajectory $I(z)$ of a point $z \in \Sigma$ is the path $t \mapsto f_t(z)$ defined on $[0, 1]$. We can define the *whole trajectory* of z as being the path

$$I^{\mathbb{Z}}(z) = \prod_{k \in \mathbb{Z}} I(f^k(z))$$

constructed by concatenation. More precisely, on every interval $[k, k + 1]$, $k \in \mathbb{Z}$, it is defined by the formula:

$$I^{\mathbb{Z}}(z) : t \mapsto f_{t-k}(f^k(z)).$$

We define the *fixed point set* and the *domain* of I as follows:

$$\text{fix}(I) = \bigcap_{t \in [0,1]} \text{fix}(f_t), \quad \text{dom}(I) = \Sigma \setminus \text{fix}(I).$$

Denote \mathcal{I} the set of identity isotopies of f . We have a preorder on \mathcal{I} defined as follows: say that $I \preceq I'$ if

- $\text{fix}(I) \subset \text{fix}(I')$;
- I' is homotopic to I relative to $\text{fix}(I)$.

Let us state two important results. The first one is due to Béguin-Crovisier-Le Roux [BeCLer] (see also [J] for a weaker version). The second can be found in [Lee1].

Theorem 4.1. *For every $I \in \mathcal{I}$, there exists $I' \in \mathcal{I}$ such that $I \preceq I'$ and such that I' is maximal for the preorder.*

Remark. An isotopy I is maximal if and only if, for every $z \in \text{fix}(f) \setminus \text{fix}(I)$, the loop $I(z)$ is not contractible in $\text{dom}(I)$. Equivalently, if we lift the isotopy $I|_{\text{dom}(I)}$ to an identity isotopy $\tilde{I} = (\tilde{f}_t)_{t \in [0,1]}$ on the universal covering space $\widetilde{\text{dom}(I)}$ of $\text{dom}(I)$, the maximality of I means that \tilde{f}_1 is fixed point free. Note that every connected component of $\widetilde{\text{dom}(I)}$ must be a topological plane.

Theorem 4.2. *If $I \in \mathcal{I}$ is maximal, then there exists a topological oriented singular foliation \mathcal{F} on M such that*

- *the singular set $\text{sing}(\mathcal{F})$ coincides with $\text{fix}(I)$;*
- *for every $z \in \text{dom}(I)$, the trajectory $I(z)$ is homotopic in $\text{dom}(I)$, relative to the ends, to a transverse path γ joining z to $f(z)$.*

We will say that \mathcal{F} is *transverse to I* . It can be lifted to a non singular foliation $\tilde{\mathcal{F}}$ on $\widetilde{\text{dom}(I)}$ which is transverse to \tilde{I} . This last property is equivalent to saying that every leaf $\tilde{\phi}$ of $\tilde{\mathcal{F}}$ is a Brouwer line of the lift \tilde{f} induced by I , as defined in Section 2.1. The path γ is uniquely defined up to equivalence: if γ_1 and γ_2 are two such paths and if $z \in \widetilde{\text{dom}(I)}$ lifts $z \in \text{dom}(I)$, then the respective lifts $\tilde{\gamma}_1, \tilde{\gamma}_2$ of γ_1, γ_2 starting at \tilde{z} join this point to $\tilde{f}(\tilde{z})$ and consequently meet the same leaves of $\tilde{\mathcal{F}}$. We will write $\gamma = I_{\mathcal{F}}(z)$ and call this path the *transverse trajectory of z* . It is defined, up to equivalence, on $[0, 1]$. For every $n \geq 1$, we will define by concatenation the path

$$I_{\mathcal{F}}^n(z) = I_{\mathcal{F}}(z)I_{\mathcal{F}}(f(z)) \cdots I_{\mathcal{F}}(f^{n-1}(z)).$$

We can also define the *whole transverse trajectory* of z as being the path

$$I_{\mathcal{F}}^{\mathbb{Z}}(z) = \prod_{k \in \mathbb{Z}} I_{\mathcal{F}}(f^k(z))$$

coinciding on $[k, k+1]$, $k \in \mathbb{Z}$, with $I_{\mathcal{F}}(f^k(z))$ after translation by $-k$. Similarly, we define

$$\tilde{I}_{\tilde{\mathcal{F}}}^n(\tilde{z}) = \tilde{I}_{\tilde{\mathcal{F}}}(\tilde{z})\tilde{I}_{\tilde{\mathcal{F}}}(\tilde{f}(\tilde{z})) \cdots \tilde{I}_{\tilde{\mathcal{F}}}(\tilde{f}^{n-1}(\tilde{z}))$$

and

$$\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z}) = \prod_{k \in \mathbb{Z}} \tilde{I}_{\tilde{\mathcal{F}}}(\tilde{f}^k(\tilde{z})).$$

Recall that a *flow-box* of $\tilde{\mathcal{F}}$ is an open disk \tilde{U} of $\widetilde{\text{dom}}(I)$ such that the foliation $\tilde{\mathcal{F}}|_{\tilde{U}}$ is homeomorphic to the foliation of \mathbb{R}^2 by verticals. The following results, easy to prove (see [LecT1]), will be useful in the article.

Proposition 4.3. *For every $\tilde{z} \in \widetilde{\text{dom}}(I)$ and every pair of integers $k_1 < k_2$ there exists a neighborhood \tilde{U} of \tilde{z} such that $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})|_{[k_1, k_2]}$ is a subpath (up to equivalence) of $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z}')|_{[k_1-1, k_2+1]}$.*

Proposition 4.4. *For every $\tilde{z} \in \widetilde{\text{dom}}(I)$ and every neighborhood \tilde{V} of \tilde{z} , there exists a flow-box $\tilde{U} \subset \tilde{V}$ containing \tilde{z} , such that for every $\tilde{z}' \in \tilde{U}$, the path $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z}')$ intersects every leaf that meets \tilde{U} .*

Remind that if f is a homeomorphism of Σ , a point z is *positively recurrent* if $z \in \omega(z)$ and *negatively recurrent* if $z \in \alpha(z)$. In the case where $z \in \alpha(z) \cap \omega(z)$, we say that z is *recurrent*. For instance, if μ is an invariant finite Borel measure on S , then μ -almost every point is recurrent. The following result is an immediate consequence of Proposition 4.3.

Proposition 4.5. *If $z \in \text{dom}(I)$ is positively recurrent, then $I_{\mathcal{F}}^{\mathbb{Z}}(z)$ is positively recurrent. If z is negatively recurrent, then $I_{\mathcal{F}}^{\mathbb{Z}}(z)$ is negatively recurrent.*

Let us state now the key lemma of [LecT1] (Proposition 20) that is the elementary brick of the forcing theory and which will be used later.

Lemma 4.6. *Suppose that there exist \tilde{z}_1, \tilde{z}_2 in $\widetilde{\text{dom}}(I)$ and positive integers n_1, n_2 such that $\tilde{I}_{\tilde{\mathcal{F}}}^{n_1}(\tilde{z}_1)$ and $\tilde{I}_{\tilde{\mathcal{F}}}^{n_2}(\tilde{z}_2)$ have an $\tilde{\mathcal{F}}$ -transverse intersection at $\tilde{I}_{\tilde{\mathcal{F}}}^{n_1}(\tilde{z}_1)(t_1) = \tilde{I}_{\tilde{\mathcal{F}}}^{n_2}(\tilde{z}_2)(t_2)$. Then there exists $\tilde{z}_3 \in \widetilde{\text{dom}}(I)$ such that $\tilde{I}_{\tilde{\mathcal{F}}}^{n_1+n_2}(\tilde{z}_3)$ is equivalent to $\tilde{I}_{\tilde{\mathcal{F}}}^{n_1}(\tilde{z}_1)|_{[0, t_1]}\tilde{I}_{\tilde{\mathcal{F}}}^{n_2}(\tilde{z}_2)|_{[t_2, n_2]}$.*

Let us give now the principal result of [LecT2]. Here, \mathcal{G} is the group of covering automorphisms of $\widetilde{\text{dom}}(I)$ and $[T]_{\mathcal{FHL}} \in \mathcal{FHL}(S)$ is the free homotopy class (in S) of a loop $\Gamma \subset \text{dom}(I)$ naturally defined by T (see Paragraph 2.5).

Theorem 4.7. *Suppose that there exists $\tilde{z} \in \widetilde{\text{dom}}(I)$, $T \in \mathcal{G} \setminus \{\text{Id}\}$ and $r \geq 1$ such that $\tilde{I}_{\tilde{\mathcal{F}}}^r(\tilde{z})$ and $T\tilde{I}_{\tilde{\mathcal{F}}}^r(\tilde{z})$ have an $\tilde{\mathcal{F}}$ -transverse intersection at $\tilde{I}_{\tilde{\mathcal{F}}}^r(\tilde{z})(a) = T(\tilde{I}_{\tilde{\mathcal{F}}}^r(\tilde{z}))(a')$ where $a' < a$. Then f admits a rotational horseshoe of type $([T]_{\mathcal{FHL}}, r)$.*

Proof. What is proved in [LecT2] is the following, where $\widehat{\text{dom}}(I) = \widetilde{\text{dom}}(I)/T$ and \hat{f} is the homeomorphism of $\widehat{\text{dom}}(I)$ induced by \tilde{f} .

There exists an \hat{f}^r -invariant compact set \hat{Y} such that

- \hat{f}^r is an extension of the Bernoulli shift $\sigma : \{1, 2\}^{\mathbb{Z}} \rightarrow \{1, 2\}^{\mathbb{Z}}$;
- the preimage of every q -periodic sequence of $\{1, 2\}^{\mathbb{Z}}$ by the factor map contains at least one q -periodic point of \hat{f}^r ;
- for every $p/q \in [0, 1] \cap \mathbb{Q}$ written in an irreducible way, there exists $\hat{z}_{p/q} \in \hat{Y}$ such that $\hat{f}^{rq}(\hat{z}_{p/q}) = T^p(\hat{z}_{p/q})$ if $\tilde{z}_{p/q} \in \widetilde{\text{dom}}(I)$ lifts $\hat{z}_{p/q}$.

The image Y of \hat{Y} by the covering projection $\hat{\pi} : \widehat{\text{dom}}(I) \rightarrow \text{dom}(I)$ is invariant by f^r . It is a topological horseshoe because $\hat{\pi}|_{\hat{Y}}$ is a semi-conjugacy from $\hat{f}^r|_{\hat{Y}}$ to $f^r|_Y$ and because every $z \in Y$ has finitely many lifts in \hat{Y} (with an uniform bound s) because \hat{Y} is compact. The loop of S naturally defined by $I^{rq}(z_{p/q})$, where $z_{p/q} = \hat{\pi}(\hat{z}_{p/q})$, belongs to $[T]_{\mathcal{F}\mathcal{H}\mathcal{L}}^p$. Moreover, the \hat{f}^r -orbit of $\hat{z}_{p/q}$ has q points because p and q are relatively prime. It projects onto the f^r -orbit of $z_{p/q}$, which has at least q/s points. So, the period of $z_{p/q}$ (for f) is at least q/s . \square

Remark. In particular, the theorem asserts the existence of a topological horseshoe, and so the positiveness of the topological entropy, in the case where there exists $z \in \text{dom}(I)$ such that $I_{\mathcal{F}}^{\mathbb{Z}}(z)$ has an \mathcal{F} -transverse self-intersection. It was proved in [LecT1] that such a situation occurs in the case where there exist two positively (or negatively) recurrent points z_1, z_2 in $\text{dom}(I)$ such that $I_{\mathcal{F}}^{\mathbb{Z}}(z_1)$ and $I_{\mathcal{F}}^{\mathbb{Z}}(z_2)$ have an \mathcal{F} -transverse intersection. For example this happens if f preserves a Borel probability measure with total support and if there exist two points z_1, z_2 in $\text{dom}(I)$ such that $I_{\mathcal{F}}^{\mathbb{Z}}(z_1)$ and $I_{\mathcal{F}}^{\mathbb{Z}}(z_2)$ have an \mathcal{F} -transverse intersection. Indeed, by Proposition 4.3, it is also the case for $I_{\mathcal{F}}^{\mathbb{Z}}(z'_1)$ and $I_{\mathcal{F}}^{\mathbb{Z}}(z'_2)$ if z'_1, z'_2 are close to z_1, z_2 respectively. But if f preserves a Borel probability measure λ with total support, then λ -almost every point is recurrent and so, the set of recurrent points is dense.

What follows, which is stronger than what is said in the previous remark, is crucial in [Lel] and will also be fundamental in our study.

Corollary 4.8. *Suppose that Σ is a closed surface and that ν_1, ν_2 are ergodic invariant probability measures. If there exists $\tilde{z}_1 \in \text{dom}(I) \cap \text{supp}(\nu_1)$ and $\tilde{z}_2 \in \text{dom}(I) \cap \text{supp}(\nu_2)$ such that $\tilde{I}_{\mathcal{F}}^{\mathbb{Z}}(\tilde{z}_1)$ and $\tilde{I}_{\mathcal{F}}^{\mathbb{Z}}(\tilde{z}_2)$ intersect \mathcal{F} -transversally, then for every neighborhood \mathcal{U} of $\text{rot}_f(\nu_1)$ in $H_1(S, \mathbb{R})$, there exists $T \in \mathcal{G} \setminus \{\text{Id}\}$ and $r \geq 1$ such $[T]/r \in \mathcal{U}$ and such that f admits a rotational horseshoe of type $([T]_{\mathcal{F}\mathcal{H}\mathcal{L}}, r)$.*

Note that this corollary can be applied in the case where $\nu_1 = \nu_2$ and some $\tilde{z} \in \text{dom}(I) \cap \text{supp}(\nu_1)$ is such that $\tilde{I}_{\mathcal{F}}^{\mathbb{Z}}(\tilde{z})$ has an \mathcal{F} -transverse self-intersection.

Proof. Let $j \in \{1, 2\}$. One knows that ν_j -almost every point z'_j satisfies the following properties:

- z'_j is recurrent;
- its orbit is dense in $\text{supp}(\nu_j)$;




FIGURE 6. The configuration of the proof of Corollary 4.8. The orange lines are leaves.

- if $\tilde{z}_j' \in \widetilde{\text{dom}}(I)$ is a lift of z_j' , then there exists a sequence $(T_{j,i})_{i \geq 0}$ in \mathcal{G} and a sequence $(n_{j,i})_{i \geq 0}$ in $\mathbb{N} \setminus \{0\}$ such that

$$\lim_{i \rightarrow +\infty} n_{j,i} = +\infty, \quad \lim_{i \rightarrow +\infty} \frac{[T_{j,i}]}{n_{j,i}} = \text{rot}_f(\nu_j), \quad \lim_{i \rightarrow +\infty} T_{j,i}^{-1} f^{n_{j,i}}(\tilde{z}_j') = \tilde{z}_j'.$$

By Proposition 4.3 and the hypothesis of the corollary we know that $\tilde{I}_{\mathcal{F}}^{\mathbb{Z}}(\tilde{z}_1')$ and $\tilde{I}_{\mathcal{F}}^{\mathbb{Z}}(\tilde{z}_2')$ intersect \mathcal{F} -transversally. So there exists $r' \in \mathbb{N} \setminus \{0\}$, $s_1, s_2 \in \mathbb{Z}$ and two lifts \tilde{z}_1'' and \tilde{z}_2'' of \tilde{z}_1' and \tilde{z}_2' such that $\tilde{I}_{\mathcal{F}}^{r'}(\tilde{f}^{s_1}(\tilde{z}_1''))$ and $\tilde{I}_{\mathcal{F}}^{r'}(\tilde{f}^{s_2}(\tilde{z}_2''))$ intersect \mathcal{F} -transversally. Denote $\tilde{z}_j'' = \tilde{f}^{s_j}(\tilde{z}_j')$. See Figure 6 for a description of the proof configuration.

By Proposition 4.3, if i is large enough then, up to equivalence, $\tilde{I}_{\mathcal{F}}^{r'}(\tilde{z}_j'')$ is a subpath of $T_{j,i}^{-1} \tilde{I}_{\mathcal{F}}^{r'+2}(\tilde{f}^{n_{j,i}-1}(\tilde{z}_j''))$. So $T_{1,i}^{-1} \tilde{I}_{\mathcal{F}}^{r'+2}(\tilde{f}^{n_{1,i}-1}(\tilde{z}_1''))$ and $\tilde{I}_{\mathcal{F}}^{r'}(\tilde{z}_2'')$ have an $\tilde{\mathcal{F}}$ -transverse intersection at $T_{1,i}^{-1} \tilde{I}_{\mathcal{F}}^{r'+2}(\tilde{f}^{n_{1,i}-1}(\tilde{z}_1''))(a) = \tilde{I}_{\mathcal{F}}^{r'}(\tilde{z}_2'')(b)$, as well as $T_{2,i}^{-1} \tilde{I}_{\mathcal{F}}^{r'+2}(\tilde{f}^{n_{2,i}-1}(\tilde{z}_2''))$ and $\tilde{I}_{\mathcal{F}}^{r'}(\tilde{z}_1'')$ have an $\tilde{\mathcal{F}}$ -transverse intersection at $T_{2,i}^{-1} \tilde{I}_{\mathcal{F}}^{r'+2}(\tilde{f}^{n_{2,i}-1}(\tilde{z}_2''))(c) = \tilde{I}_{\mathcal{F}}^{r'}(\tilde{z}_1'')(d)$ (we omit here the dependences on i, i' for brevity of notations).

Lemma 4.6 then implies that for any i, i' , there exists $\tilde{z}_3 \in \widetilde{\text{dom}}(I)$ such that $\tilde{I}_{\mathcal{F}}^{\tilde{2r'+2+n_{1,i}+n_{2,i'}}}(\tilde{z}_3)$ is equivalent to the path

$$\tilde{\gamma}_{i,i'} = T_{1,i}^{-1} \tilde{I}_{\mathcal{F}}^{r'+1+n_{1,i}}(\tilde{z}_1'')|_{[0, n_{1,i}-1+a]} \cdot \tilde{I}_{\mathcal{F}}^{r'+1+n_{2,i'}}(\tilde{z}_2'')|_{[b, r'+1+n_{2,i'}]}.$$

Consider the parameter $e \in [0, 2r' + 3 + n_{1,i} + n_{2,i'}]$ such that

$$\tilde{I}_{\mathcal{F}}^{\mathbb{Z}}(\tilde{z}_3)(e) = T_{1,i}^{-1} \tilde{I}_{\mathcal{F}}^{\mathbb{Z}}(\tilde{z}_1'')(n_{1,i} - 1 + a) = \tilde{I}_{\mathcal{F}}^{\mathbb{Z}}(\tilde{z}_2'')(b).$$

Note that if i, i' are large enough, then $n_{1,i} - 1 + a \geq d$, and $b \leq n_{2,i'} - 1 + c$. It implies that $T_{1,i} \tilde{\gamma}_{i,i'}$ has an $\tilde{\mathcal{F}}$ -transverse intersection with $T_{2,i'}^{-1} \tilde{\gamma}_{i,i'}$ at a point

$T_{1,i}\tilde{\gamma}_{i,i'}(e') = T_{2,i'}^{-1}\tilde{\gamma}_{i,i'}(e'')$, where $e' < e < e''$. So, $\tilde{\gamma}_{i,i'}$ has an $\tilde{\mathcal{F}}$ -transverse intersection with $T_{2,i'}T_{1,i}\tilde{\gamma}_{i,i'}$ at a point $\tilde{\gamma}_{i,i'}(e'') = T_{2,i'}T_{1,i}\tilde{\gamma}_{i,i'}(e')$, where $e' < e''$. By Theorem 4.7, there exists $s \geq 1$ such that f admits a rotational horseshoe of type $([T_{2,i'}T_{1,i}]_{\mathcal{FHL}}, 2r' + 2 + n_{1,i} + n_{2,i'})$. If i is large enough (i' being fixed but large enough to ensure that the above properties hold), then we have $[T_{2,i'}T_{1,i}]_{\mathcal{FHL}}/(2r' + 2 + n_{1,i} + n_{2,i'}) \in \mathcal{U}$. \square

Let us finish this quick introduction to some forcing theory tools by the following theorem of Lellouch's thesis [Lel, Théorème C]:

Theorem 4.9. *Suppose that $g \geq 2$. If $f \in \text{Homeo}_*(S)$ preserves two Borel probability measures μ_1 and μ_2 such that $\text{rot}_f(\mu_1) \wedge \text{rot}_f(\mu_2) \neq 0$, then f has a topological horseshoe. In particular, f has infinitely many periodic points.*

Moreover, if μ_1 is ergodic, then these periodic points can be supposed to have rotation vectors arbitrarily close to $\text{rot}_f(\mu_1)$ and with arbitrarily large period: for every neighbourhood \mathcal{U} of $\text{rot}_f(\mu_1)$ in $H_1(S, \mathbb{R})$, there exists a rotational horseshoe of type (κ, r) with $[\kappa]/r \in \mathcal{U}$.

Here \wedge is the *intersection form*. It is the symplectic form on $H_1(S, \mathbb{R})$ defined by the property that if Γ_1 and Γ_2 are two loops in S , then $[\Gamma_1] \wedge [\Gamma_2]$ is the algebraic intersection number between Γ_1 and Γ_2 . Equivalently, up to a multiplicative constant, it is the form induced *via* Poincaré duality by $\wedge : H^1(S, \mathbb{R}) \times H^1(S, \mathbb{R}) \rightarrow H^2(S, \mathbb{R})$.

4.2. Forcing theory in the annular covering space. We suppose now that Σ is an oriented closed surface and denote it S . We keep the other notations. We consider $T \in \mathcal{G} \setminus \text{Id}$ and a T -strip $\tilde{B} \subset \widetilde{\text{dom}}(I)$ (we suppose that T coincides with the identity on the connected components of $\text{dom}(I)$ that do not contain \tilde{B}). We fix a T -invariant line $\tilde{\gamma}_* \subset \tilde{B}$. We define

- the surface $\widehat{\text{dom}}(I) = \widetilde{\text{dom}}(I)/T$;
- the projections $\pi : \widetilde{\text{dom}}(I) \rightarrow \widehat{\text{dom}}(I)$ and $\hat{\pi} : \widehat{\text{dom}}(I) \rightarrow \text{dom}(I)$;
- the identity isotopy \hat{I} on $\widehat{\text{dom}}(I)$ lifted by \tilde{I} ;
- the lift \hat{f} of $f|_{\text{dom}(I)}$ to $\widehat{\text{dom}}(I)$ lifted by \tilde{f} ;
- the foliation $\hat{\mathcal{F}}$ on $\widehat{\text{dom}}(I)$ lifted by $\tilde{\mathcal{F}}$;
- the loop $\hat{\Gamma}_* = \pi(\tilde{\gamma}_*)$.

The complement of $\hat{\Gamma}_*$ in its connected component has two annular connected components $L(\hat{\Gamma}_*)$ and $R(\hat{\Gamma}_*)$. We denote $\hat{\infty}_L$ the common end of $\widehat{\text{dom}}(I)$ and $L(\hat{\Gamma}_*)$ and $\hat{\infty}_R$ the common end of $\widehat{\text{dom}}(I)$ and $R(\hat{\Gamma}_*)$.

We consider

- the set $\tilde{W}^{R \rightarrow L}$ of points $\tilde{z} \in \widetilde{\text{dom}}(I)$ such that $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})$ crosses \tilde{B} from the right to the left;
- the set $\tilde{W}^{L \rightarrow R}$ of points $\tilde{z} \in \widetilde{\text{dom}}(I)$ such that $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})$ crosses \tilde{B} from the left to the right;
- the set $\tilde{W}^{R \rightarrow R}$ of points $\tilde{z} \in \widetilde{\text{dom}}(I)$ such that $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})$ visits \tilde{B} on the right;
- the set $\tilde{W}^{L \rightarrow L}$ of points $\tilde{z} \in \widetilde{\text{dom}}(I)$ such that $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})$ visits \tilde{B} on the left;
- the set \tilde{W}^D of points $\tilde{z} \in \widetilde{\text{dom}}(I)$ such that $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})$ draws \tilde{B} .

Note that all these sets are invariant by \tilde{f} and by T . Note also that they are open, as a consequence of Proposition 4.3.

We define the respective projections in $\widehat{\text{dom}}(I)$

$$\hat{W}^{R \rightarrow L}, \hat{W}^{L \rightarrow R}, \hat{W}^{R \rightarrow R}, \hat{W}^{L \rightarrow L}, \hat{W}^D,$$

that are open and invariant by \hat{f} and the respective projections in $\text{dom}(I)$

$$W^{R \rightarrow L}, W^{L \rightarrow R}, W^{R \rightarrow R}, W^{L \rightarrow L}, W^D,$$

that are open and invariant by f .

Finally, we define

- the set $\hat{\infty}_R \rightarrow \hat{\infty}_L$ of points $\hat{z} \in \widehat{\text{dom}}(I)$ such that

$$\lim_{k \rightarrow -\infty} \hat{f}^k(\hat{z}) = \hat{\infty}_R, \quad \lim_{k \rightarrow +\infty} \hat{f}^k(\hat{z}) = \hat{\infty}_L;$$

- the set $\hat{\infty}_L \rightarrow \hat{\infty}_R$ of points $\hat{z} \in \widehat{\text{dom}}(I)$ such that

$$\lim_{k \rightarrow -\infty} \hat{f}^k(\hat{z}) = \hat{\infty}_L, \quad \lim_{k \rightarrow +\infty} \hat{f}^k(\hat{z}) = \hat{\infty}_R.$$

We will state some results that have been proven in [Lel] and will add some others that do not explicitly appear there. The following result has been proved in [Lel] (Proposition 2.2.12).

Lemma 4.10. *Suppose that $\nu \in \mathcal{M}(f)$ is ergodic and that ν -almost every point z has a lift $\tilde{z} \in \widehat{\text{dom}}(I)$ such that $\tilde{I}_{\tilde{f}}^{\mathbb{Z}}(\tilde{z})$ is equivalent in $+\infty$ or $-\infty$ to $\tilde{\gamma}_*$. Then there exists $a \geq 0$ ² such that $\text{rot}(\nu) = a[T]$.*

The next one also has been proved in [Lel] (Lemma 2.2.3 and Proposition 2.2.4).

Lemma 4.11. *Suppose that $\nu \in \mathcal{M}(f)$ is ergodic. We have the following:*

- (1) *if $[T] \wedge \text{rot}_f(\nu) > 0$, then $\nu(\hat{\pi}(\hat{\infty}_R \rightarrow \hat{\infty}_L)) = 1$;*
- (2) *if $[T] \wedge \text{rot}_f(\nu) < 0$, then $\nu(\hat{\pi}(\hat{\infty}_L \rightarrow \hat{\infty}_R)) = 1$.*

Let us prove now:

Lemma 4.12. *If there exists $\mu \in \mathcal{M}(f)$ with total support such that $[T] \wedge \text{rot}_f(\mu) = 0$, then every essential simple loop of $\widehat{\text{dom}}(I)$ meets its image by \hat{f} .*

Proof. Suppose that there exists an essential simple loop $\hat{\Gamma}$ such that $\hat{f}(\hat{\Gamma}) \cap \hat{\Gamma} = \emptyset$. Orient $\hat{\Gamma}$ in such a way that $\hat{\infty}_L$ is the common end of $\widehat{\text{dom}}(I)$ and $L(\hat{\Gamma})$ and $\hat{\infty}_R$ the common end of $\widehat{\text{dom}}(I)$ and $R(\hat{\Gamma})$. There is no loss of generality by supposing that $\hat{f}(\hat{\Gamma})$ is included in $L(\hat{\Gamma})$. Consider the line $\tilde{\gamma}$ of \tilde{S} that lifts $\hat{\Gamma}$. We have $\tilde{f}(\overline{L(\tilde{\gamma})}) \subset L(\tilde{\gamma})$ and more generally $\tilde{f}(\overline{L(T'(\tilde{\gamma}))}) \subset L(T'(\tilde{\gamma}))$ for every $T' \in \mathcal{G}$ because \tilde{f} commutes with T' .

If $\tilde{\gamma}'$ is an oriented line of $\widehat{\text{dom}}(I)$, recall that $\widehat{\text{dom}}(I)_{\tilde{\gamma}'}$ is the connected component of $\widehat{\text{dom}}(I)$ that contains $\tilde{\gamma}'$. Denote $\eta_{\tilde{\gamma}'}$ the function defined on $\widehat{\text{dom}}(I)_{\tilde{\gamma}'}$ that is equal to 0 on $R(\tilde{\gamma}')$, to 1 on $L(\tilde{\gamma}')$ and to 1/2 on $\tilde{\gamma}'$. Noting that $T''(\tilde{\gamma}) = T'(\tilde{\gamma})$ if $T''^{-1}T' \in \langle T \rangle$, one deduces that the notation $\tau\tilde{\gamma}$ has

²The proof given in [Lel] says that $a \geq 0$ but we will slightly improve it in Lemma 4.18 to obtain $a > 0$.

a sense for every left coset $\tau \in \mathcal{G}/\langle T \rangle$. Furthermore, if $\nu \in \mathcal{M}(f)$ is ergodic, then for ν -almost every point z , the following holds for every lift \tilde{z} of z :

$$[T] \wedge \text{rot}_f(\nu) = \lim_{n \rightarrow +\infty} \frac{1}{n} \sum_{\tau \in \mathcal{G}/\langle T \rangle} \left(\eta_{\tau\tilde{\gamma}}(\tilde{f}^n(\tilde{z})) - \eta_{\tau\tilde{\gamma}}(\tilde{z}) \right).$$

Indeed, if one considers the loop $\Gamma = \hat{\pi}(\hat{\Gamma})$ of S , then $\sum_{\tau \in \mathcal{G}/\langle T \rangle} \eta_{\tau\tilde{\gamma}}(\tilde{f}^n(\tilde{z})) - \eta_{\tau\tilde{\gamma}}(\tilde{z})$ (note that the sum is finite) is equal to the sum of the algebraic intersection numbers between all lifts of Γ with the trajectory $\tilde{I}^n(\tilde{z})$ (at least when z and $f^n(z)$ are not on Γ), meaning the algebraic intersection number between Γ and $I^n(z)$.

Observe that for every $\tau \in \mathcal{G}/\langle T \rangle$, the function $\eta_{\tau\tilde{\gamma}} \circ \tilde{f} - \eta_{\tau\tilde{\gamma}}$ is non negative on $\widehat{\text{dom}}(I)_{\tilde{\gamma}}$ and positive in the strip between $\tilde{\gamma}$ and $\tilde{f}(\tilde{\gamma})$. We deduce that for every ergodic invariant probability measure ν it holds that $[T] \wedge \text{rot}_f(\nu) \geq 0$. Moreover, we have a strict inequality if the measure of the strip between $\tilde{\gamma}$ and $\tilde{f}(\tilde{\gamma})$ is non zero for the measure $\tilde{\nu}$ that lifts ν . By using the ergodic decomposition of μ , we deduce that $[T] \wedge \text{rot}_f(\mu) > 0$, which contradicts the hypothesis. \square

Lemma 4.13. *Suppose that $\nu \in \mathcal{M}(f)$ and $\nu' \in \mathcal{M}(f)$ are ergodic and satisfy*

$$\nu(W^{R \rightarrow L} \cap W^D) = 1, \quad [T] \wedge \text{rot}_f(\nu') < 0.$$

Then one of the following assertions holds:

- for ν -almost every point z and ν' -almost every point z' , the paths $I_{\mathcal{F}}^{\mathbb{Z}}(z)$ and $I_{\mathcal{F}}^{\mathbb{Z}}(z')$ have an \mathcal{F} -transverse intersection;
- for ν -almost every point z and ν' -almost every point z' , the path $I_{\mathcal{F}}^{\mathbb{Z}}(z')$ accumulates on $I_{\mathcal{F}}^{\mathbb{Z}}(z)$.

Proof. Define three f -invariant sets W_1, W_2, W_3 as follows:

- $z' \in W_1$ if it has a lift \tilde{z}' such that $\tilde{I}_{\mathcal{F}}^{\mathbb{Z}}(\tilde{z}')$ is equivalent to $\tilde{\gamma}_*$ at $+\infty$ or at $-\infty$;
- $z' \in W_2$ if it has a lift \tilde{z}' such that $\tilde{I}_{\mathcal{F}}^{\mathbb{Z}}(\tilde{z}')$ accumulates on $\tilde{\gamma}_*$ positively or negatively;
- $z' \in W_3$ if it has a lift \tilde{z}' such that $\tilde{I}_{\mathcal{F}}^{\mathbb{Z}}(\tilde{z}')$ crosses \tilde{B} from the left to the right.

By Lemma 4.11, we know that ν' -almost every point z' has a lift $\tilde{z}' \in \widehat{\text{dom}}(I)$ that belongs to $\hat{\infty}_L \rightarrow \hat{\infty}_R$. Consequently $\nu'(W_1 \cup W_2 \cup W_3) = 1$, which implies by ergodicity of ν' that one of the sets W_1, W_2, W_3 has ν' -measure 1. By Lemma 4.10, $\nu'(W_1) \neq 1$ because $\text{rot}_f(\nu') \notin \mathbb{R}[T]$ (by the hypothesis $[T] \wedge \text{rot}_f(\nu') < 0$). If $\nu'(W_2) = 1$, then the second item of the lemma holds because for every leaf $\tilde{\phi} \subset \tilde{B}$, ν -almost every point z belongs to W_D and so has a lift $\tilde{z} \in \widehat{\text{dom}}(I)$ such that $\tilde{I}_{\mathcal{F}}^{\mathbb{Z}}(\tilde{z})$ meets $\tilde{\phi}$. By Proposition 3.1, if $\nu'(W_3) = 1$, then the first item of the lemma holds. \square

Corollary 4.14. *Suppose that $\nu \in \mathcal{M}(f)$ is ergodic and satisfies*

$$\nu(W^{R \rightarrow L} \cap W^D) = 1, \quad [T] \wedge \text{rot}_f(\nu) < 0.$$

Then, for ν -almost every point z , the path $I_{\mathcal{F}}^{\mathbb{Z}}(z)$ has an \mathcal{F} -transverse self intersection.

Proof. Let us apply Lemma 4.13 with $\nu' = \nu$ and use the fact that a recurrent transverse path does not accumulate on itself (Corollary 3.9). \square

This result is still true if $\nu(W^{R \rightarrow L} \cap W^D) = 1$ and $[T] \wedge \text{rot}_f(\nu) = 0$. More precisely we have (see [Lel], Proposition 3.3.1).

Lemma 4.15. *Suppose that $\nu \in \mathcal{M}(f)$ is ergodic and satisfies*

$$\nu(W^{R \rightarrow L} \cap W^D) = 1, \quad [T] \wedge \text{rot}_f(\nu) = 0.$$

Then $\nu(W^{L \rightarrow R}) = 1$ and for ν -almost every point z , the path $I_{\mathcal{F}}^{\mathbb{Z}}(z)$ has an \mathcal{F} -transverse self intersection.

Remark. The conclusion $\nu(W^{L \rightarrow R}) = 1$ is not explicitly stated in [Lel], Proposition 3.3.1. But, as explained by the author at the beginning of the proof, it is the key point that permits to get the second conclusion. The first condition says that there are points “that go up”, which implies by the second condition, that there are points “that go down”. We have a situation very similar to the one that occurs under the hypothesis of Corollary 4.14, but more subtle arguments of ergodic theory are needed.

Lemma 4.16. *Suppose that there exist $\lambda \in \mathcal{M}(f)$ such that $\text{supp}(\lambda) = S$. If $\nu \in \mathcal{M}(f)$ is ergodic and satisfies*

$$\nu(W^{R \rightarrow L} \cap W^D) = 1, \quad [T] \wedge \text{rot}_f(\nu) > 0,$$

then there exists $\nu' \in \mathcal{M}(f)$ ergodic, such that one of the following assertions holds:

- *for ν -almost every point z and ν' -almost every point z' , the paths $I_{\mathcal{F}}^{\mathbb{Z}}(z)$ and $I_{\mathcal{F}}^{\mathbb{Z}}(z')$ have an \mathcal{F} -transverse intersection;*
- *for ν -almost every point z and ν' -almost every point z' , the path $I_{\mathcal{F}}^{\mathbb{Z}}(z')$ accumulates on $I_{\mathcal{F}}^{\mathbb{Z}}(z)$.*

Proof. By hypothesis $W^{R \rightarrow L} \cap W^D$ is a non empty invariant open set and so we have

$$\lambda(W^{R \rightarrow L} \cap W^D) > 0.$$

Suppose first that $[T] \wedge \text{rot}_f(\lambda_{W^{R \rightarrow L} \cap W^D}) \leq 0$. Using the ergodic decomposition of $\lambda_{W^{R \rightarrow L} \cap W^D}$, we deduce that there exists $\nu' \in \mathcal{M}(f)$ ergodic such that $\nu'(W^{R \rightarrow L} \cap W^D) = 1$ and $[T] \wedge \text{rot}_f(\nu') \leq 0$. If $[T] \wedge \text{rot}_f(\nu') < 0$, we can apply Lemma 4.13 and so the conclusion of Lemma 4.16 holds. If $[T] \wedge \text{rot}_f(\nu') = 0$ we know that $\nu'(W^{L \rightarrow R}) = 1$ by Lemma 4.15 and so the first item of the conclusion of Lemma 4.16 holds thanks to Proposition 3.1.

Suppose now that $[T] \wedge \text{rot}_f(\lambda_{W^{R \rightarrow L} \cap W^D}) > 0$. From the equalities

$$[T] \wedge \text{rot}_f(\lambda) = 0$$

and

$$\text{rot}_f(\lambda_{\text{fix}(I)}) = 0 \text{ if } \lambda(\text{fix}(I)) \neq 0,$$

we deduce that

$$\lambda(\text{dom}(I) \setminus (W^{R \rightarrow L} \cap W^D)) > 0$$

and

$$[T] \wedge \text{rot}_f(\lambda_{\text{dom}(I) \setminus (W^{R \rightarrow L} \cap W^D)}) < 0.$$

Using the ergodic decomposition of $\lambda_{\text{dom}(I) \setminus (W^{R \rightarrow L} \cap W^D)}$, we deduce that there exists $\nu' \in \mathcal{M}(f)$ such that $[T] \wedge \text{rot}_f(\nu') < 0$. Here again we refer to Lemma 4.13 to ensure that the conclusion of Lemma 4.16 holds. \square

Let us conclude this section with a new result that will be useful for our purpose.

Proposition 4.17. *Suppose that $\nu \in \mathcal{M}(f)$ and $\nu' \in \mathcal{M}(f)$ are ergodic and that for ν -almost every point $z \in \text{dom}(I)$ and ν' -almost every point $z' \in \text{dom}(I)$, the path $I_{\mathcal{F}}^{\mathbb{Z}}(z')$ accumulates on $I_{\mathcal{F}}^{\mathbb{Z}}(z)$, then $\text{rot}_f(\nu) \wedge \text{rot}_f(\nu') \neq 0$.*

Proof. There is no loss of generality by supposing that $I_{\mathcal{F}}^{\mathbb{Z}}(z')$ accumulates positively on $I_{\mathcal{F}}^{\mathbb{Z}}(z)$. By Proposition 3.2, there exists a transverse simple loop $\Gamma_* \subset \Sigma$ such that

- $I_{\mathcal{F}}^{\mathbb{Z}}(z')$ is equivalent to the natural lift of Γ_* ;
- the union B of leaves met by Γ_* is an open annulus of S ;
- if $\tilde{\gamma}_*$ is a lift of Γ_* to $\widetilde{\text{dom}(I)}$, then for ν -almost every point $z \in \text{dom}(I)$, there is a lift $\tilde{z} \in \widetilde{\text{dom}(I)}$ such that $\tilde{I}_{\mathcal{F}}^{\mathbb{Z}}(\tilde{z})$ meets $\partial \tilde{B}^L$;
- for every $\tilde{\phi} \subset \partial \tilde{B}^L$ it holds that $\tilde{B} \subset R(\tilde{\phi})$.

The point z can be chosen recurrent and so every leaf of \mathcal{F} met by $I_{\mathcal{F}}^{\mathbb{Z}}(z)$ is met infinitely many often in the past and in the future. In particular, $I_{\mathcal{F}}^{\mathbb{Z}}(z)$ goes in and out of B infinitely many times, but it never enters in B on the left because $\tilde{B} \subset R(\tilde{\phi})$ for every $\tilde{\phi} \subset \partial \tilde{B}^L$. We deduce that every lift $\tilde{z} \in \tilde{B}$ of z crosses \tilde{B} from the right to the left. So, referring to the notations of the whole section, we have $\nu(W^{R \rightarrow L}) = 1$.

Lemma 4.18. *Let ν' be an f -invariant ergodic probability measure such that $\nu'(\text{dom}(I)) = 1$. Suppose that there is some deck transformation $T \in \mathcal{G} \setminus \{\text{Id}\}$ and a T -strip \tilde{B} projecting on an open annulus B of S such that ν' -almost every point $z' \in \text{dom}(I)$ satisfies $I_{\mathcal{F}}^{\mathbb{Z}}(z') \subset B$. Then there exists $a > 0$ such that $\text{rot}_f(\nu') = a[T]$.*

Proof. By Lemma 4.10 there exists $a \geq 0$ such that $\text{rot}_f(\nu') = a[T]$. We need to prove that $a \neq 0$. Let $U' \subset B$ be a topological open disk such that $\nu'(U') \neq 0$. We can suppose that U' is a flow-box that satisfies the conclusion of Proposition 4.4. Write $\varphi'_{U'} : U' \rightarrow U'$ for the first return map of f and $\tau'_{U'} : U' \rightarrow \mathbb{N} \setminus \{0\}$ for the time of first return map, which are defined ν' -almost everywhere on U' . Note that $\nu'|_{U'}$ is an ergodic invariant measure for $\varphi'_{U'}$. Fix a lift $\tilde{U}' \subset \tilde{B}$ of U' . For every point $z \in U'$ such that $\tau'_{U'}(z)$ exists, denote \tilde{z} the lift of z that is in \tilde{U}' and $\delta_{U'}(z)$ the integer such that $\tilde{f}^{\tau'_{U'}(z)}(\tilde{z}) \in T^{\delta(z)}\tilde{U}'$. One gets a map $\delta_{U'} : U' \rightarrow \mathbb{Z}$ defined ν' -almost everywhere on U' . Remind that a map $\rho_{U'} : U' \rightarrow H_1(S, \mathbb{Z})$ has been defined in the introduction and that $\rho_{U'}(z) = \delta(z)[T]$. Note also that $\delta(z) > 0$. The measure ν' being ergodic, by Kac's theorem one knows that

$$\int_{U'} \tau_{U'} d\nu' = \nu' \left(\bigcup_{k \geq 0} f^k(U') \right) = \nu' \left(\bigcup_{k \in \mathbb{Z}} f^k(U') \right) = 1,$$

and consequently that $\tau_{U'}^*(z) = 1/\nu'(U')$ for ν' -almost every point $z \in U'$, where $\tau_{U'}^*$ and $\rho_{U'}^*$ has been defined in (1) (page 3). Furthermore, for ν' -almost every point $z \in U'$, it holds that

$$\text{rot}_f(\nu') = \text{rot}_f(z) = \rho_{U'}^*(z)/\tau_{U'}^*(z) = \nu'(U')\rho_{U'}^*(z) = \left(\int_{U'} \delta(z) d\nu'(z) \right) [T].$$

Observe now that $\int_{U'} \delta(z) d\nu'(z) > 0$. This proves the lemma. \square

To prove Proposition 4.17 it remains to prove that $\text{rot}_f(\nu') \wedge \text{rot}_f(\nu) > 0$ which would lead to the result with Theorem 4.9. Let $U \subset B$ be a topological open disk such that $\nu(U) \neq 0$ and that is a flow-box that satisfies the conclusion of Proposition 4.4. Perturbing Γ_* and reducing U if necessary, one can suppose that $U \cap \Gamma_* = \emptyset$. Write $\varphi_U : U \rightarrow U$ for the first return map of f and $\tau_U : U \rightarrow \mathbb{N} \setminus \{0\}$ for the time of first return map, which are defined ν -almost everywhere on U . We will define a function $\delta_U : U \rightarrow \mathbb{Z}$ in a different way. For every point $z \in U$ such that $\tau_U(z)$ exists, set $m = \tau_U(z)$ and consider the set

$$X_z = \{t \in [0, m] \mid I_{\mathcal{F}}^t(z) \subset U\}.$$

Suppose first that $X_z \neq [0, \tau_U(z)]$. Then denote $(J_\xi)_{\xi \in \Xi}$ the family of connected components of X_z . One component J_{ξ^-} can be written $J_{\xi^-} = [0, b_{\xi^-})$, one component J_{ξ^+} can be written $J_{\xi^+} = (a_{\xi^+}, m]$ and the remaining components can be written $J_\xi = (a_\xi, b_\xi)$. Consider such a component J_ξ . The path $I_{\mathcal{F}}^m(z)$ can be lifted to a path $\tilde{I}_{\mathcal{F}}^m(\tilde{z})$ (the lift depending on ξ) such that $\tilde{I}_{\mathcal{F}}^m(\tilde{z})(a_\xi, b_\xi) \subset \tilde{B}$. By assumptions, one knows that $\tilde{I}_{\mathcal{F}}^m(\tilde{z})(a_\xi) \in \partial \tilde{B}^R$ and we set

$$\delta_\xi = \begin{cases} 0 & \text{if } \tilde{I}_{\mathcal{F}}^m(\tilde{z})(b_\xi) \in \partial \tilde{B}^R, \\ 1 & \text{if } \tilde{I}_{\mathcal{F}}^m(\tilde{z})(b_\xi) \in \partial \tilde{B}^L. \end{cases}$$

In the first situation $\tilde{I}_{\mathcal{F}}^m(\tilde{z})|_{[a_\xi, b_\xi]}$ visits \tilde{B} on the right, in the second one it crosses \tilde{B} for the right to the left. Note that there are finitely many $\xi \in \Xi$ such that $\delta_\xi = 1$ because there are finitely many $\xi \in \Xi$ such that $I_{\mathcal{F}}^m(z)([a_\xi, b_\xi]) \cap \Gamma_* \neq \emptyset$. Indeed, $\tilde{\gamma}_*$ is contained in \tilde{B} , while each such $I_{\mathcal{F}}^m(z)([a_\xi, b_\xi])$ meets $\partial \tilde{B}$; the conclusion follows by a compactness argument.

The path $I_{\mathcal{F}}^m(z)$ can be lifted to a path $\tilde{I}_{\mathcal{F}}^m(\tilde{z})$ such that $\tilde{I}_{\mathcal{F}}^m(\tilde{z})([0, b_{\xi^-})) \subset \tilde{B}$. Set

$$\delta_{\xi^-} = \begin{cases} 1/2 & \text{if } \tilde{I}_{\mathcal{F}}^m(\tilde{z})(b_{\xi^-}) \in \partial \tilde{B}^L, \\ -1/2 & \text{if } \tilde{I}_{\mathcal{F}}^m(\tilde{z})(b_{\xi^-}) \in \partial \tilde{B}^R. \end{cases}$$

Finally, set $\delta_{\xi^+} = 1/2$. Observe now that we have

$$[\Gamma_*] \wedge \rho_U(z) = \delta_U(z),$$

where ρ_U is defined page 3, and

$$\delta_U(z) = \begin{cases} \sum_{\xi \in \Xi} \delta_i & \text{if } X_z \neq [0, \tau_U(z)], \\ 0 & \text{if } X_z = [0, \tau_U(z)]. \end{cases}$$

The function δ_U is non negative but does not vanishes almost ν_U -everywhere because $I_{\mathbb{Z}}^{\mathbb{Z}}(z)$ does not stay in B for ν -almost every point. So, we have

$$[\Gamma] \wedge \text{rot}_f(\nu) = \nu(U)[\Gamma] \wedge \rho_U^*(z) = \int_U \delta_U(z) d\nu(z) > 0.$$

□

Remark. Using Lellouch's techniques [Lel, Section 3.4], one can more generally show that if z and z' are recurrent points (not necessarily trajectories of typical points for ergodic measures) and if $I_{\mathcal{F}}^{\mathbb{Z}}(z')$ accumulates on $I_{\mathcal{F}}^{\mathbb{Z}}(z)$, then f has a topological horseshoe³. However, we will not use this property in the sequel.

5. PROOF OF THE MAIN THEOREM

We suppose in this section that the hypotheses of Theorem A are satisfied. We consider an oriented closed surface S of genus $g \geq 2$ and a homeomorphism f of S isotopic to the identity that preserves a Borel probability measure λ with total support such that $\text{rot}_f(\lambda) = s\rho$, with $\rho \in H_1(S, \mathbb{Z}) \setminus \{0\}$ and $s \in \mathbb{R}$. We keep the notations of the article. We consider a Borel probability measure ν , invariant by f and ergodic. We consider a neighborhood \mathcal{U} of $\text{rot}_f(\nu)$ in $H_1(S, \mathbb{R})$ and want to prove that there exists a homotopical interval of rotation (κ, r) such that $[\kappa]/r \in \mathcal{U}$.

There is no loss of generality by supposing that f is not the identity map; in this case one can consider a maximal isotopy I of f by Theorem 4.1 with non empty domain. By Theorem 4.2, one can find a non singular foliation \mathcal{F} on $\text{dom}(I)$ transverse to I . Remind that:

- $\widetilde{\text{dom}}(I)$ is the universal covering space of $\text{dom}(I)$;
- $\text{dom}(I)_X$ is the connected component of $\widetilde{\text{dom}}(I)$ that contains a given connected set $X \subset \widetilde{\text{dom}}(I)$;
- $\tilde{\pi} : \widetilde{\text{dom}}(I) \rightarrow \text{dom}(I)$ is the covering projection;
- \mathcal{G} is the group of covering automorphism of $\tilde{\pi}$;
- $[T] \in H_1(S, \mathbb{Z})$ is the homology class of a loop $\Gamma \subset \text{dom}(I)$ associated to $T \in \mathcal{G}$;
- \tilde{I} is the lift of $I|_{\text{dom}(I)}$ to $\widetilde{\text{dom}}(I)$ that starts from the identity;
- \tilde{f} is the lift of $f|_{\text{dom}(I)}$ to $\widetilde{\text{dom}}(I)$ that is the end point of \tilde{I} ;
- $\tilde{\mathcal{F}}$ is the lift of \mathcal{F} to $\widetilde{\text{dom}}(I)$;
- $I_{\mathcal{F}}^{\mathbb{Z}}(z)$ is the whole \mathcal{F} -transverse trajectory of a point $z \in \text{dom}(I)$;
- $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})$ is the whole $\tilde{\mathcal{F}}$ -transverse trajectory of a point $\tilde{z} \in \widetilde{\text{dom}}(I)$.

Suppose first that $\text{rot}_f(\nu) \wedge \text{rot}_f(\lambda) \neq 0$. Using the ergodic decomposition of λ , we deduce that there exists $\nu' \in \mathcal{M}(f)$ ergodic such that $\text{rot}_f(\nu) \wedge \text{rot}_f(\nu') \neq 0$. By Theorem 4.9, we know that $f|_{\text{dom}(I)}$ has a rotational topological horseshoe of type (κ, r) with $[\kappa]/r \in \mathcal{U}$. If $\Gamma \subset \text{dom}(I)$ is a loop associated to T , then for every $p/q \in [0, 1]$ written in an irreducible way, there exists a periodic point $z \in \text{dom}(I)$ of period rq such that $I^{rq}(z)$ is freely homotopic to $[\Gamma]^p$ in $\text{dom}(I)$: it is freely homotopic to $[\Gamma]^p$ in S . Hence, f

³Be careful, in this case we do not have that $I_{\mathcal{F}}^{\mathbb{Z}}(z')$ and $I_{\mathcal{F}}^{\mathbb{Z}}(z)$ intersect \mathcal{F} -transversally.

has a homotopical interval of rotation of type (κ, r) such that $[\kappa]/r \in \mathcal{U}$, and the conclusion of Theorem A holds.

It remains to study the case where $\text{rot}_f(\nu) \wedge \text{rot}_f(\lambda) = 0$.

Lemma 5.1. *Suppose that $\text{rot}_f(\nu) \wedge \text{rot}_f(\lambda) = 0$. There exists $T \in \mathcal{G} \setminus \{\text{Id}\}$ satisfying $[T] \wedge \text{rot}_f(\lambda) = 0$ and a T -strip \tilde{B} such that ν -almost every point $z \in \text{dom}(I)$ has a lift \tilde{z} such that $\tilde{I}_{\tilde{f}}^{\mathbb{Z}}(\tilde{z})$ draws \tilde{B} . Moreover if \mathcal{U} is a neighborhood of $\text{rot}_f(\nu)$, one can suppose that there exists $r \geq 1$ such that $[T]/r \in \mathcal{U}$.*

Proof. Fix $z_0 \in \text{supp}(\nu) \cap \text{dom}(I)$ and a lift $\tilde{z}_0 \in \widetilde{\text{dom}(I)}$ of z_0 . One can find a topological open disk $U \subset \text{dom}(I)$ containing z_0 such that the connected component \tilde{U} of $\tilde{\pi}^{-1}(U)$ containing \tilde{z}_0 is a flow-box that satisfies the conclusion of Proposition 4.4. Write $\varphi_U : U \rightarrow U$ for the first return map of f and $\tau_U : U \rightarrow \mathbb{N} \setminus \{0\}$ for the time of first return map, which are defined ν -almost everywhere on U . Note that $\nu|_U$ is an ergodic invariant measure of φ_U . Remind that a map $\rho_U : U \rightarrow H_1(S, \mathbb{Z})$ has been defined in the introduction. For every point $z \in U$ such that $\tau_U(z)$ exists, denote \tilde{z} the preimage of z by $\tilde{\pi}$ that is in \tilde{U} and $\delta_U(z)$ the automorphism such that $\tilde{f}^{\tau_U(z)}(\tilde{z}) \in \delta_U(z)(\tilde{U})$. One gets a map $\delta_U : U \rightarrow \mathcal{G}$ defined ν -almost everywhere on U such that $\rho_U(z) = [\delta_U(z)]$. The measure ν being ergodic, one knows that

$$\int_U \tau_U d\nu = \nu \left(\bigcup_{k \geq 0} f^k(U) \right) = \nu \left(\bigcup_{k \in \mathbb{Z}} f^k(U) \right) = 1,$$

and consequently that $\tau_U^*(z) = 1/\nu(U)$ for ν -almost every point $z \in U$, where τ_U^* and ρ_U^* has been defined in (1) (page 3). Furthermore, for ν -almost every point $z \in U$, it holds that

$$\int_U \rho_U(z) d\nu(z) = \nu(U) \rho_U^*(z) = \text{rot}_f(z) = \text{rot}_f(\nu),$$

which implies that

$$\int_U \rho_U(z) \wedge \text{rot}_f(\lambda) d\mu(z) = \nu(U) \rho_U^*(z) \wedge \text{rot}_f(\lambda) = \text{rot}_f(\nu) \wedge \text{rot}_f(\lambda) = 0.$$

By Atkinson's Theorem [At], one knows that if $\varepsilon > 0$ is fixed, then for $\nu|_U$ almost every point z , there exists $n \geq 1$ such that

$$\left| \sum_{k=0}^{n-1} \rho_U(\varphi_U^k(z)) \wedge \text{rot}_f(\lambda) \right| < \varepsilon.$$

As observed by Lellouch [Lel], we can slightly improve this result: for $\nu|_U$ almost every point z , it holds that

$$\liminf_{n \rightarrow +\infty} \left| \sum_{k=0}^{n-1} \rho_U(\varphi_U^k(z)) \wedge \text{rot}_f(\lambda) \right| = 0.$$

So, if we fix a norm $\| \cdot \|$ on $H_1(S, \mathbb{R})$ and $\eta > 0$, we can find $z_1 \in \text{supp}(\mu) \cap U$ and $n \geq 1$ such that (recall that $\text{rot}_f(\lambda) = s\rho$, with $\rho \in H_1(S, \mathbb{Z}) \setminus \{0\}$ and

$s \in \mathbb{R}$)

$$\left| \sum_{k=0}^{n-1} \rho_U(\varphi_U^k(z_1)) \wedge \text{rot}_f(\lambda) \right| < s,$$

and such that

$$\left\| \frac{1}{n} \sum_{k=0}^{n-1} \rho_U(\varphi_U^k(z_1)) - \text{rot}_f(\nu) \right\| < \eta.$$

Every number $\rho_U(\varphi_U^k(z_1)) \wedge \text{rot}_f(\lambda)$ belonging to $s\mathbb{Z}$ we deduce that

$$\sum_{k=0}^{n-1} \rho_U(\varphi_U^k(z_1)) \wedge \text{rot}_f(\lambda) = 0.$$

Set

$$r = \sum_{0 \leq k < n} \tau_U(\varphi_U^k(z_1))$$

and denote \tilde{z}_1 the lift of z_1 that belongs to \tilde{U} . The automorphism T such that $\tilde{f}^r(\tilde{z}_1) \in T(\tilde{U})$ can be written

$$T = T_{n-1} \circ \cdots \circ T_1,$$

where T_k is an automorphism conjugated to $\delta_U(\varphi_U^k(z_1))$, so we have

$$[T] = \sum_{0 \leq k < n} [\delta_U(\varphi_U^k(z_1))].$$

Consequently, it holds that

$$[T] \wedge \text{rot}_f(\lambda) = 0, \quad \|[T]/r - \text{rot}_f(\nu)\| < \eta.$$

Note that we have $\tilde{f}^r(\tilde{z}_1) \in T(\tilde{U})$ if \tilde{z}_1 is the lift of z_1 that belongs to \tilde{U} . The property of \tilde{U} stated in Proposition 4.4 tells us that $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z}_1)$ intersects every leaf that meets \tilde{U} and every leaf that meets $T(\tilde{U})$. So, there is subpath $\tilde{\gamma}_1$ of $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z}_1)$ that joins $\phi_{\tilde{z}_1}$ to $T(\phi_{\tilde{z}_1})$. Of course we have $T \neq \text{Id}$. Moreover $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z}_1)$ draws the T -strip \tilde{B} defined by the line $\tilde{\gamma}_*$ obtained by concatenating⁴ the paths $T^k(\tilde{\gamma}_1)$, $k \in \mathbb{Z}$. As explained before, Proposition 4.3 tells us that the set W^D of points $z \in U$ that have a lift \tilde{z} such that $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})$ draws \tilde{B} is open. It is T -invariant and contains $z_1 \in \text{supp}(\nu)$. The measure ν being ergodic, it holds that $\nu(W^D) = 1$. \square

Proof of Theorem A. Let us summarize in which cases the results we have already proved allow us to get Theorem A. Recall that the sets W^* are defined in Paragraph 4.2.

- If $\nu(W^{R \rightarrow R} \cap W^D) = 1$ or $\nu(W^{L \rightarrow L} \cap W^D) = 1$, then by Proposition 3.1 for ν -almost every point z , the path $I_{\mathcal{F}}^{\mathbb{Z}}(z)$ has an \mathcal{F} -transverse self intersection; this allows to apply Corollary 4.8 and to get a suitable rotational horseshoe.
- If $\nu(W^{R \rightarrow L} \cap W^D) = 1$, there are three cases:

⁴Strictly speaking one has to modify the path γ_1 lifted by $\tilde{\gamma}_1$ to be able to concatenate $T^k(\tilde{\gamma}_1)$ with $T^{k+1}(\tilde{\gamma}_1)$: it is sufficient to move it along the leaves so that the last endpoint of $\tilde{\gamma}_1$ with the first endpoint of $T(\tilde{\gamma}_1)$ coincide.

- If $[T] \wedge \text{rot}_f(\nu) < 0$, then one can apply Corollary 4.14 which shows that for ν -almost every point z , the path $I_{\mathcal{F}}^{\mathbb{Z}}(z)$ has an \mathcal{F} -transverse self intersection; this allows to apply Corollary 4.8 and to get a suitable rotational horseshoe.
- If $[T] \wedge \text{rot}_f(\nu) = 0$, then one can apply Lemma 4.15 which shows that for ν -almost every point z , the path $I_{\mathcal{F}}^{\mathbb{Z}}(z)$ has an \mathcal{F} -transverse self intersection; as before this allows to apply Corollary 4.8 and to get a suitable rotational horseshoe.
- If $[T] \wedge \text{rot}_f(\nu) > 0$, then one can apply Lemma 4.16. It tells us that there exists an ergodic invariant probability measure ν' such that for ν -almost every point z and ν' -almost every point z' , either the paths $I_{\mathcal{F}}^{\mathbb{Z}}(z)$ and $I_{\mathcal{F}}^{\mathbb{Z}}(z')$ have an \mathcal{F} -transverse intersection, or the path $I_{\mathcal{F}}^{\mathbb{Z}}(z')$ accumulates on $I_{\mathcal{F}}^{\mathbb{Z}}(z)$. In the first case one can apply Corollary 4.8 to get a suitable rotational horseshoe. In the second case Proposition 4.17 tells us that $\text{rot}_f(\nu) \wedge \text{rot}_f(\nu') \neq 0$. Lellouch's Theorem 4.9 then gives us a suitable rotational horseshoe.
- The case $\nu(W^{L \rightarrow R} \cap W^D) = 1$ is identical to the case $\nu(W^{R \rightarrow L} \cap W^D) = 1$.

In all these cases the existence of a suitable homotopical interval of rotation is due to the presence of a rotational topological horseshoe. To get Theorem A it remains to study a last case where the existence of a suitable homotopical interval of rotation will have another reason. One can write $T = T'^m$, $m \geq 1$, where $T' \in \mathcal{G}$ is irreducible. The following proposition will permit us to finish the proof of Theorem A. Indeed, let \mathcal{U} be a neighborhood of $\text{rot}_f(\nu)$ in $H_1(S, \mathbb{R})$. One can find $p_0/q_0 \in (0, a)$ written in an irreducible way such that $p_0[T']/q_0 \in \mathcal{U}$. By Proposition 5.2, for every $p/q \in [0, 1]$ written in an irreducible way, there exists $\tilde{z}_{p/q}$ such that $\tilde{f}^{qq_0}(\tilde{z}) = T'^{pp_0}(\tilde{z})$. The image $z_{p/q} = \tilde{\pi}(\tilde{z}_{p/q}) \in S$ is fixed by f^{qq_0} and the loop of S defined by $I^{qq_0}(z_{p/q})$ belongs to $[T']_{\mathcal{F}\mathcal{H}\mathcal{L}}^{pp_0}$. Denote $q' = qq_0/s$ the period of $z_{p/q}$. There exists $R \in \mathcal{G}$ such that $\tilde{f}^{q'}(\tilde{z}_{p/q}) = R(\tilde{z}_{p/q})$. We deduce that $T'^{pp_0}(\tilde{z}_{p/q}) = \tilde{f}^{qq_0}(\tilde{z}_{p/q}) = R^s(\tilde{z}_{p/q})$. It implies that $T'^{pp_0} = R^s$. The group $\langle T', R \rangle$ being a free group, it must be infinite cyclic. We deduce that R is a power of T' because T' is irreducible and so s divides pp_0 and qq_0 . The integers p_0 and q_0 being relatively prime, it holds that $s \gcd(s, p_0)^{-1} \gcd(s, q_0)^{-1}$ is an integer. Moreover it is relatively prime with p_0 and with q_0 . So it divides p and q . These integers being relatively prime, we have $s = \gcd(s, p_0) \gcd(s, q_0) \leq p_0 q_0$ and hence the period $q' = qq_0/r = s$ of $z_{p/q}$ satisfies $q' \geq q/p_0$. We deduce that $([T']_{\mathcal{F}\mathcal{H}\mathcal{L}}^{p_0}, q_0, p_0)$ is a homotopical interval of rotation. \square

Proposition 5.2. *If the sets*

$$W^{R \rightarrow L} \cap W^D, \quad W^{L \rightarrow R} \cap W^D, \quad W^{R \rightarrow R} \cap W^D, \quad W^{L \rightarrow L} \cap W^D$$

are ν -null sets, then there exists $a > 0$ such that :

- *one has $\text{rot}_f(\nu) = a[T']$;*
- *for every $p/q \in [0, a) \cap \mathbb{Q}$, written in an irreducible way, there exists \tilde{z} such that $\tilde{f}^q(\tilde{z}) = T'^p(\tilde{z})$.*

Proof. Recall that $T = T'^m$, where $m \geq 1$. Let us begin by proving that \tilde{B} is invariant by T' . It is sufficient to prove that for every $n > 0$ we have $\overline{L(T'^n \tilde{\phi})} \subset L(\tilde{\phi})$. If $L(\tilde{\phi}) \subset L(T'^n \tilde{\phi})$, then for every $k \geq 1$ we have $L(T'^{nk} \tilde{\phi}) \subset L(T'^{n(k+1)} \tilde{\phi})$ and so we deduce that $L(\tilde{\phi}) \subset L(T'^{nm} \tilde{\phi})$, which contradicts the inclusion $\overline{L(T'^{nm} \tilde{\phi})} \subset L(\tilde{\phi})$. If $L(\tilde{\phi}) \cap L(T'^n \tilde{\phi}) = \emptyset$, then $L(\tilde{\phi})$ is disjoint from its image by T'^n . The map T'^n being fixed point free, Brouwer Translation Theorem [Br] tells us that $L(\tilde{\phi})$ is disjoint from its image by T'^{nm} , which contradicts the inclusion $\overline{L(T'^{nm} \tilde{\phi})} \subset L(\tilde{\phi})$. Similarly, if $R(\tilde{\phi}) \cap R(T'^n \tilde{\phi}) = \emptyset$, then $\overline{R(\tilde{\phi})}$ is disjoint from its image by T'^{nm} , which contradicts the inclusion $\overline{R(\tilde{\phi})} \subset R(T'^{nm} \tilde{\phi})$. The only remaining case is the case where $\overline{L(T'^n \tilde{\phi})} \subset L(\tilde{\phi})$.

In the following instead of seeing \tilde{B} as a \tilde{T} -strip, we will see it as a T' -strip: one can choose $\tilde{\gamma}_*$ to be invariant by T' and suppose that $\tilde{\gamma}_*(t) = T' \gamma_*(t)$ for every $t \in \mathbb{R}$.

By construction of \tilde{B} we know that $\nu(W^D) = 1$. So, ν -almost every point $z \in \text{dom}(I)$ is recurrent and has a lift $\tilde{z} \in \widetilde{\text{dom}(I)}$ such that $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})$ is equivalent to $\tilde{\gamma}_*$ at $+\infty$ or $-\infty$. Indeed if $z \in W^D$ is recurrent and if \tilde{z} was a lift of z such that $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})$ accumulates on $\tilde{\gamma}_*$, then there would exist $k \in \mathbb{Z}$ such that $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})$ accumulates on $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(T'^k(\tilde{z}))$. It is impossible because z is recurrent and so has no self-accumulation by Corollary 3.9. Hence, $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})$ does not accumulate on $\tilde{\gamma}_*$, and by the hypothesis of the proposition it cannot go out of \tilde{B} both before and after it draws \tilde{B} . This implies that it has to be equivalent to $\tilde{\gamma}_*$ at $+\infty$ or $-\infty$.

In fact we can be more precise: if there are $a < a'$ and $b \in \mathbb{R}$ such that $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})|_{[a, a']}$ is equivalent to $\tilde{\gamma}_*|_{[b, b+1]}$, then either $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})|_{[a, +\infty)}$ is equivalent to a subpath of $\tilde{\gamma}_*$ (and equivalent to $\tilde{\gamma}_*$ at $+\infty$ but we will not use this property) or $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})|_{(-\infty, a']}$ is equivalent to a subpath of $\tilde{\gamma}_*$. From this we will deduce the following lemma.

Lemma 5.3. *The transverse path $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})$ is equivalent to $\tilde{\gamma}_*$. Moreover there is neighborhood \tilde{U} of \tilde{z} such that if the orbit of \tilde{z} meets $R\tilde{U}$ for some $R \in \mathcal{G}$, then R is a power of T' .*

Proof. Let us treat the case where $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})|_{[a, +\infty)}$ is equivalent to a subpath of $\tilde{\gamma}_*$, the other case being identical.

Suppose that $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})$ is not equivalent to $\tilde{\gamma}_*$. Then, as we have already seen that it cannot accumulate in $\tilde{\gamma}_*$, this means that there exists $b < a$, $b \in \mathbb{Z}$, such that $\tilde{I}_{\tilde{\mathcal{F}}}^b(\tilde{z}) \notin \tilde{B}$.

By recurrence of the point z , there exists a sequence of integers $n_k \rightarrow -\infty$, and a sequence of deck transformations $(R_k)_{k \in \mathbb{N}} \in \mathcal{G}$ such that $R_k \tilde{f}^{n_k}(\tilde{z})$ tends to z ; in particular for any k large enough:

- $\tilde{\gamma}_*|_{[b, b+1]}$ is equivalent to a subpath of $R_k \tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})|_{[n_k + a - 1, +\infty)}$ (and in particular this path draws \tilde{B});
- $R_k \tilde{I}_{\tilde{\mathcal{F}}}^{n_k + b}(\tilde{z}) \notin \tilde{B}$.

By the same reasoning as before the lemma, we deduce that either the trajectory $R_k \tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})|_{[n_k + a + 1, +\infty)}$ is equivalent to a subpath of $\tilde{\gamma}_*$, or $R_k \tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})|_{(-\infty, n_k + a' - 1]}$

is equivalent to a subpath of $\tilde{\gamma}_*$. By the second point above, the second situation is impossible. Hence, $R_k \tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})|_{[n_k+a+1, +\infty)}$ is equivalent to a subpath of $\tilde{\gamma}_*$.

In particular, this implies that $R_k \tilde{\gamma}_*$ is equivalent at $+\infty$ to $\tilde{\gamma}_*$. By Lemma 3.3, this implies that $R_k \tilde{\gamma}_* \cap \tilde{\gamma}_* \neq \emptyset$; more precisely it implies that for any n large enough, $R_k \tilde{\gamma}_* \cap \tilde{\gamma}_*|_{[b+n, b+n+1)} \neq \emptyset$, hence that $R_k \tilde{\gamma}_* \cap \tilde{\gamma}_*$ is infinite. This implies that $R_k \tilde{\gamma}_* = \tilde{\gamma}_*$, in other words $R_k = T'^{i_k}$ for some $i_k \in \mathbb{Z}$.

We deduce that $T'^{i_k} \tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})|_{[n_k+a+1, +\infty)}$ is equivalent to a subpath of $\tilde{\gamma}_*$, equivalently (as $\tilde{\gamma}_*$ is T' -invariant), for any $k \in \mathbb{N}$, the path $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})|_{[n_k+a+1, +\infty)}$ is equivalent to a subpath of $\tilde{\gamma}_*$.

This proves that $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})$ is equivalent to a subpath of $\tilde{\gamma}_*$. As it cannot accumulate in $\tilde{\gamma}_*$, this proves that $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})$ is equivalent to $\tilde{\gamma}_*$.

To get the second part of the lemma, consider a neighborhood \tilde{U} of \tilde{z} such that for every $\tilde{z}' \in \tilde{U}$, the path $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z}')$ draws $\tilde{\gamma}_*$. If $f^k(\tilde{z}) \in R\tilde{U}$, $R \in \mathcal{G}$, then $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})$ draws $R(\tilde{\gamma}_*)$. We deduce that $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})$ is equivalent to $R\tilde{\gamma}_*$. What was done above tells us that $R \in \langle T' \rangle$. \square

Now, let us consider

- the connected component $\widetilde{\text{dom}}(I)_{\tilde{\gamma}_*}$ of $\widetilde{\text{dom}}(I)$ that contains $\tilde{\gamma}_*$,
- the quotient space $\widehat{\text{dom}}(I) = \widetilde{\text{dom}}(I)/T$,
- the foliation $\hat{\mathcal{F}}$ of $\widehat{\text{dom}}(I)$ lifted by $\tilde{\mathcal{F}}$,
- the covering projection $\hat{\pi} : \widehat{\text{dom}}(I) \rightarrow \text{dom}(I)$,
- the annulus $\widehat{\text{dom}}(I)_{\tilde{\gamma}_*} = \widetilde{\text{dom}}(I)_{\tilde{\gamma}_*}/T'$,
- the universal covering projection $\pi : \widetilde{\text{dom}}(I)_{\tilde{\gamma}_*} \rightarrow \widehat{\text{dom}}(I)_{\tilde{\gamma}_*}$.

Lemma 5.4. *It holds that ν -almost every point z has a lift in $\widetilde{\text{dom}}(I)_{\tilde{\gamma}_*}$ that is positively recurrent and has a rotation number $a > 0$ (in the annulus). Moreover we have $\text{rot}_f(\nu) = a[T']$.*

Proof. We know that ν -almost every point z is positively recurrent and has a lift \tilde{z} in $\widetilde{\text{dom}}(I)_{\tilde{\gamma}_*}$ that draws \tilde{B} . We have seen in Lemma 5.3 that $\tilde{I}_{\tilde{\mathcal{F}}}^{\mathbb{Z}}(\tilde{z})$ is equivalent to $\tilde{\gamma}_*$ and that there exists a neighborhood \tilde{U} of \tilde{z} such that if the orbit of \tilde{z} meets $R\tilde{U}$, for some $R \in \mathcal{G}$, then R is a power of T' . Using the fact that z is recurrent, we deduce that $\hat{z} = \pi(\tilde{z})$ is positively recurrent. By the argument given in the proof of Lemma 4.18, we deduce that z has rotation number $a > 0$. Moreover we have $\text{rot}_f(\nu) = a[T]$. \square

Now there are two cases to consider. The first case is the case where the stabilizer of $\widetilde{\text{dom}}(I)_{\tilde{\gamma}_*}$ is generated by T' and the second case is when it is larger. In the first case, $\hat{\pi}$ sends homeomorphically $\widehat{\text{dom}}(I)_{\tilde{\gamma}_*}$ onto a connected component of $\text{dom}(I)$. Moreover, the frontier of this annulus is made of contractible fixed points of f . In the second case, $\hat{\pi}$ sends $\widehat{\text{dom}}(I)_{\tilde{\gamma}_*}$ onto a hyperbolic surface whose universal covering space is $\widetilde{\text{dom}}(I)_{\tilde{\gamma}_*}$ and the group of covering automorphisms is the stabilizer of $\widetilde{\text{dom}}(I)_{\tilde{\gamma}_*}$ in \mathcal{G} . In both cases, there exists an extension $\widetilde{\widehat{\text{dom}}}(I)_{\tilde{\gamma}_*}$ of $\widehat{\text{dom}}(I)_{\tilde{\gamma}_*}$ obtained by blowing

at least one end e with a circle $\hat{\Gamma}_e$ and \hat{f} extends to a homeomorphism $\widehat{\tilde{f}}$ of $\widehat{\text{dom}(I)_{\tilde{\gamma}_*}}$ (see Paragraph 2.4). Furthermore, the rotation number(s) induced on the added circle(s) by the lift of $\widehat{\tilde{f}}$ that extends \tilde{f} are equal to 0.

By Lemma 5.4, there exist positively recurrent points with rotation number $a > 0$ where $\text{rot}_f(\nu) = a[T']$. Consequently, according to Theorem 2.1 that can be applied thanks to Lemma 4.12, for every rational number $p/q \in (0, a)$, written in an irreducible way, there exists a point \tilde{z} such that $\tilde{f}^q(\tilde{z}) = T^p(\tilde{z})$. As \tilde{f} also has a fixed point by Lefschetz index theorem, this means that f has a homotopical interval of rotation of type (κ, r) such that $[\kappa]/r \in \mathcal{U}$. \square

REFERENCES

- [Ad] S. ADDAS ZANATA: On properties of the vertical rotation interval for twist mappings. II. *Qual. Theory Dyn. Syst.* **4** (2003), no. 2, 125–137.
- [AdT] S. ADDAS-ZANATA, F.A. TAL: On periodic points of area preserving torus homeomorphisms. *Far East J. Dyn. Syst.* **9** (2007), no. 3, 371–378.
- [At] G. ATKINSON: Recurrence of co-cycles and random walks. *J. London Math. Soc.* (2) **13** (1976), 486–488.
- [BeCLer] F. BÉGUIN, S. CROVISIER, F. LE ROUX: Fixed point sets of isotopies on surfaces. *J. Eur. Math. Soc. (JEMS)* **22** (2020), no. 6, 197–2046.
- [Bi] G.-D. BIRKHOFF: An extension of Poincaré’s last geometric theorem. *Acta Math.* **47** (1926), no. 4 297–311.
- [Br] L. E. J. BROUWER : Beweis des ebenen Translationssatzes, *Math. Ann.*, **72** (1912), 37-54.
- [CPoPrZ] D. CRISTOFARO-GARDINER, D. POMERLEANO, R. PRASAD, B. ZHANG A note in U -cyclic elements in monopole Floer homology, *arXiv* 2110.13844v2 (2022).
- [CPrZ] D. CRISTOFARO-GARDINER, R. PRASAD, B. ZHANG Periodic Floer homology and the smooth closing lemma for area-preserving surface diffeomorphisms, *arXiv* 2110.02925 (2022).
- [D] E. DOEFF: Rotation measures for homeomorphisms of the torus homotopic to a Dehn twist. *Ergodic Theory Dynam. Systems* **17** (1997), no. 3, 575–591.
- [EH] O. EDTMAIR, M. HUTCHINGS PFH spectral invariants and C^∞ -closing lemmas, *arXiv* 2110.02463v4 (2022).
- [F] J. FRANKS: Generalizations of the Poincaré-Birkhoff theorem. *Ann. of Math.*(2) **128** (1988), no. 1, 139–151.
- [FH] J. FRANKS, M. HANDEL: Periodic points of Hamiltonian surface diffeomorphisms. *Geometry and Topology* **7** (2003), 713–756.
- [G] V. GINZBURG: The Conley conjecture. *Annals of Math.* (2) **172** (2010), no. 2, 1127–1180.
- [H] M.-E. HAMSTROM. Homotopy groups of the space of homeomorphisms on a 2-manifold, *Illinois J. Math.*, **10** (1966), 563-573.
- [J] O. JAULENT: Existence d’un feuilletage positivement transverse à un homéomorphisme de surface. *Ann. Inst. Fourier (Grenoble)* **64** (2014), no. 4, 1441–1476.
- [K] M. KAC: On the notion of recurrence in discrete stochastic processes *Bull. Amer. Math. Soc* **53** (1947), no. 10, 1002–1010.
- [Lec1] P. LE CALVEZ: Une version feuilletée équivariante du théorème de translation de Brouwer. *Publ. Math. Inst. Hautes Études Sci.* **102**, (2005) 1-98.
- [Lec2] P. LE CALVEZ: Periodic orbits of Hamiltonian homeomorphisms of surfaces. *Duke Math. J.* **133**, (2006) no. 1, 125–184.
- [Lec3] P. LE CALVEZ: Conservative surface homeomorphisms with finitely many periodic points. *J. Fixed Point Theory Appl.* **24**, (2022) no. 2, 36 pp.
- [LecSa] P. LE CALVEZ, M. SAMBARINO: Homoclinic orbits for area preserving diffeomorphisms of surfaces. *Ergodic Theory Dynam. Systems* **42**, (2022), no. 3, 1122–1165.

- [LecT1] P. LE CALVEZ, F. TAL: Forcing theory for transverse trajectories of surface homeomorphisms. *Invent. Math.* **212**, (2018), no. 2, 619–729.
- [LecT2] P. LE CALVEZ, F. TAL: Topological horseshoes for surface homeomorphisms. *Duke Math. J.* **171** (2022), no. 12, 2519–2626.
- [Lel] G. LELLOUCH: Sur les ensembles de rotation des homéomorphismes de surface en genre ≥ 2 . *Mém. Soc. Math. Fr. (N.S.)*, to appear.
- [Math] J. MATHER: Topological proofs of some purely topological consequences of Caratheodory’s Theory of prime ends. in Selected Studies, TH. M. Rassias, G. M. Rassias, eds, *North Holland*, **168** (1982), 225–255.
- [Mats1] S. MATSUMOTO: Rotation sets of surface homeomorphisms. *Bol. Soc. Brasil. Mat. (N.S.)*, **28** (1997), no. 1, 225–255.
- [Mats2] S. MATSUMOTO: Arnold conjecture for surface homeomorphisms. Proceedings of the French-Japanese Conference ”Hyperspace Topologies and Applications” (La Bussière, 1997). *Topology Appl.*, **104** (2000), no. 1-3, 191–214.
- [OI1] F. OLIVEIRA: On the generic existence of homoclinic points. *Ergodic Theory Dynam. Systems* **7** (1987), no. 4, 567–595.
- [OI2] F. OLIVEIRA: On the C^∞ genericity of homoclinic orbits. *Nonlinearity* **13** (2000), no. 3, 653–662.
- [OxU] J. C. OXTOBY, S. M. ULAM Measure-preserving homeomorphisms and metrical transitivity. *Ann. of Math. (2)* **42** (1941), 874–920.
- [PaPotSa] A. PASSEGGI, R. POTRIE, M. SAMBARINO: Rotation intervals and entropy on attracting annular continua. *Geom. Topol.* **22** (2018), no. 4, 2145–2186.
- [Pi] D. PIXTON: Planar homoclinic point. *J. Differential Equations* **44** (1982), no. 3, 365–382.
- [Pol] M. POLLICOTT: Rotation sets for homeomorphisms and homology. *Trans. Amer. Math. Soc.*, **331** (1992), no. 2, 881–894.
- [Pr] R. PRASAD: Periodic points of rational area-preserving homeomorphisms. Manuscript, (2023).
- [R] C. ROBINSON: Closing stable and unstable manifolds on the two sphere. *Proc. Amer. Math. Soc.* **41** (1973), 299–303.
- [Sc] S. SCHWARTZMAN: Asymptotic cycles. *Ann. of Math.* **66** (1957), no. 2, 270–284.
- [W] H. WEISS: A remark on papers by Pixton and Oliveira: Genericity of symplectic diffeomorphisms of S^2 with positive topological entropy. *J. Stat. Phys.* **80** (1995), no. 1-2, 481–485.

INSTITUT DE MATHÉMATIQUES DE JUSSIEU-PARIS RIVE GAUCHE, IMJ-PRG, SORBONNE UNIVERSITÉ, UNIVERSITÉ PARIS-DIDEROT, CNRS, F-75005, PARIS, FRANCE
Email address: pierre-antoine.guiheneuf@imj-prg.fr

INSTITUT DE MATHÉMATIQUES DE JUSSIEU-PARIS RIVE GAUCHE, IMJ-PRG, SORBONNE UNIVERSITÉ, UNIVERSITÉ PARIS-DIDEROT, CNRS, F-75005, PARIS, FRANCE & INSTITUT UNIVERSITAIRE DE FRANCE
Email address: patrice.le-calvez@imj-prg.fr

CMAT, FACULTAD DE CIENCIAS, UNIVERSIDAD DE LA REPÚBLICA, URUGUAY.
Current address: Igua 4225 esq. Mataojo. Montevideo, Uruguay.
Email address: alepasseggi@cmat.edu.uy