



HAL
open science

Quasi-two-dimensional turbulence

Alexandros Alexakis

► **To cite this version:**

Alexandros Alexakis. Quasi-two-dimensional turbulence. *Reviews of Modern Plasma Physics*, 2023, 7 (1), pp.31. 10.1007/s41614-023-00134-3 . hal-04268090

HAL Id: hal-04268090

<https://hal.science/hal-04268090>

Submitted on 2 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Quasi-two-dimensional Turbulence

Alexandros Alexakis
Laboratoire de Physique de l'Ecole normale supérieure, ENS,
Université PSL, CNRS, Sorbonne Université, Université de Paris,
F-75005 Paris, France
alexakis@phys.ens.fr

November 2, 2023

Abstract

Many fluid-dynamical systems met in nature are *quasi-two-dimensional*: they are constrained to evolve in approximately two dimensions with little or no variation along the third direction. This has a drastic effect in the flow evolution because the properties of three dimensional turbulence are fundamentally different from those of two dimensional turbulence. In three-dimensions energy is transferred on average towards small scales, while in two dimensions energy is transferred towards large scales. Quasi-two-dimensional flows thus stand in a crossroad, with two-dimensional motions attempting to self-organize and generate large scales while three dimensional perturbations cause disorder, disrupting any large scale organization. Where is energy transferred in such systems? It has been realised recently that in fact the two behaviors can coexist with a simultaneous transfer of energy both to large and to small scales. How the cascade properties change as the variations along the third direction are suppressed has lead to discovery of different regimes or *phases* of turbulence of unexpected richness in behavior. Here, recent discoveries on such systems are reviewed. It is described how the transition from three-dimensional to two-dimensional flows takes place, the different phases of turbulence met and the nature of the transitions from one phase to the other. Finally, the implications these new discoveries have on different physical systems are discussed.

1 Introduction

We live in a world of (at least) three spatial dimensions. However, for some physical systems a two dimensional description appears to provide a suitable approximation for the dynamics involved. This happens when one dimension is highly compactified like in thin films or due to some other physical mechanisms, like rotation or strong magnetic fields, that prevents variations along one dimension. In fluid dynamics such a dimension-reduction is met in a variety of systems from molecular to astrophysical scales. At the smallest scales, such a reduction is met in the dynamics of electrons in ultra-pure materials such as graphene that display two dimensional hydrodynamic behavior [77, 3, 10]. In an equally exotic case, two-dimensional flows are met in fluids of light [1, 36]. Two dimensional flows have also been observed in Bose-Einstein condensates in liquid Helium [93, 46, 52]. Furthermore, two-dimensional dynamics are shown to be a good approximation for the motion of bacteria in films or thin layers [104, 62, 122, 121] and for the flow in soap films [72, 91, 120, 54]. Increasing in scale, plasma flows in the presence of strong magnetic fields, such as in Tokamak devices are also known to display close to two-dimensional dynamics [125, 41]. Finally, at the planetary scale, flows are very often constrained to two dimensional motions due to the presence of rotation, stratification and geometrical constraints [129, 21, 56, 100].

In the systems above, despite being close to two-dimensional, variations along the third direction can not always be disregarded, and some times play an important role in the energy balance relation. This is in particular true for the turbulent case where the dynamics of the flow drastically differ in two and in three dimensions. Turbulence is realised both in three and in two dimensions when the Reynolds number Re (the ratio of the viscous time scale to the to eddy turn-over time) attains large values. In three dimensional turbulence, the self-interaction between eddies generates smaller and smaller eddies transferring energy towards the smallest scales where it is dissipated effectively by viscosity, independently of how small it is. In two dimensions, the opposite behavior is observed. Eddies self-organize to generate larger eddies and thus transfer energy to larger scales. Unlike the three-dimensional case, in two dimensions, because energy is transferred upscale, viscosity is not efficient at dissipating it and energy piles up at the largest scale of the system. The dynamics of flows in three and in two dimensions are thus fundamentally different.

This brings out the question: what happens when a flow is only approximately two-dimensional and the flow lies between the two extreme situations? On the one hand, if two dimensional motions dominate energetically one can expect that flow dynamics will not be far from those of two dimensions and the transfer of energy will be towards the large scales. On the other hand, the inverse transfer of energy can be proved only if *enstrophy* (the mean square of vorticity) is conserved exactly, which is only true in two dimensional flows. Three dimensional perturbations, even if subdominant, will break enstrophy conservation and the upscale transfer of energy is questioned. This question for the behavior of quasi-two dimensional flows has been around for a long time [25]. What has been revealed in the last years is that a hybrid state of turbulence exists where transfer to larger and to smaller scales coexist, building what is referred to as a *bidirectional* or *split* cascade.

This review deals with when, how and in what sense can a flow transition from three dimensional to two dimensional behavior as a control parameter is varied. We will give a phenomenological description of the different *phases* of turbulence that are observed and describe their behavior close to the *critical points* where *phase transitions* are observed. The richness of behaviors observed in quasi-two-dimensional turbulence surpassed all expectations leading to the discovery in the last years of new phenomena that challenge our mathematical and physical understanding of fluid turbulence. These new discoveries are reviewed and categorized in order to give direction to future research. Furthermore, the appearance of quasi-two-dimensional turbulence in a variety of different systems is also reviewed by mentioning key works in the different systems. Finally, an attempt is made to note the many questions that are open in the field, stressing the need for further studies. Throughout the text, the mathematical formalism is made as light as possible in favor of readability and give references for more formal descriptions.

The rest of this review is structured as follows. In section 2, three and two dimensional turbulence are briefly reviewed. Then in section 3, thin layer turbulence is discussed and a detailed description of the transition from three-dimensional (3D) to two-dimensional (2D) dynamics is given. In section 4, some recent results in other quasi-2D systems like rotating and magneto-hydrodynamic flows are given. In section 5 conclusions are drawn and open questions in the field are presented.

2 Two and three dimensional turbulence

Turbulence in three and two dimensions has been a subject of study for well over a century, earning itself the title of the last unsolved problem of classical physics. Some excellent books on three dimensional turbulence can be found in [39, 84, 27, 26] while reviews on three and two dimensional turbulence can be found in [6, 130, 107, 16, 81]. The present review is limited in presenting some basic results that are indispensable for the discussion that follows and refer the reader to the former mentioned reviews for any further information.

2.1 Three dimensional turbulence

In its simplest form, turbulence in three and two dimensions of a unit density fluid, is described by the evolution of the incompressible velocity field \mathbf{u} that follows the Navier-Stokes equation

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} = -\nabla P + \nu \nabla^2 \mathbf{u} - \alpha \bar{\mathbf{u}} + \mathbf{f} \quad (1)$$

where P is the pressure imposing the incompressibility condition $\nabla \cdot \mathbf{u} = 0$, ν is the viscosity and \mathbf{f} is an external force assumed to act on a lengthscale ℓ_f and inject energy at a rate $\mathcal{I}_\mathcal{E}$. The term $-\alpha \bar{\mathbf{u}}$ (where $\bar{\mathbf{u}}$ stands for the vertically averaged velocity) is an additional drag term that models the effect of boundaries and is added here to make contact later on with 2D turbulence. We will consider as domain an orthogonal box of dimensions $L \times L \times H$ with H being the height along the direction which fluctuations are suppressed. For simplicity, we consider here only periodic boundary conditions. For a given functional form of the forcing and non-dimensionalizing using ℓ_f and $\mathcal{I}_\mathcal{E}$ the resulting non-dimensional numbers of the system are (i) the Reynolds number is given by $Re = \frac{\mathcal{I}_\mathcal{E}^{1/3} \ell_f^{4/3}}{\nu}$ (ii) the large scale Reynolds number $Re_\alpha = \frac{\mathcal{I}_\mathcal{E}^{1/3}}{\ell_f^{2/3} \alpha}$ (iii) the length scale ratio L/ℓ_f and (iv) the normalized height H/ℓ_f . In what follows all quantities will be non dimensionalized using the forcing length-scale ℓ_f and the energy injection rate $\mathcal{I}_\mathcal{E}$ (thus setting $\ell_f = 1$ and $\mathcal{I}_\mathcal{E} = 1$).

In three dimensions, the inviscid unforced system 1 for smooth flows, conserves two quadratic invariants. The first is energy $\mathcal{E} = \frac{1}{2} \langle |\mathbf{u}|^2 \rangle$ where the angular brackets stand for volume average. The second is helicity $\mathcal{H} = \frac{1}{2} \langle \mathbf{u} \cdot \mathbf{w} \rangle$ (where $\mathbf{w} = \nabla \times \mathbf{u}$ is the vorticity) that is not going to be discussed here but refer the reader to [6, 89]. In the presence of a forcing and dissipation the following energy balance relation holds

$$\mathcal{I}_\mathcal{E} = \epsilon_\nu + \epsilon_\alpha \quad (2)$$

where $\mathcal{I}_\mathcal{E} = \langle \mathbf{u} \cdot \mathbf{f} \rangle$ is the time and volume averaged energy injection rate, $\epsilon_\nu = \nu \langle |\nabla \mathbf{u}|^2 \rangle$ is the energy dissipation rate due to viscosity and $\epsilon_\alpha = \alpha \langle |\bar{\mathbf{u}}|^2 \rangle$ is the energy dissipation rate due to the drag term.

To express the notion of scale we are going to use the Fourier transformed fields $\hat{\mathbf{u}}_{\mathbf{k}}$ such that

$$\hat{\mathbf{u}}_{\mathbf{k}} = \left\langle e^{-i\mathbf{k}\cdot\mathbf{x}} \mathbf{u} \right\rangle \quad \mathbf{u} = \sum_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{x}} \hat{\mathbf{u}}_{\mathbf{k}} \quad (3)$$

where $\ell = 1/k$ gives a natural definition of scale. Using this we can define the spherically averaged energy spectrum as

$$E(k) = \frac{L}{2} \sum_{k \leq |\mathbf{q}| < k+1/L} |\hat{\mathbf{u}}_{\mathbf{q}}|^2 \quad (4)$$

where the sum is over all wavenumbers \mathbf{q} that satisfy $k \leq |\mathbf{q}| < k+1/L$. Note that we have normalized $E(k)$ by L so that it has dimensions of energy per unit of wavenumber.

It was the pioneering work of last century that led to the understanding that it is the flux of energy through scales that controls the statistical properties of high Re flows. Early work of Kolmogorov [57] argued for the existence of a constant flux of energy in scale space from the large to the small dissipative scales such that in the infinite Re, Re_α limit (in our notation)

$$\epsilon_\nu = \mathcal{I}_\mathcal{E}, \quad \text{and} \quad \epsilon_\alpha = 0 \quad (5)$$

so that all of the injected energy arrives at small scales and it is dissipated by viscosity. Further assuming that this process is self-similar in scale, led to the prediction of the famous Kolmogorov energy spectrum

$$E(k) = c_K \epsilon_\nu^{2/3} k^{-5/3}, \quad k_f < k < k_\nu \quad (6)$$

for wavenumbers larger than $k_f = 1/\ell_f$ where c_K is a non-dimensional order one constant. His prediction is valid up to what is now known as the Kolmogorov wavenumber $k_\nu = 1/\ell_\nu = \epsilon_\nu^{1/4}/\nu^{3/4} \simeq k_f Re^{-3/4}$. For wavenumbers smaller than k_f since there is no flux of energy these scales are expected to reach a *thermal equilibrium* state with an equipartition of energy among modes leading to

$$E(k) = c_T \mathcal{I}_\mathcal{E}^{2/3} \ell_f^{11/3} k^2, \quad k < k_f \quad (7)$$

where c_T is an other non-dimensional constant [24, 7, 49].

Later research showed that the forward cascade is not self-similar and the distributions of velocity differences develop stronger tails as smaller scales are examined. As a result there is a small correction to the exponent of the energy spectrum. This phenomenon referred to as *intermittency* has been the study of numerous studies in order to quantify and understand these corrections, that however is not going to be discussed here.

2.2 Two dimensional turbulence

Despite the fact that two dimensional flows obey the same equation as three dimensional flows they do not display the same dynamical properties. In two dimensions the Navier-Stokes equation 1 can be written in terms of the out-of-plane vorticity as

$$\partial_t w + \mathbf{u} \cdot \nabla w = \nu \nabla^2 w - \alpha w + f_w \quad (8)$$

where $w = \mathbf{e}_z \cdot \mathbf{w}$ (where \mathbf{e}_z is the unit vector along z taken to be the out-of-plane direction), $f_w = \mathbf{e}_z \cdot \nabla \times \mathbf{f}$ and we have added the drag force $-\alpha w$ often met in two dimensional systems due to bottom friction. There are two quadratic invariants in two dimensions the energy $\mathcal{E} = \frac{1}{2} \langle |\mathbf{u}|^2 \rangle$ and the enstrophy $\Omega = \frac{1}{2} \langle |\mathbf{w}|^2 \rangle$. Their balance relations read

$$\mathcal{I}_\mathcal{E} = \epsilon_\nu + \epsilon_\alpha, \quad \mathcal{I}_\Omega = \eta_\nu + \eta_\alpha \quad (9)$$

where $\mathcal{I}_\mathcal{E}$ is again the energy injection rate, $\mathcal{I}_\Omega = \langle w f_w \rangle \simeq k_f^2 \mathcal{I}_\mathcal{E}$ is the enstrophy injection rate. The energy dissipation terms at small and large scales are given by $\epsilon_\nu = \nu \langle |\mathbf{w}|^2 \rangle$ and $\epsilon_\alpha = \alpha \langle |\mathbf{u}|^2 \rangle$ respectively and the enstrophy dissipation rates by $\eta_\nu = \nu \langle |\nabla \mathbf{w}|^2 \rangle$ and $\eta_\alpha = \alpha \langle |\mathbf{w}|^2 \rangle$.

It was first realized by Onsager [79] using a point vortex model that in two-dimensions negative ‘‘temperature’’ states can exist where the flow self-organizes to generate large scale structures. Later on, the work of Kraichnan, Leith and Batchelor [59, 60, 65, 12], led to what is known as the KLB dual cascade picture of two dimensional turbulence. Their work argued that a constant forward flux of energy is incompatible with a constant flux of enstrophy. In spectral space the energy spectrum $E(k)$ is related to the enstrophy spectrum $E_\Omega(k) = k^2 E(k)$. This implies that a constant forward energy flux would imply an ever larger flux of enstrophy that is inconsistent with the enstrophy conservation. As a result, the only possibility is that enstrophy cascades *forward* towards the small scales while energy cascades *inversely* in the large scales. Therefore at infinite domains and in the infinite Re, Re_α limit

$$\epsilon_\alpha \simeq \mathcal{I}_\mathcal{E}, \quad \epsilon_\nu \simeq 0 \quad \text{and} \quad \eta_\alpha \simeq 0, \quad \eta_\nu \simeq \mathcal{I}_\Omega. \quad (10)$$

The presence of this dual cascade picture has been verified in numerical simulations [17, 66, 67] and experiments [20, 119, 55]. Same arguments as in the three dimensional cascade lead to the prediction of an $E(k) \propto k^{-5/3}$




Figure 1: Visualisation of vorticity for the turbulent state $L \gg L_\alpha$ (left) and for the condensate state $L_\alpha \gg L$ (right). In both cases the forcing scale ℓ_f was 80 times smaller than the domain size L .

energy spectrum for the range of wavenumbers that an inverse cascade of energy is present $k_\alpha \ll k \ll k_f$ while the steeper spectrum $E(k) \propto k^{-3}$ is predicted for the range of wavenumbers $k_f \ll k \ll k_\nu$ that a forward enstrophy cascade is observed. Here $k_\nu = \eta_\nu^{1/6}/\nu^{1/2} \simeq k_f Re^{1/2}$ while $k_\alpha = \epsilon_\alpha^{1/2}/\alpha^{3/2} = k_f Re_\alpha^{-3/2}$. Note that the viscous wavenumber k_ν has a different scaling in 2D than 3D. We have also introduced a new cut-off wavenumber k_α (and lengthscale $L_\alpha = 1/k_\alpha$) such that wavenumbers with $k < k_\alpha$ are severely damped by the drag force. The inverse cascade of energy is shown not to be intermittent [14] unless energy is injected in a fractal set [103].

Unlike the the small scale viscous cut off $\ell_\nu = 1/k_\nu$ that can become arbitrarily small as $Re \rightarrow \infty$ the large scale cut of $L_\alpha = 1/k_\alpha$ is limited by the domain size L . Thus while for $L_\alpha \ll L$ the aforementioned inverse cascade and power laws will be present, when $L \ll L_\alpha$ the inverse cascade will reach the domain size scale before the drag coefficient α becomes effective. In the latter case energy that arrives at the large scales will pile-up at the domain size in what is known as a spectral *condensate* [32]. In square periodic domains this condensate takes the form of two large counter rotating vortexes with large enough amplitude for the dissipation to balance the energy arriving from the small scales. This leads to the estimate for the energy of the condensate to be

$$\mathcal{E} \simeq \frac{\mathcal{I}_\mathcal{E}}{\alpha + \nu L^2}. \quad (11)$$

This energy can be very large in particular in the $\alpha = 0$ case that we will examine later on. The cascade picture described before is then altered and steeper power-laws are observed. The amplitude of the velocity fluctuations is so large that it brings the system to a quasi-equilibrium state that merits various equilibrium statistical approaches [79, 61, 92, 78, 18, 64, 40, 112]. Figure 1 shows two vorticity visualisations of two-dimensional flows such that on the left $L_\alpha < L$ while on the right $L_\alpha \gg L$. The condensate state has been realized in various experiments [20, 38]. The statistical properties of the inverse cascade case $L_\alpha \ll L$ and the condensate case $L_\alpha \gg L$ are so different that we are going to treat them separately for the quasi-2D case.

3 Thin Layer turbulence

Turbulent flows confined in thin layers is perhaps the simplest and most intuitive flows that can display split energy cascades. It is thus an optimal choice for the study of quasi-two-dimensional flows. Furthermore, such anisotropic domains are very relevant for the atmosphere whose horizontal direction is of the order of 1000 km, while the pressure scale height of the order of 10 km. As such they have been extensively studied in the literature.

Here we will discuss how the cascade properties of thin layer turbulence change as the layer height is varied in the limit of large Re, Re_α . We will distinguish between two cases: (i) cases that the drag is efficient at absorbing the large scale energy that arrives in the presence of an inverse cascade so that $L_\alpha \ll L$ and (ii) cases that the a large scale drag is very weak or absent so that a condensate is formed so that $L_\alpha \gg L$. In the first case we will quantify the forward and inverse flux of energy using ϵ_α and ϵ_ν (measured in units of $\mathcal{I}_\mathcal{E}$) that express the fraction of energy that is transferred in the large and the small scales respectively. In the infinite Re, Re_α limit $\epsilon_\alpha = 0, \epsilon_\nu = 1$ for three dimensional turbulence while $\epsilon_\alpha = 1, \epsilon_\nu = 0$ for two dimensional turbulence.

In the condensate case we will typically consider $\alpha = 0$ so that $\epsilon_\alpha, \epsilon_\nu$ are not suitable to quantify the state of the flow. Instead we will use the energy of the largest scale modes \mathcal{E}_C defined as

$$\mathcal{E}_C = \frac{1}{2} \sum_{|\mathbf{k}| < c/L} |\hat{\mathbf{u}}_{\mathbf{k}}|^2 \quad (12)$$

with $c > 2\pi$ an order one number. Finally we will also consider the energy of 3D fluctuations as

$$\mathcal{E}_{3D} = \frac{1}{2} \langle |\mathbf{u} - \bar{\mathbf{u}}|^2 \rangle. \quad (13)$$

In what follows we describe the behavior of $\epsilon_\alpha, \epsilon_\nu, \mathcal{E}_C, \mathcal{E}_{3D}$ as the height of the layer is varied for the turbulent and the condensate case.

3.1 From 3D turbulence to 2D turbulence

First, the case $L_\alpha \ll L$ (that no condensate forms) is considered. The first work that made the remark that flows in thin layers can cascade energy to both large and small scales was [102]. Since then a series of more systematic works followed that measured the fraction of energy that cascaded to that large scales as the height was varied [23, 75, 13, 109, 87] whose results will be described in this section.

We begin the presentation of thin layer turbulence by considering first a layer of height H much larger than the forcing scale and then gradually decreasing it. For $H \gg \ell_f$ the flow displays three dimensional turbulence with only forward cascade observed at scales ℓ smaller than ℓ_f forming a $k^{-5/3}$ energy spectrum. At scales larger than ℓ_f energy is expected to reach a thermal equilibrium state with equipartition of energy among modes. An argument can be made that if one considers horizontal scales $\ell_\perp \gg H \gg \ell_f$ where the flow is constrained to move primarily in two dimensions an inverse cascade can build up. However the primary interactions at such scales are not with same scale two dimensional eddies but rather directly with the forcing scale modes that are more energetic and act as a turbulent diffusion. Thus even at these scales the cascade will be strictly forward. This is a conjecture however that needs to be verified.

The strictly forward cascade behavior changes as smaller layer heights are considered. Eventually a critical height H_{3D} is reached such that for heights $H < H_{3D}$ a new phase of turbulence appears that a bidirectional cascade is present. Numerical simulations indicate that the fraction of the energy that cascades inversely ϵ_α is gradually increasing from zero as a power-law [13, 109]

$$\epsilon_\alpha \propto (H_{3D} - H)^{\beta_1} \quad (14)$$

where β_1 is measured to be close to unity but its precise value has yet to be determined. The presence of this critical point is not trivial nor fully understood. The only evidence we have are from numerical simulations [13, 109] that however can be questioned because unavoidably they suffer from limited resolution. Further investigations would be required both numerical and theoretical to investigate this point and conclude on its presence and on the universality class of this transition.

As H is decreased further than H_{3D} the fraction of energy that cascades to the small scales is decreased. For $H \ll H_{3D}$ all scales $\ell > H$ have a two dimensional behavior with an inverse cascade of energy and a forward cascade of enstrophy. This however does not imply that no energy arrives at scales close to the height $\ell \sim H$. Along with the enstrophy cascade some energy has to be transported to the small scales as well, so at scale H the flux of energy is $\epsilon_\nu \propto \eta_\nu H^2$ [15]. This energy is then transported to even smaller scales $\ell < H$ by three dimensional interactions. Note that at the limit $H \rightarrow 0$ the fraction of energy transported to the small scales also goes to zero $\epsilon_\nu \rightarrow 0$ and 2D behavior is recovered. Given that $\eta_\nu \sim \mathcal{I}_\Omega \simeq \mathcal{I}_\mathcal{E} / \ell_f^2$ implies that

$$\epsilon_\nu \propto H^2 \quad (15)$$

This behavior was first predicted in [15] and verified using a shell model. This scaling however has never been tested using numerical simulations and it is something future research needs to confirm.

The scaling in 15 does not continue for arbitrarily small H . If $H \sim \ell_\nu$ then a new transition is observed towards a third phase of turbulence where all three dimensional perturbations are damped out. In particular in the case that the forcing is two-dimensional, the transition to exactly 2D behavior, occurs in a critical way: there is a second critical height H_{2D} such that for all $H < H_{2D}$ the flow becomes exactly two dimensional. A very interesting dynamical behavior is observed for layer heights slightly larger than H_{2D} . If we denote as \mathcal{E}_{3D} the energy contained in 3D modes alone then it is measured that close to this new critical height \mathcal{E}_{3D} scales like

$$\mathcal{E}_{3D} \propto (H - H_{2D})^{\beta_2} \quad (16)$$

where β_2 is an exponent larger than one [13, 8]. This exponent is linked to the fact that as H_{2D} is approached from above 3D fluctuations grow or decay with growth that is randomly distributed in space and time. As a result not only the amplitude of 3D perturbations depends on $H - H_{2D}$ but also the fraction of area that is occupied

by them. In [8] this behavior was linked to the universality class observed in the presence of multiplicative noise in extended systems [108, 48] that is related to the KPZ equation [53]. This leads to the prediction that $\beta_2 \simeq 1.7 \dots$ compatible with present data but further work is needed to draw firm conclusions.

Finally for $H \ll \ell_\nu$ all three dimensional fluctuations are severely damped and the flow can be proven to become exactly two dimensional [44, 42] and thus all properties mentioned in section 2.2 are recovered. A sum-up of all the different phases of thin layer turbulence and their transitions in the $L_\alpha \ll L$ case is sketched in fig 2 and reported in table 1.




Figure 2: A sketch of the qualitative behavior of ϵ_α (top panel, blue line), ϵ_ν (top panel red line) and \mathcal{E}_{3D} (bottom panel red line) as a function of the layer height H .

3.2 From 3D turbulence to a 2D condensate

In finite domains when the drag coefficient is very small (or even zero) even a weak inverse cascade will lead to the formation of a large scale condensate at long times. The study of condensates is costly with numerical simulations due to the very long times that are involved. For this reason such studies always come after studies without the condensate. Nonetheless various numerical studies of thin layer turbulence exist in the literature [109, 76]. On the contrary to the numerical simulations, laboratory experiments do not have such time limitations and various studies of thin layer flows have been investigated in the literature although with limited separation between the forcing scale and the domain height [123, 124, 38, 99].

As in the previous section we begin with a layer that has a height H much larger than the forcing scale $H \gg \ell_f$ so that no inverse cascade and no condensate is formed, and gradually reduce this height. As H is reduced beyond the critical height H_{3D} (discussed in the previous section) a weak inverse cascade starts to build up leading to the formation of a condensate of energy \mathcal{E}_C . If the drag coefficient α is finite the increase of \mathcal{E}_C is gradual. For $\alpha = 0$ however even a small inverse cascade can lead to a large value of \mathcal{E}_C . In fact studies very close to H_{3D} showed that the transition to the condensate state is discontinuous [109]. In more detail it was shown that for H larger than H_{3D} the large scale energy remained small $\mathcal{E}_C = \mathcal{O}(1/L^2)$, but as soon as it became slightly smaller than H_{3D} a condensate formed and \mathcal{E}_C jumped to a finite value $\mathcal{E}_C = \mathcal{O}(1)$. Furthermore, it was shown in [109] that if H was gradually increased again to values larger than H_{3D} the condensate state remained with $\mathcal{E}_C = \mathcal{O}(1)$ up to some second critical value $H'_{3D} > H_{3D}$. Thus for values of H in the range $H_{3D} < H < H'_{3D}$ two steady states (two different attractors) exist for the same value of H and a hysteresis diagram was constructed [109] as shown in fig.3.

Further investigations of this system [115] revealed that if run for a long time there are random jumps from one attractor to the other. The time distribution of these random jumps follows an exponential distribution indicating a memory-less process [115]. It was also shown that as the Reynolds number and the domain size are increased this range of H where both attractors are stable increases [29]. This bistability is thus a behavior that is expected to survive in the large Re limit.

As H is decreased significantly from H_{3D} so that $\ell_f \gg H$ the condensate becomes stable and its amplitude depends on the principal mechanism that saturates the inverse cascade. If the Reynolds number is moderate, viscous dissipation will provide the main saturation mechanism and $\mathcal{E}_C \propto \mathcal{I}_\mathcal{E} L^2 / \nu$ [109]. For large Reynolds numbers however a new mechanism for saturation is present that originates from an eddy-viscosity effect due




Figure 3: A sketch of the qualitative behavior of \mathcal{E}_C (blue line), and \mathcal{E}_{3D} (red line) as a function of the layer height H in the condensate state.

to the small 3D eddies at scales $\ell < H$. These 3D eddies extract energy from the condensate scales. A flux loop mechanism thus is present in which energy injected at ℓ_f moves up scale to the condensate scale L through 2D motions and then back to smaller scales $\ell < H$ through interactions with the forcing scale 3D eddies. The condensate energy then in this case is not inversely proportional to viscosity but rather reaches a viscosity independent scaling $\mathcal{E}_C \propto (\mathcal{I}_\mathcal{E} \ell_f)^{2/3}$ [76, 109]. A simple model that captures this behavior was proposed in [109].

At even smaller layer heights, such that $H \sim \ell_\nu$, a new transition appears again towards a state where 3D perturbations are damped as in the previous section. The case of the condensate however is significantly different from the turbulent case. While in the former turbulence was unstructured and uniform in space allowing 3D perturbations to grow anywhere in the domain, in the condensate state the flow is still chaotic but self-organized in coherent structures of high concentration of vorticity and strain in small regions of space. In [98, 68] the evolution of infinitesimal 3D perturbations in a two dimensional flow in the condensate state were followed. This study revealed that \mathcal{E}_{3D} in this limit followed a random behavior with long periods of decay and very short periods of very large exponential increase. The periods of increase were shown to appear when the extreme of vorticity or the strain of a flow crossed a certain threshold. The statistics of these extremes were studied in detail recently in [94]. Using a point vortex model that was coupled to point like 3D perturbations in [113] showed that the instantaneous growth-rate of the energy of the 3D perturbations displayed power-law distributions that were linked to the power-law distribution of strain in space. As such the logarithm of the energy of the perturbations followed a Lévy random walk that explains the sudden jumps in the growth of energy observed in [98]. This lead to a new type of intermittency the Lévy-On-Off intermittency that was described and studied in detail in [114, 116]. This also leads to new power-law behavior for the 3D energy

$$\mathcal{E}_{3D} \propto (H - H_{2D}^*)^{\beta_3} \quad (17)$$

where H_{2D}^* is the critical value of H below which all 3D perturbation are damped and β_3 is an other exponent that depends on the Lévy noise parameters (see [114, 116]). This behavior suggested by the model has yet to be confirmed by direct numerical simulations.

Finally, for $H \ll \ell_\nu$ all 3D perturbations are dumped and the systems recovers its exact two dimensional behavior. The different stages of thin layer condensates are shown in figure 3 and summed up in table 1 along with the turbulent case.

4 Other examples of Quasi-2D flows

Similar transitions to the ones observed in thin layer turbulence are expected to be found in other quasi-two-dimensional flows. However due to the increased numerical cost or the technical difficulties in constructing laboratory experiments has limited their study. In what follows we mention a few of these systems focusing on the additional phenomena that are present.

$H&L$	Turbulent cascade $L_\alpha \ll L$	Condensate $L_\alpha \gg L$
$H > \ell_f$	3D Forward Cascade of Energy $\epsilon_\alpha = 0, \epsilon_\nu = 1$	No condensate $\mathcal{E}_C \propto L^{-2}$
$H = H_{3D} \sim \ell_f$	Criticality $\epsilon_\alpha \propto (H_{3D} - H)^{\beta_1}$ $\beta_1 \simeq 1.$	Discontinuous transition hysteresis and “rare event” jumps
$\ell_f \gg H \gg \ell_\nu$	2D and 3D Bidirectional cascade $\epsilon_\nu \propto H^2$	Flux-loop condensate $\mathcal{E}_C \propto 1$
$H \sim \ell_\nu$	Extended Multiplicative noise $\mathcal{E}_{3D} \propto (H - H_{2D})^{\beta_2}$ $\beta_2 \sim 1.7$	Lévy flight on-off Intermittency $\mathcal{E}_{3D} \propto (H - H_{2D}^*)^{\beta_3}$
$H \ll \ell_\nu$	2D Inverse energy cascade $\epsilon_\alpha = 1, \epsilon_\nu = 0$	2D condensate $\mathcal{E}_C \propto L^2/\nu$

Table 1: Table with the different phases and critical behaviors observed in thin layer turbulence. The left column gives the range of H , the middle gives the characteristics of the flow for the turbulent case and the right column gives the characteristics of the flow for the condensate state.

4.1 Rotating turbulence

Rotating turbulence is perhaps the simplest system after thin layers that displays quasi-2D behavior and split cascades. It deals with turbulence in a rotating reference frame of rotation Ω quantified here by the Rossby number $Ro = \mathcal{I}_E^{1/3}/(\Omega\ell_f)^{2/3}$. The transition from 3D turbulence to a 2D inverse cascade has been investigated in [30] and [82]. The critical layer height H_{3D} was shown to increase with the rotation rate from the weak rotation value $H_{3D} \propto \ell_f$ to a value that was shown to increase as

$$H_{3D} \propto \ell_f/Ro. \quad (18)$$

This scaling was confirmed in [110] using an asymptotic model of the rotating Navier-Stokes equation for fast-rotating turbulence within highly elongated domains.

Although, rotating turbulence appeared to have the same phenomenological description as in thin layers for the 3D to 2D transition in the large Ro and small Ro limit, at intermediate values of Ro a new state of turbulence was discovered in [31] at which the flow formed a *crystal* of co-rotating vortexes at scales larger than the forcing scale. This state of turbulence occurs close to the H_{3D} transition height at which the co-rotating 2D vortexes are stable while counter-rotating 2D vortexes decay due to 3D fluctuations. The remaining co-rotating vortexes form crystals as have been seen in point vortex models [9] and experiments on magnetized electron columns [37]. This new state of turbulence is supported by a flux loop mechanism and was shown to be metastable that reduced to an inverse cascade if a strong perturbation is applied. Such vortex crystals have been observed in the North pole of Jupiter by the Juno spacecraft mission [2] and have been interpreted by quasi-geostrophic dynamics in a curved domain [101]. The results of [31] could provide an alternative explanation. We note that in both models a segregation between co-rotating and counter rotating vortexes has lead to the formation of the crystal.

Condensates in rotating turbulence have been studied in [96, 5, 128] where a discontinuous transition and hysteresis were also found close to H_{3D} , however the significant larger complexity of rotating turbulence did not allow for a thorough investigation. However, recently new experimental platforms have been build that have been able to quantify the formation of large scales in rotating turbulence and disentangle the forward and inverse transfers as well as the presence of wave turbulence [63, 22, 43, 69, 127, 58, 19, 73]. These experiments are very suitable in studying the long term behavior of rotating turbulence close to critical points and future work is expected to verify numerical observations as well as reveal new interesting physics.

4.2 Stratified turbulence

By stratified we refer to turbulence in the presence of gravity $-\mathbf{e}_z g$ and a mean stable density gradient $S = -\rho_0^{-1} d\rho/dz$. Unlike rotation, stratification works against the two dimensionalization of turbulence by leading to the formation of strong vertical gradients, despite the fact that it suppresses vertical motions it. In [105] it was

shown that the critical height H_{3D} that the inverse cascade appears is decreased with stratification as

$$H_{3D} \propto \ell_f Fr, \tag{19}$$

where $Fr = \mathcal{I}_\varepsilon^{1/3} / (gS)^{1/2} \ell_f^{2/3}$ is the Froude number. As a result in strongly stratified flows deviations from two dimensional turbulence can appear at much thinner layers than in the absence of stratification.

Little or no work has been done for the other critical points in strongly stratified turbulence or in the presence of a condensate.

4.3 Rotating and stratified turbulence

Rotating and stratified turbulence provides perhaps the simplest model of a dry atmosphere and various studies have been devoted to its cascade properties directly from the Navier-Stokes [11, 71, 88, 51] or from reduced models [126]. It combines both the effect of two-dimensionalization of rotation and the suppression of vertical motions of stratification. The resulting complexity however is larger than the sum of its parts. In [111] using an asymptotic model for strong rotation and long boxes it was shown that in this limit the parameter space split in three different phases, one with no inverse cascade, one rotating dominated regime with an inverse cascade due to a two-dimensionalization and a third strongly stratified regime with an inverse cascade corresponding to quasi-geostrophic dynamics. It seems thus the phase space of rotating and stratified turbulence is far more complex than anticipated and a careful exploration of it and its asymptotic limits is needed.

4.4 Convection

Although convection involves inherently 3D motions in the presence of strong rotation or vertical confinement it can lead to quasi-2D behavior. There have been numerous reports in the last years of an inverse cascade and the formation of coherent large scale quasi-2D vortices in rotating Rayleigh-Benard convection that coexists with 3D eddies that extract energy from the unstable stratification [50, 35, 70]. The behavior of the large scale vortices form resemble the ones observed in thin layer turbulence and rotating turbulence displaying a discontinuous transitions and a hysteresis [34, 28].

Finally, inverse cascade has been reported in [118] for very horizontally extended domains even in the absence of rotation.

4.5 Magnetohydrodynamic turbulence

Like strong rotation for ordinary fluids a strong uniform magnetic field is also capable of bi-dimensionalizing electrically conducting fluids. Within the Magneto-Hydro-Dynamic (MHD) approximation two particular limits are of interest. First, the high conductivity limit (corresponding to the large magnetic Reynolds number limit) relevant to astrophysical plasmas and tokamak devices. The transition to a two-dimensional behavior in this limit has been investigated in [4, 106] see also the review of [80]. Second, for low magnetic Reynolds numbers (low conductivity) the magnetic field acts as a damping mechanism for all velocity fluctuations varying along the direction of the magnetic field. This so called quasi-static MHD with relation to the transition to 2D flow has been studied in [33, 90, 117]. This limit is suitable for liquid metals and has been investigated in some innovating experiments [85, 86, 124, 125, 45].

Just like in the previous examples in MHD transitions from a three dimensional forward cascade to a bi-directional cascade and two dimensional cascade are observed. However the presence of additional invariants both in 3D and 2D make the phase space of MHD turbulence sufficiently more complex such that even in pure 2D MHD such transitions can be observed [97, 95].

4.6 Quantum turbulence

Quantum turbulence refers to turbulent flow of quantum fluids, where vortex are quantized. There are various models used to describe such flows that is beyond the purpose of this work to describe. We will however mention the experiments in [93, 52, 47] where quasi-two dimensionalization in super-fluids is observed and the formation of large scale vortices. Numerically such bi-dimensionalization has been observed in thin superfluid layers [74] using the Gross-Pitaevskii equation and in the two fluid description of superfluid turbulence in the presence of a counter-flow [83].

5 Conclusions

If something is kept from this review, it should be the plethora of new dynamical phenomena that appear in quasi-2D turbulence when it is pushed to the right limits. To begin with, even in the simplest case of a thin layer flow, different phases of turbulence are observed in the infinite Re limit. Unlike the commonly accepted

expectations for homogeneous and isotropic turbulence for which a universal behavior is expected, in the presence of confinement and anisotropy distinct phases of turbulence are present such that in one phase there is large scale energy transfer and self-organization while in the other disorder and efficient energy dissipation. These phase are separated by critical points that display continuous or discontinuous phase transitions. Near these critical points, novel dynamical behaviors are observed that include the appearance of hysteresis diagrams and new critical exponents that are summed up in the table 1. As the complexity of the system is increased further new phases are discovered like the vortex-crystal meta-stable phase observed in rotating turbulence.

Recent research has only scratched the surface of these new phenomena and a lot of further work is required. In particular, investigating the behavior of the flows close to criticality and classifying them in universality classes is a much needed direction for the field. Theoretical, numerical and experimental investigations need to proceed in parallel in this direction in order to establish a clear and quantitative understanding of these transitions.

Furthermore, expanding the system complexity including more physical effects would allow to make contact with with physical and industrial systems. Here, only idealized situations were examined with a well defined injection and dissipation scale. Reality is far more complex, with forcing mechanisms that span a wide range of scales (like convection, planetary scale baroclinic instabilities, etc) and large scale dissipation much more complex than the linear drag force assumed here. However, if we can not understand this behavior in the idealised models what hope do we have to understand the more complex physical systems. Progress in both directions and a connection between the the idealized and more physical case would be required in order to obtain accurate predicting models.

Finally I would like to note that, we live in a world that climate gradually changes and an atmosphere belongs in the wider class of quasi-two-dimensional flows. It is thus important to understand how this system responds to variations of parameters. For this reason studies of fundamental questions along the directions reviewed in this work are imperative.

A peer-reviewed version of this article can be found at *Reviews of Modern Plasma Physics volume 7, Article number: 31 (2023)* <https://doi.org/10.1007/s41614-023-00134-3>

References

- [1] Myrann Abobaker, Wei Liu, Tangui Aladjidi, Alberto Bramati, and Quentin Glorieux. Inverse energy cascade in two-dimensional quantum turbulence in a fluid of light. *arXiv preprint arXiv:2211.08441*, 2022.
- [2] Alberto Adriani, Alessandro Mura, G Orton, C Hansen, Francesca Altieri, ML Moriconi, J Rogers, G Eichstädt, T Momary, Andrew P Ingersoll, et al. Clusters of cyclones encircling jupiter’s poles. *Nature*, 555(7695):216–219, 2018.
- [3] Amit Aharon-Steinberg, Tobias Völkl, Arkady Kaplan, Arnab K Pariari, Indranil Roy, Tobias Holder, Yotam Wolf, Alexander Y Meltzer, Yuri Myasoedov, Martin E Huber, Binghai Yan, Gregory Falkovich, Leonid S. Levitov, Markus Hücker, and Eli Zeldov. Direct observation of vortices in an electron fluid. *arXiv preprint arXiv:2202.02798*, 2022.
- [4] Alexandros Alexakis. Two-dimensional behavior of three-dimensional magnetohydrodynamic flow with a strong guiding field. *Physical Review E*, 84(5):056330, 2011.
- [5] Alexandros Alexakis. Rotating taylor–green flow. *Journal of Fluid Mechanics*, 769:46–78, 2015.
- [6] Alexandros Alexakis and Luca Biferale. Cascades and transitions in turbulent flows. *Physics Reports*, 767:1–101, 2018.
- [7] Alexandros Alexakis and Marc-Etienne Brachet. On the thermal equilibrium state of large-scale flows. *Journal of Fluid Mechanics*, 872:594–625, 2019.
- [8] Alexandros Alexakis, François Pétrélis, Santiago J Benavides, and Kannabiran Seshasayanan. Symmetry breaking in a turbulent environment. *Physical Review Fluids*, 6(2):024605, 2021.
- [9] Hassan Aref and Dmitri L Vainchtein. Point vortices exhibit asymmetric equilibria. *Nature*, 392(6678):769–770, 1998.
- [10] Denis A Bandurin, Andrey V Shytov, Leonid S Levitov, Roshan Krishna Kumar, Alexey I Berdyugin, Moshe Ben Shalom, Irina V Grigorieva, Andre K Geim, and Gregory Falkovich. Fluidity onset in graphene. *Nature communications*, 9(1):1–8, 2018.
- [11] Peter Bartello. Geostrophic adjustment and inverse cascades in rotating stratified turbulence. *Journal of Atmospheric Sciences*, 52(24):4410–4428, 1995.
- [12] George K Batchelor. Computation of the energy spectrum in homogeneous two-dimensional turbulence. *The Physics of Fluids*, 12(12):II–233, 1969.

- [13] Santiago Jose Benavides and Alexandros Alexakis. Critical transitions in thin layer turbulence. *Journal of Fluid Mechanics*, 822:364–385, 2017.
- [14] Guido Boffetta, A Celani, and M Vergassola. Inverse energy cascade in two-dimensional turbulence: Deviations from gaussian behavior. *Physical Review E*, 61(1):R29, 2000.
- [15] Guido Boffetta, Filippo De Lillo, and Stefano Musacchio. Shell model for quasi-two-dimensional turbulence. *Physical Review E*, 83(6):066302, 2011.
- [16] Guido Boffetta, Robert E Ecke, et al. Two-dimensional turbulence. *Annual review of fluid mechanics*, 44(1):427–451, 2012.
- [17] Guido Boffetta and S Musacchio. Evidence for the double cascade scenario in two-dimensional turbulence. *Physical Review E*, 82(1):016307, 2010.
- [18] Freddy Bouchet and Antoine Venaille. Statistical mechanics of two-dimensional and geophysical flows. *Physics reports*, 515(5):227–295, 2012.
- [19] Maxime Brunet, Basile Gallet, and Pierre-Philippe Cortet. Shortcut to geostrophy in wave-driven rotating turbulence: the quartetic instability. *Physical Review Letters*, 124(12):124501, 2020.
- [20] D Byrne, H Xia, and M Shats. Robust inverse energy cascade and turbulence structure in three-dimensional layers of fluid. *Physics of Fluids*, 23(9):095109, 2011.
- [21] David Byrne and Jun A Zhang. Height-dependent transition from 3-d to 2-d turbulence in the hurricane boundary layer. *Geophysical research letters*, 40(7):1439–1442, 2013.
- [22] Antoine Campagne, Basile Gallet, Frédéric Moisy, and Pierre-Philippe Cortet. Direct and inverse energy cascades in a forced rotating turbulence experiment. *Physics of Fluids*, 26(12):125112, 2014.
- [23] Antonio Celani, Stefano Musacchio, and Dario Vincenzi. Turbulence in more than two and less than three dimensions. *Physical Review Letters*, 104(18):184506, 2010.
- [24] Vassilios Dallas, Stephan Fauve, and Alexandros Alexakis. Statistical equilibria of large scales in dissipative hydrodynamic turbulence. *Physical review letters*, 115(20):204501, 2015.
- [25] Sergei D Danilov and D Gurarie. Quasi-two-dimensional turbulence. *Physics-Uspekhi*, 43(9):863, 2000.
- [26] Peter A Davidson, Yukio Kaneda, Keith Moffatt, and Katapalli R Sreenivasan. *A voyage through turbulence*. Cambridge University Press, 2011.
- [27] Peter Alan Davidson. *Turbulence: an introduction for scientists and engineers*. Oxford university press, 2015.
- [28] Xander M de Wit, Andrés J Aguirre Guzmán, Herman JH Clercx, and Rudie PJ Kunnen. Discontinuous transitions towards vortex condensates in buoyancy-driven rotating turbulence. *Journal of Fluid Mechanics*, 936:A43, 2022.
- [29] Xander M de Wit, Adrian van Kan, and Alexandros Alexakis. Bistability of the large-scale dynamics in quasi-two-dimensional turbulence. *Journal of Fluid Mechanics*, 939, 2022.
- [30] Enrico Deusebio, Guido Boffetta, Erik Lindborg, and Stefano Musacchio. Dimensional transition in rotating turbulence. *Physical Review E*, 90(2):023005, 2014.
- [31] P Clark Di Leoni, Alexandros Alexakis, L Biferale, and M Buzzicotti. Phase transitions and flux-loop metastable states in rotating turbulence. *Physical Review Fluids*, 5(10):104603, 2020.
- [32] G. Falkovich. Inverse cascade and wave condensate in mesoscale atmospheric turbulence. *Physical Review Letters*, 69:3173–3176, 1992.
- [33] Benjamin Favier, Fabien S Godeferd, Claude Cambon, Alexandre Delache, and Wouter JT Bos. Quasi-static magnetohydrodynamic turbulence at high reynolds number. *Journal of Fluid Mechanics*, 681:434–461, 2011.
- [34] Benjamin Favier, Céline Guervilly, and Edgar Knobloch. Subcritical turbulent condensate in rapidly rotating rayleigh–bénard convection. *Journal of Fluid Mechanics*, 864:R1, 2019.
- [35] Benjamin Favier, Lara J Silvers, and Michael RE Proctor. Inverse cascade and symmetry breaking in rapidly rotating boussinesq convection. *Physics of Fluids*, 26(9):096605, 2014.
- [36] Tiago D Ferreira, Vicente Rocha, Duarte Silva, Ariel Guerreiro, and Nuno A Silva. Towards the experimental observation of turbulent regimes and the associated energy cascades with paraxial fluids of light. *New Journal of Physics*, 24(11):113050, 2022.
- [37] KS Fine, AC Cass, WG Flynn, and CF Driscoll. Relaxation of 2d turbulence to vortex crystals. *Physical review letters*, 75(18):3277, 1995.

- [38] Nicolas Francois, Hua Xia, Horst Punzmann, and Michael Shats. Inverse energy cascade and emergence of large coherent vortices in turbulence driven by faraday waves. *Physical Review Letters*, 110(19):194501, 2013.
- [39] Uriel Frisch and Andreĭ Nikolaevich Kolmogorov. *Turbulence: the legacy of AN Kolmogorov*. Cambridge university press, 1995.
- [40] Anna Frishman and Corentin Herbert. Turbulence statistics in a two-dimensional vortex condensate. *Physical Review Letters*, 120(20):204505, 2018.
- [41] Akihide Fujisawa. A review of zonal flow experiments. *Nuclear Fusion*, 49(1):013001, 2008.
- [42] Basile Gallet. Exact two-dimensionalization of rapidly rotating large-reynolds-number flows. *Journal of Fluid Mechanics*, 783:412–447, 2015.
- [43] Basile Gallet, Antoine Campagne, P-P Cortet, and Frédéric Moisy. Scale-dependent cyclone-anticyclone asymmetry in a forced rotating turbulence experiment. *Physics of Fluids*, 26(3):035108, 2014.
- [44] Basile Gallet and Charles R Doering. Exact two-dimensionalization of low-magnetic-reynolds-number flows subject to a strong magnetic field. *Journal of Fluid Mechanics*, 773:154–177, 2015.
- [45] Basile Gallet, Johann Herault, Claude Laroche, François Pétrélis, and Stéphan Fauve. Reversals of a large-scale field generated over a turbulent background. *Geophysical & Astrophysical Fluid Dynamics*, 106(4-5):468–492, 2012.
- [46] Guillaume Gauthier, Matthew T Reeves, Xiaoquan Yu, Ashton S Bradley, Mark A Baker, Thomas A Bell, Halina Rubinsztein-Dunlop, Matthew J Davis, and Tyler W Neely. Giant vortex clusters in a two-dimensional quantum fluid. *Science*, 364(6447):1264–1267, 2019.
- [47] Guillaume Gauthier, Matthew T Reeves, Xiaoquan Yu, Ashton S Bradley, Mark A Baker, Thomas A Bell, Halina Rubinsztein-Dunlop, Matthew J Davis, and Tyler W Neely. Giant vortex clusters in a two-dimensional quantum fluid. *Science*, 364(6447):1264–1267, 2019.
- [48] Walter Genovese, Miguel A Munoz, and José M Sancho. Nonequilibrium transitions induced by multiplicative noise. *Physical Review E*, 57(3):R2495, 1998.
- [49] Jean-Baptiste Gorce and Eric Falcon. Statistical equilibrium of large scales in three-dimensional hydrodynamic turbulence. *Physical Review Letters*, 129(5):054501, 2022.
- [50] Céline Guervilly, David W Hughes, and Chris A Jones. Large-scale vortices in rapidly rotating rayleigh–bénard convection. *Journal of Fluid Mechanics*, 758:407–435, 2014.
- [51] Corentin Herbert, Raffaele Marino, Duane Rosenberg, and Annick Pouquet. Waves and vortices in the inverse cascade regime of stratified turbulence with or without rotation. *Journal of Fluid Mechanics*, 806:165–204, 2016.
- [52] Shaun P Johnstone, Andrew J Groszek, Philip T Starkey, Christopher J Billington, Tapio P Simula, and Kristian Helmerston. Evolution of large-scale flow from turbulence in a two-dimensional superfluid. *Science*, 364(6447):1267–1271, 2019.
- [53] Mehran Kardar, Giorgio Parisi, and Yi-Cheng Zhang. Dynamic scaling of growing interfaces. *Physical Review Letters*, 56(9):889, 1986.
- [54] Hamid Kellay and Walter I Goldburg. Two-dimensional turbulence: a review of some recent experiments. *Reports on Progress in Physics*, 65(5):845, 2002.
- [55] Douglas H Kelley and Nicholas T Ouellette. Spatiotemporal persistence of spectral fluxes in two-dimensional weak turbulence. *Physics of Fluids*, 23(11):115101, 2011.
- [56] Gregory P King, Jur Vogelzang, and Ad Stoffelen. Upscale and downscale energy transfer over the tropical pacific revealed by scatterometer winds. *Journal of Geophysical Research: Oceans*, 120(1):346–361, 2015.
- [57] Andrey Nikolaevich Kolmogorov. The local structure of turbulence in incompressible viscous fluid for very large reynolds numbers. *Cr Acad. Sci. URSS*, 30:301–305, 1941.
- [58] Itamar Kolvin, Kobi Cohen, Yuval Vardi, and Eran Sharon. Energy transfer by inertial waves during the buildup of turbulence in a rotating system. *Physical Review Letters*, 102(1):014503, 2009.
- [59] Robert H Kraichnan. Inertial ranges in two-dimensional turbulence. *The Physics of Fluids*, 10(7):1417–1423, 1967.
- [60] Robert H Kraichnan. Inertial-range transfer in two-and three-dimensional turbulence. *Journal of Fluid Mechanics*, 47(3):525–535, 1971.
- [61] Robert H Kraichnan. Statistical dynamics of two-dimensional flow. *Journal of Fluid Mechanics*, 67(1):155–175, 1975.

- [62] Hüseyin Kurtuldu, Jeffrey S Guasto, Karl A Johnson, and Jerry P Gollub. Enhancement of biomixing by swimming algal cells in two-dimensional films. *Proceedings of the National Academy of Sciences*, 108(26):10391–10395, 2011.
- [63] Cyril Lamriben, Pierre-Philippe Cortet, and Frédéric Moisy. Direct measurements of anisotropic energy transfers in a rotating turbulence experiment. *Physical review letters*, 107(2):024503, 2011.
- [64] Jason Laurie, Guido Boffetta, Gregory Falkovich, Igor Kolokolov, and Vladimir Lebedev. Universal profile of the vortex condensate in two-dimensional turbulence. *Physical Review Letters*, 113(25):254503, 2014.
- [65] Cecil E Leith. Diffusion approximation for two-dimensional turbulence. *The Physics of Fluids*, 11(3):671–672, 1968.
- [66] Douglas K Lilly. Numerical simulation of two-dimensional turbulence. *The Physics of Fluids*, 12(12):II–240, 1969.
- [67] Douglas K Lilly. Numerical simulation studies of two-dimensional turbulence: I. models of statistically steady turbulence. *Geophysical Fluid Dynamics*, 3(4):289–319, 1972.
- [68] Chandra Shekhar Lohani, Suraj Kumar Nayak, and Kannabiran Seshasayanan. Effect of confinement on the transition from 2d to 3d fast rotating flows. *arXiv preprint arXiv:2305.12255*, 2023.
- [69] Nathanaël Machicoane, Frédéric Moisy, and Pierre-Philippe Cortet. Two-dimensionalization of the flow driven by a slowly rotating impeller in a rapidly rotating fluid. *Physical Review Fluids*, 1(7):073701, 2016.
- [70] Stefano Maffei, Mitchell J Krouss, Keith Julien, and Michael A Calkins. On the inverse cascade and flow speed scaling behaviour in rapidly rotating rayleigh–bénard convection. *Journal of Fluid Mechanics*, 913:A18, 2021.
- [71] Raffaele Marino, Pablo Daniel Mininni, Duane Rosenberg, and Annick Pouquet. Inverse cascades in rotating stratified turbulence: fast growth of large scales. *EPL (Europhysics Letters)*, 102(4):44006, 2013.
- [72] BK Martin, XL Wu, WI Goldberg, and MA Rutgers. Spectra of decaying turbulence in a soap film. *Physical Review Letters*, 80(18):3964, 1998.
- [73] Eduardo Monsalve, Maxime Brunet, Basile Gallet, and Pierre-Philippe Cortet. Quantitative experimental observation of weak inertial-wave turbulence. *Physical Review Letters*, 125(25):254502, 2020.
- [74] Nicolás P Müller, Marc-Etienne Brachet, Alexandros Alexakis, and Pablo D Mininni. Abrupt transition between three-dimensional and two-dimensional quantum turbulence. *Physical Review Letters*, 124(13):134501, 2020.
- [75] Stefano Musacchio and Guido Boffetta. Split energy cascade in turbulent thin fluid layers. *Physics of Fluids*, 29(11):111106, 2017.
- [76] Stefano Musacchio and Guido Boffetta. Condensate in quasi-two-dimensional turbulence. *Physical Review Fluids*, 4(2):022602, 2019.
- [77] Boris N Narozhny. Hydrodynamic approach to two-dimensional electron systems. *La Rivista del Nuovo Cimento*, pages 1–76, 2022.
- [78] A Naso, Pierre-Henri Chavanis, and Bérengere Dubrulle. Statistical mechanics of two-dimensional euler flows and minimum enstrophy states. *The European Physical Journal B*, 77(2):187–212, 2010.
- [79] Lars Onsager. Statistical hydrodynamics. *Il Nuovo Cimento (1943-1954)*, 6:279–287, 1949.
- [80] Sean Oughton, William H Matthaeus, and Pablo Dmitruk. Reduced mhd in astrophysical applications: Two-dimensional or three-dimensional? *The Astrophysical Journal*, 839(1):2, 2017.
- [81] Rahul Pandit, Debarghya Banerjee, Akshay Bhatnagar, Marc Brachet, Anupam Gupta, Dhruvadya Mitra, Nairita Pal, Prasad Perlekar, Samriddhi Sankar Ray, Vishwanath Shukla, et al. An overview of the statistical properties of two-dimensional turbulence in fluids with particles, conducting fluids, fluids with polymer additives, binary-fluid mixtures, and superfluids. *Physics of fluids*, 29(11):111112, 2017.
- [82] Tiago Pestana and Stefan Hickel. Regime transition in the energy cascade of rotating turbulence. *Physical Review E*, 99(5):053103, 2019.
- [83] Juan Ignacio Polanco and Giorgio Krstulovic. Counterflow-induced inverse energy cascade in three-dimensional superfluid turbulence. *Physical Review Letters*, 125(25):254504, 2020.
- [84] Stephen B Pope and Stephen B Pope. *Turbulent flows*. Cambridge university press, 2000.
- [85] Alban Pothérat and Rico Klein. Why, how and when mhd turbulence at low rm becomes three-dimensional. *Journal of Fluid Mechanics*, 761:168–205, 2014.
- [86] Alban Pothérat, Joël Sommeria, and René Moreau. An effective two-dimensional model for mhd flows with transverse magnetic field. *Journal of Fluid Mechanics*, 424:75–100, 2000.

- [87] Basile Pujol, Adrian van Kan, and Alexandros Alexakis. Role of the forcing dimensionality in thin-layer turbulent energy cascades. *Physical Review Fluids*, 5(6):064610, 2020.
- [88] Annick Pouquet, Raffaele Marino, Pablo Daniel Mininni, and Duane Rosenberg. Dual constant-flux energy cascades to both large scales and small scales. *Physics of Fluids*, 29(11):111108, 2017.
- [89] Annick Pouquet and Nobumitsu Yokoi. Helical fluid and (hall)-mhd turbulence: a brief review. *Philosophical Transactions of the Royal Society A*, 380(2219):20210087, 2022.
- [90] K Sandeep Reddy and Mahendra K Verma. Strong anisotropy in quasi-static magnetohydrodynamic turbulence for high interaction parameters. *Physics of Fluids*, 26(2):025109, 2014.
- [91] Michael Rivera, Peter Vorobieff, and Robert E Ecke. Turbulence in flowing soap films: velocity, vorticity, and thickness fields. *Physical Review Letters*, 81(7):1417, 1998.
- [92] Raoul Robert and Joel Sommeria. Statistical equilibrium states for two-dimensional flows. *Journal of Fluid Mechanics*, 229:291–310, 1991.
- [93] Sang Won Seo, Bumsuk Ko, Joon Hyun Kim, and Yong-il Shin. Observation of vortex-antivortex pairing in decaying 2d turbulence of a superfluid gas. *Scientific reports*, 7(1):1–8, 2017.
- [94] Kannabiran Seshasayanan. Spatial extreme values of vorticity and velocity gradients in two-dimensional turbulent flows. *arXiv preprint arXiv:2301.09900*, 2023.
- [95] Kannabiran Seshasayanan and Alexandros Alexakis. Critical behavior in the inverse to forward energy transition in two-dimensional magnetohydrodynamic flow. *Physical Review E*, 93(1):013104, 2016.
- [96] Kannabiran Seshasayanan and Alexandros Alexakis. Condensates in rotating turbulent flows. *Journal of Fluid Mechanics*, 841:434–462, 2018.
- [97] Kannabiran Seshasayanan, Santiago Jose Benavides, and Alexandros Alexakis. On the edge of an inverse cascade. *Physical Review E*, 90(5):051003, 2014.
- [98] Kannabiran Seshasayanan and Basile Gallet. Onset of three-dimensionality in rapidly rotating turbulent flows. *Journal of Fluid Mechanics*, 901, 2020.
- [99] Michael Shats, David Byrne, and Hua Xia. Turbulence decay rate as a measure of flow dimensionality. *Physical Review Letters*, 105(26):264501, 2010.
- [100] Lia Siegelman, Patrice Klein, Andrew P Ingersoll, Shawn P Ewald, William R Young, Annalisa Bracco, Alessandro Mura, Alberto Adriani, Davide Grassi, Christina Plainaki, et al. Moist convection drives an upscale energy transfer at jovian high latitudes. *Nature Physics*, 18(3):357–361, 2022.
- [101] Lia Siegelman, William R Young, and Andrew P Ingersoll. Polar vortex crystals: Emergence and structure. *Proceedings of the National Academy of Sciences*, 119(17):e2120486119, 2022.
- [102] Leslie M Smith, Jeffrey R Chasnov, and Fabian Waleffe. Crossover from two-to three-dimensional turbulence. *Physical Review Letters*, 77(12):2467, 1996.
- [103] George Sofiadis, Ioannis E Sarris, and Alexandros Alexakis. Inducing intermittency in the inverse cascade of two-dimensional turbulence by a fractal forcing. *Physical Review Fluids*, 8(2):024607, 2023.
- [104] Andrey Sokolov, Igor S Aranson, John O Kessler, and Raymond E Goldstein. Concentration dependence of the collective dynamics of swimming bacteria. *Physical Review Letters*, 98(15):158102, 2007.
- [105] Alessandro Sozza, Guido Boffetta, P Muratore-Ginanneschi, and Stefano Musacchio. Dimensional transition of energy cascades in stably stratified forced thin fluid layers. *Physics of Fluids*, 27(3):035112, 2015.
- [106] Nicolás Eduardo Sujovolsky and Pablo Daniel Mininni. Tridimensional to bidimensional transition in magnetohydrodynamic turbulence with a guide field and kinetic helicity injection. *Physical Review Fluids*, 1(5):054407, 2016.
- [107] Patrick Tabeling. Two-dimensional turbulence: a physicist approach. *Physics reports*, 362(1):1–62, 2002.
- [108] Yuhai Tu, G Grinstein, and MA Munoz. Systems with multiplicative noise: critical behavior from kpz equation and numerics. *Physical Review Letters*, 78(2):274, 1997.
- [109] Adrian van Kan and Alexandros Alexakis. Condensates in thin-layer turbulence. *Journal of Fluid Mechanics*, 864:490–518, 2019.
- [110] Adrian van Kan and Alexandros Alexakis. Critical transition in fast-rotating turbulence within highly elongated domains. *Journal of Fluid Mechanics*, 899:A33, 2020.
- [111] Adrian van Kan and Alexandros Alexakis. Energy cascades in rapidly rotating and stratified turbulence within elongated domains. *Journal of Fluid Mechanics*, 933:A11, 2022.

- [112] Adrian van Kan, Alexandros Alexakis, and Marc Brachet. Geometric microcanonical theory of two-dimensional truncated euler flows. *Philosophical Transactions of the Royal Society A*, 380(2226):20210049, 2022.
- [113] Adrian van Kan, Alexandros Alexakis, and Marc-Etienne Brachet. Intermittency of three-dimensional perturbations in a point-vortex model. *Physical Review E*, 103(5):053102, 2021.
- [114] Adrian van Kan, Alexandros Alexakis, and Marc-Etienne Brachet. Lévy on-off intermittency. *Physical Review E*, 103(5):052115, 2021.
- [115] Adrian van Kan, Takahiro Nemoto, and Alexandros Alexakis. Rare transitions to thin-layer turbulent condensates. *Journal of Fluid Mechanics*, 878:356–369, 2019.
- [116] Adrian van Kan and François Pétrélis. $1/f$ noise and anomalous scaling in lévy noise-driven on-off intermittency. *Journal of Statistical Mechanics: Theory and Experiment*, 2023(1):013204, 2023.
- [117] Mahendra K Verma. Anisotropy in quasi-static magnetohydrodynamic turbulence. *Reports on Progress in Physics*, 80(8):087001, 2017.
- [118] Philipp P Vieweg, Janet D Scheel, Rodion Stepanov, and Jörg Schumacher. Inverse cascades of kinetic energy and thermal variance in three-dimensional horizontally extended turbulent convection. *Physical Review Research*, 4(4):043098, 2022.
- [119] Alexandra von Kameke, Fernández Huhn, Guillermo Fernández-García, Alberto P Muñozuri, and Vicente Pérez-Muñozuri. Double cascade turbulence and richardson dispersion in a horizontal fluid flow induced by faraday waves. *Physical Review Letters*, 107(7):074502, 2011.
- [120] P Vorobieff, M Rivera, and RE Ecke. Soap film flows: Statistics of two-dimensional turbulence. *Physics of Fluids*, 11(8):2167–2177, 1999.
- [121] Da Wei, Yaochen Yang, Xuefeng Wei, Ramin Golestanian, Ming Li, Fanlong Meng, and Yi Peng. Scaling transition of active turbulence from two to three dimensions. *arXiv preprint arXiv:2307.15720*, 2023.
- [122] Xiao-Lun Wu and Albert Libchaber. Particle diffusion in a quasi-two-dimensional bacterial bath. *Physical Review Letters*, 84(13):3017, 2000.
- [123] H Xia, M Shats, and Gregory Falkovich. Spectrally condensed turbulence in thin layers. *Physics of Fluids*, 21(12):125101, 2009.
- [124] Hua Xia, David Byrne, Gregory Falkovich, and Michael Shats. Upscale energy transfer in thick turbulent fluid layers. *Nature Physics*, 7(4):321–324, 2011.
- [125] Hua Xia and MG Shats. Inverse energy cascade correlated with turbulent-structure generation in toroidal plasma. *Physical Review Letters*, 91:155001, 2003.
- [126] Jin-Han Xie. Downscale transfer of quasigeostrophic energy catalyzed by near-inertial waves. *Journal of Fluid Mechanics*, 904:A40, 2020.
- [127] Ehud Yarom and Eran Sharon. Experimental observation of steady inertial wave turbulence in deep rotating flows. *Nature Physics*, 10(7):510–514, 2014.
- [128] Naoto Yokoyama and Masanori Takaoka. Hysteretic transitions between quasi-two-dimensional flow and three-dimensional flow in forced rotating turbulence. *Physical Review Fluids*, 2(9):092602, 2017.
- [129] Roland Young and Peter L Read. Forward and inverse kinetic energy cascades in jupiter’s turbulent weather layer. *Nature Physics*, 13(11):1135–1140, 2017.
- [130] Ye Zhou. Turbulence theories and statistical closure approaches. *Physics Reports*, 935:1–117, 2021.