



HAL
open science

Low regularity solutions to the logarithmic Schrodinger equation

Rémi Carles, Masayuki Hayashi, Tohru Ozawa

► **To cite this version:**

Rémi Carles, Masayuki Hayashi, Tohru Ozawa. Low regularity solutions to the logarithmic Schrodinger equation. 2023. hal-04268646v1

HAL Id: hal-04268646

<https://hal.science/hal-04268646v1>

Preprint submitted on 2 Nov 2023 (v1), last revised 10 Nov 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

LOW REGULARITY SOLUTIONS TO THE LOGARITHMIC SCHRÖDINGER EQUATION

RÉMI CARLES, MASAYUKI HAYASHI, AND TOHRU OZAWA

ABSTRACT. We consider the logarithmic Schrödinger equation, in various geometric settings. We show that the flow map can be uniquely extended from H^1 to L^2 , and that this extension is Lipschitz continuous. Moreover, we prove the regularity of the flow map in intermediate Sobolev spaces.

1. INTRODUCTION

We consider the Cauchy problem associated to the logarithmic Schrödinger equation

$$i\partial_t u + \Delta u + \lambda u \ln(|u|^2) = 0, \quad u|_{t=0} = \varphi, \quad (1.1)$$

with $x \in \Omega$, $\lambda \in \mathbb{R}$. The standard occurrence for Ω is \mathbb{R}^d . In view of the framework considered in numerical simulations (see e.g. [3]), the domain Ω that we consider may be:

- The whole space $\Omega = \mathbb{R}^d$,
- A half space $\Omega = \mathbb{R}_+^d$, with zero Dirichlet boundary condition, $u|_{\partial\Omega} = 0$,
- A smooth domain $\Omega \subset \mathbb{R}^d$ with bounded boundary; (1.1) is then considered with zero Dirichlet boundary condition, and the Laplacian Δ is understood to be the self-adjoint realization in $H^{-1}(\Omega)$ with the domain $D(\Delta) = H_0^1(\Omega)$ (see e.g. [13, Chapter 2]),
- The torus $\Omega = \mathbb{T}^d = \mathbb{R}^d/\mathbb{Z}^d$.

In all cases, there is no restriction on the space dimension $d \geq 1$. Formally, the mass and energy are independent of time:

$$M(u(t)) = \|u(t)\|_{L^2(\Omega)}^2,$$

$$E(u(t)) = \|\nabla u(t)\|_{L^2(\Omega)}^2 - \lambda \int_{\Omega} |u(t,x)|^2 (\ln(|u(t,x)|^2) - 1) dx.$$

The equation (1.1) was introduced in [5] for quantum mechanics, and has attracted the interest of physicists from various fields ever since (see e.g. [7, 32, 20, 26, 25, 31, 33, 34]). From the mathematical point of view, the first study goes back to [12], where the Cauchy problem was investigated in the case $\Omega = \mathbb{R}^d$. The main results in [12] we mention here are:

- Theorem 1.2 b): if $\varphi \in L^2(\mathbb{R}^d)$, then (1.1) has a unique weak solution $u \in C(\mathbb{R}, L^2(\mathbb{R}^d))$ in the sense of Brezis [6].

- Theorem 2.1: if $\lambda > 0$, and $\varphi \in H^1(\mathbb{R}^d)$ is such that $|\varphi|^2 \ln(|\varphi|^2) \in L^1(\mathbb{R}^d)$, then (1.1) has a unique solution $u \in C(\mathbb{R}, H^1(\mathbb{R}^d))$. In addition, $|u(t)|^2 \ln(|u(t)|^2) \in L^1(\mathbb{R}^d)$ for all $t \in \mathbb{R}$, and the mass and energy of u are independent of time.

The goal of this paper is mostly to revisit the first statement above. The weak solution in Theorem 1.2 b) is obtained as a limit of the sequence of strong solutions, and the existence of this limit is guaranteed through the maximal monotone theory. For the convenience of the reader, more details are provided in Appendix A.

The mathematical study of (1.1) has been considered since [12], regarding both the Cauchy problem ([16, 9, 23]) and the dynamical properties of the solutions (e.g. [10, 2, 9, 18, 19]). More recently, the Cauchy problem was revisited by the second and third authors in [24], where strong global solutions are provided in a constructive way, in each of the following functional settings:

- $\varphi \in H^1(\mathbb{R}^d)$ for $\lambda \neq 0$,
- Energy space: $\varphi \in W_1 := \{f \in H^1(\mathbb{R}^d), |f|^2 \ln(|f|^2) \in L^1(\mathbb{R}^d)\}$ for $\lambda \neq 0$,
- H^2 energy space: $\varphi \in W_2 := \{f \in H^2(\mathbb{R}^d), f \ln(|f|^2) \in L^2(\mathbb{R}^d)\}$ for $\lambda > 0$.

We emphasize that unlike in the case of more standard power-like nonlinearities for Schrödinger equations, due to the singularity of the logarithm at the origin, several questions regarding the Cauchy problem (1.1) are unclear, such as:

- The propagation of higher regularity: if $\varphi \in H^3$, can we guarantee that the solution u remains in H^3 , even locally in time?
- Is there a minimal regularity for a “reasonable” notion of solution?

In this paper, we focus on the second question. When $\Omega = \mathbb{R}^d$, (1.1) is invariant under Galilean transformations

$$u(t, x) \mapsto e^{iv \cdot x - i|v|^2 t} u(t, x - 2vt) \quad \text{for } v \in \mathbb{R}^d,$$

which leave the $L^2(\mathbb{R}^d)$ norm invariant, so it may be expected that like for nonlinear Schrödinger equations with pure power nonlinearities, the flow map fails to be uniformly continuous on $H^s(\mathbb{R}^d)$ when $s < 0$ ([28, 14, 15]). See also [15, 8, 30] for stronger ill-posedness results. We note, however, that in the case of the KdV equation the flow map was extended continuously to the level of $H^{-1}(\mathbb{R})$ ([29]), even though it has been known that the flow map cannot be uniformly continuous on $H^s(\mathbb{R})$ when $s < -3/4$. Similar well-posedness pictures can also be seen in the cubic NLS equation and the modified KdV equation ([22]).

Note that there is no natural scaling associated to (1.1), when $\Omega = \mathbb{R}^d$, of the form $u_\kappa(t, x) = \kappa^\alpha u(\kappa^\beta t, \kappa^\gamma x)$, which leaves the equation invariant. Yet, another invariance, rather unique for nonlinear Schrödinger equations, shows that the size of the initial data does not affect the Cauchy problem, nor the dynamical properties of the solution: if u solves (1.1), then for any $z \in \mathbb{C}$,

$$u_z(t, x) = zu(t, x)e^{i\lambda t \ln|z|^2}$$

solves the same equation, with initial datum $z\varphi$. This unusual invariance also shows that regardless of the function spaces considered, the flow map cannot be C^1 at the origin, due to the above oscillating factor: it is at most Lipschitz continuous.

The main result of this paper is that the flow map defined on $H^1(\Omega)$ can be uniquely extended on $L^2(\Omega)$, this extension is Lipschitz continuous, and preserves possible intermediate Sobolev regularity:

Theorem 1.1. *The equation (1.1) is globally well-posed in $L^2(\Omega)$ in the following sense: The H^1 solution map Φ is uniquely extended to $L^2(\Omega)$, and for $\varphi \in L^2(\Omega)$, $u = \Phi(\varphi) \in C(\mathbb{R}, L^2(\Omega))$ satisfies*

$$i\partial_t u + \Delta u + \lambda u \ln(|u|^2) = 0 \quad \text{in } H^{-2}(\omega)$$

for any open sets $\omega \Subset \Omega$ and all $t \in \mathbb{R}$, with $u|_{t=0} = \varphi$. Moreover, Φ is Lipschitz continuous:

$$\|\Phi(\varphi)(t) - \Phi(\psi)(t)\|_{L^2} \leq e^{2|\lambda t|} \|\varphi - \psi\|_{L^2}$$

for any $\varphi, \psi \in L^2(\Omega)$ and all $t \in \mathbb{R}$. If in addition $\varphi \in H^s(\Omega)$ for some $s \in (0, 1)$, then $\Phi(\varphi) \in C(\mathbb{R}, H^s(\Omega))$.

Our contributions in this paper can be summarized as follows.

1. *Meaning of L^2 solutions revisited.* We show that logarithmic nonlinearities make sense for general functions belonging to $L^\infty((-T, T), L^2(\Omega))$ for $T > 0$, which in particular gives the meaning of solutions to (1.1) in the distribution sense (see Lemma 3.1 and Remark 3.2). This differs from [12] in that it gives the meaning to L^2 solutions regardless of how the solution is constructed or independently of limiting procedures.

2. *GWP in L^2 independently of the maximal monotone theory.* We construct L^2 solutions as an extension of the solution map on H^1 while preserving L^2 Lipschitz flow in [12] (see Lemma 3.3 below). The formulation of the global well-posedness is inspired from recent results on completely integrable systems ([29, 22, 21]). Note that the L^2 Lipschitz flow is a natural consequence of the remarkable inequality

$$\left| \operatorname{Im} \left[\overline{(z_1 - z_2)} (z_1 \log(|z_1|) - z_2 \log(|z_2|)) \right] \right| \leq |z_1 - z_2|^2 \quad \text{for all } z_1, z_2 \in \mathbb{C}, \quad (1.2)$$

which was discovered in [12, Lemme 1.1.1]. It may be common to [12, Theorem 1.2 b)] that L^2 solutions are constructed as a limit of sequences of strong solutions, but our construction would be regarded to be a more direct consequence of (1.2) and it is independent of the maximal monotone theory.

3. *Intermediate Sobolev regularity.* This regularity result is obtained as a new application of the inequality (1.2). By using the Sobolev norm based on the difference quotient, we can effectively utilize (1.2) to obtain an a priori estimates on H^s for all $s \in (0, 1)$.

The rest of the paper is organized as follows. In Section 2, we recall the construction of the H^1 solution map, from [24]. In Section 3, we show that this map

can be uniquely extended to L^2 , as a Lipschitz continuous map. In Section 4, we show that the intermediate Sobolev regularity is propagated by this map.

Notation. We sometimes use the abbreviated notation such as

$$C_T(X) = C([-T, T], X), \quad L_T^\infty(X) = L^\infty((-T, T), X)$$

for $T > 0$ and a Banach/Fréchet space X . For open sets $\omega, \Omega \subset \mathbb{R}^d$, we write $\omega \Subset \Omega$ if $\bar{\omega} \subset \Omega$ and $\bar{\omega}$ is compact, where $\bar{\omega}$ is the closure of ω in \mathbb{R}^d .

According to [17], we define the fractional Sobolev spaces H^s . For a general open set $\Omega \subset \mathbb{R}^d$, the fractional Sobolev spaces $H^s(\Omega)$ for $s \in (0, 1)$ are defined via the norm

$$\|f\|_{H^s(\Omega)}^2 = \|f\|_{L^2(\Omega)}^2 + \iint_{\Omega \times \Omega} \frac{|f(x+y) - f(x)|^2}{|y|^{d+2s}} dx dy. \quad (1.3)$$

We define $H_0^s(\Omega)$ by the closure of $C_c^\infty(\Omega)$ in the norm $\|\cdot\|_{H^s(\Omega)}$ and $H^{-s}(\Omega)$ by the dual space of $H_0^s(\Omega)$ for $s \in (0, 1)$. When $\Omega = \mathbb{R}^d$, the Sobolev space via the norm (1.3) coincides with Bessel potential spaces endowed with the norm

$$\|f\|_{H^s(\mathbb{R}^d)}^2 = \left\| (1 + 4\pi^2|\xi|^2)^s \hat{f}(\xi) \right\|_{L^2(\mathbb{R}^d)}^2 \quad \text{for } s \in \mathbb{R}, \quad (1.4)$$

where $\hat{f}(\xi)$ is the Fourier transform defined by

$$\hat{f}(\xi) = \int_{\mathbb{R}^d} f(x) e^{-2\pi i x \cdot \xi} dx \quad \text{for } \xi \in \mathbb{R}^d.$$

Similarly to Bessel potential spaces on \mathbb{R}^d , the Sobolev spaces $H^s(\mathbb{T}^d)$ on the torus are defined via the norm

$$\|f\|_{H^s(\mathbb{T}^d)}^2 = \sum_{n \in \mathbb{Z}^d} (1 + 4\pi^2|n|^2)^s |\hat{f}(n)|^2 \quad \text{for } s \in \mathbb{R}, \quad (1.5)$$

where $\hat{f}(n)$ is the discrete Fourier transform defined by

$$\hat{f}(n) = \int_{\mathbb{T}^d} f(x) e^{-2\pi i x \cdot n} dx \quad \text{for } n \in \mathbb{Z}^d.$$

We use $A \lesssim B$ to denote the inequality $A \leq CB$ for some constant $C > 0$. The dependence of C is usually clear from the context and we often omit this dependence. We may sometimes write $A \lesssim_* B$ to clarify the dependence of the implicit constant.

2. GLOBAL H^1 SOLUTIONS

In this section we review global H^1 solutions to (1.1). Here let $\Omega \subset \mathbb{R}^d$ be a general domain and let $\varphi \in H_0^1(\Omega)$. Following the strategy of [24], we regularize (1.1) like in [3]: for $\varepsilon > 0$, we consider approximate solutions u^ε to

$$i\partial_t u^\varepsilon + \Delta u^\varepsilon + 2\lambda u^\varepsilon \ln(|u^\varepsilon| + \varepsilon) = 0, \quad u_{|t=0}^\varepsilon = \varphi. \quad (2.1)$$

As the nonlinearity is now smooth, with moderate growth at infinity, the equation (2.1) has a unique, global solution (see e.g. [11, Chapter 3], [24, Section 2.1]),

$$u^\varepsilon \in C(\mathbb{R}, H_0^1(\Omega)) \cap C^1(\mathbb{R}, H^{-1}(\Omega)).$$

Uniqueness and regularity in time in the case of (2.1) follow from the generalization of the inequality (1.2), generalized successively in [3] and [24]. We state the most general version:

Lemma 2.1 ([24, Lemma A.1]). *For all $z_1, z_2 \in \mathbb{C}$, $\varepsilon_1, \varepsilon_2 \geq 0$,*

$$\begin{aligned} |\operatorname{Im}[(\overline{z_1 - z_2})(z_1 \ln(|z_1| + \varepsilon_1) - z_2 \ln(|z_2| + \varepsilon_2))]| &\leq |z_1 - z_2|^2 \\ &+ |\varepsilon_1 - \varepsilon_2| \times |z_1 - z_2|. \end{aligned}$$

The uniform energy estimate

$$\|\nabla u^\varepsilon(t)\|_{L^2}^2 \leq e^{4|\lambda t|} \|\nabla \varphi\|_{L^2}^2,$$

is easily obtained by differentiating the left hand side in time and invoking Gronwall's lemma. Combined this with Lemma 2.1, one can prove that $\{u^\varepsilon\}$ forms a Cauchy sequence in $C_T(L_{\text{loc}}^2(\Omega))$ as $\varepsilon \downarrow 0$ for any $T > 0$. Thus, we obtain the following result.

Theorem 2.2 (From Theorem 4.1 in [24]). *For any $\varphi \in H_0^1(\Omega)$, there exists a unique solution $u \in C(\mathbb{R}, H_0^1(\Omega))$ to (1.1), in the sense of*

$$i\partial_t u + \Delta u + \lambda u \ln(|u|^2) = 0 \quad \text{in } H^{-1}(\omega),$$

for all $\omega \in \Omega$ and all $t \in \mathbb{R}$, and with $u|_{t=0} = \varphi$.

The whole argument holds true for the torus \mathbb{T}^d (not considered in [24]), in the same way, and we obtain:

Theorem 2.3. *For any $\varphi \in H^1(\mathbb{T}^d)$, there exists a unique solution $u \in C(\mathbb{R}, H^1(\mathbb{T}^d))$ to (1.1), in the sense of*

$$i\partial_t u + \Delta u + \lambda u \ln(|u|^2) = 0 \quad \text{in } H^{-1}(\mathbb{T}^d),$$

for all $t \in \mathbb{R}$, and with $u|_{t=0} = \varphi$.

3. THE CAUCHY PROBLEM IN L^2

In this section we construct strong L^2 solutions to (1.1). For convenience of notation, here we only consider the case where Ω is a general domain in \mathbb{R}^d . In the case of the torus \mathbb{T}^d , the same argument still works if we replace $H_0^1(\Omega)$ by $H^1(\mathbb{T}^d)$.

The first task in order to consider (1.1) with initial data $\varphi \in L^2(\Omega)$ is to clarify in what sense the L^2 solution satisfies the equation.

Lemma 3.1. *Let $u \in L_T^\infty(L^2(\Omega))$ for $T > 0$. Then, for any small $\varepsilon > 0$ and all $\omega \in \Omega$, the nonlinear term satisfies $u \ln(|u|^2) \in L_T^\infty(H^{-\varepsilon}(\omega))$.*

Proof. We note that

$$|u \ln(|u|^2)| \lesssim |u|^{1-\delta} + |u|^{1+\delta},$$

for any $\delta \in (0, 1)$. Writing, for ψ a test function supported in $\omega \Subset \Omega$,

$$\begin{aligned} |\langle u \ln |u|^2, \psi \rangle| &\lesssim \int_{\omega} |u|^{1-\delta} |\psi| + \int_{\omega} |u|^{1+\delta} |\psi| \\ &\lesssim \|u\|_{L^{p'(1-\delta)}}^{1-\delta} \|\psi\|_{L^p} + \|u\|_{L^{q'(1+\delta)}}^{1+\delta} \|\psi\|_{L^q}, \end{aligned}$$

and considering $p'(1-\delta) = 2$, that is $p = \frac{2}{1-\delta}$, and $q = \frac{2}{1+\delta}$, we get

$$\begin{aligned} |\langle u \ln(|u|^2), \psi \rangle| &\lesssim \|u\|_{L^2}^{1-\delta} \|\psi\|_{L^{\frac{2}{1-\delta}}(\omega)} + \|u\|_{L^2}^{1+\delta} \|\psi\|_{L^{\frac{2}{1+\delta}}(\omega)} \\ &\lesssim \|u\|_{L^2}^{1-\delta} \|\psi\|_{L^2(\omega)} + \|u\|_{L^2}^{1+\delta} \|\psi\|_{H^{d\delta/2}(\omega)}, \end{aligned}$$

where we have used the Sobolev embedding in the last inequality.¹ Therefore, we deduce that $u \ln(|u|^2) \in H^{-d\delta/2}(\omega)$, which proves the result. \square

Remark 3.2. If $u \in L_T^\infty(L^2(\Omega))$, then the equation (1.1) makes sense in the sense of

$$i\partial_t u + \Delta u + \lambda u \ln(|u|^2) = 0 \quad \text{in } H^{-2}(\omega),$$

for any $\omega \Subset \Omega$ and a.e. t . In particular, it gives the meaning of the equation in the distribution sense for any $u \in L_T^\infty(L^2(\Omega))$.

We now recall the following important lemma:

Lemma 3.3 ([12, Lemma 2.2.1]). *Assume that $u, v \in C_T(H_0^1(\Omega))$ satisfies (1.1) in the distribution sense. Then, we have*

$$\|u(t) - v(t)\|_{L^2} \leq e^{2|\lambda t|} \|u(0) - v(0)\|_{L^2} \quad \text{for } t \in [-T, T]. \quad (3.1)$$

Proof. For completeness, we give a proof of this result. To simplify the presentation, we consider only the case $\Omega = \mathbb{R}^d$. We set

$$M := \max \left\{ \|u\|_{C_T(H^1)}, \|v\|_{C_T(H^1)} \right\}.$$

We note that u, v satisfy the equation in the sense of

$$i\partial_t u + \Delta u + \lambda u \ln(|u|^2) = 0 \quad \text{in } H^{-1}(B_R) \quad (3.2)$$

for any $R > 0$ and for all $t \in \mathbb{R}$, where B_R is the open ball of radius R with center at the origin of \mathbb{R}^d . Take a function $\zeta \in C_c^\infty(\mathbb{R}^d)$ satisfying

$$\zeta(x) = \begin{cases} 1 & \text{if } |x| \leq 1, \\ 0 & \text{if } |x| \geq 2, \end{cases} \quad 0 \leq \zeta(x) \leq 1 \quad \text{for all } x \in \mathbb{R}^d.$$

¹More precisely, we first consider the zero extension of ψ , and then apply the Sobolev embedding $H^{d\delta/2}(\mathbb{R}^d) \subset L^{\frac{2}{1-\delta}}(\mathbb{R}^d)$.

We set $\zeta_R(\cdot) = \zeta(\cdot/R)$ for $R > 0$. It follows from (3.2) and Lemma 2.1 (with $\varepsilon_1 = \varepsilon_2 = 0$) that

$$\begin{aligned} \frac{d}{dt} \|\zeta_R(u - v)\|_{L^2}^2 &= 2 \operatorname{Im} \langle i\zeta_R^2 \partial_t(u - v), u - v \rangle_{H^{-1}(B_{2R}), H_0^1(B_{2R})} \\ &= 2 \operatorname{Im} (\nabla(\zeta_R^2) \nabla(u - v), u - v)_{L^2} \\ &\quad - 4\lambda \operatorname{Im} (\zeta_R^2 (u \ln |u| - v \ln |v|), u - v)_{L^2} \\ &\leq \frac{C(M)}{R} + 4|\lambda| \|\zeta_R(u - v)\|_{L^2}^2. \end{aligned}$$

Integrating the last inequality over $[0, t]$, and applying Gronwall's lemma,

$$\|\zeta_R(u - v)(t)\|_{L^2}^2 \leq e^{4|\lambda|t} \left(\|u(0) - v(0)\|_{L^2}^2 + \frac{C(M)}{R} T \right),$$

for all $t \in (-T, T)$. Applying Fatou's lemma,

$$\|u(t) - v(t)\|_{L^2}^2 \leq \liminf_{R \rightarrow \infty} \|\zeta_R(u - v)(t)\|_{L^2}^2 \leq e^{4|\lambda|t} \|u(0) - v(0)\|_{L^2}^2,$$

which proves (3.1). \square

Remark 3.4. For the proof of Lemma 3.3, we need to assume H^1 solutions to give a sense of the duality product. Note, however, that (3.1) remains meaningful for L^2 solutions.

Now we take $\varphi \in L^2(\Omega)$ as the initial data. Take $\{\varphi_n\} \subset H_0^1(\Omega)$ such that $\varphi_n \rightarrow \varphi$ in $L^2(\Omega)$. We know from Theorem 2.2 that there exists a unique solution $u_n \in C(\mathbb{R}, H_0^1(\Omega))$ of (1.1) with $u_n(0) = \varphi_n$. Then, it follows from (1.1) that $\{u_n\}$ forms a Cauchy sequence in $L_{\text{loc}}^\infty(\mathbb{R}, L^2(\Omega))$. Therefore we deduce that there exists $u \in C(\mathbb{R}, L^2(\Omega))$ such that

$$u_n \rightarrow u \quad \text{in } L_{\text{loc}}^\infty(\mathbb{R}, L^2(\Omega)).$$

The rest of the proof consists in verifying that u is an L^2 solution in the above sense.

Lemma 3.5. *For any $\omega \Subset \Omega$ and for any $\varepsilon > 0$, we have*

$$g(u_n) \rightarrow g(u) \quad \text{in } L_{\text{loc}}^\infty(\mathbb{R}, H^{-\varepsilon}(\omega)),$$

where we have set $g(z) = z \ln(|z|^2)$.

Proof. We take a function $\theta \in C_c^1(\mathbb{C}, \mathbb{R})$ satisfying

$$\theta(z) = \begin{cases} 1 & \text{if } |z| \leq 1/2, \\ 0 & \text{if } |z| \geq 1, \end{cases} \quad 0 \leq \theta(z) \leq 1 \quad \text{for } z \in \mathbb{C}.$$

We set

$$g_1(u) = \theta(u)g(u), \quad g_2(u) = (1 - \theta(u))g(u).$$

We note that for $\alpha \in (0, 1)$,

$$|g_1(z) - g_1(w)| \lesssim_\alpha |z - w|^\alpha, \quad (3.3)$$

$$|g_2(z) - g_2(w)| \lesssim (\ln^+ |z| + \ln^+ |w|) |z - w|, \quad (3.4)$$

for any $z, w \in \mathbb{C}$. It follows from (3.3) that

$$g_1(u_n) \rightarrow g_1(u) \quad \text{in } L_{\text{loc}}^\infty(\mathbb{R}, L^2(\omega)). \quad (3.5)$$

Regarding the convergence of $g_2(u_n)$, we use the argument in the proof of Lemma 3.1. For $\psi \in C_c^1(\Omega)$ and any $\delta > 0$, we obtain from (3.4) that

$$\begin{aligned} \left| \int (g_2(u_n) - g_2(u))\psi \right| &\lesssim \int (|u_n|^\delta + |u|^\delta) |u_n - u| |\psi| \\ &\lesssim \left(\|u_n\|_{L^2}^\delta + \|u\|_{L^2}^\delta \right) \|u_n - u\|_{L^2} \|\psi\|_{L^{\frac{2}{1-\delta}}}, \end{aligned}$$

where we have used Hölder inequality with the exponent relation

$$\frac{\delta}{2} + \frac{1}{2} + \frac{1-\delta}{2} = 1.$$

Thus, we obtain

$$\left| \int (g_2(u_n) - g_2(u))\psi \right| \leq C(\|\varphi\|_{L^2}) \|u_n - u\|_{L^2} \|\psi\|_{H^\varepsilon},$$

where we take $\varepsilon = d\delta/2$ by the Sobolev embedding. Therefore, we deduce that

$$\|g_2(u_n) - g_2(u)\|_{H^{-\varepsilon}} \lesssim \|u_n - u\|_{L^2},$$

which implies that

$$g_2(u_n) \rightarrow g_2(u) \quad \text{in } L_{\text{loc}}^\infty(\mathbb{R}, H^{-\varepsilon}(\Omega)). \quad (3.6)$$

Hence, the result follows from (3.5) and (3.6). \square

We recall that u_n satisfies

$$i\partial_t u_n + \Delta u_n + \lambda u_n \ln(|u_n|^2) = 0 \quad \text{in } H^{-1}(\omega), \quad (3.7)$$

for all $\omega \in \Omega$. We now fix $\omega \in \Omega$, and take $\psi \in C_c^\infty(\omega)$ and $\phi \in C_c^1(\mathbb{R})$. It follows from (3.7) that

$$\begin{aligned} \int_{\mathbb{R}} (iu_n, \psi)_{L^2} \phi'(t) dt &= - \int_{\mathbb{R}} \langle i\partial_t u_n, \psi \rangle_{H^{-1}, H_0^1} \phi(t) dt \\ &= \int_{\mathbb{R}} \left((u_n, \Delta \psi)_{L^2} + \langle \lambda g(u_n), \psi \rangle_{H^{-1}, H_0^1} \right) \phi(t) dt. \end{aligned}$$

Passing to the limit as $n \rightarrow \infty$, we obtain from Lemma 3.5 that

$$\begin{aligned} \int_{\mathbb{R}} (iu, \psi)_{L^2} \phi'(t) dt &= \int_{\mathbb{R}} \left((u, \Delta \psi)_{L^2} - \langle \lambda g(u), \psi \rangle_{H^{-1}, H_0^1} \right) \phi(t) dt \\ &= \int_{\mathbb{R}} \langle \Delta u - \lambda g(u), \psi \rangle_{H^{-2}, H_0^2} \phi(t) dt. \end{aligned}$$

It is easily verified from this formula that

$$u \in C(\mathbb{R}, L^2(\Omega)) \cap C^1(\mathbb{R}, H^{-2}(\omega)),$$

for any $\omega \in \Omega$, and u satisfies (3.7) for any $\omega \in \Omega$ and all $t \in \mathbb{R}$.

The L^2 solution just constructed can be regarded as a unique extension of the solution map in H^1 . We define the solution map from H^1 initial data by

$$\Phi : H_0^1(\Omega) \ni \varphi \mapsto u \in C(\mathbb{R}, H_0^1(\Omega)). \quad (3.8)$$

For any $\varphi \in L^2(\Omega)$, we take a sequence $\{\varphi_n\} \subset H_0^1(\Omega)$ such that $\varphi_n \rightarrow \varphi$ in $L^2(\Omega)$ and define

$$\Phi(\varphi) = \lim_{n \rightarrow \infty} \Phi(\varphi_n) \in C(\mathbb{R}, L^2(\Omega)).$$

From the above discussion, $\Phi(\varphi)$ yields an L^2 solution of (1.1). We note from (3.1) that $\Phi(\varphi)$ is defined independently of the approximate sequence $\varphi_n \rightarrow \varphi$. Therefore the solution map (3.8) is uniquely extended from $H_0^1(\Omega)$ to $L^2(\Omega)$, and hence the first part of Theorem 1.1 follows.

4. INTERMEDIATE SOBOLEV REGULARITY

To prove the last part of Theorem 1.1, we show that the flow map associated to (2.1) propagates H^s regularity for $s \in (0, 1)$, uniformly in $\varepsilon \in (0, 1]$. For domains $\Omega \subset \mathbb{R}^d$ the fractional Sobolev spaces space $H^s(\Omega)$ may be defined either by real/complex interpolation between $L^2(\Omega)$ and $H_0^1(\Omega)$. When $\Omega = \mathbb{R}^d$, it is well known that the Bessel potential spaces by (1.4) are characterized by complex interpolation as

$$H^s(\mathbb{R}^d) = [L^2(\mathbb{R}^d), H^1(\mathbb{R}^d)]_s, \quad s \in (0, 1).$$

In the first three cases considered for Ω in the introduction (whole space, half space, or smooth bounded domain), [1, Theorem 7.48] states that the fractional Sobolev spaces defined by real interpolation are equivalent to the ones equipped with the norm (1.3). Regarding the last case $\Omega = \mathbb{T}^d$, it is also known that a similar equivalence relation holds as follows.

Lemma 4.1 ([4, Proposition 1.3]). *Let $s \in (0, 1)$. Then, for $f \in H^s(\mathbb{T}^d)$ we have the relation*

$$\|f\|_{H^s(\mathbb{T}^d)}^2 \sim \|f\|_{L^2(\mathbb{T}^d)}^2 + \iint_{\mathbb{T}^d \times [-\frac{1}{2}, \frac{1}{2}]^d} \frac{|f(x+y) - f(x)|^2}{|y|^{d+2s}} dx dy.$$

We denote the approximate nonlinearity by

$$g_\varepsilon(z) = 2z \ln(|z| + \varepsilon) \quad \text{for } \varepsilon \in (0, 1].$$

For domains $\Omega \subset \mathbb{R}^d$, we differentiate $\|u^\varepsilon(t)\|_{H^s}^2$ with respect to time, where we recall the H^s norm defined by (1.3). Then, in view of the conservation of mass,

we obtain

$$\begin{aligned}
& \frac{d}{dt} \|u^\varepsilon(t)\|_{H^s(\Omega)}^2 \\
&= 2 \operatorname{Re} \iint_{\Omega \times \Omega} \overline{(u^\varepsilon(t, x+y) - u^\varepsilon(t, x))} \partial_t (u^\varepsilon(t, x+y) - u^\varepsilon(t, x)) \frac{dx dy}{|y|^{d+2\alpha}} \\
&= -\operatorname{Im} \iint_{\Omega \times \Omega} \overline{(u^\varepsilon(t, x+y) - u^\varepsilon(t, x))} \Delta (u^\varepsilon(t, x+y) - u^\varepsilon(t, x)) \frac{dx dy}{|y|^{d+2\alpha}} \\
&\quad - 2\lambda \operatorname{Im} \iint_{\Omega \times \Omega} \overline{(u^\varepsilon(t, x+y) - u^\varepsilon(t, x))} (g_\varepsilon(u^\varepsilon(t, x+y)) - g_\varepsilon(u^\varepsilon(t, x))) \frac{dx dy}{|y|^{d+2\alpha}}.
\end{aligned}$$

The first term on the right hand side is zero by integration by parts in x . For the second term, by applying Lemma 2.1 we obtain

$$\begin{aligned}
& \frac{d}{dt} \|u^\varepsilon(t)\|_{H^s(\Omega)}^2 \\
&\leq 2|\lambda| \iint_{\Omega \times \Omega} \left| \operatorname{Im} \left[\overline{(u^\varepsilon(t, x+y) - u^\varepsilon(t, x))} (g_\varepsilon(u^\varepsilon(t, x+y)) - g_\varepsilon(u^\varepsilon(t, x))) \right] \right| \frac{dx dy}{|y|^{d+2\alpha}} \\
&\leq 4|\lambda| \iint_{\Omega \times \Omega} |u^\varepsilon(t, x+y) - u^\varepsilon(t, x)|^2 \frac{dx dy}{|y|^{d+2\alpha}} \leq 4|\lambda| \|u^\varepsilon(t)\|_{H^s(\Omega)}^2.
\end{aligned}$$

Therefore, by Gronwall's lemma we deduce

$$\|u^\varepsilon(t)\|_{H^s(\Omega)}^2 \leq e^{4|\lambda t|} \|\varphi\|_{H^s(\Omega)}^2 \quad \text{for all } t \in \mathbb{R}. \quad (4.1)$$

For the torus \mathbb{T}^d , a similar estimates is obtained through Lemma 4.1 as

$$\|u^\varepsilon(t)\|_{H^s(\mathbb{T}^d)}^2 \leq e^{C|\lambda t|} \|\varphi\|_{H^s(\mathbb{T}^d)}^2 \quad \text{for all } t \in \mathbb{R} \quad (4.2)$$

for some $C > 0$ independent of ε and t .

In view of the construction presented in Section 3, it follows from (4.1), (4.2), and limiting procedures that $\Phi(\varphi) \in C_w(\mathbb{R}, H^s(\Omega))$ when $\varphi \in H^s(\Omega)$. Applying the argument of [27, Remarks (c)], one can improve the regularity in time as $\Phi(\varphi) \in C(\mathbb{R}, H^s(\Omega))$ (see the proof of [24, Lemma 2.11] for more details). This completes the proof of the last part of Theorem 1.1.

APPENDIX A. WEAK SOLUTIONS IN THE SENSE OF BREZIS

In this section, we state the content of [12, Theorem 1.2 b)] in a mostly self-contained manner. The construction of their L^2 weak solutions depend on the maximal monotone theory. We define the nonlinear operator A by

$$Au = -i\Delta u - i\lambda u \ln(|u|^2)$$

with the domain

$$D(A) = \{u \in H_{\text{loc}}^1(\mathbb{R}^d) \cap L^2(\mathbb{R}^d) : \Delta u + \lambda u \ln(|u|^2) \in L^2(\mathbb{R}^d)\}.$$

From [12, Theorem 1.1] we know that $A + 2|\lambda|I$ is maximal monotone in $L^2(\mathbb{R}^d)$. Note that the inequality (1.2) is used to show the monotonicity of A . We now rewrite the equation (1.1) as

$$\frac{du}{dt} + (A + 2|\lambda|)u = 2|\lambda|u, \quad u(0) = \varphi. \quad (\text{A.1})$$

The definition of weak solutions in the sense of Brezis is given as follows.

Definition A.1 ([6, Definition 3.1]). *Let A be a maximal monotone operator on the Hilbert space H . Assume $f \in L^1((0, T), H)$ for some $T > 0$. We say that $u \in C([0, T], H)$ is a weak solution of the equation $\frac{du}{dt} + Au = f$ if there exists sequences $f_n \in L^1((0, T), H)$ and $u_n \in C([0, T], H)$ such that u_n satisfies $\frac{du_n}{dt} + Au_n = f_n$ for a.e. $t \in (0, T)$, $f_n \rightarrow f$ in $L^1((0, T), H)$, and $u_n \rightarrow u$ in $C([0, T], H)$.*

The authors in [12] apply [6, Theorem 3.17] to the equation (A.1) and prove that there exists a unique L^2 weak solution in the sense of Definition A.1. We note that in the proof of [6, Theorem 3.17] the inequality (26) therein plays a key role in guaranteeing both the existence and uniqueness of solutions, and that this inequality is a consequence of monotonicity of the operator.

ACKNOWLEDGMENTS

The authors would like to thank Thierry Cazenave for clearly explaining Theorem 1.2 b) in [12], and Nobu Kishimoto for pointing out the reference [4].

R.C. is partially supported by Centre Henri Lebesgue, program ANR-11-LABX-0020-0.

M.H. is supported by JSPS KAKENHI Grant Number JP22K20337 and by the Italian MIUR PRIN project 2020XB3EFL.

T.O. is supported by JSPS KAKENHI Grant Numbers 18KK073 and 19H00644.

A CC-BY public copyright license has been applied by the authors to the present document and will be applied to all subsequent versions up to the Author Accepted Manuscript arising from this submission.

REFERENCES

- [1] R. A. Adams, *Sobolev Spaces*, Pure and Applied Mathematics, Vol. 65, Academic Press, New York-London, 1975.
- [2] A. H. Ardila, Orbital stability of Gausson solutions to logarithmic Schrödinger equations, *Electron. J. Differential Equations* (2016), Paper No. 335, 9 pp.
- [3] W. Bao, R. Carles, C. Su, and Q. Tang, Error estimates of a regularized finite difference method for the logarithmic Schrödinger equation, *SIAM J. Numer. Anal.* **57** (2019), 657–680.
- [4] A. Bényi and T. Oh, The Sobolev inequality on the torus revisited, *Publ. Math. Debrecen* **83** (2013), 359–374.
- [5] I. Białyński-Birula and J. Mycielski, Nonlinear wave mechanics, *Ann. Physics* **100** (1976), 62–93.
- [6] H. Brézis, *Opérateurs Maximaux Monotones et Semi-Groupes de Contractions dans les Espaces de Hilbert*, North-Holland Mathematics Studies, No. 5, New York, 1973.

- [7] H. Buljan, A. Šiber, M. Soljačić, T. Schwartz, M. Segev, and D. Christodoulides, Incoherent white light solitons in logarithmically saturable noninstantaneous nonlinear media, *Phys. Rev. E* **68** (2003), 036607.
- [8] R. Carles, E. Dumas, and C. Sparber, Geometric optics and instability for NLS and Davey-Stewartson models, *J. Eur. Math. Soc. (JEMS)* **14** (2012), 1885–1921.
- [9] R. Carles and I. Gallagher, Universal dynamics for the defocusing logarithmic Schrödinger equation, *Duke Math. J.* **167** (2018), 1761–1801.
- [10] T. Cazenave, Stable solutions of the logarithmic Schrödinger equation, *Nonlinear Anal.* **7** (1983), 1127–1140.
- [11] T. Cazenave, *Semilinear Schrödinger Equations*, Courant Lecture Notes in Mathematics, vol. 10, New York University Courant Institute of Mathematical Sciences, New York, 2003.
- [12] T. Cazenave and A. Haraux, Équations d'évolution avec non linéarité logarithmique, *Ann. Fac. Sci. Toulouse Math. (5)* **2** (1980), 21–51.
- [13] T. Cazenave and A. Haraux, *An Introduction to Semilinear Evolution Equations*, Oxford Lecture Series in Mathematics and its Applications, vol. 13, The Clarendon Press Oxford University Press, New York, 1998.
- [14] M. Christ, J. Colliander, and T. Tao, Asymptotics, frequency modulation, and low regularity ill-posedness for canonical defocusing equations, *Amer. J. Math.* **125** (2003), 1235–1293.
- [15] M. Christ, J. Colliander, and T. Tao, Ill-posedness for nonlinear Schrödinger and wave equations, preprint (2003), [arXiv:math/0311048].
- [16] P. d'Avenia, E. Montefusco, and M. Squassina, On the logarithmic Schrödinger equation, *Commun. Contemp. Math.* **16** (2014), 1350032, 15 pp.
- [17] E. Di Nezza, G. Palatucci, and E. Valdinoci, Hitchhiker's guide to the fractional Sobolev spaces, *Bull. Sci. Math.* **136** (2012), 521–573.
- [18] G. Ferriere, The focusing logarithmic Schrödinger equation: analysis of breathers and nonlinear superposition, *Discrete Contin. Dyn. Syst.* **40** (2020), 6247–6274.
- [19] G. Ferriere, Existence of multi-solitons for the focusing logarithmic non-linear Schrödinger equation, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **38** (2021), 841–875.
- [20] T. Hansson, D. Anderson, and M. Lisak, Propagation of partially coherent solitons in saturable logarithmic media: A comparative analysis, *Phys. Rev. A* **80** (2009), 033819.
- [21] B. Harrop-Griffiths, R. Killip, M. Ntekoume, and M. Visan, Global well-posedness for the derivative nonlinear Schrödinger equation in $L^2(\mathbb{R})$, preprint (2022), [arXiv:2204.12548].
- [22] B. Harrop-Griffiths, R. Killip, and M. Visan, Sharp well-posedness for the cubic NLS and mKdV in $H^s(\mathbb{R})$, preprint (2020), [arXiv:2003.05011].
- [23] M. Hayashi, A note on the nonlinear Schrödinger equation in a general domain, *Nonlinear Anal.* **173** (2018), 99–122.
- [24] M. Hayashi and T. Ozawa, The Cauchy problem for the logarithmic Schrödinger equation revisited, preprint (2023), [arXiv:2309.01695].
- [25] E. F. Hefter, Application of the nonlinear Schrödinger equation with a logarithmic inhomogeneous term to nuclear physics, *Phys. Rev. A* **32** (1985), 1201–1204.
- [26] E. S. Hernandez and B. Remaud, General properties of Gausson-conserving descriptions of quantal damped motion, *Physica A* **105** (1980), 130–146.
- [27] T. Kato and C. Y. Lai, Nonlinear evolution equations and the Euler flow, *J. Funct. Anal.* **56** (1984), 15–28.
- [28] C. Kenig, G. Ponce, and L. Vega, On the ill-posedness of some canonical dispersive equations, *Duke Math. J.* **106** (2001), 617–633.
- [29] R. Killip and M. Visan, KdV is well-posed in H^{-1} , *Ann. of Math.* **190** (2019), 249–305.
- [30] N. Kishimoto, A remark on norm inflation for nonlinear Schrödinger equations, *Commun. Pure Appl. Anal.* **18** (2019), 1375–1402.

- [31] W. Krolikowski, D. Edmundson, and O. Bang, Unified model for partially coherent solitons in logarithmically nonlinear media, *Phys. Rev. E* **61** (2000), 3122–3126.
- [32] S. D. Martino, M. Falanga, C. Godano, and G. Lauro, Logarithmic Schrödinger-like equation as a model for magma transport, *Europhys. Lett.* **63** (2003), 472–475.
- [33] K. Yasue, Quantum mechanics of nonconservative systems, *Annals Phys.* **114** (1978), 479–496.
- [34] K. G. Zloshchastiev, Logarithmic nonlinearity in theories of quantum gravity: Origin of time and observational consequences, *Grav. Cosmol.* **16** (2010), 288–297.

UNIV RENNES, CNRS, IRMAR - UMR 6625, F-35000 RENNES, FRANCE
Email address: Remi.Carles@math.cnrs.fr

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PISA, LARGO BRUNO PONTECORVO, 5
56127 PISA, ITALY

WASEDA RESEARCH INSTITUTE FOR SCIENCE AND ENGINEERING, WASEDA UNIVERSITY,
TOKYO 169-8555, JAPAN

Email address: masayuki.hayashi@dm.unipi.it

DEPARTMENT OF APPLIED PHYSICS, WASEDA UNIVERSITY, TOKYO 169-8555, JAPAN

Email address: txozawa@waseda.jp