



HAL
open science

Package 'ExGaussEstim'

Yousri Slaoui, Abir El Haj, Alandra Zakkour, Caroline Bordes, Cyril Perret,
Jean Dumoncel

► **To cite this version:**

Yousri Slaoui, Abir El Haj, Alandra Zakkour, Caroline Bordes, Cyril Perret, et al.. Package 'ExGauss-
Estim'. /, 2023. hal-04269036

HAL Id: hal-04269036

<https://hal.science/hal-04269036v1>

Submitted on 3 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Public Domain

Package ‘ExGaussEstim’

October 6, 2023

Title Quantile Maximization Likelihood Estimation and Bayesian
Ex-Gaussian Estimation

Version 0.1.2

Description Presents two methods to estimate the parameters 'mu', 'sigma', and 'tau' of an ex-Gaussian distribution. Those methods are Quantile Maximization Likelihood Estimation ('QMLE') and Bayesian. The 'QMLE' method allows a choice between three different estimation algorithms for these parameters : 'neldermead' ('NEMD'), 'fmin-search' ('FMIN'), and 'nlminb' ('NLMF'). For more details about the methods you can refer at the following list: Brown, S., & Heathcote, A. (2003) <[doi:10.3758/BF03195527](https://doi.org/10.3758/BF03195527)>; McCormack, P. D., & Wright, N. M. (1964) <[doi:10.1037/h0083285](https://doi.org/10.1037/h0083285)>; Van Zandt, T. (2000) <[doi:10.3758/BF03214357](https://doi.org/10.3758/BF03214357)>; El Haj, Aret, C. (2021) <[doi:10.19139/soic-2310-5070-1251](https://doi.org/10.19139/soic-2310-5070-1251)>; Gilks, W. R., Best, N. G., & Tan, K. K. C. (1995) <[doi:10.2307/2986138](https://doi.org/10.2307/2986138)>.

License GPL-2

Encoding UTF-8

LazyLoad true

RoxygenNote 7.2.3

Imports pracma, stats, nloptr, invgamma, dlm, fitdistrplus,
gamlss.dist

NeedsCompilation no

Suggests testthat (>= 3.0.0)

Config/testthat/edition 3

Author Yousri SLAOUI [aut] (<<https://orcid.org/0000-0001-5295-3311>>),
Abir EL HAJ [aut, ctb] (<<https://orcid.org/0000-0003-3210-9097>>),
Alandra ZAKKOUR [aut],
Caroline BORDES [aut, ctb],
Cyril PERRET [aut] (<<https://orcid.org/0000-0002-4552-9093>>),
Jean DUMONCEL [aut, cre] (<<https://orcid.org/0000-0003-0789-7458>>)

Maintainer Jean DUMONCEL <jean.dumoncel@univ-poitiers.fr>

Repository CRAN

Date/Publication 2023-10-06 07:20:02 UTC

R topics documented:

| | |
|------------------------------|----------|
| BayesianExgaussian | 2 |
| QMLEEstim | 3 |
| Index | 5 |

| | |
|--------------------|--------------------------------------|
| BayesianExgaussian | <i>Bayesian Ex-gaussian Estimate</i> |
|--------------------|--------------------------------------|

Description

Estimates the mu, sigma, and tau parameters of an ex-Gaussian distribution using a bayesian method.

Usage

```
BayesianExgaussian(n, x, nSamples = 5000, Ti = 2500)
```

Arguments

| | |
|----------|--|
| n | the data size |
| x | the data. Must be a vector, with no missing values |
| nSamples | number of Samples |
| Ti | burn-in |

Value

BayesianExgaussian() returns an object "theta" which is a list with components: estimates of mu, sigma, and tau

References

- Brown, S., & Heathcote, A. (2003). QMLE: Fast, robust, and efficient estimation of distribution functions based on quantiles. *Behavior Research Methods, Instruments, & Computers*, **35**, 485-492.
- McCormack, P. D., & Wright, N. M. (1964). The positive skew observed in reaction time distributions. *Canadian Journal of Psychology/Revue canadienne de psychologie*, **18**, 43-51.
- Van Zandt, T. (2000). How to fit a response time distribution. *Psychonomic Bulletin & Review*, **7**, 424-465.
- El Haj, A., Slaoui, Y., Solier, C., & Perret, C. (2021). Bayesian Estimation of The Ex-Gaussian distribution. *Statistics, Optimization & Information Computing*, **9(4)**, 809-819.
- Gilks, W. R., Best, N. G., & Tan, K. K. C. (1995). Adaptive rejection Metropolis sampling within Gibbs sampling. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, **44**, 455-472.

Examples

```
library(gamlss.dist)
set.seed(2703)
data<-rexGAUS(n=100, mu = 500, sigma = 150, nu = 100)
BayesianExgaussian(n = 100, x = data)
```

QMLEEstim

*Ex-Gaussian Quantile Maximum Likelihood Estimate***Description**

Estimates the mu, sigma, and tau parameters of an ex-Gaussian distribution. 3 different algorithms can be used : neldermead ('NEMD'), fminsearch ('FMIN') and nlminb ('NLMI').

Usage

```
QMLEEstim(y, func)
```

Arguments

| | |
|------|--|
| y | the data. Must be a vector with no missing values |
| func | the function selected for the estimation. 'NEMD' for neldermead, 'FMIN' for fminsearch, and 'NLMI' for nlminb. |

Value

QMLEEstim() returns an object "valEstim" which is a list with components: estimates of mu, sigma, and tau

References

- Brown, S., & Heathcote, A. (2003). QMLE: Fast, robust, and efficient estimation of distribution functions based on quantiles. *Behavior Research Methods, Instruments, & Computers*, **35**, 485-492.
- McCormack, P. D., & Wright, N. M. (1964). The positive skew observed in reaction time distributions. *Canadian Journal of Psychology/Revue canadienne de psychologie*, **18**, 43-51.
- Van Zandt, T. (2000). How to fit a response time distribution. *Psychonomic Bulletin & Review*, **7**, 424-465.
- El Haj, A., Slaoui, Y., Solier, C., & Perret, C. (2021). Bayesian Estimation of The Ex-Gaussian distribution. *Statistics, Optimization & Information Computing*, **9(4)**, 809-819.
- Gilks, W. R., Best, N. G., & Tan, K. K. C. (1995). Adaptive rejection Metropolis sampling within Gibbs sampling. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, **44**, 455-472.

Examples

```
library(gamlss.dist)
set.seed(2703)
data<-rexGAUS(n=100, mu = 500, sigma = 150, nu = 100)
QMLEEstim(data, 'NEMD')
```

Index

BayesianExgaussian, [2](#)

QMLEstim, [3](#)