



**HAL**  
open science

# Solutan echinoderms from the Lower Ordovician of the Montagne Noire (France): new data and palaeobiogeographic implications

Christophe Dupichaud, Bertrand Lefebvre, Martina Nohejlová

## ► To cite this version:

Christophe Dupichaud, Bertrand Lefebvre, Martina Nohejlová. Solutan echinoderms from the Lower Ordovician of the Montagne Noire (France): new data and palaeobiogeographic implications. *Estonian Journal of Earth Sciences*, 2023, 72 (1), pp.26. 10.3176/earth.2023.80 . hal-04271100

**HAL Id: hal-04271100**

**<https://hal.science/hal-04271100v1>**

Submitted on 5 Nov 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## SHORT COMMUNICATION

**Solutan echinoderms from the Lower Ordovician of the Montagne Noire (France):  
new data and palaeobiogeographic implications**Christophe Dupichaud<sup>a\*</sup>, Bertrand Lefebvre<sup>a</sup> and Martina Nohejlová<sup>b</sup><sup>a</sup> Univ Lyon, Univ Lyon 1, ENSL, CNRS, LGL-TPE, F-69622, Villeurbanne, France<sup>b</sup> Czech Geological Survey, Klárov 3, Praha 1, 118 21 Czech Republic\* Corresponding author, [christophe.dupichaud@univ-lyon1.fr](mailto:christophe.dupichaud@univ-lyon1.fr)

Received ..., accepted ..., available online ...

**Abstract.** Solutans are relatively common members of echinoderm assemblages in the Saint-Chinian Formation (upper Tremadocian) of the Montagne Noire (France). The revision of all available material confirms that most specimens can be assigned to *Minervae cystis vidali*, which is here reconstructed for the first time. The occurrence of *Plasiacystis mobilis* evidenced based on a single, well-preserved dististele. The interpretation of a small-sized individual as a putative dendrocystitid is rejected: it corresponds to a juvenile specimen of *M. vidali*. Early Ordovician solutans from the Montagne Noire partly fill the gap between Laurentian mid–late Cambrian syringocrinids and Avalonian-Gondwanan Early–Mid Ordovician taxa.

**Keywords:** Echinodermata, Montagne Noire, Ordovician, Soluta, Tremadocian.

**INTRODUCTION**

In spite of some recent reports (e.g. Zhu et al. 2016), very few Furongian–Early Ordovician echinoderm faunas are known worldwide from this critical interval at the transition between the Cambrian Explosion and the Great Ordovician Biodiversification (Smith 1988). With the Fillmore Formation in western Utah, USA (Sprinkle & Guensburg 1995) and the lower part of the Fezouata Formation in the Central Anti-Atlas, Morocco (Lefebvre et al. 2016), the Saint-Chinian Formation in Montagne Noire, France, is one of the few lithostratigraphic units yielding abundant and diverse Tremadocian echinoderm remains (Thoral 1935, Vizcaïno & Lefebvre 1999, Vizcaïno et al. 2001). This situation probably results from their remarkable three-dimensional preservation in aluminosiliceous concretions and also from nearly 150 years of intensive, continuous sampling (Vizcaïno et al. 2001 and references therein). Echinoderm assemblages from the Saint-Chinian Formation comprise over 30 taxa belonging to eight classes. In terms of abundance and diversity, stylophorans (cornutes and mitrates) are the dominant group. The Saint-Chinian Formation also comprises some of the earliest representatives of the subphylum Asterozoa and of the class Crinoidea. This unit also yielded edrioasteroids, eocrinoids, rhombiferans and solutans (see Vizcaïno & Lefebvre 1999, Vizcaïno et al. 2001 and references therein).

Along with eocrinoids, glyptocystitid rhombiferans and stylophorans, solutans are typical members of the cosmopolitan, low-diversity, Furongian–Early Ordovician echinoderm assemblages (Sumrall et al. 1997, Lefebvre et al. 2013, Zhu et al. 2016). Late Cambrian and younger solutans were vagile members of the benthos, generally interpreted as probable detritus-feeders, sweeping soft substrates with their unique feeding appendage (e.g. Kolata 1973, Noailles et al. 2014, Nohejlová & Lefebvre 2022). In the Lower Ordovician of the Montagne Noire, a single species of solutans has been described so far (*Minervae cystis vidali*) from the concretion-bearing shales of the Saint-Chinian Formation (Thoral 1935, Ubaghs 1970). Based on the examination of all available material (historical collections and additional specimens collected since Ubaghs' 1970 seminal description of *M. vidali*), the aims of this paper are (1) to revise the morphology and produce the first reconstruction of *Minervae cystis*; (2) to describe the first occurrence of *Plasiacystis mobilis* in the Lower Ordovician of southern France; and (3) to discuss the palaeobiogeographic affinities of the Montagne Noire solutans.


**MATERIALS AND METHODS**

The study material includes 42 solutan specimens collected in the Lower Ordovician of the Montagne Noire (southern France) and catalogued in the following public collections: Natural History Museum, London (BMNHUK: Miquel collection), Musée du Biterrois, Béziers (MBB: Griffé collection), Muséum National d'Histoire Naturelle, Paris (MNHN: Courtessole-Griffé collection), Lyon 1 University, Villeurbanne (UCBL.FSL: Lignièrès, Kundura, Villebrun, and Vizcaïno collections), and Montpellier University, Montpellier (UM: Azaïs, Roquefort, Thoral, and Villebrun collections). All specimens are from the Saint-Chinian Formation (late Tremadocian, *Euloma filacovi* Zone; see Vizcaïno & Álvaro 2003). In the Saint-Chinian area, this 500 m thick formation consists of relatively monotonous dark concretion-bearing siltstones, with fine sandstone intercalations (Vizcaïno et al. 2001, Vizcaïno & Álvaro 2003). The Saint-Chinian Formation has yielded a particularly diverse fossiliferous assemblage typical of

moderately deep, well-oxygenated shelf environments, below storm wave base, and comprising bivalves, brachiopods, cephalopods, conulariids, echinoderms, gastropods, graptolites, hyolithids, machaeridians, and trilobites (Thoral 1935, Vizcaïno et al. 2001).

All fossils are preserved three-dimensionally, as empty negative moulds, in aluminosiliceous concretions. So as to reveal their original aspect, solutans were cast with latex, and coated with ammonium chloride (NH<sub>4</sub>Cl) for drawing and photographic purposes. Observations and camera-lucida drawings were made with a Zeiss StEREO Discovery.V8 stereomicroscope binocular, and photographs with a Canon 5DSR camera equipped with a MP-E 65 mm macro-lens. Digital drawings were done under the software Krita 5.0.2.


## RESULTS



**Fig. 1.** Solutans from the Saint-Chinian Formation (upper Tremadocian), Montagne Noire (Hérault, France); latex casts of original specimens; **A–E**, *Minervaecystis vidali* (Thoral, 1935); **A–B**, cluster of four individuals (part and counterpart), BMNHUK.E.23695 (Miquel coll.), Cassagnoles; **A**, nearly complete individual (in lower aspect) in the centre of the cluster, with fully articulated brachiole and long, strongly curved, twisted dististele; **B**, same individuals (in upper aspect); **C**, well-preserved individual (in lower aspect) with strongly curved, twisted dististele; UCBL.FSL.713595 (Kundurá coll.), Prades-sur-Vernazobre; **D**, fully articulated theca (in upper aspect) with brachiole and well-preserved anal pyramid, right of homoiostele insertion; lectotype, UM.ACI.634 (Villebrun coll.), Saint-Chinian; **E**, juvenile individual, UM.ACI.636 (Thoral coll.), Prades-sur-Vernazobre; **F–G**, *Plasiacystis mobilis* Prokop & Petr, 2003; latex casts of the dististele (part and counterpart) of the single known specimen (MBB.GG.6, Griffé coll.), Saint-Chinian; note the presence of a well-defined spike-shaped tubercle on one of the two sides (**F**).

The re-examination of Thoral's (1935) original type series and of over 30 additional specimens (Fig. 1A–E) did not bring any significant additional morphological information to the very detailed description of *Minervaecystis vidali* provided by Ubaghs (1970). However, this abundant material made it possible to elaborate the first detailed, specimen-based reconstruction of this solutan (Fig. 2). Tentative reconstructions of *M. vidali* were already provided by Lefebvre et al. (2012, fig. 2.1) and Rozhnov (2022, fig. 4h), but they were simply combining two original camera-lucida drawings figured in Ubaghs (1970, figs 39.1 and 41.1). Moreover, the observation of all available individuals of *M. vidali* indicates that, when it is preserved, their twisted dististele is not straight, but consistently curved towards the right (Fig. 1A, C). *Minervaecystis* shares with *Pahvanticystis* and *Plasiacystis* the possession of a non-apical, laterally inserted brachiole, a relatively large, inflated, ovoid theca, and a flattened, twisted dististele (Lefebvre & Lerosey-Aubril 2018). *Minervaecystis* differs from *Pahvanticystis* in the more posterior insertion of its brachiole, a proportionately longer proxistele (comprising over 20 tetramerous rings), and the morphology of its dististele (narrower, and with lateral comb-like spikes). *Minervaecystis* mostly differs from *Plasiacystis* in the possession of a proxistele consisting of numerous, regularly arranged tetramerous rings (this region is made of numerous,

unorganized, imbricate, scale-like plates in *P. mobilis*), and of an elongate, stem-like dististele, instead of a short, broad, paddle-like distal structure (Prokop & Petr 2003, Lefebvre et al. 2012).



**Fig. 2.** Reconstruction of *Minervaecystis vidali* (Thoral, 1935), Saint-Chinian Formation (upper Tremadocian), Montagne Noire (France).

Lefebvre & Lerosey-Aubril (2018) questioned the possibility that the smallest known individual assigned to *Minervaecystis* (UM.ACI.636) and figured by both Thoral (1935, pl. 11 fig. 2) and Ubaghs (1970, pl. 16 fig. 5) could represent one of the oldest known dendrocystitid solutans. Their suggestion relied on the quadrangular aspect of the theca and, mostly, on the location and morphology of the periproct (relatively large, rounded, low structure formed by several elongate tapered plates forming a well-defined anal rosette). However, the re-examination of the original specimen (Fig. 1E) does not support this interpretation, but confirms its identification as a juvenile individual of *M. vidali* (Thoral 1935, Ubaghs 1970). Its unusual aspect results from the combination of a theca preserved as a hollow internal mould, with two appendages (brachiolo and homoistele) occurring as three-dimensional external moulds. Consequently, contrary to most other specimens of *Minervaecystis*, the periproct of UM.ACI.636 is visible in internal view. Its regular arrangement into a well-defined, closed rosette suggests that the prominent, cone-shaped (external) morphology of the periproct observed in most individuals is possibly preservational and/or results from its ability to open outwards (valvular structure). As in all other specimens of *Minervaecystis*, the brachiolo is inserted laterally, away from the apex and on the antanal side of the theca (i.e. opposite the periproct).

Although the occurrence of *Plasiacystis mobilis* was regularly mentioned in the Lower Ordovician of the Montagne Noire (e.g. Lefebvre et al. 2012, 2016), no remains of this solutan have been formally described or illustrated so far from this region. The detailed re-examination of historical collections made it possible to confirm the presence of *P. mobilis*, based on a single, well-preserved, isolated dististele (MBB.GG.6, part and counterpart) collected in the Saint-Chinian Formation. Its short and broad, bean-shaped, flattened morphology (Fig. 1F–G) is typical of *Plasiacystis*, and identical to that of the dististele in Czech specimens of *P. mobilis* (see Prokop & Petr 2003, Lefebvre et al. 2012). The distally incomplete paddle-like appendage of MBB.GG.6 consists of two unequal series of wide, flattened plates, with those forming the concave lateral edge about twice longer than those on the opposite, convex margin. The dististele slightly increases in thickness along the longitudinal suture running along opposite, left and right, sets of plates. In MBB.GG.6, a well-defined tubercle (spike) occurs on one side of the dististele. It cannot be formally excluded that some poorly preserved solutan thecae traditionally assigned to *Minervaecystis* may indeed belong to *Plasiacystis*.

## DISCUSSION

Their abundant and almost continuous fossil record from the lower Drumian to the upper Floian of (modern) North America (Alabama, Nevada and Utah) suggests that solutans very likely originated in Laurentia (Zamora et al. 2013a, Lefebvre and Lerosey-Aubril 2018), possibly from a gogiid-like ancestor (Parsley 1997). Yet undescribed solutans from the Sandu Formation (Jiangshanian) of South China represent the earliest known occurrence of this clade outside of Laurentia (Zamora et al. 2013b, Zhu et al. 2016). In this context, the late Tremadocian solutans from the Montagne Noire correspond, along with coeval occurrences from the Central Anti-Atlas (Morocco), to the earliest known record of this class in high-latitude Gondwanan regions (Lefebvre et al. 2013, 2016). The two taxa occurring in the Lower Ordovician of southern France (*Minervaecystis* and *Plasiacystis*) are closely related to Cambrian syringocrinids from Laurentia (*Castericystis*, and mostly *Pahvanticystis*; see Lefebvre & Lerosey-Aubril 2018). Although *Minervaecystis vidali* is so far considered as endemic to the Montagne Noire, very similar isolated homoiosteles sp. from the Whipple Cave Formation (upper Furongian) of Nevada (Ubaghs 1963, Sumrall et al. 1997) have been tentatively assigned to this genus (*Minervaecystis?* sp.), and yet undescribed remains of minervaecystids are also known from the Wah Wah Formation (late Floian) of Utah (Sprinkle & Guensburg 1993). In sharp contrast with the Montagne Noire, where it is extremely rare, *Plasiacystis mobilis* is particularly abundant and widespread in the Lower Ordovician of Morocco, with occurrences ranging from the late Tremadocian to the late Floian (Lefebvre et al. 2016, Dupichaud & Lefebvre 2022, Saleh et al. 2022). The latter solutan is remarkably long time-ranging, as it is also present in the Darriwilian of Shropshire, UK (Hope Shale; Lefebvre et al. 2012) and of the Prague Basin, Czech Republic (Šarka Formation; Prokop & Petr 2003, Lefebvre et al. 2012).

## CONCLUSIONS

Solutans were relatively common members of late Tremadocian benthic communities in the southern Montagne Noire (France). The revision of all available specimens from the Saint-Chinian Formation made it possible to confirm the occurrence of two distinct taxa: (1) the most abundant one, *Minervaecystis vidali*, is here reconstructed for the first time, with new informations on the morphology of its periproct; (2) *Plasiacystis mobilis*, which is particularly rare in the Lower Ordovician of the Montagne Noire, is described and figured for the first time; and (3) the occurrence of putative dendrocystitids is not confirmed. The Montagne Noire solutan assemblage documents a key evolutionary transition, with a mixture of plesiomorphic morphologies inherited from the worldwide expansion of syringocrinids outside of Laurentia in the late Cambrian (*Minervaecystis*), and the onset of new, highly derived morphotypes in Avalonia and Gondwana (*Plasiacystis*) in the Early and Mid Ordovician. During the Darriwilian, the persistence and diversification of solutans in the latter two geographic areas fuelled their expansion to Baltica and Laurentia (where they had disappeared at the end of the Floian) in the Late Ordovician.

**Acknowledgements.** This study is a contribution to the IGCP Project 735 "Rocks and the Rise of Ordovician Life" and was supported by the PHC Barrande Project n°49290RD. The authors are particularly grateful to \_\_\_\_\_ and \_\_\_\_\_ for their helpful and constructive remarks, which helped improving the MS, and also to Vincent Perrier, for assistance with photographs.

## REFERENCES

- Dupichaud C. and Lefebvre B. 2022. Solutan echinoderms from the Fezouata Shale Lagerstätte (Lower Ordovician, Morocco): diversity, exceptional preservation and palaeobiogeographic implications. *Abstract book*, Second Annual Meeting of IGCP 735, Cadi Ayyad University, Marrakesh, *October 19-20 2022*, 23–28.
- Kolata, D. R. 1973. *Scalenocystites strimplei*, a new Middle Ordovician belemnocystitid solute from Minnesota. *Journal of Paleontology*, **47**, 969–974.
- Lefebvre, B. and Lerosey-Aubril, R. 2018. Laurentian origin of solutan echinoderms: new evidence from the Guzhangian (Cambrian Series 3) Weeks Formation of Utah, USA. *Geological Magazine*, **155**, 1190–1204.
- Lefebvre, B., Derstler, K. and Sumrall, C. D. 2012. A reinterpretation of the solutan *Plasiacystis mobilis* (Echinodermata) from the Middle Ordovician of Bohemia. *Zoosymposia*, **7**, 287–306.
- Lefebvre, B., Sumrall, C. D., Shroat-Lewis, R. A., Reich, M., Webster, G. D., Hunter, A. W., Nardin, E., Rozhnov, S. V., Guensburg, T. E., Touzeau, A., Noailles, F. and Sprinkle, J. 2013. Palaeobiogeography of Ordovician echinoderms. In *Early Palaeozoic Biogeography and Palaeogeography* (Harper, D. A. T and Servais, T., eds). *Geological Society London, Memoirs*, **38**, 173–198.
- Lefebvre, B., Allaire, N., Guensburg, T. E., Hunter, A. W., Kouraïss, K., Martin, E., Nardin, E., Noailles, F., Pittet, B., Sumrall, C. D. and Zamora, S. 2016. Palaeoecological aspects of the diversification of echinoderms in the Lower Ordovician of central Anti-Atlas, Morocco. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **460**, 97–121.
- Noailles, F., Lefebvre, B. and Kašička, L. 2014. A probable case of heterochrony in the solutan *Dendrocystites* Barrande, 1887 (Echinodermata: Blastozoa) from the Upper Ordovician of the Prague Basin (Czech Republic) and a revision of the family Dendrocystitidae Bassler, 1938. *Bulletin of Geosciences*, **89**, 451–476.

- Nohejlová M. and Lefebvre B. 2022. Late Ordovician solutan echinoderms from the western Tafilalt, Morocco. In *The Great Ordovician Biodiversification Event: Insights from the Tafilalt Biota, Morocco* (Hunter, A.W., Álvaro, J.J., Lefebvre, B., Van Roy, P. and Zamora, S., eds). *Geological Society, London, Special Publications*, **485**, 523–539.
- Parsley, R. L. 1997. The echinoderm classes Stylophora and Homoiostelea: non Calcichordata. *Paleontological Society Papers*, **3**, 225–248.
- Prokop R. V. and Petr, V. 2003. *Plasiacystis mobilis*, gen. et sp. n., a strange "carpoid" (Echinodermata, ?Homoiostelea: Soluta) in the Bohemian Ordovician (Czech Republic). *Sborník Národního muzea (B: Přírodní vědy) [= Acta Musei Nationalis Pragae (B: Natural History)]*, **59**, 151–162.
- Rozhnov, S. V. 2022. Solutans (echinoderms): Evolution frozen between torsion and pentaradiality. *Paleontological Journal*, **56**, 1306–1321.
- Saleh F., Vaucher R., Vidal M., El Hariri K., Laibl L., Daley A. C., Gutiérrez-Marco J. C., Candela Y., Harper D. A. T., Ortega-Hernandez J., Ma X., Rida A., Vizcaíno D. and Lefebvre B. 2022. New fossil assemblages from the Early Ordovician Fezouata Biota. *Scientific Reports*, **12**:20773, 1–10.
- Smith, A. B. 1988. Patterns of diversification and extinction in Early Palaeozoic echinoderms. *Palaeontology*, **31**, 799–828.
- Sprinkle, J. and Guensburg, T. E. 1993. Echinoderm biostratigraphy. In *Paleozoic Biochronology of the Great Basin, Western United States* (Taylor, M. E., ed.). *Open Field Report*, **598**, 61–63.
- Sprinkle, J. and Guensburg, T. E. 1995. Origin of echinoderms in the Paleozoic Evolutionary Fauna: the role of substrates. *Palaios*, **10**, 437–453.
- Sumrall, C. D., Sprinkle, J. and Guensburg, T. E. 1997. Systematics and paleoecology of Late Cambrian echinoderms from the western United States. *Journal of Paleontology*, **71**, 1091–1109.
- Thoral, M. 1935. *Contribution à l'étude paléontologique de l'Ordovicien inférieur de la Montagne Noire et révision sommaire de la faune cambrienne de la Montagne Noire*. Imprimerie de la Charité, Montpellier, 362 pp.
- Ubaghs, G. 1963. *Cothurnocystis* Bather, *Phyllocystis* Thoral and an undetermined member of the order Soluta (Echinodermata, Carpoidea) in the uppermost Cambrian of Nevada. *Journal of Paleontology*, **37**, 1133–1142.
- Ubaghs, G. 1970. *Les échinodermes carpoïdes de l'Ordovicien inférieur de la Montagne Noire (France)*. Editions du CNRS, Paris, 112 pp.
- Vizcaíno, D. and Álvaro, J. J. 2003. Adequacy of the Early Ordovician trilobite record in the southern Montagne Noire (France): biases for biodiversity documentation. *Transactions of the Royal Society of Edinburgh, Earth Sciences*, **93**, 393–401.
- Vizcaíno, D. and Lefebvre, B. 1999. Les échinodermes du Paléozoïque inférieur de Montagne Noire : biostratigraphie et paléodiversité. *Geobios*, **32**, 353–364.
- Vizcaíno, D., Álvaro, J. J. and Lefebvre, B. 2001. The Lower Ordovician of the southern Montagne Noire. *Annales de la Société Géologique du Nord*, **8**, 213–220.
- Zamora S., Lefebvre, B., Alvaro, J. J., Clausen, S., Elicki, O., Fatka, O., Jell, P., Kouchinsky, A., Lin, J. P., Nardin, E., Parsley, R. L., Rozhnov, S. V., Sprinkle, J., Sumrall, C. D., Vizcaíno, D. and Smith, A. B. 2013a. Cambrian echinoderm diversity and palaeobiogeography. In *Early Palaeozoic Biogeography and Palaeogeography* (Harper, D. A. T. and Servais, T., eds). *Geological Society of London, Memoirs*, **38**, 157–171.
- Zamora, S., Zhu, X. and Lefebvre, B. 2013b. A new Furongian (Cambrian) echinoderm-Lagerstätte from the Sandu Formation (South China). *Cahiers de Biologie Marine*, **54**, 565–569.
- Zhu, X., Peng, S., Zamora, S., Lefebvre, B. and Chen, G. 2016. Furongian (upper Cambrian) Guole Konservat-Lagerstätte from South China. *Acta Geologica Sinica*, **90**, 801–808.