



HAL
open science

Insights into stylophoran anatomy and taphonomy based on an exceptionally preserved mitrate from the Lorraine Group (Upper Ordovician) of New York, USA

Thomas Boisset, Bertrand Lefebvre, Rich Mooi, Andreas Kroh, Viola Winkler, Jérôme Adrien, Markus J. Martin

► To cite this version:

Thomas Boisset, Bertrand Lefebvre, Rich Mooi, Andreas Kroh, Viola Winkler, et al.. Insights into stylophoran anatomy and taphonomy based on an exceptionally preserved mitrate from the Lorraine Group (Upper Ordovician) of New York, USA. 11th European Conference on Echinoderms, Bertrand Lefebvre; Thomas Saucède, Oct 2023, Villeurbanne, France. hal-04271127

HAL Id: hal-04271127

<https://hal.science/hal-04271127>

Submitted on 6 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



ECE 11

Lyon, France

ABSTRACT BOOK

11th European Conference on Echinoderms (ECE11)

October 16–20, 2023

Lyon, France

Insights into stylophoran anatomy and taphonomy based on an exceptionally preserved mitrate from the Lorraine Group (Upper Ordovician) of New York, USA

Thomas BOISSET¹, Bertrand LEFEBVRE¹, Rich MOOI², Andreas KROH³,
Viola WINKLER⁴, Jérôme ADRIEN⁵ & Markus J. MARTIN⁶

¹ UMR CNRS 5276 LGLTPE, Université Lyon 1, Villeurbanne, France. thomas.boisset@etu.univ-lyon1.fr, bertrand.lefebvre@univ-lyon1.fr.

² Department of Invertebrate Zoology and Geology, California Academy of Sciences, San Francisco, California, USA. rmooi@calacademy.org

³ Department of Geology & Palaeontology, Naturhistorisches Museum Wien, Vienna, Austria. andreas.kroh@nhm.at

⁴ Central Research Laboratories, Naturhistorisches Museum Wien, Vienna, Austria. viola.winkler@nhm.at

⁵ UMR CNRS 5510 Laboratoire MATEIS, INSA de Lyon, Villeurbanne, France. jerome.adrien@insa-lyon.fr

⁶ Watertown, New York, USA. triarthrus@gmail.com

Several levels of the Lorraine Group (Upper Ordovician) in upstate New York (USA) have yielded low-diversity, exceptionally preserved, pyritized invertebrate assemblages dominated by the trilobite *Triarthrus eatoni*. Sedimentological and taphonomic features suggest dysoxic bottom-water conditions, with limited transport and rapid burial by distal turbidites (Farrell *et al.*, 2009, 2011). Echinoderms are extremely rare in these strata. Here we report, for the first time, the occurrence of the anomalocystitid mitrate *Enoploura* in the Konservat-Lagerstätte of Beecher's Trilobite Bed. A pyritized specimen of this stylophoran was CT-scanned and three-dimensionally reconstructed. The mitrate is laterally compressed, but its 3D-rendering provided several insights into its internal anatomy and taphonomy. The recurved position of the single feeding appendage (aulacophore) probably results from ligament-induced, post mortem contraction. This posture and the collapse of one lateral series of cover plates indicate that the individual was probably not buried alive. Nevertheless, a portion of the distal aulacophore shows clear evidence of exceptionally preserved soft parts (ambulacral system) in between two sets of slightly open cover plates and the underlying ossicles. One of the most intriguing features of this specimen is its close association with a sinuous, elongated, pyritized trace fossil, which enters the stylophoran through the mouth and disappears into the proximal aulacophore. In marked contrast with other skeletal parts of the specimen (theca and distal part of the aulacophore), the proximal rings of the aulacophore are heavily disrupted and disarticulated. Proximal rings are usually the most decay-resistant skeletal regions in stylophorans (Saleh *et al.*, in press). Therefore, close association of this disrupted region with a trace fossil that penetrates it suggests the action of an unknown infaunal scavenger. Proximal rings of the aulacophore, which housed powerful muscles in life, were likely very attractive to scavengers who could detect freshly killed benthos in the first stages of decay.

References

- Farrell, Ú.C., Martin, M.J., Hagadorn, J.W., Whiteley, T. & Briggs, D.E.G. 2009. Beyond Beecher's Trilobite Bed: widespread pyritization of soft tissues in the Late Ordovician Taconic foreland basin. *Geology*, **37**, 907–910.
- Farrell, Ú.C., Briggs, D.E.G. & Gaines, R.R. 2011. Paleoecology of the olenid trilobite *Triarthrus*: new evidence from Beecher's Trilobite Bed and other sites of pyritization. *Palaios*, **26**, 730–742.
- Saleh, F., Lefebvre, B., Dupichaud, C., Martin, E.L.O., Nohejlová M. & Spaccesi L. In press. Skeletal elements controlled soft-tissue preservation in echinoderms from the Early Ordovician Fezouata Biota. *Geobios*.