



HAL
open science

A new multiarmed asteroid from the Lower Devonian of Morocco

Frank A. Raquet, Daniel B. Blake, Bertrand Lefebvre

► **To cite this version:**

Frank A. Raquet, Daniel B. Blake, Bertrand Lefebvre. A new multiarmed asteroid from the Lower Devonian of Morocco. 11th European Conference on Echinoderms, Bertrand Lefebvre; Thomas Saucède, Oct 2023, Villeurbanne, France. hal-04271143

HAL Id: hal-04271143

<https://hal.science/hal-04271143v1>

Submitted on 5 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



ECE 11

Lyon, France

ABSTRACT BOOK

11th European Conference on Echinoderms (ECE11)

October 16–20, 2023

Lyon, France

A new multiarmed asteroid from the Lower Devonian of Morocco

Frank A. RAQUET¹, Daniel B. BLAKE² & Bertrand LEFEBVRE³

¹ Sulzbachtal, Germany. f.raquet@aol.de

² Department of Geology, University of Illinois, Urbana, Illinois, USA. dblake@illinois.edu

³ UMR CNRS 5276 LGLTPE, Université Lyon 1, Villeurbanne, France. bertrand.lefebvre@univ-lyon1.fr

Deviation from the classical pentamerous symmetry through the possession of supernumerary arms and associated ambulacra is relatively common in extant Asterozoa: five families (out of 34) are exclusively multiarmed, and nine include both multiarmed and five-armed taxa (Hotchkiss, 2000). In contrast, Palaeozoic multiarmed asteroids are rare, with only 11 occurrences reported so far (Herringshaw *et al.*, 2007a). Their earliest occurrences include *Lepidaster grayi* from the Wenlock of England, an undescribed starfish from the Ludlow of Australia (Herringshaw *et al.*, 2007a), and two Emsian taxa from the Hunsrück Shale of Germany (*Helianthaster rhenanus* and *Palaeosolaster gregoryi*; Blake, 2009).

Here, we report a new occurrence of multiarmed asteroid, based on about half of the oral surface of a single specimen from the Emsian of the Erfoud area (eastern Anti-Atlas, Morocco). Although incomplete and weathered, the morphology of this specimen, which originally possessed approximately 20 elongate arms, departs from that of coeval forms from the Emsian of Germany. The new Moroccan asteroid is tentatively assigned to the family Lepidasteridae, which includes the two multiarmed genera *Lepidaster* (Silurian of England) and *Michiganaster* (Middle Devonian of Michigan), as well as the five-armed *Lepidactis* (Silurian of England and, possibly, Middle Ordovician of western France) (Herringshaw *et al.*, 2007a, 2007b; Blake *et al.*, 2016).

References

- Blake, D.B. 2009. Re-evaluation of the Devonian family Helianthasteridae Gregory, 1899 (Asterozoa: Echinodermata). *Paläontologische Zeitschrift*, **83**, 293–308.
- Blake, D.B., Guensburg, T.E. & Lefebvre, B. 2016. New Early Paleozoic Asterozoa (Echinodermata) from the Armorican Massif, France, and the Western United States. *Annales de Paléontologie*, **102**, 161–181.
- Herringshaw, L.G., Smith, M.P. & Thomas, A.T. 2007a. Evolutionary and ecological significance of *Lepidaster grayi*, the earliest multiradiate starfish. *Zoological Journal of the Linnean Society*, **150**, 743–754.
- Herringshaw, L.G., Thomas, A.T. & Smith, M.P. 2007b. Starfish diversity in the Wenlock of England. *Palaeontology*, **50**, 1211–1229.
- Hotchkiss, F.H.C. 2000. On the number of rays in starfish. *American Zoologist*, **40**, 340–354.